

Article

Not peer-reviewed version

The Association between Coffee and Tea Consumption at Midlife and Risk of Dementia Later in Life: The HUNT Study

Denise Abbel , Bjørn Olav Åsvold , [Marit Kolberg](#) , Geir Selbæk , [Raymond Noordam](#) ,
[Håvard Kjesbu Skjellegrind](#) *

Posted Date: 20 April 2023

doi: 10.20944/preprints202304.0627.v1

Keywords: coffee consumption; tea consumption; mild cognitive impairment; dementia; Alzheimer's disease; nutrition; epidemiology; apolipoprotein E; MCI



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

The Association between Coffee and Tea Consumption at Midlife and Risk of Dementia Later in Life: The HUNT Study

Denise Abbel ^{1,2,3}, Bjørn Olav Åsvold ^{1,3,4}, Marit Kolberg ⁵, Geir Selbæk ^{6,7,8}, Raymond Noordam ² and Håvard Kjesbu Skjellegrind ^{1,9,10,*}

¹ HUNT Research Centre, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Levanger, Norway

² Department of Internal Medicine, Section of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands

³ K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Trondheim, Norway

⁴ Department of Endocrinology, Clinic of Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

⁵ Center for Oral Health Services and Research Mid-Norway (TkMidt), Trondheim, Norway

⁶ Norwegian National Centre for Aging and Health, Vestfold Hospital Trust, Tønsberg, Norway

⁷ Department of Geriatric Medicine, Oslo University Hospital, Oslo, Norway

⁸ Faculty of Medicine, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

⁹ General Practice Research Unit, Department of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

¹⁰ Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway

* Correspondence: havard.skjellegrind@ntnu.no

Abstract: Background: Coffee and tea consumption are putative protective factors for dementia, but studies so far have shown inconsistent results. We aimed to investigate whether consumption of tea and different types of coffee at midlife are associated with dementia later in life, and whether sex or ApoE4 influence such association. Methods: We included 7381 participants from the Norwegian HUNT Study. Self-reported questionnaires assessed daily consumption of coffee and tea at baseline. After 22 years, individuals 70 years or older were assessed for cognitive impairment. Results: Compared to daily consumption of 0-1 cups of coffee, daily consumption of ≥ 8 cups of boiled coffee was associated with increased dementia risk in women (OR:1.83, 95% CI: 1.10-3.04, $p=0.03$) and daily consumption of 4-5 cups of other types of coffee was associated with a decrease in dementia risk in men (OR 0.48, 95% CI: 0.32-0.72, $p=0.05$). An association between boiled coffee and dementia risk was only found in ApoE4 non-carriers. Differences on sex or ApoE4 carrier status were not confirmed in interaction analysis. Tea consumption was not associated with dementia risk. Conclusion: The type of coffee might play a role in the direction of the association between coffee-drinking habits and dementia later in life.

Keywords: coffee consumption; tea consumption; mild cognitive impairment; dementia; Alzheimer's disease; nutrition; epidemiology; apolipoprotein E

1. Introduction

Globally, around 57.4 million people are currently living with dementia increasing to an estimated 152.8 million by 2050 [1]. Alzheimer's Disease (AD), which accounts for approximately 60-70% of all dementia cases [2], is characterized by amyloid- β plaques, neurofibrillary tangles, and neuroinflammation. Moreover, mild cognitive impairment (MCI) is a possible prodromal condition of dementia, some individuals with MCI will eventually progress to dementia.

No disease-modifying treatment for cognitive decline or dementia is established yet. Prevention is therefore of great importance. A previous study [3] theorized that 40% of dementia cases could be prevented or delayed by avoiding several modifiable risk factors, including physical inactivity,

alcohol and social isolation. Diet could be another promising strategy for dementia prevention [4]. Coffee and tea are two of the most commonly consumed beverages worldwide and contain many bioactive plant compounds [5]. Previous studies have shown that regular caffeine intake, which is found in both coffee and tea, reduced neuroinflammation, amyloid- β levels and reversed cognitive impairment in animal models [6,7]. One of the mechanisms of action of caffeine may be the competitive antagonism of adenosine receptors [8]. Nevertheless, evidence from observational studies researching the association between coffee and tea consumption and dementia risk is conflicting. While some studies conclude that increased coffee and tea consumption separately or in combination may be associated with a lower dementia risk [9–11], other studies report no association [12]. Most studies had a follow-up time of less than ten years. Pre-clinical Alzheimer's disease may influence dietary habits through loss of initiative and social retraction. As the pathologic processes leading to clinical AD may have a preclinical phase of 10-15 years [13] studies with a shorter follow-up period are at risk for reverse causation bias.

While most studies are focused on universal risk reduction for the general population, assessing risk factors, including sex and genetics, could possibly enable optimal risk reduction by offering highly tailored approaches for dementia prevention. The most well-known genetic risk factor for Alzheimer's Disease is the Apolipoprotein E ϵ 4 (ApoE4) allele [14]. ApoE4 carriers are at increased risk for dementia and the environment might play a role in the translation of this genetic risk into the disease. This gene thus possibly influences associations between prevention interventions and dementia risk [15].

Lastly, little is known about the different brewing methods in the association between coffee and dementia development [16]. A coffee filter retains the lipids cafestol and kahweol and consumption of unfiltered coffee could raise low-density lipoprotein (LDL)-cholesterol blood levels. This may increase the risk of cardiovascular diseases (CVD) and thus dementia [17–19].

In the current study, we used data from the Trøndelag Health (HUNT) Study, a longitudinal cohort study with a mean follow-up period of 21.80 years, to examine whether the consumption of tea and different types of coffee at midlife are associated with long-term risk of AD, MCI and all-cause dementia and to also assess the potential impact of sex and ApoE4 carrier status on this association.

2. Materials and Methods

2.1. Study population and design

Since 1984, the HUNT Study, a population-based cohort study, has been conducted in four surveys. HUNT1 (1984-1986), HUNT2 (1995-1997), HUNT3 (2006-2008) and HUNT4 (2017-2019) invited all residents in the Nord-Trøndelag region of Norway aged over 20 years. As part of HUNT4, a specific study on aging, was performed, HUNT 70+. Individuals aged 70 years or older both community-dwelling and living in long-term care institutions were invited to participate. The current study included the participants of HUNT2 in 1995-1997 who subsequently participated in the HUNT4 70+ study in 2017-2019.

2.2. Exposure assessment

Coffee and tea consumption were assessed during HUNT2 using a self-reported questionnaire. All participants were asked to write down the number of cups of boiled coffee, other types of coffee and tea they consumed daily.

2.3. Outcome assessment

Details of dementia and cognition ascertainment are published elsewhere [20]. In brief, dementia and MCI were diagnosed based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) criteria [21]. A consensus process was applied, using all available relevant information and at least two research and clinical experts from a group of nine specialists in neurology, geriatrics and/or old age psychiatry independently assessed the cognitive status. When dementia was

diagnosed, subtypes were defined based on clinical symptoms and categorized into: AD, vascular dementia (VaD), Lewy body dementia (LBD), frontotemporal dementia (FTD), mixed dementia, other specified dementia, or unspecified dementia. The main outcome of this study was overall dementia risk. Due to the low sample sizes of dementia subtypes other than AD, secondary outcomes were limited to MCI and AD risk.

2.4. Covariates

The following covariates, collected at baseline at HUNT2, were included: sex, age, marital status, educational attainment, smoking, physical activity (PA), marital status, alcohol consumption, history of CVD, history of diabetes mellitus (DM), body mass index (BMI) and ApoE4 carrier status. A more detailed description of the data collection in HUNT2 has been described previously [22]. In brief, information on sex, age, educational attainment, marital status, smoking, alcohol, PA, myocardial infarction, stroke and diabetes was obtained using self-reported questionnaires. PA was assessed by self-reporting on weekly hours of light and vigorous PA (none/less than 1 hours/1-2 hours/ ≥ 3 hours). Based on the answers, we calculated the metabolic equivalent (MET) hours per week. Less than 1 hour per week was considered as 0.5 hours (h) per week, 1-2 h/week as 1.5 h/week and ≥ 3 h/week as 3.5 h/week. For light PA, the amount of PA hours was multiplied by 2.5 METs and for vigorous PA, PA hours were multiplied by 7 METs. Additionally, MET-h/week were grouped into: ≤ 8.3 MET-h/week, 8.4–16.6 MET-h/week and >16.6 MET-h/week.

During HUNT2, clinical measurements determined weight and height and was used to calculate BMI (kg/m^2). Venous blood was drawn, DNA was extracted from whole blood and used for genotyping. ApoE4 carrier status was determined using the two single nucleotide polymorphisms (SNPs) rs7412 and rs429358.

2.5. Statistical analyses

All statistical analyses were performed in STATA, version 17.0, and figures were made in R, version 4.0.3. Baseline characteristics of the study participants were reported as mean (standard deviation) for continuous variables and number (percentage) for categorical variables. We used logistic regression models to estimate odds ratios (ORs) and 95% confidence intervals (95% CI) of all-cause dementia according to different types of coffee consumption (0-1/2-3/4-5/6-7/ ≥ 8 cups daily) and tea consumption (0/1/2/ ≥ 3 cups daily). These main analyses were adjusted according to the following three models: model 1 included sex (female/male), and age (years); model 2 was further adjusted for marital status (unmarried/married/widow(er), divorced or separated), educational attainment (primary school/high school/college or university), smoking (never /previous/current daily smoker), alcohol consumption (total units/week), PA ($\leq 8.3/8.4\text{--}16.6/>16.6$ MET h/week), and tea /coffee consumption and model 3 was additionally adjusted for BMI ($<25/25\text{--}29.9/30\text{--}34.9/\geq 35$ kg/m^2), history of CVD (none/history myocardial infarction and/or stroke), and diabetes (yes/no). The analyses to test the association between the different types of coffee and dementia risk was adjusted according to model 4, that consisted of model 2 and was additionally adjusted for the alternative type of coffee (0-1/2-3/4-5/6-7/ ≥ 8 cups daily). To test the robustness of our findings on the type of coffee, we excluded participants that drank the alternative type of coffee and performed sensitivity analyses. Tests for linear trends in the OR across categories of exposure were performed using coffee and tea consumption categories as continuous variables.

Furthermore, multinomial logistic regression analyses estimated the relative risk (RR) and 95% CI for MCI and AD according to tea and different types of coffee consumption, adjusted according to model 4. In these analyses, other types of dementia were excluded.

To assess potential interaction effects, subgroup analyses according to model 4 were carried out according to the two potential effect modifiers, sex (female/men) and ApoE4 carrier status (negative/positive) to assess the dementia risk by coffee and tea consumption. For the ApoE4 analyses, we excluded participants with missing information on ApoE4 carrier status. We used the likelihood ratio test (LRT) to calculate the p value for interaction, contrasting maximally-adjusted models with and without the cross-product term of the stratifying variable with coffee or tea

consumption in the model. To account for survival bias when assessing possible sex differences, we additionally performed sensitivity analysis where we ran the model, excluding participants older than 80 years at HUNT4.

2.6. Ethics

This project was approved by the Regional Committee for Medical and Health Research Ethics in Norway (456138) and the HUNT study was approved by the Norwegian Data Protection Authority. Participation was voluntary and all participants provided written informed consent.

3. Results

3.1. Participants

In total, 8758 people participated in both in HUNT2 and HUNT4 70+ study (Fig. 1) with a mean follow-up time of 21.8 years. Participants with missing data on coffee consumption, tea consumption, dementia diagnosis or covariates (n=1377) were excluded, resulting in 7381 individuals available for main analyses. For the ApoE4 sub-analyses and the analyses on AD/MCI risk, boiled coffee and other types of coffee we respectively excluded participants with missing data on the ApoE4 carrier status (n=368), participants with other causes of cognitive impairment apart from MCI and AD (n=417), participants who drank other types of coffee (n=4561) and participants who drank boiled coffee (n=3368).




Figure 1. Flowchart of participant selection.

In the total study sample, 985 (13.4%) cases of dementia were identified during HUNT 70+ (Table 1). The mean age of the participants was 55.9 years (SD 6.2) at baseline and the sample contained 3432

(46.5%) men. In general, people who solely completed primary school tended to drink more cups of coffee, compared to people with college or university education. Moreover, smokers tended to drink more cups of coffee than nonsmokers and participants who were less physically active (≤ 8.3 MET-h/week) tended to drink more cups of coffee and fewer cups of tea compared to physically active participants (>16.6 MET-h/week). There was an inverse relationship between coffee and tea intake.

Table 1. Characteristics of study sample examining the possible association between coffee and dementia risk, collected at HUNT2 and HUNT4.

	Coffee consumption (cups/day)						Tea consumption (cups/day)			
	Overall	0-1	2-3	4-5	6-7	≥ 8	0	1	2	≥ 3
Total study population, n		580	1770	2584	1248	1199	4413	1132	1184	652
	7381									
Sex, male	3432 (46.5)	286 (49.3)	768 (43.4)	1099 (42.5)	594 (47.6)	685 (57.1)	2229 (50.5)	497 (43.9)	411 (34.7)	295 (45.2)
Age	55.85 \pm 6.20	56.22 \pm 6.19	56.57 \pm 6.56	56.18 \pm 6.43	55.40 \pm 5.96	54.39 \pm 5.36	55.76 \pm 6.21	55.83 \pm 6.31	56.22 \pm 6.47	55.90 \pm 5.96
Educational attainment										
Primary school	2976 (40.3)	162 (27.9)	633 (35.8)	1104 (42.7)	532 (42.6)	545 (45.5)	1918 (43.5)	363 (32.1)	454 (38.3)	241 (37.0)
High school	2655 (36.0)	222 (38.3)	642 (36.3)	891 (34.5)	452 (36.2)	448 (37.4)	1578 (35.8)	425 (37.5)	432 (36.5)	220 (33.7)
College/university	1750 (23.7)	196 (33.8)	495 (28.0)	589 (22.8)	264 (21.2)	206 (17.2)	917 (20.8)	344 (30.4)	298 (25.2)	191 (29.3)
Marital status										
Unmarried	310 (4.2)	42 (7.2)	70 (4.0)	104 (4.0)	47 (3.8)	47 (3.9)	197 (4.5)	42 (3.7)	41 (3.5)	30 (4.6)
Married	6097 (82.6)	462 (79.7)	1457 (82.3)	2137 (82.7)	1057 (84.7)	984 (82.1)	3635 (82.4)	945 (83.5)	984 (83.1)	533 (81.7)
Widow(er)/divorced/separated		76 (13.1)	243 (13.7)	343 (13.3)	144 (11.5)	168 (14.0)	581 (13.2)	145 (12.8)	159 (13.4)	89 (13.7)
	974 (13.2)									
Tea consumption (cups/day)										
0	4413 (59.8)	181 (31.2)	721 (40.7)	1595 (61.7)	935 (74.9)	981 (81.8)				
1	1132 (15.3)	103 (17.8)	436 (24.6)	398 (15.4)	120 (9.6)	75 (6.3)				
2	1184 (16.0)	119 (20.5)	397 (22.4)	438 (17.0)	139 (11.1)	91 (7.6)				
≥ 3	652 (8.8)	177 (30.5)	216 (12.2)	153 (5.9)	54 (4.3)	52 (4.3)				
Coffee consumption (cups/day)										
0-1	580 (7.9)						181 (4.1)	103 (9.1)	119 (10.1)	177 (27.1)
2-3	1770 (24.0)						721 (16.3)	436 (38.5)	397 (33.5)	216 (33.1)
4-5	2584 (35.0)						1595 (36.1)	398 (35.2)	438 (37.0)	153 (23.5)
6-7	1248 (16.9)						935 (21.2)	120 (10.6)	139 (11.7)	54 (8.3)
≥ 8	1199 (16.2)						981 (22.2)	75 (6.6)	91 (7.7)	52 (8.0)
BMI (kg/m ²)										
< 25	2449 (33.2)	223 (38.4)	613 (34.6)	853 (33.0)	383 (30.7)	377 (31.4)	1392 (31.5)	407 (36.0)	403 (34.0)	247 (37.9)
25-29.99	3759 (51.0)	272 (46.9)	888 (50.2)	1317 (51.0)	671 (53.8)	611 (51.0)	2286 (51.8)	575 (50.8)	597 (50.4)	301 (46.2)
30-34.99	939 (12.7)	61 (10.5)	210 (11.9)	333 (12.9)	163 (13.1)	172 (14.3)	597 (13.5)	113 (10.0)	146 (12.3)	83 (12.7)
≥ 35	234 (3.2)	24 (4.1)	59 (3.3)	81 (3.1)	31 (2.5)	39 (3.3)	138 (3.1)	37 (3.3)	38 (3.2)	21 (3.2)
Alcohol (units/week)	1.71 \pm 2.22	1.27 \pm 2.13	1.61 \pm 2.19	1.68 \pm 2.13	1.78 \pm 2.07	2.06 \pm 2.58	1.78 \pm 2.36	1.68 \pm 2.02	1.50 \pm 1.94	1.65 \pm 2.12
PA (MET-h/week)										
≤ 8.3	3930 (53.2)	303 (52.2)	908 (51.3)	1363 (52.7)	672 (53.8)	684 (57.0)	2439 (55.3)	558 (49.3)	618 (52.2)	315 (48.3)
8.3-16.6	2273 (30.8)	168 (29.0)	575 (32.5)	812 (31.4)	393 (31.5)	325 (27.1)	1302 (29.5)	383 (33.8)	369 (31.2)	219 (33.6)
>16.6	1178 (16.0)	109 (18.8)	287 (16.2)	409 (15.8)	183 (14.7)	190 (15.8)	672 (15.2)	191 (16.9)	197 (16.6)	118 (18.1)
Smoking										
Never	3305 (44.8)	402 (69.3)	1028 (58.1)	1186 (45.9)	441 (35.3)	248 (20.7)	1682 (38.1)	636 (56.2)	638 (53.9)	349 (53.5)
Previous	2579 (34.9)	133 (22.9)	571 (32.3)	943 (36.5)	481 (38.5)	451 (37.6)	1608 (36.4)	351 (31.0)	395 (33.4)	225 (34.5)
Current	1497 (20.3)	45 (7.8)	171 (9.7)	455 (17.6)	326 (26.1)	500 (41.7)	1123 (25.4)	145 (12.8)	151 (12.8)	78 (12.0)
DM, yes	141 (1.9)	14 (2.4)	32 (1.8)	49 (1.9)	29 (2.3)	17 (1.4)	69 (1.6)	25 (2.2)	31 (2.6)	16 (2.5)
CVD, at least one		11 (1.9)	43 (2.4)	59 (2.3)	26 (2.1)	40 (3.3)	120 (2.7)	25 (2.2)	20 (1.7)	14 (2.1)

	179 (2.4)										
ApoE4 carrier status, positive	2101 (30.0)	182 (33.1)	480 (28.7)	721 (29.4)	368 (31.1)	350 (30.3)	1255 (29.8)	315 (29.1)	348 (31.2)	183 (30.0)	
Cognitive status											
No CI	3840 (55.1)	331 (61.1)	927 (55.8)	1344 (55.1)	636 (53.6)	602 (53.0)	2230 (53.7)	636 (59.0)	617 (55.2)	357 (58.1)	
MCI	2552 (36.7)	177 (32.7)	584 (35.2)	896 (36.7)	448 (37.8)	447 (39.4)	1571 (37.8)	358 (33.2)	412 (36.9)	211 (34.4)	
Dementia all causes	985 (13.4)	71 (12.2)	259 (14.6)	343 (13.3)	163 (13.1)	149 (12.4)	610 (13.8)	136 (12.0)	155 (13.1)	84 (12.9)	
AD	572 (8.2)	34 (6.3)	150 (9.0)	200 (8.2)	102 (8.6)	86 (7.6)	354 (8.5)	84 (7.8)	88 (7.9)	46 (7.5)	

Values are expressed as mean \pm standard deviation or number (%). Abbreviations: AD = Alzheimer's disease, BMI = body mass index; CI = cognitive impairment; CVD = cardiovascular diseases, DM = diabetes mellitus, MCI = mild cognitive impairment, PA = physical activity.

3.2. Coffee and tea consumption and dementia risk

In the regression analysis only adjusted for age and sex, daily consumption of ≥ 8 cups of coffee was associated with an increased risk of developing dementia (Model 1: OR:1.45; 95% CI 1.05-2.01, p-value for trend=0.02), compared to daily consumption of 0-1 cup of coffee. (Table 2). Upon further adjustment for confounders, the association between coffee and dementia attenuated (Model 2: OR: 1.11; 95% CI 0.78-1.57, p-value for trend=0.81). Results for adjustment model 2 and 3 were similar. We found no other associations between general coffee consumption and dementia risk, nor was tea consumption associated with dementia risk.

Table 2. OR (95% CI) of dementia by consumption of coffee, stepwise adjusted. .

Coffee consumption (cups/day)							Tea consumption (cups/day)				
0-1		2-3	4-5	6-7	≥8	p-value for trend	0	1	2	≥3	p-value for trend
Model 1	Ref	1.16	1.09	1.26	1.45	0.02	Ref	0.81	0.84	0.91	0.11
		(0.86-1.57)	(0.81 - 1.46)	(0.91 - 1.73)	(1.05 - 2.01)			(0.66-1.01)	(0.69-1.03)	(0.70-1.18)	
Model 2	Ref	1.12	0.95	1.06	1.11	0.81	Ref	0.92	0.92	1.02	0.71
		(0.82-1.52)	(0.70-1.30)	(0.75-1.48)	(0.78 - 1.57)			(0.74-1.15)	(0.75-1.14)	(0.77-1.34)	
Model 3	Ref	1.11	0.94	1.04	1.09	0.90	Ref	0.92	0.92	1.01	0.65
		(0.81 - 1.51)	(0.69-1.28)	(0.74-1.46)	(0.77-1.54)			(0.74-1.16)	(0.74-1.14)	(0.76-1.33)	

Model 1: Results were adjusted for participation age (years) and sex (female/men). Model 2: Results were additionally adjusted for educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), coffee/tea consumption, exercise (MET-h/week score) and alcohol consumption (total units/week). Model 3: Results were further adjusted for diabetes (yes/no), CVD (none/at least one) and BMI (<25, 25-29.9, 30-34.9, ≥ 35).

3.3. Type of coffee and dementia risk

According to adjustment model 2, daily consumption of ≥ 6 cups of boiled coffee was associated with an increased risk of dementia (6-7 cups: OR:1.38, 95%CI: 1.05-1.81, ≥ 8 cups: OR: 1.46, 95% CI:1.08-1.96, p-value for trend<0.01), compared to consumption of 0-1 cups of boiled coffee a day (Table 3). After further adjusting for other types of coffee, this association disappeared (6-7 cups: OR=1.19, 95% CI 0.86-1.66, ≥ 8 cups: OR=1.26, 95% CI 0.88-1.80, p-value for trend=0.17) and associations between other consumption levels of boiled coffee and dementia were not found. In adjustment models 2 and 4, only the daily consumption of 4-5 cups of other types of coffee was associated with reduced dementia risk compared to drinking 0-1 cups daily (model 2: OR: 0.67, 95%CI: 0.54-0.82, p-value for trend<0.01, model 4: OR=0.71 95% CI 0.54-0.92, p-value for trend=0.24). The results according to

adjustment model 4 were similar to adjustment model 3 and the sensitivity analyses models where we excluded participants who drank the alternative type of coffee (results not shown).

Table 3. OR (95% CI) of dementia risk by consumption of boiled and other types of coffee consumption, adjusted according to model 2 and 4. .

			Coffee consumption (cups/day)					p-value for trend
			0-1	2-3	4-5	6-7	≥8	
Boiled coffee	Model 2	n	4284	1123	1006	511	457	<0.01
		Ref		1.14 (0.93-1.41)	1.14 (0.92-1.41)	1.38 (1.05-1.81)	1.46 (1.08-1.96)	
	Model 4	n	3148	1447	1540	639	607	0.17
		Ref		1.01 (0.79-1.30)	1.00 (0.75-1.30)	1.19 (0.86-1.66)	1.26 (0.88-1.80)	
Other type coffee	Model 2	n	3148	1447	1540	639	607	<0.01
		Ref		0.96 (0.79-1.18)	0.67 (0.54-0.82)	0.80 (0.60-1.07)	0.86 (0.64-1.15)	
	Model 4	n	3148	1447	1540	639	607	0.24
		Ref		1.01 (0.80-1.28)	0.71 (0.54-0.92)	0.86 (0.61-1.21)	0.93 (0.65-1.32)	

Model 2: Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), tea consumption (0/1/2/≥3), exercise (MET-h/week score) and alcohol consumption (total units/week). Model 4: Results were adjusted according to model 2 and additionally adjusted for the alternative coffee type consumption (0-1/2-3/4-5/6-7/≥8).

3.4. Coffee, tea and the risk of MCI and AD

After adjusting according to model 4, daily consumption of 4-7 cups of boiled coffee was associated with an increased risk of MCI, compared to 0-1 cup of boiled coffee (Table 4) (4-5 cups: RR:1.41; 95% CI 1.16-1.70, 6-7 cups: RR:1.29 95% CI 1.01-1.63, p-value for trend=0.04). Daily consumption of other types of coffee was not associated with MCI risk. Additionally, ≥6 cups of boiled coffee daily was associated with an increased risk of AD compared to daily consumption of 0-1 cups of boiled coffee (6-7 cups: RR: 1.85 95% CI 1.22-2.81, ≥8 cups: RR=1.65 95% CI 1.03-2.53, p-value for trend<0.01) and other types of coffee were not associated with AD risk. Daily tea consumption was also not associated with the risk of MCI or AD (Appendix: Table A1). Results for all analyses were similar to adjustment model 3 (results not shown).

Table 4. RR (95% CI) of MCI and AD risk by coffee consumption, adjusted according to model 4.

			Coffee consumption (cups/day)					p-value for trend
			0-1	2-3	4-5	6-7	≥8	
MCI	Boiled coffee	n	1405	369	407	199	172	0.04
		Ref		1.03 (0.87-1.22)	1.41 (1.16-1.70)	1.29 (1.01-1.63)	1.11 (0.86-1.43)	
	Other type of coffee	n	1138	462	521	212	219	0.87
		Ref		0.98 (0.83-1.16)	0.97 (0.81-1.15)	0.94 (0.75-1.18)	1.01 (0.80-1.28)	
AD	Boiled coffee	n	279	94	97	59	43	
		Ref						

		Ref	1.15 (0.83-1.58)	1.37 (0.96-1.95)	1.85 (1.22-2.81)	1.65 (1.03-2.53)	<0.01
Other type of coffee	n	295	109	92	41	35	
		Ref	1.14 (0.83-1.55)	0.79 (0.56-1.12)	1.00 (0.64-1.56)	1.03 (0.64-1.67)	0.71

Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), tea consumption (0/1/2/≥3), exercise (MET-h/week score) and alcohol consumption (total units/week) and the alternative coffee type consumption (0-1/2-3/4-5/6-7/≥8).

3.5. Sex differences

After adjusting according to model 4, consumption of ≥8 cups of boiled coffee daily was associated with an increase in dementia risk in women compared to drinking 0-1 cups of boiled coffee daily (OR:1.83, 95% CI: 1.10-3.04, p-value for trend=0.03) and boiled coffee was not associated with dementia risk in men (p-value for trend=0.66). There was little statistical evidence that the association differed between women and men (p-value for interaction=1.00) (Figure 2). Additionally, we found that daily consumption of 4-5 cups of other types of coffee was associated with a lower dementia risk in men (OR 0.48, 95% CI: 0.32-0.72, p-value for trend=0.05), but other types of coffee were not associated with dementia risk in women (p-value for trend=0.84). Again, there was little statistical evidence that the association differed between women and men (p-value for interaction=0.41). Adjusting for ApoE4 carrier status did not change the results (results not shown). Also, when limiting the participants' age at outcome measurement from 70 to 80 years, the sex differences in the direction of the association between coffee and dementia risk remained (Appendix: Table A2). Lastly, in women and men, tea was not associated with dementia risk, and there was little statistical evidence that the association differed between women and men (p-value for interaction=0.45) (Appendix: Table A3).




Figure 2. OR (95% CI) of dementia risk by type of coffee consumption and sex, adjusted according to model 4. (a) women; (b) men. Compared to the 0-1 cup/day category. Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), tea consumption (0/1/2/≥3), exercise (MET-h/week score) and alcohol consumption (total units/week) and the alternative coffee type consumption (0-1/2-3/4-5/6-7/≥8).

3.6. ApoE4 carrier status

In ApoE4 non-carriers, 6-7 cups of boiled coffee was associated with an increased dementia risk, compared to consumption of 0-1 cups of boiled coffee (OR=1.53 95% CI: 1.00-2.32, p-value for

trend=0.10. This association was not found in ApoE4 carriers (OR=1.02 95% CI: 0.47-1.48. p-value for trend=0.57) (Table 5). There was little statistical evidence that the association differed by ApoE4 carrier status (p-value for interaction=0.30). Other types of coffee were not associated with dementia risk in neither ApoE4 subgroups (p-value for interaction=0.55). Lastly, in both ApoE4 subgroups, tea was not associated with dementia risk (p-value for interaction=0.92) (Appendix: Table A4).

Table 5. OR (95% CI) of AD risk by type of coffee consumption and ApoE4, adjusted according to model 4.

			Coffee consumption (cups/day)					p-value for trend
			0-1	2-3	4-5	6-7	≥8	
Boiled Coffee	Negative	n	2844	753	675	330	310	0.10
		Ref		1.15 (0.84-1.58)	1.15 (0.81-1.64)	1.53 (1.00-2.32)	1.31 (0.82-2.09)	
	Positive	n	1227	314	281	156	123	0.57
		Ref		0.87 (0.56-1.35)	0.80 (0.44-1.46)	1.02 (0.47-1.48)	1.45 (0.79-2.66)	
Other type coffee	Negative	n	2070	983	1031	417	411	0.19
		Ref		1.14 (0.84-1.55)	0.77 (0.54-1.09)	0.81 (0.51-1.29)	0.86 (0.54-1.39)	
	Positive	n	912	392	433	29	181	0.84
		Ref		0.78 (0.51-1.20)	0.61 (0.39-0.96)	0.85 (0.48-1.52)	1.05 (0.58-1.89)	

Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), tea consumption (0/1/2/≥3), exercise (MET-h/week score) and alcohol consumption (total units/week) and the alternative coffee type consumption (0-1/2-3/4-5/6-7/≥8).

4. Discussion

This study, following 7381 participants with a mean follow-up time of 21.8 years, investigated the possible association between coffee and tea consumption and dementia risk. The main findings were as follows: 1.After adjusting for age and sex, general coffee consumption was associated with an increased dementia risk, however, upon further adjustment for confounders, the association disappeared; 2. Consumption of 4-7 cups of boiled coffee a day were associated with an increased risk of MCI, and ≥6 cups of boiled coffee daily was associated with an increased risk of AD. Consumption of other types of coffee was not associated with MCI and AD risk; 3. Daily consumption of 4-5 cups of other types of coffee was associated with reduced dementia risk in men, but not in women. The daily consumption of ≥8 cups of boiled coffee was associated with an increased dementia risk in women, but not in men. Daily consumption of 6-7 cups of boiled coffee was associated with an increased dementia risk in ApoE4 non-carriers, but not in ApoE4 carriers. These suggestive differences by sex and ApoE4 carrier status were however not substantiated by statistical evidence for interaction; 4. Tea was not associated with dementia, MCI or AD risk.

Previous studies showed inconsistent results concerning coffee consumption and dementia risk. A cohort study that followed 13757 participants for eight years, found that participants who had a daily intake of 2-2.9 cups and >3 cups of coffee had the lowest HRs for dementia incidence (0.69, 95% CI: 0.48-0.98 and 0.53, 95% CI: 0.31-0.89), compared to participants that consumed 0 cups of coffee daily [10]. On the other hand, a study including 3498 men with a follow-up period of 25 years, found no significant association between coffee intake during midlife on the risk of cognitive impairment and dementia [12]. In addition, two studies reported only short-term associations between coffee

intake and cognition and no long-term effects were observed [23,24]. A possible explanation for the inconsistent results in previous studies could be the lack of differentiation between the types of coffee. Our study distinguished boiled and other types of coffee from the association between coffee and dementia risk. We showed that, in general, boiled coffee seems to be associated with an increased risk of developing dementia, while consumption of other types of coffee may be associated with a decrease in dementia risk. Although our study lacks information on the specifics of the other type of coffee, we assume that the other type of coffee mainly consisted of filtered coffee. Filtered coffee has been associated with lower cardiovascular risks, compared to non-coffee drinkers and boiled coffee drinkers [25]. Boiled coffee, which is non-filtered and includes coffee lipids, increases the low-density lipoprotein LDL-cholesterol and thus the risk of atherosclerosis and cardiovascular diseases (CVD) [17,26]. CVD are associated with an increased risk of future dementia [18,19]. Although boiled coffee is not widely used anymore, current popular methods, including French press, Turkish and Greek coffee have comparable amounts of cafestol and kahweol, due to the absence of a paper filter [27]. Espresso contains intermediate amounts of cafestol and kahweol. Although causality cannot be established, this study highlights that coffee might affect dementia risk and may therefore be used as a cost-effective complement to prevention strategies.

The results of this current study suggest that women had an increased dementia risk when drinking higher amounts of boiled coffee, while men had a dementia risk reduction when drinking other types of coffee. The direction of the associations and significance remained after excluding participants aged over 80 years at HUNT4. These suggestive differences were however not substantiated by statistical evidence for interaction. Previous literature showed conflicting findings on sex differences in the association between coffee and dementia risk. While some studies showed that coffee consumption was more strongly associated with a reduction in dementia risk in men [10], others showed that coffee consumption was especially associated with a dementia risk reduction in women [9,28]. Although evidence favoring possible causality cannot be established through the observational nature of our study, we can speculate about the possible mechanisms behind the possible sex differences in the association between coffee and dementia risk. The antioxidant effect of coffee might be more beneficial in men than in women. Both human and animal studies have reported that oxidative stress levels were higher in men than in women [29,30] and Ishizaka, Yamakado [31] found that coffee consumption, in a graded matter, was associated with decreased oxidative stress levels in men, but not in women. If men truly benefit more from coffee consumption, this may balance out the negative effects of boiled coffee and could explain why we found that boiled coffee was associated with an increased dementia risk in women, but not in men. More research is needed to determine the sex differences in coffee consumption and the possible influence of this on dementia development.

Moreover, our results showed that in ApoE4 non-carriers, boiled coffee was associated with an increased dementia risk, while this association was not found in ApoE4 carriers. These suggestive differences were however not substantiated by statistical evidence for interaction. ApoE4 is an apolipoprotein that plays a major role in the transport of cholesterol, as it is a LDL-receptor ligand [32]. A population-based cohort study found that increased levels of LDL-cholesterol were associated with increased risk of AD in ApoE4 non-carriers, but not in ApoE4 carriers [33] and another cohort study showed that higher LDL-cholesterol was associated with decreased verbal memory in ApoE4 non-carriers, but did not in ApoE4 carriers [34]. ApoE4 non-carriers therefore might be more at risk for dementia when drinking the lipids that are found in boiled coffee, but the precise underlying mechanisms remain unknown.

We found no association between tea consumption and MCI, AD and dementia risk. However, tea drinking was relatively rare in our population and led to decreased statistical power. In line with our results, the Finnish Cardiovascular Risk Factor, Aging and Dementia (CAIDE) study, reported that only coffee, but not tea, was associated with dementia risk [35]. However, another study, with a follow-up period of 11.4 years, concluded that tea, separately or in combination with coffee, was associated with a decrease in dementia risk [11]. Compared to non-drinkers, 2-3 cups of coffee or tea were associated with a 28% lower dementia. The type and amount of possible beneficial components

differ per tea type [36]. In our study, there was no information available on the type of tea. Black tea dominates the market in Norway [37] and it is thus plausible that the majority of our tea-drinking participants drank black tea. Several studies showed that green tea was associated with a reduced dementia risk, while black tea was not [38,39].

The precise protective mechanisms of non-boiled coffee against dementia are unknown. The antagonist binding of caffeine to adenosine receptors, which stimulates cholinergic neurons (Fredholm et al., 1999), is a possible pathway that may explain the association. Through the competitive antagonist binding of adenosine receptors, caffeine reduced neuroinflammation and neurodegeneration in rats [7,40]. Moreover, the antioxidants in coffee might influence the risk of dementia by reducing oxidative stress [41]. Lastly, coffee consumption has been associated with reduced risk of several diseases including Parkinson's disease [42,43], diabetes [44,45] and cardiovascular disease [25,46]. Previously mentioned comorbidities are risk factors for dementia and decreasing the risk for comorbidities may mediate the association between coffee and tea and dementia risk. In our study, we did not find significant changes between adjustment model 2 and the model 3, which was additionally adjusted for comorbidities. However, as the participants had a mean age of 55.9 (SD=6.3) at baseline, the prevalence of comorbidities at baseline was low.

Our study has several strengths. Firstly, while most previous epidemiological studies had shorter follow-up periods, this current study has a mean follow-up period of 21.8 years, making our study less prone to reverse causation bias. Other strengths of this current study are the participation of both home-dwelling and institutionalized elderly, controls without dementia who were identified through diagnostic tests and ability to adjust for lots of confounders. Several limitations should also be taken into account when interpreting our results. Firstly, when assessing coffee consumption, the question remains what the alternative is. One must note that all participants drank something. Coffee may be more or less harmful than other drinks and are also related to other dietary habits. Therefore, our analyses may not completely reflect the true association. Additionally, covariates and consumption of coffee and tea were surveyed at baseline and may not reflect long-term drinking habits. Lifestyle factors are dynamic. For example, during the follow-up period, most participants retired, which may lead to major changes in daily habits. Moreover, our results have wide CIs, representing uncertainty of the effect sizes. Also, we had limited statistical power to assess interactions. Although our participation rates were relatively high, non-participants had lower socio-economic status and higher mortality than individuals that participated in the HUNT Study [47,48]. This may limit the generalizability of our findings. Moreover, this study population was predominantly Caucasian, so our results may be less representative for populations with other ethnic backgrounds. Additionally, although a systematic approach was used to diagnose people with dementia, we did not have access to biomarker data for the diagnosis of dementia and AD. Lastly, when researching older adults above the age of 70, the chance of competing risk from death and participation bias based on cognitive status are unavoidable constraints.

To conclude, consumption of 4-7 cups of boiled coffee a day was associated with an increased risk of MCI, and ≥ 6 cups of boiled coffee daily was associated with an increased risk of AD. Consumption of other types of coffee was not associated with MCI or AD risk. Moreover, boiled coffee was associated with increased dementia risk in women and ApoE4 non-carriers, consumption of other types of coffee was associated with decreased dementia risk in men and tea consumption was not associated with dementia risk. These suggestive differences in sex and ApoE4 carrier status were however not substantiated by statistical evidence for interaction. Since coffee and tea are among the most widely consumed beverages worldwide and cognitive decline is common among older adults, even minor potential health benefits or risks associated with coffee and tea consumption could have a significant effect on public health. Long-term intervention studies are needed to disentangle to possible relationship between coffee intake, dementia risk, type of coffee, sex and ApoE4 carrier status.

Author Contributions: Conceptualization, D.A., B.O.Å., G.S. and H.K.S.; methodology, D.A., B.O.Å., G.S., M.K. and H.K.S.; formal analysis D.A., B.O.Å. and H.K.S.; investigation D.A., B.O.Å., M.K. and H.K.S.; resources G.S., M.K. and H.K.S.; writing—original draft preparation, D.A.; writing—review and editing, D.A., B.O.Å., M.K.,

G.S., R.N. and H.K.S.; visualization, D.A.; supervision, B.O.Å, R.N. and H.K.S.; project administration, R.N. and H.K.S.; funding acquisition, D.A., M.K. and H.K.S.;

Funding: This research was funded by Alzheimer Nederland, Jo Kolk Studiefonds, Leiden University Fund (LUF), Center for Oral Health Services and Research Mid-Norway (TkMidt), Trondheim, Norway and General Practice Research Unit, Department of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), Trondheim, Norway. .

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the regional committee on research ethics (REK approval #456138).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Restrictions apply to the availability of these data. Data was obtained from HUNT Databank. HUNT data are available for research projects approved by a Norwegian Regional Committee for Medical Research Ethics (REK). .

Acknowledgments: The Trøndelag Health Study (HUNT) is a collaboration between HUNT Research Centre (Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology NTNU), Trøndelag County Council, Central Norway Regional Health Authority, and the Norwegian Institute of Public Health.



Conflicts of Interest: The authors declare no conflict of interest and the funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Appendix A

Table A1. RR (95% CI) of MCI and AD risk by tea consumption, adjusted according to model 2.

			0	1	2	≥3	p-value for trend
MCI	n		1571	358	412	211	0.91
		Ref		0.91 (0.78-1.06)	1.07 (0.93-1.25)	0.95 (0.78-1.16)	
AD	n		354	84	88	46	0.68
		Ref		0.94 (0.71-1.25)	0.90 (0.68-1.19)	1.01 (0.71-1.45)	

Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), coffee consumption (0-1/2-3/4-5/6-7/≥8), exercise (MET-h/week score) and alcohol consumption (total units/week).

Table A2. OR (95% CI) of dementia risk by type of coffee consumption and sex for people aged 70-80 years at outcome measurement, adjusted according to model 2.

			Coffee consumption (cups/day)					p-value for trend
			0-1	2-3	4-5	6-7	≥8	
Boiled Coffee	Female	n	1578	403	358	183	161	<0.01
		Ref		1.01 (0.57-1.80)	1.28 (0.69-2.36)	1.49 (0.73-3.05)	2.68 (1.35-5.33)	
	Male	n	1454	353	304	174	185	0.47
		Ref		0.47 (0.25-0.85)	1.02 (0.60-1.77)	0.52 (0.25-1.09)	0.78 (0.40-1.53)	

Other type coffee	Female	n	1083	577	582	254	187	
			Ref	1.26 (0.75- 2.14)	0.94 (0.52- 1.71)	1.06 (0.50- 2.25)	1.59 (0.75- 3.40)	0.48
	Male	n	1019	444	494	220	293	
			Ref	0.78 (0.46- 1.32)	0.49 (0.28- 0.87)	0.57 (0.28- 1.14)	0.64 (0.34- 1.21)	0.10

Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), tea consumption (0/1/2/≥3), exercise (MET-h/week score) and alcohol consumption (total units/week) and the alternative coffee type consumption (0-1/2-3/4-5/6-7/≥8).

Table A3. OR (95% CI) of dementia risk by tea consumption and sex, adjusted according to model 2.

Sex		0	1	2	≥3	p-value for trend
Female	n	2184	635	773	357	
		Ref	0.96 (0.71- 1.29)	0.97 (0.74- 1.28)	1.33 (0.92- 1.92)	0.39
Male	n	2229	497	411	295	
		Ref	0.89 (0.63- 1.26)	0.92 (0.65- 1.30)	0.72 (0.47- 1.10)	0.15

Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), coffee consumption (0-1/2-3/4-5/6-7/≥8), exercise (MET-h/week score) and alcohol consumption (total units/week).

Table A4. OR (95% CI) of AD risk by tea consumption and ApoE4 carriers status, adjusted according to model 2.

ApoE4 carrier status		0	1	2	≥3	p-value for trend
Negative	n	2950	767	769	42	
		Ref	0.98 (0.74- 1.30)	0.95 (0.72- 1.26)	0.98 (0.69- 1.41)	0.79
Positive	n	1255	315	348	183	
		Ref	0.85 (0.57- 1.28)	0.91 (0.63- 1.32)	1.16 (0.71- 1.87)	0.95

Results were adjusted for participation age (years) and sex (female/men), educational attainment (primary school/high school/ college or university), marital status (unmarried/married/ widow(er), divorced or separated), smoking status (never/ previous/currently daily smoker), coffee consumption (0-1/2-3/4-5/6-7/≥8), exercise (MET-h/week score) and alcohol consumption (total units/week).

References

1. Global Burden of Disease Dementia Forecasting Collaborators, *Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019*. Lancet Public Health, 2022. 7(2): p. e105-e125.

2. World Health Organization. *Dementia*. Fact Sheets 2022 [cited 2022 02-11]; Available from: <https://www.who.int/news-room/fact-sheets/detail/dementia>.

3. Livingston, G., et al., *Dementia prevention, intervention, and care: 2020 report of the Lancet Commission*. Lancet, 2020. 396(10248): p. 413-446.

4. Middleton, L.E. and K. Yaffe, *Targets for the prevention of dementia*. J Alzheimers Dis, 2010. 20(3): p. 915-24.

5. Gomez-Ruiz, J.A., D.S. Leake, and J.M. Ames, *In vitro antioxidant activity of coffee compounds and their metabolites*. J Agric Food Chem, 2007. 55(17): p. 6962-9.

6. Cao, C., et al., *Caffeine suppresses amyloid-beta levels in plasma and brain of Alzheimer's disease transgenic mice*. J Alzheimers Dis, 2009. **17**(3): p. 681-97.
7. Ullah, F., et al., *Caffeine prevents d-galactose-induced cognitive deficits, oxidative stress, neuroinflammation and neurodegeneration in the adult rat brain*. Neurochem Int, 2015. **90**: p. 114-24.
8. Fredholm, B.B., et al., *Actions of caffeine in the brain with special reference to factors that contribute to its widespread use*. Pharmacol Rev, 1999. **51**(1): p. 83-133.
9. Chen, J.Q.A., et al., *Associations Between Caffeine Consumption, Cognitive Decline, and Dementia: A Systematic Review*. J Alzheimers Dis, 2020. **78**(4): p. 1519-1546.
10. Matsushita, N., et al., *Association of coffee, green tea, and caffeine with the risk of dementia in older Japanese people*. J Am Geriatr Soc, 2021. **69**(12): p. 3529-3544.
11. Zhang, Y., et al., *Consumption of coffee and tea and risk of developing stroke, dementia, and poststroke dementia: A cohort study in the UK Biobank*. PLOS Medicine, 2021. **18**(11): p. e1003830.
12. Gelber, R.P., et al., *Coffee intake in midlife and risk of dementia and its neuropathologic correlates*. J Alzheimers Dis, 2011. **23**(4): p. 607-15.
13. Villemagne, V.L., Burnham, S., Bourgeat, P., Brown, B., Ellis, K. A., Salvado, O., Szoek, C., S.L. Macaulay, Martins, R., Maruff, P., Ames, D., Rowe, C. C., Masters, C. L., Australian, and B. Imaging, *Amyloid beta deposition, neurodegeneration, and cognitive decline in sporadic Alzheimer's disease: a prospective cohort study*. Lancet Neurol, 2013. **12**(4): p. 357-367.
14. Burke, J.R. and A.D. Roses, *Genetics of Alzheimer's disease*. International Journal of Neurology, 1991: p. 25-26:41-51.
15. Norwitz, N.G., et al., *Precision Nutrition for Alzheimer's Prevention in ApoE4 Carriers*. Nutrients, 2021. **13**(4).
16. Wu, L., D. Sun, and Y. He, *Coffee intake and the incident risk of cognitive disorders: A dose-response meta-analysis of nine prospective cohort studies*. Clin Nutr, 2017. **36**(3): p. 730-736.
17. Schoeneck, M. and D. Iggman, *The effects of foods on LDL cholesterol levels: A systematic review of the accumulated evidence from systematic reviews and meta-analyses of randomized controlled trials*. Nutr Metab Cardiovasc Dis, 2021. **31**(5): p. 1325-1338.
18. Clair, L., et al., *Cardiovascular disease and the risk of dementia: a survival analysis using administrative data from Manitoba*. Can J Public Health, 2022. **113**(3): p. 455-464.
19. Farnsworth Von Cederwald, B., et al., *Association of Cardiovascular Risk Trajectory With Cognitive Decline and Incident Dementia*. Neurology, 2022: p. 10.1212/WNL.000.
20. GjØra, L., et al., *Current and Future Prevalence Estimates of Mild Cognitive Impairment, Dementia, and Its Subtypes in a Population-Based Sample of People 70 Years and Older in Norway: The HUNT Study*. J Alzheimers Dis, 2021. **79**(3): p. 1213-1226.
21. Black, D.W. and J.E. Grant, *The essential companion to the diagnostic and statistical manual of mental disorders, fifth edition* 2014: American Psychiatric Publishing
22. Holmen, J., Midthjell, K., KrØger, Ø., Langhammer, A., Lingaas Holmen, T., Bratberg, G., Vatten, L., & Lund-Larsen, P. , *The Nord-TrØndelag Health Study 1995-97 (HUNT 2): Objectives, contents, methods and participation*. Norsk Epidemiologi, 2003. **13**(1): p. 19-32.
23. Araújo, L.F., et al., *Association of Coffee Consumption with MRI Markers and Cognitive Function: A Population-Based Study*. J Alzheimers Dis, 2016. **53**(2): p. 451-61.
24. Mirza, S.S., et al., *Coffee consumption and incident dementia*. Eur J Epidemiol, 2014. **29**(10): p. 735-41.
25. Ding, M., et al., *Long-term coffee consumption and risk of cardiovascular disease: a systematic review and a dose-response meta-analysis of prospective cohort studies*. Circulation, 2014. **129**(6): p. 643-59.
26. Tverdal, A., et al., *Coffee consumption and mortality from cardiovascular diseases and total mortality: Does the brewing method matter?* Eur J Prev Cardiol, 2020. **27**(18): p. 1986-1993.
27. Urgert, R., van der Weg, G., Kosmeijer-Schuil, T. G., van de Bovenkamp, P., Hovenier, R., & Katan, M. B., *Levels of the Cholesterol-Elevating Diterpenes Cafestol and Kahweol in Various Coffee Brews*. Journal of Agricultural and Food Chemistry, 1995. **43**(8): p. 2167-2172.
28. Johnson-Kozlow, M., et al., *Coffee consumption and cognitive function among older adults*. Am J Epidemiol, 2002. **156**(9): p. 842-50.
29. Barp, J., et al., *Myocardial antioxidant and oxidative stress changes due to sex hormones*. Braz J Med Biol Res, 2002. **35**(9): p. 1075-81.
30. Ide, T., et al., *Greater oxidative stress in healthy young men compared with premenopausal women*. Arterioscler Thromb Vasc Biol, 2002. **22**(3): p. 438-42.
31. Ishizaka, Y., et al., *Relationship between coffee consumption, oxidant status, and antioxidant potential in the Japanese general population*. Clin Chem Lab Med, 2013. **51**(10): p. 1951-9.
32. Mahley, R.W., *Apolipoprotein E: cholesterol transport protein with expanding role in cell biology*. Science, 1988. **240**(4852): p. 622-30.
33. Hall, K., et al., *Cholesterol, APOE genotype, and Alzheimer disease: an epidemiologic study of Nigerian Yoruba*. Neurology, 2006. **66**(2): p. 223-7.

34. Wroolie, T., et al., *Effects of LDL Cholesterol and Statin Use on Verbal Learning and Memory in Older Adults at Genetic Risk for Alzheimer's Disease*. J Alzheimers Dis, 2020. **75**(3): p. 903-910.
35. Eskelinen, M.H., et al., *Midlife coffee and tea drinking and the risk of late-life dementia: a population-based CAIDE study*. J Alzheimers Dis, 2009. **16**(1): p. 85-91.
36. Almajano, M.P., Carbó, R., Jiménez, J. A. L., & Gordon, M. H. , *Antioxidant and antimicrobial activities of tea infusions*. Food chemistry, 2008. **108**(1): p. 55-63.
37. Beresniak, A., et al., *Relationships between black tea consumption and key health indicators in the world: an ecological study*. BMJ Open, 2012. **2**(6): p. e000648.
38. Xu, H., et al., *Gender differences in the protective effects of green tea against amnesic mild cognitive impairment in the elderly Han population*. Neuropsychiatr Dis Treat, 2018. **14**: p. 1795-1801.
39. Noguchi-Shinohara, M., et al., *Consumption of green tea, but not black tea or coffee, is associated with reduced risk of cognitive decline*. PLoS One, 2014. **9**(5): p. e96013.
40. Kolahdouzan, M. and M.J. Hamadeh, *The neuroprotective effects of caffeine in neurodegenerative diseases*. CNS Neurosci Ther, 2017. **23**(4): p. 272-290.
41. Luca, M., A. Luca, and C. Calandra, *The Role of Oxidative Damage in the Pathogenesis and Progression of Alzheimer's Disease and Vascular Dementia*. Oxid Med Cell Longev, 2015. **2015**: p. 504678.
42. Nie, J., et al., *Independent and Joint Associations of Tea Consumption and Smoking with Parkinson's Disease Risk in Chinese Adults*. J Parkinsons Dis, 2022.
43. Ross, G.W., *Association of Coffee and Caffeine Intake With the Risk of Parkinson Disease*. JAMA, 2000. **283**(20): p. 2674.
44. Adu, M.D., et al., *Association between non-tea flavonoid intake and risk of type 2 diabetes: the Australian diabetes, obesity and lifestyle study*. Food Funct, 2022. **13**(8): p. 4459-4468.
45. Mirmiran, P., et al., *Long-term effects of coffee and caffeine intake on the risk of pre-diabetes and type 2 diabetes: Findings from a population with low coffee consumption*. Nutr Metab Cardiovasc Dis, 2018. **28**(12): p. 1261-1266.
46. Shin, S., et al., *Coffee and tea consumption and mortality from all causes, cardiovascular disease and cancer: a pooled analysis of prospective studies from the Asia Cohort Consortium*. Int J Epidemiol, 2022. **51**(2): p. 626-640.
47. Krokstad, S., et al., *Cohort Profile: the HUNT Study, Norway*. Int J Epidemiol, 2013. **42**(4): p. 968-77.
48. Åsvold, B.O., et al., *Cohort Profile Update: The HUNT Study, Norway*. Int J Epidemiol, 2022.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.