

# Invasion of Chronic Lymphocytic Leukemia (CLL/SLL) blood-stream tumor cells by borrelia spirochetes

Alan B. MacDonald MD

Molecular Interrogation of Naples Laboratory; Naples Florida 34114

**Abstract:** A sixty-six-year-old woman with a strong family history of breast cancer noted gradual onset of axillary lymph node enlargement in 2019. Axillary lymph node biopsies disclosed Lymph node involvement by Chronic Lymphocytic Leukemia (CLL/SLL). Immunophenotype analysis of nodal tissue and of bloodstream leukemic cells in immunohistochemistry and flow cytometry and next generation molecular diagnostics confirmed CLL/SLL. Replicate absolute peripheral blood lymphocyte counts in excess of  $5.0 \times 10^3/\text{microliter}$  confirmed the numerical threshold for diagnosis of Chronic Lymphocytic Leukemia in the bloodstream. FISH studies disclosed an immunoglobulin "mutation negative" CLL/SLL genotype without a light chain restriction pattern. No translocations or germline mutations were present. Borrelia spirochetes in the bloodstream were detected with simultaneous high spirochetemia of both *borrelia miyamotoi* and *borrelia burgdorferi*. FISH hybridization of leukemic lymphocytes disclosed extensive borrelia invasion of the cytoplasm and of the nucleus compartments of CLL/SLL tumor cells. Retrospective studies of the patient's archival stored blood smears obtained three years prior in 2016 at age 63 disclosed no evidence of CLL/SLL but confirmed asymptomatic bloodstream *borrelia miyamotoi* and *borrelia burgdorferi* borrelia infection. Lyme serology had been reported as negative in 2016. Additionally, retrospective FISH DNA hybridizations focused on preleukemic bloodstream Lymphocytes disclosed rare borrelia spirochetal adherence to benign blood lymphocytes in the year 2016 in the preleukemic blood smears. This case report is the very first to describe precursor chronic asymptomatic Lyme borreliosis and Miyamotoi borrelia bloodstream infections three years prior to the diagnosis of Rai Stage 1 Chronic Lymphocytic leukemia.

**Keywords:** Borrelia infection; Leukemia; burgdorferi; Miyamotoi

Introduction: Lyme borreliosis and miyamotoi borreliosis infections have diverse clinical presentations which exceed the narrow year 2022 Centers for Disease Control and Prevention list of clinical stigmata and symptoms on the CDC website. [1.] Patients who are clinically refractory to cure with antibiotic treatment for Lyme disease are estimated to constitute 20% of all physician- diagnosed Lyme borreliosis cases. Chronic infection with *borrelia burgdorferi* and *borrelia miyamotoi* includes both symptom free patients and patients with a wide spectrum of symptomatic cases from chronic pain and fatigue syndromes and neuropsychiatric illnesses to some chronic Lyme patients with suicidal ideation or death. Hematologic diseases including cutaneous small lymphocytic malignant lymphomas in Italian and European patients have been diagnosed in association with patients with concurrent *borrelia burgdorferi* infections. [2.10.] Miyamotoi borrelia infections concurrent with nodal B cell lymphomas have been diagnosed in rare patients from Europe and from the USA. [11.12.]

Dorward and colleagues documented in 1987 with the electron microscope in an in vitro laboratory model that borrelia spirochetes show a tropism for attachment to and penetration of human lymphocytes. [13.-14.] Lyme meningitis may present as a marked tumor-like excesses of lymphocytes and plasmacytoid cells which resemble malignant lymphoma in the cerebrospinal fluid and in the meninges.[15.-17.] Borrelia cutis infections may mimic lymphocytic Lymphomas of either the diffuse type or of the follicular type. Borrelia can induce the formation of "Ectopic " germinal centers in cutaneous Borrelia

lymphocytoma sites which rarely have been misinterpreted as deposits of follicular pattern lymphoma in the dermis.[18,21.]

In 2012, a commentary by Stricker and Johnson[ 22.] entitled called attention to the diagnostic shortcomings of current negative blood serology testing techniques for Lyme disease related antibodies and warned that negative Lyme serology test results in physician diagnosed cases of Lyme borreliosis might be augmented by direct microscopic detections of borrelia infections in blood and in tissue specimens in patients. Such chronic active borrelia infections without a concurrent malignant diagnosis and possibly might guide clinical thinking in select human malignancies which co-exist with chronic borrelia infections. Stricker and Johnson discussed chronic lymphocytic leukemia as a possible disease condition for research scrutiny for associated borrelia infections would be germane.[23.]

This case report begins with a previously healthy woman with previous asymptomatic mixed species borrelia bloodstream infection three years previously, who in year 2019 palpated enlarged lymph nodes in her axilla. Her family history was positive for breast cancer and remarkable for one brother with a 30 year history of CLL/SLL who is alive and well on Rituxan therapy.. Needle biopsies demonstrated increased numbers of small lymphocytes permeating bilateral axillary nodal sites as diffuse infiltrates. Immunohistochemistry confirmed a diagnosis of Chronic Lymphocytic Leukemia /Small Lymphocytic Lymphoma (CLL/SLL). FISH studies of the lymph nodes determined a “mutation negative” light chain genotype CLL/SLL. Her peripheral blood in Flow cytometry and in immunohistochemistry (IHC) nodal studies demonstrated that her leukemic cells were positive with the following markers: CD5( Dim), CD19, CD 23( bright). Kappa and Lambda stains disclosed no light chain restriction pattern. Radiology imaging confirmed enlargement of lymph nodes in bilateral axillary and scapular sites without enlargement of her spleen of liver and without lymphadenopathy in other sites. She was determined to be clinically stable, and she did not receive chemotherapy.

Three years prior to the discovery of her CLL/SLL diagnosis, in 2016 the patient had volunteered to participate in a pro bono research study directed by this author to evaluate the use of microscopy to search for *borrelia burgdorferi* in her peripheral blood. She had not received a Lyme vaccine previously and was ELISA seronegative for Lyme related antibodies in 2016. Bloodstream borrelia was identified with acridine orange staining and her absolute counts of bloodstream lymphocytes were normal in number and in morphology. She did not receive antibiotic therapy. In 2022, after being notified of the year 2019 diagnosis of CLL/SLL, the author retrieved from storage archival blood smears from 2016. DNA probes in FISH method confirmed positive hybridizations for *Borrelia* gene sequences in single separate borrelia in the bloodstream and biofilm communities diagnostic of chronic blood stream infection. Leukemic phase blood smears in 2022 also showed persistence of chronic biofilm positive double borrelia species infection. The author then commenced a search for possible invasion by borrelia of leukemic lymphocytes in the year 2022 blood films motivated by the Italian patient reports of concurrent B cell lymphomas in skin sites and concurrent *Borrelia burgdorferi* infections.

## Methods:

Blood smears : EDTA anticoagulated peripheral blood was smeared onto clean glass slides, and the slides were air dried, and heat fixed, followed by post fixation in 99% isopropyl alcohol, and again air dried and stored for use in FISH studies.

Wright Giemsa Stains : Blood smears were stained with Wright Giemsa stain according to the manufacturer’s recommendation.(Volusol Inc)

Immunohistochemistry: Monoclonal antibody CB10 [24.] which uniquely binds to *borrelia burgdorferi* Outer surface protein A (OSP A) conjugated to Horseradish peroxidase was layered over blood smears previously fixed in absolute alcohol and incubated at room temperature for 30 minutes followed by triplicate washes with PBS pH 7.0.

Diaminobenzidine chromogen ( Vector Labs catalog SK 4105 ImmPACT DAB substrate kit, peroxidase) was applied according to the manufacturer's recommendations. All positive and negative controls produced expected results.

FISH : Fluorescence in situ hybridization was completed as previously described. Briefly, single DNA probes were individually dropped onto the blood films and the slides were then flooded with 100% DMF reagent. The flooded slides were then individually heated ( Not cover slipped) to 115 Degrees F for 30 minutes and carefully observed to assure that the DNA probes in solution did not evaporate . The slides were then removed from the heating block and were cover slipped over RPMI tissue culture medium and examined under oil immersion 1000x final magnification with an LED monochromatic light source delivered through a substage darkfield condenser to detect tissue bound sites of FISH DNA probe hybridizations. The photography was completed with a 16 MP digital camera mounted on a trinocular microscope.

#### FISH DNA probe Reagents for borrelia species gene target sites

*Borrelia burgdorferi* B31 strain DNA probe for **gene bbo 0147 for the flagellin B gene of borrelia burgdorferi:**

5' - (FITC)-cacggt-TAATCTTACCAGAACTCCC-accgtg -(Dabcyl)-3' {green color}

DNA probe for gene **bb00740 inner cell membrane** [common to thirty *borrelia burgdorferi* strains (GenBank) sequences:

[5' -( Cy5) -cgcgagATATATTCAAGCAAATTCGATGA CATC-ctcgcg-(Dabcyl -)3']. (red color)

*Borrelia Miyamotoi* strain LB2001 DNA probe for **gene Miyamotoi Flagellin B** (BA1\_FlaB\_Miy\_mb3)

5' (-Cy5) -cgtccg- CGTCAGCCATAAATGCTTCCAGAAATAA-ccgagc -(BBQ650)-3' (red color)

*Borrelia miyamotoi* Strain LB2001: DNA probe for **gene GlpQ of borrelia miyamotoi** ,(BA2\_glpQ\_Miy\_MB3) (yellow color)

5' - (Cy3)-cgtcgg- AGAACATACCTTAGAAGCTAAAGCATATGCCcgagc- (BHQ\_2)-3' (yellow color)

Because the patient resided in Wisconsin, which is hyperendemic for ticks carrying *borrelia burgdorferi*, *borrelia miyamotoi*, and *babesia Duncani*, special manual microscopy of Giemsa-stained blood films was undertaken to search for babesia co-infection in peripheral blood smears. This required microscopy at 1000x oil immersion magnification to search for ring trophozoite babesia inclusions inside erythrocytes in one hundred consecutive microscopic fields of view.

Coombs method for detection antibody coated borrelia spirochetes and antibody coated borrelia biofilm communities was deployed as previously published. Coombs antiglobulin reagent labeled with a blue fluorochrome Biotium mix N stain 403 was layered over thin blood smears and incubated at room Temperature for 30 minutes, and then the slides were washed three times in PBS pH7.4. and examined in Immunofluorescence microscopy.[25.]

### Controls:

Positive Control for *borrelia burgdorferi* used pure cultures strain B31 *borrelia burgdorferi* of American Type Culture Collection ATCC 35210 in log phase culture thin smears on clean glass slides fixed in Absolute isopropyl alcohol and post alcohol heat fixed.

Positive Control for *borrelia miyamotoi* used pure cultures of *Borrelia miyamotoi* strain LB2001 thin smears fixed on clean glass slides ( a gift from Catherine Brissette PhD, School of Medicine, University of North Dakota)

Negative tissue controls used normal autopsy liver tissue sections and peripheral blood smears from normal patient volunteers.

Negative IHC controls used Antibodies to Alpha Synuclein ( StressMarq Inc) and antibodies to Cytokeratin Ae1/Ae3. ( SantaCruz Biotechnology Inc. )

Negative Controls for FISH used A Nonsense DNA probe Gn2, which by design functioned as a negative control in FISH studies was designed such that its structure does not correspond to any known DNA sequence in GenBank deposits at the NCBI NIH site.

5' (-Marina Blue NHS AmMC6)-cgcgat- TCTCCGAACGTGTCACGC-gtcgcg-(BHQ\_1)-3' {Nonsense probe}

### Results:

Blood smears from 2016 showed normal lymphocyte morphology in Wright Giemsa stains in peripheral blood lymphocytes. Rare benign blood lymphocytes demonstrated questionable borrelia like forms in adjacent to lymphocyte cell membrane sites. No evidence of spirochetes compatible with borrelia inside lymphocyte nucleus compartments was found. (Figure 1) .

Blood smears from year 2022 blood films demonstrated leukemic nuclei of larger caliber lymphocytes than those seen in the 2016 blood lymphocytes. Giemsa stains disclosed spirochetes compatible with borrelia inside leukemic cells. The full hematopathology repertoire of Immunohistochemical staining CD markers confirmed chronic lymphocytic leukemia/ small lymphocytic lymphoma. (Figure 2)

Immunohistochemistry studies in 2016 blood smears demonstrated borrelia specific protein OSP A borrelia spirochetes attached to the cell membrane of rare blood benign lymphocytes (Figure 3)

Extensive invasion of leukemic blood lymphocytes as found in 2022 smears in immunohistochemistry detection of Outer surface protein A of *borrelia burgdorferi* resident inside leukemic cells. IHC studies for *borrelia miyamotoi* specific proteins were not undertaken due to the lack of available commercially available antibodies to *borrelia miyamotoi* protein GLPQ and due to the availability of FISH DNA hybridization results for the GLPQ gene expression in the borrelia resident inside the patient's leukemic lymphocytes. (Figure 4)

FISH studies of smears from 2022 demonstrated borrelia DNA content inside live *borrelia burgdorferi* and *borrelia miyamotoi* spirochetes which have invaded leukemic lymphocytes in blood. Leukemic cells contained both *borrelia burgdorferi* (sl) group spirochetes and *borrelia miyamotoi* spirochetes. Cytoplasmic and nuclear compartments of the leukemic cells contained intact borrelia spirochetes. [Figure 5 and Figure 6]

Wright Giemsa-stained blood smears from 2022 disclosed Babesia ring trophozoites in red blood cells. The species of babesiosis was not determined. The estimated percentage of parasitized RBC was between 1% and 5% in serial counts of 1000 erythrocytes in oil immersion magnification examinations.(Figure 7)

Immunofluorescence microscopy detected blue fluorescent staining of single borrelia spirochetes coated with Coombs antibody and of biofilms of borrelia coated with Coombs antibody. (Figure 8)

All positive and negative controls in Immunohistochemistry with *borrelia burgdorferi* and with *Borrelia miyamotoi* DNA probes returned expected results. *Borrelia burgdorferi* and *borrelia miyamotoi* FISH specific DNA probes returned expected positive and negative

results with the cultured reference glass slide control material. The nonsense DNA probe returned negative hybridization results in FISH studies.

**Discussion:** Asymptomatic bloodstream *Borrelia* infections with dual species *burgdorferi* and *miyamotoi* *borrelia* were found in preleukemic blood smears from year 2016. Lymphocytes in blood smears from 2016 were normal in appearance and in total lymphocyte absolute counts in blood in 2016. Retrospective FISH studies of 2016 blood smears with DNA specific gene probes disclosed parasitic *borrelia* spirochetal hybridization signals adherent to the cell membranes of rare benign lymphocytes.

Leukemic phase asymptomatic *Borrelia* infections in year 2022 presented with dual species *burgdorferi* and *miyamotoi* *borrelia*. Double infection of lymphocytes in blood smears from 2022 with *burgdorferi* and *miyamotoi* *borrelia* species were established with FISH studies. The leukemic lymphocyte cytomorphology in 2022 demonstrated enlargement of the nuclei and an uncondensed open chromatin pattern which contrasted with the microscopy from the year 2016 blood lymphocytes whose nuclei were smaller in caliber and showed a more compact condense chromatin pattern.

The differential results from the years 2016 and 2022 FISH focused DNA hybridizations disclosed that outright penetration of the patient's tumoral lymphocytes by live *borrelia* spirochetes had occurred. These spatial and temporal leukemic and infection events overlap with a biological transformation in lymphocyte biology from a benign to a leukemic status as confirmed with Immunohistochemistry results which confirmed a phenotype of Chronic Lymphocytic leukemia. Mere coincidence of ongoing pre-existing *borrelia* bloodstream infection and the separate emergence of lymphocytic leukemia/small lymphocytic lymphoma is one possible conclusion here. However, Leukemic cell parasitism by *borrelia* infection is not a hallmark of usual infection in a neoplastic setting. Further FISH studies with *borrelia burgdorferi* and with *borrelia miyamotoi* specific DNA probes are needed in additional CLL/SLL patients who are residents of endemic *borrelia miyamotoi* and *borrelia burgdorferi* infection to search for evidence of *borrelia* infectomes inside the nuclei of lymphocytic leukemia/Lymphoma tumor cells.

The microscopic detection of simultaneous *borrelia miyamotoi* and *borrelia burgdorferi* infection of human lymphocytes is without precedent. Previous reports have described patients with active *borrelia* infections in CLL patients. These cases have been confirmed with PCR positive confirmation of *borrelia burgdorferi* DNA in biopsies from CLL/SLL patients whose illnesses include cutaneous B cell lymphomatous infiltrates. Previous authors have concluded that cutaneous B cell lymphomas with associated *borrelia* infections were mere consequences of the immune system impairments of patients by their CLL/SLL illnesses. In this case, host antibodies coated *borrelia* spirochetes in Coombs reagent testing, and this observation argues that the host serological immune system was not silenced and therefore the spatial co-localization of *borrelia* inside leukemic tumor cells merits further research study.

Miyamotoi *borrelia* infections concurrent with B cell Lymphomas in published case reports from two patients establish a second category of concurrent *borrelia* infection and B cell neoplasias. No attempt was made to evaluate the lymphoma cells for possible invasion by *borrelia miyamotoi* spirochetes

The patient presented here is distinguished by seven observational findings which have not been previously described in CLL/SLL *borrelia* infected patients .

First, In this case study, chronic asymptomatic bloodstream *borrelia miyamotoi* and *burgdorferi* infections preceded the diagnosis of CLL/SLL by three years.

Second, dual species infection with *miyamotoi* *borrelia* and *burgdorferi* group *sl borrelia* in the bloodstream of a CLL/SLL patient is a first of kind observation.

Third, outright invasion and cell membrane adherence of benign lymphocytes by *borrelia burgdorferi* has never previously been observed in vivo but has been documented in laboratory conditions.

Fourth, outright invasion of Leukemic lymphocytes by *borrelia* spirochetes of any species has never previously been observed.



Fifth, *Borrelia miyamotoi* infections have been previously identified in patients with B cell lymphoid malignancies but have never been shown to invade neoplastic B cells.

Sixth, biofilm communities of *Borrelia* spirochetes which, confirm a diagnosis of chronicity of infection, were identified in patient blood smears in her preleukemic blood in year 2016 and were identified in her leukemic blood smears in year 2022. Biofilm infections are a type of chronic infection which has been linked to oncogenic activity in several human malignancies.

Seventh, Leukemic blood smears demonstrated *Babesia* ring trophozoites inside erythrocytes in 2022 which declare emergence of *Babesia* co-infection which was not present in her 2016 blood smears.

Eighth, the *Borrelia* spirochetes in preleukemic blood smears from 2016 and in leukemic blood smears were coated with the patient's antibodies in a Coombs method test indicating that the patient's immune system generated a serological host response to *Borrelia* infections. (Figure 8)

**Acknowledgments:** Hematopathology and Molecular diagnostic testing for Chronic Lymphocytic Leukemia were completed at Froedtert Hospital, Milwaukee, Wisconsin.


## References –

1. CDC website Lyme disease case 2022 definitions: <https://ndc.services.cdc.gov/case-definitions/lyme-disease-2022/>
2. Kash N, Fink-Puches R, Cerroni L. Cutaneous manifestations of B-cell chronic lymphocytic leukemia associated with *Borrelia burgdorferi* infection showing a marginal zone B-cell lymphoma-like infiltrate. The American journal of dermatopathology. 2011 Oct 1;33(7):712-5.
3. Roggero E, Zucca E, Mainetti C, Bertoni F, Valsangiacomo C, Pedrinis E, Borisch B, Piffaretti JC, Cavalli F, Isaacson PG. Eradication of *Borrelia burgdorferi* infection in primary marginal zone B-cell lymphoma of the skin. Human pathology. 2000 Feb 1;31(2):263-8.
4. Maraspin V, Ružić-Sabljic E, Lusa L, Strle F. Course, and outcome of early Lyme borreliosis in patients with hematological malignancies. Clinical Infectious Diseases. 2015 Aug 1;61(3):42
5. Colli C, Leinweber B, Müllegger R, Chott A, Kerl H, Cerroni L. *Borrelia burgdorferi*-associated lymphocytoma cutis: clinicopathologic, immunophenotypic, and molecular study of 106 cases. Journal of cutaneous pathology. 2004 Mar;31(3):232-40.
6. Chang ET, van Roos G, Hansen M, Adami HO, Hjalgrim H. *Borrelia* infection and risk of non-Hodgkin lymphoma. BLOOD. 2008 Jun 15;111(12).
7. Tothova SM, Bonin S, Trevisan G, Stanta G. Mycosis fungoides: is it a *Borrelia burgdorferi*-associated disease? British journal of cancer. 2006 Mar;94(6):879-83.
8. Solaiman F, Afshan M, Dana S. The Challenges of Diagnosing Disseminated Lyme Disease in a Patient with CLL. Open Journal of Blood Diseases. 2012 Mar 5;2012.
9. Futuri S, Malik A, Shani D. The Challenges of Diagnosing Disseminated Lyme Disease in a Patient with CLL. Open Journal of Blood Diseases. 2012 Mar 5;2(1):14-5.
10. Ali L, Cheney R, Merzianu M. Subclinical chronic lymphocytic leukemia with atypical cutaneous presentation. Journal of
11. Gugliotta, J.L. et al "Meningoencephalitis from *Borrelia miyamotoi* in an immunocompromised patient", New England J. Medicine, 2013, 368: (3), 240-245.
12. Hovius J. W. R., et al, "A case of meningoencephalitis by the relapsing fever spirochete in Europe," Lancet 2013;382: (9892) 658
13. Dörward, D.W., et al, "Invasion and cytopathic killing of human lymphocytes by spirochetes causing Lyme disease", Clin Inf Dis 1997 25: (supp 1); S2-S8
14. Duray, P.H. et al, "Invasion of human Tissue ex vivo by *Borrelia burgdorferi*", JID, 2005, 191(10) 1747-1754
15. Schweighofer CD, Fätkenheuer G, Staib P, Hallek M, Reiser M. Lyme disease in a patient with chronic lymphocytic leukemia mimics leukemic meningoencephalitis. Oncology Research and Treatment. 2007;30(11):564-6.
16. Kalac M, Suvic-Krizanic V, Ostojic S, Kardum-Skelin I, Barsic B, Jaksic B. Central nervous system involvement of previously undiagnosed chronic lymphocytic leukemia in a patient with neuroborreliosis. International journal of hematology. 2007 May;85(4):323-5.
17. Harrer T, Geissdörfer W, Schoerner C, Lang E, Helm G. Seronegative Lyme neuroborreliosis in a patient on treatment for chronic lymphatic leukemia. Infection. 2007 Apr;35(2):110-3.
18. Beda IM, Karla EB, Almudena CC, Flavio GG, Armando RP, Patricia VF. B-cell leukemia/lymphoma and neuroborreliosis: Diagnostic and therapeutic challenges. Journal of Vector Borne Diseases. 2017 Jan 1;54(1):106.
19. Raufi A, Alsharedi M, Khelfa Y, Griswold DC, Lebowicz Y. Leukemia cutis in a patient with chronic lymphocytic leukemia presenting as bilateral helical nodules. SAGE Open Medical Case Reports. 2016 Dec;4:2050313X16683624

20. Koletsis T, Patsatsi A, Kostopoulos I, Georgiou E, Sotiriadis D. Earlobe involvement in chronic lymphocytic leukemia. JDDG: Journal der Deutschen Dermatologischen Gesellschaft. 2013 11(1):80-2
- 21.. Riede I. Borrelia infection appears as chronic lymphocytic leukemia: Therapy with Amanita phalloides and Terebinthina laricina. Br. J. Med. Med. Res. 2015;7(7):630-7.
22. Stricker and Johnson Stricker RB, Johnson L. Risk of malignancy associated with Lyme disease: Still up in the air. International journal of cancer. 2012 Dec 1;131(11):2717
- 23.. Cerroni L, Höfler G, Bäck B, Wolf P, Maier G, Kerl H. Specific cutaneous infiltrates of B-cell chronic lymphocytic leukemia (B-CLL) at sites typical for *Borrelia burgdorferi* infection. Journal of cutaneous pathology. 2002 Mar;29(3):142-7.
24. Benach, J.L., Coleman, J, Golightly, M.J., "A murine IgM monoclonal antibody binds an antigenic determinant in outer surface protein A, an immunodominant basic protein of the Lyme disease spirochete," J. Immunology, 1988, 140(10) 265-272
25. MacDonald, A.B. "Lyme ELISA negative patients with borrelia antibody coated borrelia spirochetes in their blood" , Medical and Clinical Research, 2022, 7(9); 1-4


Supplementary Data :

Supplementary File S1: Bloodstream infection with dual species of borrelia ( Burgdorferi and Miyamotoi)




Supplementary File S2:

Size Grid – 400x objective



Ten Microns per division




Dual species infection ( borrelia burgdorferi (red color – Monoclonal Ab CB10 – OspA specific) and  
Borrelia miyamotoi – blue color – Miyamotoi borrelia does not contain OspA protein  
400 x final magnification – IHC with CB10 monoclonal Ab and mixed with Acridine orange stain – blue  
color

Note: Protein studies demonstrate two populations of borrelia spirochetes; one population displays  
Red color fluorescence signals of Protein OSP A and one population which is negative for red color signals ( No con-  
tent of Protein OSP A). Miyamotoi borrelia lack Protein OSP A. Burgdorferi borrelia group (SL) universally contain pro-  
tein OSP A.

Supplementary File S3:





*Borrelia burgdorferi* spirochetes ( red color) and *borrelia miyamotoi* spirochetes (blue color)

Indicate dual species infection in this patient. Final Magnification 1000 x original.

Note : Select spirochetes display both red color and blue color signals. This is suggestive of a chimeric borrelia microbe which contains dual borrelia species gene DNA.

Supplementary File S4:

**Borrelia burgdorferi biofilm community ( Red color plaque) in peripheral blood patient E.M. IHC with Monoclonal Ab CB10 – specific for OSP A protein of burgdorferi group spirochetes.**

**Note:** a preponderance of granular type borrelia ( Red dots) populates the biofilms. Rare cylindrical type borrelia are present at the radial outer margins of each biofilm. Extracellular Protein OspA Deposits are diffusely present inside the amorphous matrix “slime layer” inside the biofilm.

**Black spaces inside the biofilm communities are “water channel networks” which provide conduits for Nutrient influxes and for “waste removal” channels for living microbes inside the biofilm.**

**Final Magnification 1000x original.**

