

Article

Not peer-reviewed version

Lipid-Associated GWAS Loci Predict Antiatherogenic Effects of Rosuvastatin in Patients with Coronary Artery Disease

Stanislav Kononov , [Iuliia Azarova](#) , [Elena Klyosova](#) , Marina Bykanova , [Mikhail Churnosov](#) , Maria Solodilova , [Alexey Polonikov](#) *

Posted Date: 19 May 2023

doi: 10.20944/preprints202305.1365.v1

Keywords: coronary artery disease; plasma lipids; carotid intima-media thickness; pharmacogenetics; personalized medicine; single nucleotide polymorphisms; lipid-lowering therapy; rosuvastatin.



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Lipid-Associated GWAS Loci Predict Antiatherogenic Effects of Rosuvastatin in Patients with Coronary Artery Disease

Stanislav Kononov ¹, Iuliia Azarova ^{2,3}, Elena Klyosova ^{3,4}, Marina Bykanova ^{4,5}, Mikhail Churnosov ⁶, Maria Solodilova ⁴ and Alexey Polonikov ^{4,7,*}

¹ Department of Internal Medicine No 2, Kursk State Medical University, 3 Karl Marx Street, 305041 Kursk, Russia

² Department of Biological Chemistry, Kursk State Medical University, 3 Karl Marx Street, 305041 Kursk, Russia

³ Laboratory of Biochemical Genetics and Metabolomics, Research Institute for Genetic and Molecular Epidemiology, Kursk State Medical University, 18 Yamskaya Street, 305041 Kursk, Russia

⁴ Department of Biology, Medical Genetics and Ecology, Kursk State Medical University, 3 Karl Marx Street, 305041 Kursk, Russia

⁵ Laboratory of Genomic Research, Research Institute for Genetic and Molecular Epidemiology, Kursk State Medical University, 18 Yamskaya Street, 305041 Kursk, Russia

⁶ Department of Medical Biological Disciplines, Belgorod State University, 85 Pobedy Street, 308015 Belgorod, Russia

⁷ Laboratory of Statistical Genetics and Bioinformatics, Research Institute for Genetic and Molecular Epidemiology, Kursk State Medical University, 18 Yamskaya Street, 305041 Kursk, Russia

* Correspondence: polonikov@rambler.ru

Abstract: We have shown that lipid-associated loci discovered by genome-wide association studies (GWAS) have pleiotropic effects on lipid metabolism, carotid intima-media thickness (CIMT), and CAD risk. Here, we investigated the impact of lipid-associated GWAS loci on the efficacy of rosuvastatin therapy in terms of changes in plasma lipid levels and CIMT. The study comprised 116 CAD patients with hypercholesterolemia. CIMT, total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and triglycerides (TG) were measured at baseline and after 6 and 12 months of follow-up, respectively. Genotyping of fifteen lipid-associated GWAS loci was performed by the MassArray-4 System. Linear regression analysis adjusted for sex, age, body mass index, and rosuvastatin dose was used to estimate the phenotypic effects of polymorphisms, and P-values were calculated through adaptive permutation tests. Over one-year therapy, a decrease in CIMT was linked to rs1689800, rs4846914, rs12328675, rs55730499, rs9987289, rs11220463, rs16942887, and rs881844 polymorphisms ($P_{\text{perm}} < 0.05$). TC change was associated with rs55730499, rs11220463, and rs6065906; LDL-C change was linked to the rs55730499, rs1689800, and rs16942887 polymorphisms; and TG change was linked to polymorphisms rs838880 and rs1883025 ($P_{\text{perm}} < 0.05$). In conclusion, the polymorphisms rs1689800, rs55730499, rs11220463, and rs16942887 were found to be predictive markers for multiple antiatherogenic effects of rosuvastatin in CAD patients.

Keywords: coronary artery disease; plasma lipids; carotid intima-media thickness; pharmacogenetics; personalized medicine; single nucleotide polymorphisms; lipid-lowering therapy; rosuvastatin

1. Introduction

Atherosclerosis is a complex disease, which pathogenesis involving numerous pathological processes increased plasma cholesterol, its deposition in the arterial wall, endothelial dysfunction and vascular remodeling. These processes are influenced both by environmental and genetic factors [1,2]. It is widely agreed that dyslipidemia plays a crucial role in the development of atherosclerosis

and coronary artery disease (CAD) [2]. Lipid metabolism abnormalities and coronary heart disease are determined by substantial effects of genetic factors [3–5]. According to the GWAS Catalog (<https://www.ebi.ac.uk/gwas/home>), approximately 1700 genome-wide association studies (GWAS) on over 450 cardiovascular phenotypes have been conducted, yielding substantial insights into the genetic etiology of cardiovascular disease and related traits [6]. Numerous GWAS including large-scale studies of the Global Lipids Genetics Consortium have discovered loci associated with plasma lipids [7–12]. GWAS were also conducted to find genetic determinants of statin pharmacogenetics, including JUPITER, the largest study of lipid-lowering effects of rosuvastatin [13–15]. Statins are the first line of lipid-lowering drugs widely used in both primary and secondary prevention of CAD. For one of the most popular statins, rosuvastatin, it was shown that it could reduce both blood lipid levels, including total cholesterol (TC), LDL-C (low-density lipoprotein cholesterol), triglycerides (TG), and carotid intima-media thickness (CIMT) [16].

Carotid intima-media thickness is a surrogate marker for the presence and progression of atherosclerosis, useful for evaluating risk and incidence of cardiovascular disease [17]. A causal relationship between LDL-C and CIMT has been observed [18], and CIMT is being considered as a marker of preclinical atherosclerosis [18,19]. A growing body of evidence has been provided to consider CIMT as a predictive marker for coronary artery atherosclerosis, its severity, and the extent of plaque burden [20–23]. Importantly, rosuvastatin may reduce not only CIMT but also atherosclerotic plaque growth [24,25]. It is impossible to claim definitely whether hypolipidemic effects of rosuvastatin are solely responsible for the regression of atherosclerosis, because pleiotropic effects of the drug may contribute to in arterial wall changes through inflammation reduction [25,26]. The mechanisms by which rosuvastatin is responsible for vascular wall changes may be unraveled by pharmacogenetic studies; however, a majority of such studies have investigated only lipid-lowering effects of the drug [13,14,27,28]. Our research team has conducted several studies on the effect of gene polymorphisms on plasma lipids and CIMT, as well as the dynamics of their changes during rosuvastatin lipid-lowering therapy [29–31]. It has been discovered that some lipid-associated GWAS loci contribute to lipid-lowering effects associated with rosuvastatin therapy and determine CIMT regression. Without a doubt, CIMT seems to be a surrogate marker of atherosclerosis in CAD patients, despite it correlates with disease severity [20–23]. However, in terms of clinical application, CIMT regression during rosuvastatin therapy may mirror vascular wall changes and be indicative of drug efficacy. The aim of our pharmacogenetic study was to identify the effects of fifteen lipid-associated GWAS loci such as rs4846914 (*GALNT2*), rs11220463 (*ST3GAL4*), rs881844 (*STARD3*), rs1689800 (*ZNF648*), rs12328675 (*COBLL1*), rs9987289 (*PPP1R3B*), rs55730499 (*LPA*), rs3136441 (*F2*), rs6065906 (*PLTP*), rs838880 (*SCARB1*), rs386000 (*LILRA3*), rs1883025 (*ABCA1*), rs3764261 (*CETP*), rs217406 (*NPC1L1*), and rs16942887 (*PSKH1*) on plasma lipids and CIMT in CAD patients taking rosuvastatin. These loci have recently been observed to be associated with CAD susceptibility and CIMT [32].

2. Materials and Methods

Study design. The present research was designed as a prospective study, in which the participants received rosuvastatin at an initial dose of 5 mg. Then, in 4 weeks, we controlled the levels of LDL-C. In cases where the levels were below the target level (1.8 mmol/l), patients continued taking the dose, which let them attain the target level. If it was not attained, the dose of rosuvastatin was gradually increased to 10, 20, and finally the maximum dose of 40 mg. The control of LDL-C levels was performed every 4 weeks after the dose increase. For the association analysis, we used the change (delta, Δ) in TC, LDL-C, TG, and CIMT values over the first 6 and 12 months of the study.

Study participants. The patient cohort comprised 116 Russian patients with the diagnosis of CAD, stable angina pectoris grade II–III. The study participants were described in detail in our previous work [31]. In brief, the cohort included 85 men (73%) and 31 postmenopausal women (27%), with the mean age of 61.0 ± 7.25 years. Approximately 58% of CAD patients experienced a myocardial infarction in the past, and 97.5% of study participants had arterial hypertension. Diagnosis of CAD was confirmed by qualified cardiologists according to the Canadian Cardiovascular Society, ECG

stress tests (treadmill test), and 24-h Holter's ECG monitoring. TC levels higher than 4.0 mmol/l and LDL-C levels higher than 1.8 mmol/l were considered as criteria for dyslipidemia.

Biochemical and ultrasound investigations were described in our previous paper [31]. Plasma lipid levels, except LDL-C, were detected using an automatic laboratory analyzer. LDL-C were calculated using Friedewald's equation. CIMT measurement was performed in B-mode using the standard method at the distal third of the common carotid artery at a distance of 1–1.5 cm proximal to the bifurcation along the posterior wall [33]. The following two parameters of carotid intima-media thickness were used in the study: mean CIMT (mean value of all measurements on the right and left sides) and maximal CIMT (the maximal value obtained from measurements on both sides with the further assessment of its change on the side where it was the highest).

Genetic analysis. Single nucleotide polymorphisms (SNPs) selected for this study have previously been found to be associated with plasma lipids in several genome-wide association studies [7–12] (GWAS Catalog (<https://www.ebi.ac.uk/gwas>, accessed date 13 March 2023)). As mentioned above, some of the SNPs have been recently found to be associated with CAD susceptibility and CIMT [32], and we are interested in investigating whether these variants determine the lipid- and CIMT-lowering effects of rosuvastatin in CAD patients. The investigated SNPs included rs4846914 (*GALNT2*), rs11220463 (*ST3GAL4*), rs881844 (*STARD3*), rs1689800 (*ZNF648*), rs12328675 (*COBLL1*), rs9987289 (*PPP1R3B*), rs55730499 (*LPA*), rs3136441 (*F2*), rs6065906 (*PLTP*), rs838880 (*SCARB1*), rs386000 (*LILRA3*), rs1883025 (*ABCA1*), rs3764261 (*CETP*), rs217406 (*NPC1L1*), and rs16942887 (*PSKH1*).

Extraction of DNA was performed from venous blood samples using phenol-chloroform method and precipitation with ethanol. Genotyping was performed using iPLEX technology on the MassARRAY 4 system (Agena Bioscience, San-Diego, CA, USA). Software MassARRAY Assay Design Suite was used to select a primer set and to design a multiplex panel for SNP genotyping. The sequences of primers are available on request.

Statistical analysis. The normality of the distribution of lipid and CIMT values was determined using the Kolmogorov-Smirnov and Shapiro-Wilk tests. As the trait distributions deviated from the normal one, the values were expressed as median and interquartile range (Me, Q1; Q3). Statistical package STATISTICA v13.0 (Statsoft, Tulsa, OK, USA) was utilized for descriptive statistics, distribution analysis, and determining the significance of lipid and CIMT changes. The significance of lipid and CIMT change during rosuvastatin therapy was tested by the Wilcoxon's matched pairs test. The distribution of genotype frequencies according to Hardy-Weinberg equilibrium was assessed with Fisher's exact test. For the association analysis, we used linear regression with adjustments for sex, age, body mass index (calculated as body mass in kilograms divided by height in meters in a square), and rosuvastatin dose, which allowed the participants to attain the target LDL-C level. Change in lipid levels was calculated as the difference between natural log-transformed on- and off-treatment levels divided by the natural log-transformed off-treatment level as described by Postmus et al. [13]. We applied the same approach for the CIMT change, but before the logarithmic transformation, the CIMT data were multiplied by 10. Empirical P-values (Pperm) were calculated through the adaptive permutation procedure. We tested three genetic models for SNP-phenotype associations, such as additive, dominant, and recessive. The PLINK v1.92 software [34] was used for all genetic calculations, including estimation of minor allele frequencies (MAF), tests for Hardy-Weinberg equilibrium (HWE), multiple regression analysis, and permutation procedures. When the p-value was less than 0.05, all of the results were declared statistically significant.

3. Results

The majority of the patients (N=110, 94.8 %) attained their target LDL-C level during rosuvastatin therapy. The change in lipid and CIMT values was assessed during 6 and 12 months of observation. The reduction of all lipid parameters ($p < 1 \times 10^{-6}$ for TC, LDL-C, and TG change during 6- and 12-month periods) was significant, except HDL-C levels ($P=0.16$ and 0.87 , respectively). Reductions in the maximum ($p < 1 \times 10^{-6}$) and mean CIMT ($P=0.025$ and $< 1 \times 10^{-5}$ for the change in 6 and 12 months, respectively) were also significant (Supplementary Table 1).

3.1. Associations of the SNPs with lipid and CIMT reduction during the 6-month therapy by rosuvastatin.

The distribution of genotypes for the majority of SNPs (except for rs16942887) was in Hardy-Weinberg equilibrium. The observed effects of SNPs on the lipid-lowering effect and CIMT regression in CAD patients taking rosuvastatin for the 6-month period are shown in Table 1. We found that plasma TC reduction was associated with two SNPs, such as rs55730499 of *LPA* (beta = 0.323, Pperm=0.0022, recessive effect) and rs6065906 of *PLTP* (beta = 0.140, Pperm=0.0135, recessive effect). Based on the beta coefficients, it can be estimated that the carriage of homozygous genotypes for the minor alleles of the above polymorphisms is associated with 32.3% and 14.0 % of TC reduction, respectively, compared with the carriers of alternative genotypes (the positive beta indicates a worse lipid reduction and consequently a worse statin response). LDL-C reduction was found to be associated with rs55730499 of the *LPA* gene (Pperm=0.0224, recessive effect), and rs1689800 of the *ZNF648* gene (Pperm=0.0493, additive effect). The rs1689800G allele is associated with a 4.6 percent reduction in plasma LDL-C levels after 6 months of rosuvastatin therapy. The effect size for the rs55730499 polymorphism of *LPA* was greater in the TT homozygotes and characterized by a 50.4% decrease in LDL-C concentration than in carriers of alternative genotypes.

Table 1. Associations of single nucleotide polymorphisms with the 6-month lipid-lowering effect (total cholesterol, low density lipoprotein cholesterol) and carotid intima-media thickness change in coronary artery disease patients taking rosuvastatin/.

Chr	Gene (SNP ID)	Effect allele	EAF	N	Total cholesterol		LDL-C		CIMT, maximum		CIMT, mean	
					Beta*	Pperm [#]	Beta*	Pperm [#]	Beta*	Pperm [#]	Beta*	Pperm [#]
1	<i>ZNF648</i> (rs1689800)	G	0.392	115	0.020	0.1786	0.046	0.0493^A	-0.084	0.0234^R	-0.021	0.2308
1	<i>GALNT2</i> (rs4846914)	G	0.388	115	0.004	0.7778	-0.005	0.8571	-0.045	0.0133^A	-0.038	0.0344^A
2	<i>COBL1</i> (rs12328675)	C	0.170	111	-0.003	0.8571	-0.003	1.0000	0.035	0.2647	0.041	0.1000
6	<i>LPA</i> (rs55730499)	T	0.056	115	0.323	0.0022^R	0.504	0.0224^R	-0.010	0.8571	0.028	0.8571
7	<i>NPC1L1</i> (rs217406)	G	0.203	115	0.017	0.2308	0.033	0.3556	-0.001	1.0000	-0.003	1.0000
8	<i>PPP1R3B</i> (rs9987289)	A	0.086	115	0.019	0.3404	-0.013	0.8571	0.041	0.3404	0.051	0.1550
9	<i>ABCA1</i> (rs1883025)	T	0.263	115	0.001	1.0000	-0.004	0.8571	-0.024	0.2982	-0.025	0.2535
11	<i>F2</i> (rs3136441)	C	0.180	102	0.011	0.6429	0.025	0.7778	0.021	0.6923	0.023	0.2982
11	<i>ST3GAL4</i> (rs11220463)	T	0.190	115	-0.031	0.1280	-0.068	0.0803	0.066	0.0159^D	0.061	0.0243^A
12	<i>SCARB1</i> (rs838880)	C	0.336	115	0.002	1.0000	0.005	1.0000	-0.014	0.7273	-0.015	0.6923
16	<i>CETP</i> (rs3764261)	A	0.147	115	0.007	0.6250	0.027	0.8571	-0.045	0.2466	-0.042	0.1900
16	<i>PSKH1</i> (rs16942887)	A	0.116	115	0.009	0.6429	0.025	0.4643	0.066	0.0421^D	0.068	0.0483^D
17	<i>STARD3</i> (rs881844)	C	0.310	115	0.003	0.8571	-0.038	0.1667	0.048	0.0086^A	0.057	0.0033^A
19	<i>LILRA3</i> (rs386000)	C	0.203	115	0.006	0.5455	-0.001	1.0000	0.015	0.7778	0.003	0.8571
20	<i>PLTP</i> (rs6065906)	C	0.160	115	0.140	0.0135^R	-0.008	1.0000	-0.022	0.6923	-0.022	0.5455

Chr, chromosome; SNP, single nucleotide polymorphism; EAF, effect allele frequency; LDL-C, low-density lipoprotein cholesterol; CIMT, carotid intima-media thickness.

*Beta for difference between the natural log-transformed on- and off-treatment values adjusted for natural log-transformed off-treatment level. The beta reflects the fraction of differential lipid- or CIMT-lowering effect in carriers versus non-carriers of the minor allele (for additive model), in carriers of the genotype according to the model (dominant, recessive) versus the carriers of alternative genotypes; a negative beta indicates a better statin response (stronger lipid or CIMT reduction), a positive beta – a worse statin response. Betas were generated using linear regression analysis with age, sex, body mass index, and rosuvastatin dose as covariates.

[#]p-values were generated using the permutation procedure. Superscript indicates genetic model: R, recessive; A, additive; D, dominant.

As it can be seen from Table 1, four lipid-associated GWAS loci were associated with both maximum and mean CIMT change: rs4846914 in *GALNT2* (Pperm=0.0133 and 0.0344 for the association with maximum and mean CIMT change, respectively), rs11220463 in *ST3GAL4* (Pperm=0.0159 and 0.0243, respectively), rs16942887 in *PSKH1* (Pperm=0.0421 and 0.0483, respectively), and rs881844 in *STARD3* (Pperm=0.0086 and 0.0033, respectively). SNP rs1689800 in the *ZNF648* gene (Pperm=0.0234) was associated with a change in the maximum CIMT. The most significant association with TC and LDL-C reduction during 6 months of rosuvastatin therapy was

found for SNP rs55730499 in *LPA*, whereas the rs881844 polymorphism of the *STARD3* gene showed the most substantial association with the regression of CIMT in CAD patients.

Table 2 shows associations between the SNPs and plasma triglycerides reduction. It has been revealed that the only SNP rs838880 of the *SCARB1* gene was associated with TG reduction (beta = 1.736, $P_{perm}=0.0478$, recessive effect) after 6 months of lipid-lowering therapy.

Table 2. Associations of single nucleotide polymorphisms with triglyceride-lowering effect in coronary artery disease patients taking rosuvastatin.

Chr	Gene (SNP ID)	Effect allele	EAF	N	6-month period		12-month period	
					Beta*	$P_{perm}^{\#}$	Beta*	$P_{perm}^{\#}$
1	ZNF648 (rs1689800)	G	0.392	114	-0.6495	0.1148	-0.1946	0.4643
1	GALNT2 (rs4846914)	G	0.388	114	0.4816	0.1919	0.05046	0.7778
2	COBLL1 (rs12328675)	C	0.170	110	-0.2948	0.6250	-0.1431	0.7778
6	LPA (rs55730499)	T	0.056	114	-0.5923	0.4242	0.1825	0.6923
7	NPC1L1 (rs217406)	G	0.203	114	0.4106	0.4118	-0.1208	0.6429
8	PPP1R3B (rs9987289)	A	0.086	114	-0.9197	0.1887	-0.7977	0.0756
9	ABCA1 (rs1883025)	T	0.263	114	-0.177	0.5789	-0.7246	0.0160^D
11	F2 (rs3136441)	C	0.180	101	-0.185	0.5789	0.02512	1.0000
11	ST3GAL4 (rs11220463)	T	0.190	114	-0.774	0.1587	3.624	0.0503^R
12	SCARB1 (rs838880)	C	0.336	114	1.736	0.0478^R	0.1063	0.6429
16	CETP (rs3764261)	A	0.147	114	0.695	0.2931	0.02338	1.0000
16	PSKH1 (rs16942887)	A	0.116	114	-0.6306	0.1439	-0.2626	0.3947
17	STARD3 (rs881844)	C	0.310	114	0.4075	0.4815	-0.2337	0.2647
19	LILRA3 (rs386000)	C	0.203	114	-0.1847	0.8571	-0.1448	0.6923
20	PLTP (rs6065906)	C	0.160	114	-0.4654	0.5789	-0.0718	0.8571

Chr, chromosome; SNP, single nucleotide polymorphism; EAF, effect allele frequency.

*Beta for difference between the natural log-transformed on- and off-treatment values adjusted for natural log-transformed off-treatment level. Betas were generated using linear regression analysis with age, sex, body mass index, and rosuvastatin dose as covariates.

[#]p-values were generated using the permutation procedure. Superscript indicates genetic model: R, recessive; A, additive; D, dominant.

3.2. Associations of SNPs with lipid and CIMT reduction during 12-month therapy by rosuvastatin

Table 3 shows associations of studied SNPs with the 12-month lipid-lowering effect and carotid intima-media thickness change in CAD patients receiving rosuvastatin.

Table 3. Associations of single nucleotide polymorphisms with the 12-month lipid-lowering effect and carotid intima-media thickness change in coronary artery disease patients taking rosuvastatin.

Chr	Gene (SNP ID)	Effect allele	EAF	N	Total cholesterol		LDL-C		CIMT, maximum		CIMT, mean	
					Beta*	$P_{perm}^{\#}$	Beta*	$P_{perm}^{\#}$	Beta*	$P_{perm}^{\#}$	Beta*	$P_{perm}^{\#}$
1	ZNF648 (rs1689800)	G	0.392	113	0.010	0.3091	0.024	0.2043	-0.093	0.0105^R	-0.036	0.0282^A
1	GALNT2 (rs4846914)	G	0.388	113	-0.004	0.7273	-0.016	0.4118	0.002	0.8571	-0.002	0.8571
2	COBLL1 (rs12328675)	C	0.170	109	-0.011	0.4643	-0.018	0.4643	0.059	0.0213^A	0.075	0.0056^D
6	LPA (rs55730499)	T	0.056	113	0.364	0.0001^R	0.367	0.0415^R	0.018	0.8571	0.414	0.0146^R
7	NPC1L1 (rs217406)	G	0.203	113	-0.002	1.0000	0.003	0.2043	0.019	0.3636	0.018	0.4516
8	PPP1R3B (rs9987289)	A	0.086	113	0.004	0.8571	-0.068	0.4118	0.072	0.0359^D	0.079	0.0109^D
9	ABCA1 (rs1883025)	T	0.263	113	0.010	0.3148	0.001	0.4643	0.011	0.8571	0.019	0.2687
11	F2 (rs3136441)	C	0.180	100	0.013	0.3478	0.018	0.2043	-0.011	0.6429	-0.005	1.0000
11	ST3GAL4 (rs11220463)	T	0.190	113	-0.181	0.0273^R	-0.027	0.4118	0.001	1.0000	0.032	0.1709
12	SCARB1 (rs838880)	C	0.336	113	-0.001	1.0000	-0.005	0.4643	-0.003	1.0000	-0.004	1.0000
16	CETP (rs3764261)	A	0.147	113	0.017	0.3478	0.021	0.2043	-0.053	0.0833	-0.043	0.0880
16	PSKH1 (rs16942887)	A	0.116	113	0.006	0.5789	0.086	0.0175^D	-0.011	0.5217	-0.004	0.7778
17	STARD3 (rs881844)	C	0.310	113	0.009	0.4375	0.010	0.8571	0.028	0.1852	0.040	0.0223^A
19	LILRA3 (rs386000)	C	0.203	113	0.019	0.2400	0.015	0.5200	0.026	0.2982	0.010	0.7778

20	<i>PLTP</i> (rs6065906)	C	0.160	113	0.013	0.4815	0.031	0.2571	-0.029	0.3478	-0.006	1.0000
----	-------------------------	---	-------	-----	-------	--------	-------	--------	--------	--------	--------	--------

Chr, chromosome; SNP, single nucleotide polymorphism; EAF, effect allele frequency; LDL-C, low-density lipoprotein cholesterol; CIMT, carotid intima-media thickness.

*Beta for difference between the natural log-transformed on- and off-treatment values adjusted for natural log-transformed off-treatment level. The beta reflects the fraction of differential lipid- or CIMT-lowering effect in carriers versus non-carriers of the minor allele (for additive model), in carriers of the genotype according to the model (dominant, recessive) versus the carriers of alternative genotypes; a negative beta indicates a better statin response (stronger lipid or CIMT reduction), a positive beta – a worse statin response. Betas were generated using linear regression analysis with age, sex, body mass index, and rosuvastatin dose as covariates.

*p-values were generated using the permutation procedure. Superscript indicates genetic model: R, recessive; A, additive; D, dominant.

We observed that two polymorphisms contributed to TC reduction: rs55730499 in *LPA* (beta = 0.364, Pperm=0.0001, recessive effect) and rs11220463 in *ST3GAL4* (beta = -0.181, Pperm=0.0273, recessive effect) after one year of lipid-lowering therapy. For the rs11220463 polymorphism, the homozygous genotype for the minor allele T was associated with an 18.1% more pronounced TC-lowering effect than in carriers of alternative genotypes. The reduction in plasma LDL-C levels was associated with rs55730499 of *LPA* (Pperm=0.0415, recessive effect) and rs16942887 of *PSKH1* (Pperm=0.0175, dominant effect).

Three SNPs were associated with both maximum and mean CIMT change: rs1689800 of *ZNF648* (better statin response; Pperm=0.0105 and 0.0282 for the association with maximum and mean CIMT change, respectively), rs12328675 of *COBLL1* (worse statin response; Pperm=0.0213 and 0.0056, respectively), and rs9987289 of *PPP1R3B* (worse statin response; Pperm=0.0359 and 0.0109, respectively). Two polymorphisms, such as rs55730499 of *LPA* (Pperm=0.0146) and rs881844 of *STARD3* (Pperm=0.0223) were associated with the mean CIMT change.

During 12 months of lipid-lowering therapy with rosuvastatin, the most significant effect on the TC reduction was found for SNP rs55730499 of *LPA*, whereas the most significant effects of rs16942887 of *PSKH1* and rs12328675 of *COBLL1* were found on the reduction of LDL-C and CIMT, respectively. Associations of TG reduction (Table 2) with genotype in a 12-month period were found for rs1883025 of the *ABCA1* gene (beta = -0.7246, Pperm= 0.0160, dominant effect) and for rs11220463 in *ST3GAL4* (beta = 3.624, Pperm=0.05, recessive effect).

4. Discussion

The present study demonstrated, for the first time, the impact of lipid-associated GWAS loci on CIMT (Figure 1) and plasma levels of TC, LDL-C, and TG in CAD patients during rosuvastatin therapy (Figure 2).




Figure 1. Associations of single nucleotide polymorphisms with carotid intima-media thickness change on rosuvastatin at 6 and 12 months of therapy.




Figure 2. Associations of single nucleotide polymorphisms with lipid- lowering effect of rosuvastatin at 6 and 12 months of therapy. TC, total cholesterol; LDL-C, low density lipoprotein cholesterol; TG, triglyceride levels.

In particular, we found associations of SNPs located at *ZNF648*, *LPA*, *ST3GAL4*, *PSKH1*, *GALNT2*, *COBLL1*, *PPP1R3B*, and *STARD3* genes with reduction of CIMT. Pharmacogenetic associations of the variants in *ZNF648*, *LPA*, *ST3GAL4*, *PSKH1*, *PLTP*, *SCARB1*, and *ABCA1* with the change in lipid levels were also found for the first time. Thus, lipid-associated GWAS loci are associated not only with lipids and the risk of coronary artery disease [32], but are also responsible for the antiatherogenic effects of rosuvastatin after 6 and 12 months of lipid-lowering therapy.

Potential mechanisms by which polymorphic variants of the investigated lipid metabolism genes can impact the antiatherogenic effects of rosuvastatin therapy are of interest. The *ZNF648* gene encodes zinc finger protein 648, which may be involved in the DNA-templated transcription [35]. In the present study rs1689800 polymorphism was associated with CIMT change. The possible mechanism of the effect of the variant on vascular wall may include the influence on oxidative stress (GLUL catabolizes glutamate, an amino acid required for glutathione synthesis), because such a mechanism is described for the nearest SNP rs10911021 located in the same genomic region [36].

The *GALNT2* gene encodes N-acetylgalactosaminyltransferase 2, an enzyme involved in O-linked glycosylation of the substrates, including those regulating lipid metabolism such as apolipoprotein C-III (APOC-III), angiopoietin-related protein 3 (ANGPTL3) and phospholipid transfer protein (PLTP); in these ways it influences HDL-C and TG metabolism [37,38]. The rs4846914G allele has been linked to atherogenic changes in plasma lipid metabolism, such as an increase in TG and a decrease in HDL-C levels [9,10]. We found no associations of this variant with changes in any plasma lipid, but this SNP was significantly associated with a better CIMT regression during a 6-month period of rosuvastatin therapy. The presence of the link with CIMT change and absence of any association with lipid level change suggest that the pharmacogenetic effect on CIMT change might be mediated through non-lipid-related mechanisms, taking into account that SNP rs4846914 is known to be associated with endothelial function, serum levels of insulin and glucose [39], as well as hypertension [40]. In particular, the latter study demonstrated higher promoter methylation of the *GALNT2* gene, higher levels of ApoB, and lower levels of ApoA1 in hypertensives with the GG genotype [40].

The *COBLL1* gene encodes a cordon-bleu WH2 repeat protein like 1, having actin monomer and cadherin binding activity [41]. In the present study, the variant rs12328675 was associated with worse CIMT regression (for the carriers of the C-allele, dominant effect) during 12 months of observation. This association was the most significant among all studied variants over a 12-month period. There were no associations with lipid levels change. The link between this polymorphism and CIMT change can be explained by the ability of the risk allele C, associated with the higher CAD risk, to form transcription factor binding sites for the factors involved in the processes of vascular inflammation regulation and angiogenesis (AP-1 (syn. JUN), SMAD2, SMAD3, SMAD4, and E2F8) [42].

The *LPA* gene encodes lipoprotein (a), which is a well-known and independent risk factor for coronary artery disease [11]. The studied rs55730499 variant is associated with Lp(a) concentrations [11], coronary artery disease risk [8], myocardial infarction risk [43], and stroke risk [44]. The T-allele is associated with higher TC levels out of therapy [43], and in our study, the TT genotype (homozygous for the minor allele) was associated with reduced drug response in terms of TC, LDL-C, and CIMT change. The association with TC change had the highest significance among all associations with TC change, and moreover, rs55730499 was the only variant studied that was associated with both TC, LDL-C, and CIMT change in plasma at the same period of observation (12 months). Based on these findings, the studied *LPA* variant can be considered the first of all studied variants to be used as a predictor for rosuvastatin therapy personalization.

The *PPP1R3B* gene encodes protein phosphatase 1 regulatory subunit 3B, a key protein in hepatic glycogen metabolism [12]. The rs9987289 variant is associated with plasma TC and LDL-C levels [9,12], but was not tested for carotid atherosclerosis traits or the pharmacogenetics of rosuvastatin. In our study, the presence of the minor A-allele was associated with a worse CIMT response to rosuvastatin. Such an effect could be explained by the association of the studied SNP with inflammatory markers, such as C-reactive protein [45], known for its influence on cardiovascular risk and atherosclerosis progression [46], and also by the link of the variant with metabolic syndrome [47] and glucose levels [48], taking into account the known role of hyperglycemia in the dysfunction of the endothelium [49,50].

The *ABCA1* gene encodes phospholipid-transporting ATPase ABCA1, which provides the efflux of intracellular cholesterol to apolipoproteins and the formation of nascent high-density lipoproteins [51]. The minor T-allele of the rs1883025 SNP is associated in GWAS with an anti-atherogenic phenotype: lower TC [9], LDL-C [52], TG [53], and HDL-C levels [52], and with a better response to statin therapy ([54], the particular drug and lipid are not specified). In the present study, we confirmed a better response to statin therapy in terms of triglyceride reduction in carriers of the minor T-allele (the dominant effect of the SNP).

The *ST3GAL4* gene encodes CMP-N-acetylneuraminate-beta-galactosamide-alpha-2,3-sialyltransferase 4, an enzyme involved in the terminal sialylation of glycoproteins and glycolipids. A beta-galactoside alpha2-3 sialyltransferase takes part in hemostasis (sialylation of plasma von Willebrand factor) and the inflammatory process (selectin-mediated rolling and adhesion of leukocytes during extravasation) [55]. The studied SNP in this gene, rs11220463, is associated with both blood lipids (TC and LDL-C) [53] and inflammation (CRP levels) [56]. In the present study, we have found pharmacogenetic associations of the studied variant with TC and CIMT reduction during therapy. The association with CIMT can be possibly explained by the involvement of the product of *ST3GAL4* in inflammatory processes, taking into account the known association of systemic inflammation (assessed by the systemic immune-inflammatory index) with CIMT [57]. For rs11220463 the association with inflammation was found in terms of CRP levels [56], and CRP is known to be associated with CIMT [58,59], but this association is not proven to be causal [58].

The *PSKH1* gene encodes serine/threonine-protein kinase H1, involved in intracellular protein trafficking and pre-mRNA processing [60]. For the rs16942887 variant, we found associations with LDL-C and CIMT changes on rosuvastatin therapy. There are no reported associations in the literature between this variation and changes in LDL-C and CIMT when taking rosuvastatin. The possible mechanism of the effect of SNP is difficult to predict because of the lack of information on the mechanisms of lipid and vascular influence of *PSKH1*.

The *STARD3* gene encodes StAR-related lipid transfer protein 3, which mediates cholesterol transport from the endoplasmic reticulum to endosomes [61]. In the literature, the rs881844 polymorphism in *STARD3* was known to be associated with plasma lipids, including TC and HDL-C [43,53]. In the present study we didn't find any associations with changes in plasma lipid levels on the rosuvastatin therapy, however, this SNP showed the strongest effects on CIMT changes in both 6- and 12-month periods. The effect on the vascular wall can be explained by the involvement of *STARD3* in the regulation of cholesterol-dependent inflammation and sensitivity to proinflammatory cytokines [62].

The *PLTP* gene encodes phospholipid transfer protein, which is involved in the transfer of phospholipids and free cholesterol from LDLs and VLDLs into HDLs [63], as well as the uptake of cholesterol from peripheral cells [64]. Taking into account the function of the gene product, it was rather logical to find the influence of rs6065906 in *PLTP* on the lipid-lowering effect of rosuvastatin in terms of TC reduction in the present study, which was not reported before.

Thus, the molecular mechanisms underlying the effects of the studied loci in terms of lipid and CIMT reduction remains unknown, but taking into account the biological functions of the proteins, these genes possess pleiotropic effects on biological and pathological processes including lipid metabolism, inflammation, leukocyte adhesion to the endothelium, endothelial dysfunction, and glucose metabolism. Further experimental studies are needed to explain the functional effects of polymorphisms. Many of the studied loci have been associated with HDL-C levels in GWAS. However, because of non-significant HDL-C change on the therapy, observed in the present study, we didn't test the SNPs for associations with HDL-C change. This is the limitation of our study, but we can't say it's a big omission, because one of the main phenotypes studied here was CIMT, which is known to have a causal relationship with LDL-C and with other factors but not with HDL-C itself [18].

5. Conclusions

In the present study, associations of both TC (and/or LDL-C) and CIMT changes at the same period of follow-up (6 or 12 months) were observed for the polymorphisms rs55730499 of *LPA* (after 12 months of therapy) and rs1689800 of *ZNF648* (after 6 months of therapy). In general (including 6- and 12-month periods), four polymorphisms namely rs1689800 in *ZNF648* (LDL-C and CIMT), rs55730499 in *LPA* (TC, LDL-C, and CIMT), rs11220463 in *ST3GAL4* (TC and CIMT), and rs16942887 in *PSKH1* (LDL-C and CIMT) were associated with a decrease in atherogenic lipids and CIMT. There were no TG change-associated loci, that were also associated with CIMT change. A half of eight GWAS-associated lipid loci, such as rs4846914 of *GALNT2*, rs12328675 of *COBLL1*, rs9987289 of *PPP1R3B*, and rs881844 of *STARD3*, that were linked to CIMT changes in our study were not associated with any changes in plasma lipids during rosuvastatin therapy. Polymorphisms rs881844 of *STARD3* and rs12328675 of *COBLL1* were most strongly associated with CIMT regression, whereas rs55730499 of *LPA* – with the reduction of plasma TC and rs16942887 of *PSKH1* – with LDL-C change. The observed associations should be replicated in independent pharmacogenetic studies with a larger number of hyperlipidemic patients. The successful validation of these associations may provide novel genetic markers for the personalization of rosuvastatin therapy in the monitoring of the drug effectiveness in patients with atherogenic changes in lipid metabolism.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Table S1: Change in lipid and carotid intima-media thickness values during rosuvastatin therapy. Supplementary Table 1. Change in lipid and carotid intima-media thickness values during rosuvastatin therapy.

Author Contributions: Conceptualization, A.P.; methodology, A.P., S.K., M.A., M.C.; software, S.K. and A.P.; validation, S.K., A.P.; formal analysis, S.K., M.A.; genetic investigation, Iu.A., E.K., and M.B.; resources, A.P.; data curation, S.K. and A.P.; writing—original draft preparation, S.K.; writing—review and editing, A.P.; visualization, S.K.; supervision, A.P.; project administration, A.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received funding from Kursk State Medical University.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Ethics Committee of Kursk State Medical University (11 May, 2015).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data supporting reported results are available upon request.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Gupta, RM; Schnitzler, G; Fang, S; Lee-Kim, VS; Barry, A. Multiomic Analysis and CRISPR Perturbation Screens Identify Endothelial Cell Programs and Novel Therapeutic Targets for Coronary Artery Disease. *Arterioscler Thromb Vasc Biol.* 2023. doi: 10.1161/ATVBAHA.123.318328.
- Blackburn, H. Invited commentary: 30-year perspective on the seven countries study. *Am J Epidemiol.* 2017, 185(11), 1143–1147.
- McPherson, R; Tybjaerg-Hansen, A. Genetics of Coronary Artery Disease. *Circ Res.* 2016, 118(4), 564–578. doi: 10.1161/CIRCRESAHA.115.306566.
- Churilin, M.I.; Kononov, S.I.; Mal, G.S.; Polonikov, A.V.; Lazarenko, V.A. Lipid metabolism genes and predisposition to ischemic heart disease. *Medical News of North Caucasus*, 2019, 14(2), 401–407.
- Mutlu, AS; Duffy, J; Wang, MC. Lipid metabolism and lipid signals in aging and longevity. *Dev Cell.* 2021, 56(10), 1394–1407. doi: 10.1016/j.devcel.2021.03.034.
- Walsh, R.; Jurgens, S.J.; Erdmann, J.; Bezzina, C.R. Genome-wide association studies of cardiovascular disease. *Physiol Rev.* 2023. doi: 10.1152/physrev.00024.2022.
- Willer, C.J.; Schmidt, E.M.; Sengupta, S.; Peloso, G.M.; Gustafsson, S.; Kanoni, S.; Ganna, A.; Chen, J.; Buchkovich, M.L.; Mora, S.; et al. Discovery and refinement of loci associated with lipid levels. *Nat. Genet.* 2013, 45, 1274–1283.
- Nikpay, M.; Goel, A.; Won, H.H.; Hall, L.M.; Willenborg, C.; Kanoni, S.; Saleheen, D.; Kyriakou, T.; Nelson, C.P.; Hopewell, J.C.; et al. A comprehensive 1000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat. Genet.* 2015, 47, 1121–1130.
- Teslovich, T.M.; Musunuru, K.; Smith, A.V.; Edmondson, A.C.; Stylianou, I.M.; Koseki, M.; Pirruccello, J.P.; Ripatti, S.; Chasman, D.I.; Willer, C.J.; et al. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature.* 2010, 466, 707–713.
- Kathiresan, S.; Melander, O.; Guiducci, C.; Surti, A.; Burt, N.P.; Rieder, M.J.; Cooper, G.M.; Roos, C.; Voight, B.F.; Havulinna, A.S.; et al. Six new loci associated with blood low-density lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans. *Nat. Genet.* 2008, 40, 189–197.
- Mack, S.; Coassin, S.; Rueedi, R.; Yousri, N.A.; Seppälä, I.; Gieger, C.; Schönherr, S.; Forer, L.; Erhart, G.; Marques-Vidal, P.; et al. A genome-wide association meta-analysis on lipoprotein (a) concentrations adjusted for apolipoprotein (a) isoforms. *J. Lipid Res.* 2017, 58(9), 1834–1844.
- Zhang, Y.; Gan, W.; Tian, C.; Li, H.; Lin, X.; Chen, Y. Association of PPP1R3B polymorphisms with blood lipid and C-reactive protein levels in a Chinese population. *J. Diabetes.* 2013, 5(3), 275–281.
- Postmus, I.; Trompet, S.; Deshmukh, H.A.; Arsenault B.J.; Avery C.L.; Bis J.C.; Chasman D.I.; de Keyser C.E.; Deshmukh H.A.; Evans D.S.; et al. Pharmacogenetic meta-analysis of genome-wide association studies of LDL cholesterol response to statins. *Nat. Commun.* 2014, 5, 5068.
- Chasman, D.I.; Giulianini, F.; MacFadyen J., Barratt B.J., Nyberg F., Ridker P.M. Genetic determinants of statin-induced low-density lipoprotein cholesterol reduction: the Justification for the Use of Statins in Prevention: an Intervention Trial Evaluating Rosuvastatin (JUPITER) trial. *Circ. Cardiovasc. Genet.* 2012, 5(2), 257–264.
- Chasman, D.I., Giulianini, F., Demler, O.V., Udler, M.S. Pleiotropy-Based Decomposition of Genetic Risk Scores: Association and Interaction Analysis for Type 2 Diabetes and CAD. *Am J Hum Genet.* 2020, 106(5), 646–658. doi: 10.1016/j.ajhg.2020.03.011.
- Crouse, J.R. 3rd, Raichlen, J.S., Riley, W.A.; Evans G.W.; Palmer M.K.; O'Leary D.H.; Grobbee D.E.; Bots M.L.; METEOR Study Group. Effect of rosuvastatin on progression of carotid intima-media thickness in low-risk individuals with subclinical atherosclerosis: the METEOR Trial. *JAMA.* 2007, 297(12), 1344–1353.
- Nezu, T; Hosomi, N; Aoki, S; Matsumoto M. Carotid Intima-Media Thickness for Atherosclerosis. *J Atheroscler Thromb.* 2016, 23(1), 18–31. doi: 10.5551/jat.31989.

18. Shah, S; Casas, J.P., Drenos, F.; Whittaker, J.; Deanfield, J.; Swerdlow, D.I.; Holmes, M.V.; Kivimaki, M.; Langenberg, C.; Wareham, N.; et al. Causal relevance of blood lipid fractions in the development of carotid atherosclerosis: mendelian randomization analysis. *Circ. Cardiovasc Genet.* 2013, 6(1), 63–72.
19. Vargas, JD; Manichaikul, A; Wang, X.Q.; Rich, S.S.; Rotter, J.I.; Post, W.S.; Polak, J.F.; Budoff, M.J., Bluemke, D.A. Detailed analysis of association between common single nucleotide polymorphisms and subclinical atherosclerosis: The Multi-ethnic Study of Atherosclerosis. *Data Brief.* 2016, 7, 229-242. doi: 10.1016/j.dib.2016.01.048.
20. Guaricci, A.I.; Arcadi, T.; Brunetti, ND; Maffei, E.; Montrone, D.; Martini, C.; De Luca, M.; De Rosa, F.; Cocco, D.; Midiri, M.; et al. Carotid intima media thickness and coronary atherosclerosis linkage in symptomatic intermediate risk patients evaluated by coronary computed tomography angiography. *Int J Cardiol.* 2014, 176(3), 988-993. doi: 10.1016/j.ijcard.2014.08.141.
21. Amato, M.; Montorsi, P.; Ravani, P., et al. Carotid intima-media thickness by B-mode ultrasound as surrogate of coronary atherosclerosis: correlation with quantitative coronary angiography and coronary intravascular ultrasound findings. *Eur Heart J.* 2007, 28, 2094–101.
22. Graner M, Varpula M, Kahri J et al. Association of carotid intima-media thickness with angiographic severity and extent of coronary artery disease. *Am. J. Cardiol.* 97(5), 624–629 (2006).
23. Matsushima Y, Kawano H, Koide Y, Baba T, Toda G, Seto S, Yano K. Relationship of carotid intima-media thickness, pulse wave velocity, and ankle brachial index to the severity of coronary artery atherosclerosis. *Clin Cardiol.* 2004 Nov;27(11):629-34. doi: 10.1002/clc.4960271110.
24. Ballantyne CM, Raichlen JS, Nicholls SJ et al. Effect of rosuvastatin therapy on coronary artery stenoses assessed by quantitative coronary angiography: a study to evaluate the effect of rosuvastatin on intravascular ultrasound-derived coronary atheroma burden. *Circulation* 117(19), 2458–2466 (2008).
25. Puri R, Libby P, Nissen SE et al. Long-term effects of maximally intensive statin therapy on changes in coronary atheroma composition: insights from SATURN. *Eur. Heart J. Cardiovasc. Imaging.* 15(4), 380–388 (2014).
26. Rumyantsev NA, Kukes VG, Kazakov RE, Rumyantsev AA, Sychev DA. Ispol'zovanie farmakogeneticheskogo testirovaniia dlia predotvrashcheniia nezhelatel'nykh lekarstvennykh reaktsii pri terapii statinami [Use of pharmacogenetic testing to prevent adverse drug reactions during statin therapy]. *Ter Arkh.* 2017;89(1):82-87. Russian. doi: 10.17116/terarkh201789182-87.
27. Soko ND, Masimirembwa C, Dandara C. Pharmacogenomics of Rosuvastatin: A Glocal (Global + Local) African Perspective and Expert Review on a Statin Drug. *OMICS* 20(9), 498–509 (2016).
28. Alfonsi JE, Hegele RA, Gryn SE. Pharmacogenetics of lipid-lowering agents: precision or indecision medicine? *Curr. Atheroscler. Rep.* 18(5), 24 (2016).
29. Churilin, M.I., Kononov, S.I., Luneva, Y.V., ...Solodilova, M.A., Polonikov, A.V. Polymorphisms of Intracellular Cholesterol Transporters Genes: Relationship to Blood Lipid Levels, Carotid Intima-Media Thickness, and the Development of Coronary Heart Disease. *Russian Journal of Genetics*, 2020a, 56(2), pp. 234–241.
30. Churilin, M.I. , Kononov, S.I., Luneva, Y.V., ...Polonikov, A.V., Kazanov, V.A. Apolipoprotein E gene polymorphisms: A relationship with the risk of coronary artery disease and the effectiveness of lipid-lowering therapy with rosuvastatin. *Cardiovascular Therapy and Prevention (Russian Federation)*, 2020b, 19(1), pp. 17–23.
31. Kononov S, Mal G, Azarova I, Klyosova E, Bykanova M, Churnosov M, Polonikov A. Pharmacogenetic loci for rosuvastatin are associated with intima-media thickness change and coronary artery disease risk. *Pharmacogenomics.* 2022 Jan;23(1):15-34. doi: 10.2217/pgs-2021-0097.
32. Lazarenko V, Churilin M, Azarova I, Klyosova E, Bykanova M, Ob'edkova N, Churnosov M, Bushueva O, Mal G, Povetkin S, Kononov S, Luneva Y, Zhabin S, Polonikova A, Gavrilenko A, Saraev I, Solodilova M, Polonikov A. Comprehensive Statistical and Bioinformatics Analysis in the Deciphering of Putative Mechanisms by Which Lipid-Associated GWAS Loci Contribute to Coronary Artery Disease. *Biomedicines.* 2022 Jan 25;10(2):259. doi: 10.3390/biomedicines10020259.
33. Pignoli P, Tremoli E, Poli A, Oreste P, Paoletti R. Intimal plus medial thickness of the arterial wall: a direct measurement with ultrasound imaging. *Circulation* 74(6), 1399–1406 (1986).
34. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007 Sep;81(3):559-75. doi: 10.1086/519795.

35. Gaudet P, Livstone MS, Lewis SE, Thomas PD. Phylogenetic-based propagation of functional annotations within the Gene Ontology consortium. *Brief Bioinform.* 2011 Sep;12(5):449-62. doi: 10.1093/bib/bbr042.
36. Shahid SU, Shabana, Humphries S. The SNP rs10911021 is associated with oxidative stress in coronary heart disease patients from Pakistan. *Lipids Health Dis.* 2018 Jan 5;17(1):6. doi: 10.1186/s12944-017-0654-8.
37. Khetarpal SA, Schjoldager KT, Christoffersen C, Raghavan A, Edmondson AC, Reutter HM, Ahmed B, Ouazzani R, et al. Loss of Function of GALNT2 Lowers High-Density Lipoproteins in Humans, Nonhuman Primates, and Rodents. *Cell Metab.* 2016 Aug 9;24(2):234-45. doi: 10.1016/j.cmet.2016.07.012.
38. Zilmer M, Edmondson AC, Khetarpal SA, Alesi V, Zaki MS, Rostasy K, Madsen CG, et al. Novel congenital disorder of O-linked glycosylation caused by GALNT2 loss of function. *Brain.* 2020 Apr 1;143(4):1114-1126. doi: 10.1093/brain/awaa063.
39. Chen J, Guan L, Liu H, Liu Q, Fan P, Bai H. GALNT2 Gene Variant rs4846914 Is Associated with Insulin and Insulin Resistance Depending on BMI in PCOS Patients: a Case-Control Study. *Reprod Sci.* 2021 Apr;28(4):1122-1132. doi: 10.1007/s43032-020-00380-7.
40. Zhang X, Zhao H, Zhang J, Han D, Zheng Y, Guo X, He D, Guo J, Wang Y. Gene environment interaction of GALNT2 and APOE gene with hypertension in the Chinese Han Population. *Biomed Mater Eng.* 2015;26 Suppl 1:S1977-83. doi: 10.3233/BME-151501.
41. Guo Z, Neilson LJ, Zhong H, Murray PS, Zanivan S, Zaidel-Bar R. E-cadherin interactome complexity and robustness resolved by quantitative proteomics. *Sci Signal.* 2014 Dec 2;7(354):rs7. doi: 10.1126/scisignal.2005473.
42. Churilin MI. Association of RS12328675 COBLL1 polymorphism with coronary heart disease and intermediate phenotypes of atherosclerosis: validation study in Central Russia. *Research Results in Biomedicine.* 2020;6(2):209-218. Russian. DOI: 10.18413/2658-6533-2020-6-2-0-5.
43. Sakaue S, Kanai M, Tanigawa Y, Karjalainen J, Kurki M, Koshihara S, Narita A, Konuma T, Yamamoto K, Akiyama M, Ishigaki K, Suzuki A, Suzuki K, Obara W, Yamaji K, Takahashi K, Asai S, et al. A cross-population atlas of genetic associations for 220 human phenotypes. *Nat Genet.* 2021 Oct;53(10):1415-1424. doi: 10.1038/s41588-021-00931-x.
44. Mishra A, Malik R, Hachiya T, Jürgenson T, Namba S, Posner DC, Kamanu FK, Koido M, Le Grand Q, Shi M, He Y, Georgakis MK, Caro I, Krebs K, Liaw YC, Vaura FC, Lin K, et al. Stroke genetics informs drug discovery and risk prediction across ancestries. *Nature.* 2022 Nov;611(7934):115-123. doi: 10.1038/s41586-022-05165-3. Epub 2022 Sep 30. Erratum in: *Nature.* 2022 Dec;612(7938):E7.
45. Wojcik GL, Graff M, Nishimura KK, Tao R, Haessler J, Gignoux CR, Highland HM, Patel YM, Sorokin EP, Avery CL, Belbin GM, Bien SA, Cheng I, et al. Genetic analyses of diverse populations improves discovery for complex traits. *Nature.* 2019 Jun;570(7762):514-518. doi: 10.1038/s41586-019-1310-4.
46. Sethwala AM, Goh I, Amerena JV. Combating Inflammation in Cardiovascular Disease. *Heart Lung Circ.* 2021 Feb;30(2):197-206. doi: 10.1016/j.hlc.2020.09.003.
47. Lind L. Genome-Wide Association Study of the Metabolic Syndrome in UK Biobank. *Metab Syndr Relat Disord.* 2019 Dec;17(10):505-511. doi: 10.1089/met.2019.0070.
48. Chen J, Spracklen CN, Marenne G, Varshney A, Corbin LJ, Luan J, Willems SM, Wu Y, Zhang X, Horikoshi M, Boutin TS, Mägi R, Waage J, Li-Gao R, Chan KHK, Yao J, Anasanti MD, Chu AY, et al. The trans-ancestral genomic architecture of glycemic traits. *Nat Genet.* 2021 Jun;53(6):840-860. doi: 10.1038/s41588-021-00852-9.
49. Meza CA, La Favor JD, Kim DH, Hickner RC. Endothelial Dysfunction: Is There a Hyperglycemia-Induced Imbalance of NOX and NOS? *Int J Mol Sci.* 2019 Aug 2;20(15):3775. doi: 10.3390/ijms20153775.
50. Clyne AM. Endothelial response to glucose: dysfunction, metabolism, and transport. *Biochem Soc Trans.* 2021 Feb 26;49(1):313-325. doi: 10.1042/BST20200611.
51. Krimbou L, Denis M, Haidar B, Carrier M, Marcil M, Genest J Jr. Molecular interactions between apoE and ABCA1: impact on apoE lipidation. *J Lipid Res.* 2004 May;45(5):839-48. doi: 10.1194/jlr.M300418-JLR200.
52. Huang QQ, Sallah N, Dunca D, Trivedi B, Hunt KA, Hodgson S, Lambert SA, Arciero E, Wright J, Griffiths C, Trembath RC, Hemingway H, et al. Transferability of genetic loci and polygenic scores for cardiometabolic traits in British Pakistani and Bangladeshi individuals. *Nat Commun.* 2022 Aug 9;13(1):4664. doi: 10.1038/s41467-022-32095-5.
53. Hoffmann TJ, Theusch E, Haldar T, Ranatunga DK, Jorgenson E, Medina MW, Kvale MN, Kwok PY, Schaefer C, Krauss RM, Iribarren C, Risch N. A large electronic-health-record-based genome-wide study of serum lipids. *Nat Genet.* 2018 Mar;50(3):401-413. doi: 10.1038/s41588-018-0064-5.

54. Wu Y, Byrne EM, Zheng Z, Kemper KE, Yengo L, Mallett AJ, Yang J, Visscher PM, Wray NR. Genome-wide association study of medication-use and associated disease in the UK Biobank. *Nat Commun*. 2019 Apr 23;10(1):1891. doi: 10.1038/s41467-019-09572-5.
55. Mondal N, Buffone A Jr, Stolfi G, Antonopoulos A, Lau JT, Haslam SM, Dell A, Neelamegham S. ST3Gal-4 is the primary sialyltransferase regulating the synthesis of E-, P-, and L-selectin ligands on human myeloid leukocytes. *Blood*. 2015 Jan 22;125(4):687-96. doi: 10.1182/blood-2014-07-588590.
56. Ligthart S, Vaez A, Hsu YH; Inflammation Working Group of the CHARGE Consortium; PMI-WG-XCP; LifeLines Cohort Study; Stolk R, Uitterlinden AG, Hofman A, Alizadeh BZ, Franco OH, Dehghan A. Bivariate genome-wide association study identifies novel pleiotropic loci for lipids and inflammation. *BMC Genomics*. 2016 Jun 10;17:443. doi: 10.1186/s12864-016-2712-4.
57. Ozbeyaz NB, Gokalp G, Algul E, Kilic P, Saricam O, Aydinylmaz F, Guliyev I. Could Systemic Inflammation in Healthy Individuals With Obesity Indicate Subclinical Atherosclerosis? *Angiology*. 2022 Apr 28;33197221089375. doi: 10.1177/00033197221089375.
58. Kivimäki M, Lawlor DA, Smith GD, Kumari M, Donald A, Britton A, Casas JP, Shah T, Brunner E, Timpson NJ, Halcov JP, Miller MA, Humphries SE, Deanfield J, Marmot MG, Hingorani AD. Does high C-reactive protein concentration increase atherosclerosis? The Whitehall II Study. *PLoS One*. 2008 Aug 20;3(8):e3013. doi: 10.1371/journal.pone.0003013.
59. Moran CA, Sheth AN, Mehta CC, Hanna DB, Gustafson DR, Plankey MW, Mack WJ, Tien PC, French AL, Golub ET, Quyyumi A, Kaplan RC, Ofotokun I. The association of C-reactive protein with subclinical cardiovascular disease in HIV-infected and HIV-uninfected women. *AIDS*. 2018 May 15;32(8):999-1006. doi: 10.1097/QAD.0000000000001785.
60. Brede G, Solheim J, Prydz H. PSKH1, a novel splice factor compartment-associated serine kinase. *Nucleic Acids Res*. 2002 Dec 1;30(23):5301-9. doi: 10.1093/nar/gkf648.
61. Wilhelm LP, Wendling C, Védie B, Kobayashi T, Chenard MP, Tomasetto C, Drin G, Alpy F. STARD3 mediates endoplasmic reticulum-to-endosome cholesterol transport at membrane contact sites. *EMBO J*. 2017 May 15;36(10):1412-1433. doi: 10.15252/embj.201695917.
62. Li L, Liu Y, Liu X, Zheng N, Gu Y, Song Y, Wang X. Regulatory roles of external cholesterol in human airway epithelial mitochondrial function through STARD3 signalling. *Clin Transl Med*. 2022 Jun;12(6):e902. doi: 10.1002/ctm2.902.
63. Zhang M, Zhai X, Li J, Albers JJ, Vuletic S, Ren G. Structural basis of the lipid transfer mechanism of phospholipid transfer protein (PLTP). *Biochim Biophys Acta Mol Cell Biol Lipids*. 2018 Sep;1863(9):1082-1094. doi: 10.1016/j.bbalip.2018.06.001.
64. Albers JJ, Vuletic S, Cheung MC. Role of plasma phospholipid transfer protein in lipid and lipoprotein metabolism. *Biochim Biophys Acta*. 2012 Mar;1821(3):345-57. doi: 10.1016/j.bbalip.2011.06.013.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.