

Article

The In Silico Sugarcane Genome-Encoded MicroRNA and Target Network Prediction for Targeting the SCMV

Muhammad Aleem Ashraf ^{1,†,2,*}, Xiaoyan Feng ^{1,†}, Hira Ghaffar ^{2,†}, Judith K Brown ³, Imran Shahid ⁴, Naitong Yu ¹ and Shuzhen Zhang ^{1,*}

¹ Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, China; fengxiaoyan@itbb.org.cn (X.F.); yunaitong@163.com (N.Y.)

² Institute of Biological Sciences, Faculty of Natural and Applied Sciences, Khwaja Fareed University of Engineering and Information Technology, Rahim Yar Khan 64200, Pakistan; hiraghaffar823@gmail.com

³ School of Plant Sciences, The University of Arizona, AZ85721 Tucson, USA; jbrown@ag.arizona.edu

⁴ Department of Pharmacology and Toxicology, Faculty of Medicine, Umm Al-Qura University, Makkah 21955, Saudi Arabia; iyshahid@uqu.edu.sa

* Correspondence: ashraf.muhammad.aleem@gmail.com (M.A.A.); zhangshuzhen@itbb.org.cn (S.Z.)

† These authors contributed equally to this work.

Abstract: Sugarcane mosaic virus (SCMV) is a deleterious pathogen which causes widespread Sugarcane mosaic disease (SCMD) and is classified in the genus *Potyvirus* (*Potyviridae*), disseminated by the aphid vector. RNA interference (RNAi)-mediated antiviral innate immunity is a key biological process and antiviral defence system to interfere with viral genomes for controlling plant pathogens. The current study aims to analyze sugarcane (*Saccharum officinarum* L. and *Saccharum* spp.) locus-derived microRNAs (sof-miRNAs/ssp-miRNAs) with predicted potential for targeting the SCMV +ssRNA-encoded mRNAs, using 'five algorithms' approach. The ultimate goal in this research is to mobilize the in silico endogenous predicted sof-miRNAs/ssp-miRNAs to trigger RNAi catalytic pathway experimentally and generate sugarcane cultivars for evaluating potential antiviral resistance monitoring capability and capacity for SCMV. Experimentally validated mature sugarcane (*S. officinarum*, $2n = 8X = 80$) and (*S. spp.*, $2n = 100-120$) sof-miRNAs/ssp-miRNAs ($n = 28$) were acquired for alignment with the SCMV genome. Of the 28 targeting mature locus-derived sof-miRNAs/ssp-miRNAs investigated, one sugarcane miRNA homolog, sof-miR159c, was concluded to localize potential binding site at genomic nucleotide site 3847 targeting CL ORF of SCMV. In order to validate target prediction accuracy, whether the sugarcane sof-miRNA/ssp-miRNA might bind predicted SCMV mRNA target(s), we created an integrated Circos plot. Genome-wide in-silico-predicted miRNA-mediated target gene regulatory network validated interactions that warrant in vivo analysis. The current work provides valuable evidence and biological material for generating SCMV-resistant sugarcane varieties.

Keywords: *potyvirus*; in silico tools; sugarcane mosaic virus; miRNA; RNA interference

1. Introduction

Sugarcane (*Saccharum officinarum*) is a vigorous tropical and sub-tropical pertinent economically important, long-duration, biofuel cash crop, enriched with high energy roughage and also a source of agro-industrial residue [1-3]. The octaploid sugarcane (*S. officinarum*) genome ($2n = 80$; $x = 10$) [4,5] which is referred as "noble" cane and genome of sugarcane species and cultivars have been assembled, drafted and re-sequenced [6-11]. Sugarcane mosaic virus disease (SCMV) is highly transmissible and pathogenic potyvirus that cause sugarcane mosaic virus disease (SCMD) [12,13]. Potyviruses are observed to be spread by a common sap sucking vector—aphid species complex [14]. Innovative approaches are still needed to enhance the sugarcane productivity [15]. The genome of SCMV composed of a +ss RNA molecule 9575 nucleotides in length encoding a single large polyprotein. The genome polyprotein precursor was predicted to undergo

cleavage resulting ten functional proteins, P1, HC-Pro, P3, 6 K1, CI, 6 K2, VPg, NIa, NIb and CP [16-19].

In plants, microRNAs (miRNA) are endogenously expressed small (19-25 nucleotides), evolutionary conserved, non-coding (NC)-ss RNA molecules [20]. In higher plants, biogenesis and transcription of miRNA gene (*MIR*) is governed by RNA polymerase II which is further transcribed into single-standard polycistronic *primary* transcripts (pri-miRNAs). They govern a multitude of biological process in plants regulating gene expression, cell growth, development, differentiation and host-virus interactions [21,22]. The miRNA-mediated RNAi is a post-transcriptional gene silencing mechanism providing antimicrobial innate immunity regulating host-virus interaction for restriction of inhibition of virus infection [23].

Artificial miRNA-mediated (amiRNA) technology is an alternative, safe approach based on engineering miRNA gene to control viral infection in plants [24]. RNAi-based amiRNA construct has been deployed in research to create antiviral resistance in plants against plant viral species such as tomato [25,26], cucumber [27], rice [28] and cotton [29]. Mature miRNAs in sugarcane genome have been predicted, identified, isolated, analyzed, and validated for evaluation of host-virus interactions, gene regulation and was associated with abiotic and biotic stresses [30-40]. Recently, experimental validation of 35 conserved mature sugarcane genome-encoded, high-confidence sof-miRNAs/ssp-miRNAs and further deposition in the miRBase database were reported.

An integrative multi-network approach based on evaluation of SCMV infection, deployed to identify target-binding sites of sugarcane genome-encoded sof-miRNAs/ssp-miRNAs in the SCMV genome. Identification of several homologous amiRNAs for the creation of transgenic sugarcane cultivars—resistant to SCMV is the key objective in this study. The predicted sugarcane genome-encoded sof-miRNAs/ssp-miRNAs were further evaluated to understand complex sugarcane host plant–SCMV potyviral interactions for identification of novel antiviral-targets.

2. Materials and Methods

2.1. Sugarcane MicroRNAs and SCMV Genome Data Retrieval and Processing

Experimentally validated high-confidence mature sugarcane microRNAs (sof-miRNA156-sof-miR11892/ssp-miR156-ssp-1432) (Accession ID: MIMAT0001656-MIMAT0001671/ MIMAT0020291-MIMAT0020290) and (Saccharum sp.-microRNAs) (ssp-miR166-ssp-miR1432) (Accession ID: MIMAT0030451-MIMAT0020290) (Table S1) were retrieved from the from the miRNA registry (miRBase, version 22) [41]. The full-length SCMV +ssRNA genome sequence (9575 bases) (Accession number KY548506) was acquired from the NCBI GenBank database [42].

2.2. Potential Targets of Sugarcane MicroRNAs in SCMV Genome

Predicting effective microRNA-mRNA binding sites is an initial step for understanding microRNA-regulated gene regulatory networks. The accuracy of miRNA target site prediction can be influenced by several factors, such as the specificity and sensitivity of the algorithm, the choice of reference sequence, and the length of the target sequence. To predict miRNA-mRNA target sites computationally, various in silico methods are generated for effective silencing. A computational approach refers to the use of multiple computational methods, algorithms, or tools to analyze and interpret biological data. This approach involves combining different types of publicly available in silico algorithms, miRanda[43,44], RNA22[45,46], TAPIR[47], psRNATarget[48,49] and RNAhybrid [50] (Table 1).

Table 1. Different features and parameters of algorithms applied for miRNA target predictions.

Algorithms	Features	Organism	Parameters	Source
------------	----------	----------	------------	--------

miRanda	Seed-based interaction, multiple target sites, free energy of miRNA-mRNA duplex, conservation	Human, rat, fly, worms	Score threshold= 140, Free energy=-20 Kcal/mol, Gap open penalty=-9.00, Gap extend penalty=-4.00	http://www.microrna.org/ (retrieved 14 August 2019)
RNA22	Pattern recognition, folding energy, heteroduplex	Human, mouse, fly and worms	Number of paired-up bases= 12, Sensitivity (63%), Specificity (61%), Folding energy=-15 Kcal/mol	https://cm.jefferson.edu/rna22/Interactive/ (retrieved on 22 June 2019)
TAPIR	Sees pairing, target site accessibility, multiple sites	Plants	Free energy ratio=0.2 Score= 9	http://bioinformatics.psb.ugent.be/webtools/tapir (retrieved on 25 June 2021)
psRNATarget	Complementarity scoring, multiple target sites, translation inhibition	Plants	Expectation Score= 6.5, Penalty for G:U pair= 0.5 HSP size= 19 Penalty for opening gap= 2	https://www.zhaolab.org/psRNATarget/analysis?function=2 (accessed on 26 May 2022)
RNAhybrid	Seed pairing and free energy	Any	Free energy=-20 Kcal/mol, Hit per target= 1	http://bibiserv.techfak.uni-bielefeld.de/rnahybrid (accessed on 26 May 2022)


2.3. Statistical Analysis

The miRNA-mRNA target prediction biological data were further processed. Graphical representations of miRNA data were prepared using R-language [51].

3. Results

3.1. Prediction and Analysis of Sugarcane MicroRNAs Targeting SCMV Genome

An integrative computational approach to identify possible interactions of high-confidence target sites of sugarcane mature miRNAs located in the SCMV positive-sense single-stranded (+ssRNA) genome from among the 28 sugarcane miRNAs (sof-miRNAs/ssp-miRNAs) revealed sof-miRNAs/ssp-miRNAs—derived MIR genes at high proportion of sugarcane miRNA gene loci [33,52-55]. The predicted SCMV +ssRNA encoded mRNA sequences were localized hypothetically best sof-miRNAs/ssp-miRNAs—annealing sites predicted by miRanda algorithm (19 miRNA-mRNA target pairs) and RNA22 (15 sugarcane sof-miRNAs/ssp-miRNAs and 20 loci). The TAPIR identified 7 binding sites of sugarcane mature sof-miRNA-target/ssp-miRNA-target pairs. Twenty nine sugarcane miRNAs targeting thirty three attachment sites were identified by psRNATarget. RNAhybrid predicted 28 high-probability binding sites of sugarcane miRNAs in the SCMV genomic RNA sequence (Figure 1 and Figure 2) (File S1) (Table S2).



The potyviral first protease protein P1 encoded by P1 ORF (149-847) (698 bases), the least conserved, hypervariable, modulates host responses and essential for replication of viral +ssRNA genome [56,57]. Host adaptation is a key process for virus genome evolution [58,59]. P1 is also related to virus-host adaptation [60]. The miRanda and RNA22 algorithms predicted bindings of two sof-miRNAs: sof-miR168 (a, b) at nucleotide positions 547 and 846, respectively as shown in (**Figure 2A-2B**). The sof-miRNA168a was targeted at nucleotide position 406, by the TAPIR algorithm (**Figure 2C**). No sof-miRNA/ssp-miRNA was predicted targeting the P1 region, by the psRNATarget and RNAhybrid algorithms (**Figure 2D-2E**) (**File S1**) (**Table S2-S3**).




Figure 2. Individual sugarcane sof-miRNA/ssp-miRNA and their predicted high-confidence binding sites in the SCMV genome were predicted based on ‘five algorithms’ approach. (A) miRNA-sites were detected by miRanda. (B) Several miRNA-target sites were detected by RNA22. (C) TAPIR identified sugarcane miRNA-binding sites. (D) psRNATarget predicted several binding sites of sugarcane miRNAs. (E) Prediction of miRNA-sites by RNAhybrid. (F) Union plot representing all predicted binding sites detected by all the algorithms used. Multiple copies of miRNA target binding sites were represented by colored dots. Targeted genes of SCMV were indicated by different colors.

3.3. Sugarcane miRNAs Targeting HC-Pro

The HC-Pro ORF (848-2227 nt) encodes a multifunctional, non-structural dimeric—helper component—proteinase. It has been reported as a viral suppressor. Enhanced expression by fusion of P1, symptoms development and viral replication are the key functions [61-66]. The miRanda and RNA22 algorithms predicted target site of sof-miR168 (a, b) at nucleotide position 1827. Both the algorithms binding of sof-miR168a at nt position 1296 also (Figure 2A-2B).

TAPIR predicted attachment site of sof-miRNA159e at locus 1159 (**Figure 2C**). The psRNATarget algorithm detected binding of ssp-miR444 (a, b, c-3p) at nt positions (1058 and 1763) (**Figure 2D**). The RNAhybrid algorithm predicted—sof-miR159c, sof-miR168 (a, b), ssp-miR444 (a, b, c-3p) and ssp-miR1432 at nucleotide positions 1830, 1296, 1818, 1057 and 1316, respectively (**Figure 2E**) (**File S1**) (**Table S2-S3**).

3.4. Sugarcane miRNAs Targeting P3

The P3 ORF (2228-3268 nt) encodes a P3 membrane-associated protein participating directly in the genomic RNA replication mechanism of SCMV. It is also involved in potential cell-to-cell spread (movement and transport) and is responsible to determine host-range and symptoms [67-69]. The miRanda algorithm predicted binding of sof-miRNAs: sof-miR168 (a, b) at nucleotide position 2562 (**Figure 2A**). No sof-miRNA/ssp-miRNA was predicted targeting the P3 by the RNA22 and TAPIR algorithms (**Figure 2B-2C**). Potential target sites of sof-miR167 (a, b), sof-miR168a, ssp-miR437c, ssp-miR444 (a,b, c-3p) at nucleotide positions 2427, 2971, 2981 and 2367, respectively, were detected by psRNATarget (**Figure 2D**). Further, RNAhybrid identified sof-miR167(a, b), ssp-miR437b—target sites at nt positions 2699 and 2416, respectively (**Figure 2E**) (**File S1**) (**Table S2-S3**).

3.5. Sugarcane miRNAs Targeting 6K1

The 6K1 ORF (3269-3469 nucleotides) encoding a 6K1 protein which functions viral genome replication. It mediates cell-to-cell movement, controlling defense mechanism and gene regulation. It is a key component of 6K2-induced viral replication complex (VRC), and regulation [70,71]. The 6K1 had the least number of predicted sugarcane sof-miRNAs ssp-miR444c-3p at nucleotide position 3441, by the psRNATarget algorithm (**Figure 2D**) (**File S1**) (**Table S2-S3**).

3.6. Sugarcane miRNAs Targeting CI

The CI ORF (3470-5383 nt) encodes a multifunctional cylindrical inclusion (CI) protein that is essential for ATP-binding and RNA helicase activity [72-74]. CI was targeted by two miRNAs: sof-miR396, ssp-miR166 at nt positions 3634, 4178 respectively, as indicated by miRanda (**Figure 2A**). RNA22 predicted two miRNAs: sof-miR159c, ssp-miR444b at nt positions 3730, 5311 respectively (**Figure 2B**). Further, TAPIR predicted three sugarcane miRNAs: sof-miR159c, ssp-miR437a and ssp-miR1128 at nucleotide positions 3847, 4869 and 4534, respectively (**Figure 2C**). The psRNATarget identified seven miRNAs: sof-miR159 (a, b, c, d, e), ssp-miR444b, ssp-miR1432 at nt positions 3847, 3992 and 3980, respectively (**Figure 2D**). Five miRNA-binding sites were detected by RNAhybrid: sof-miR396 (start site 5016), sof-miR408e (3633), ssp-miR166 (3714), ssp-miR437a (4868) and ssp-miR1128 (4533) (**Figure 2E**) (**File S1**) (**Table S2-S3**).

3.7. Sugarcane miRNAs Targeting 6K2

The potyviral 6K2 (5384-5542 nt) encoded 6K2 multifunctional protein, induces formation of RE-derived complexes and develop resistance against drought [75,76]. The RNA22 identified five sugarcane sof-miRNAs: sof-miR408 (a, b, c, d, e) on locus position 5538 (**Figure 2B**) (**File S1**) (**Table S2-S3**).

3.8. Sugarcane miRNAs Targeting NIa-VPg

The potyviruses NIa-VPg ORF (5543-6109 nt) encodes viral genome-linked protein (VPg) which functions as virulence determinant and genome translation [77-81]. It also involved in replication, translation and movement [82-84]. The RNA22 and TAPIR predicted binding of ssp-miR444c-3p on locus position 5552 (**Figure 2B-2C**). The psRNATarget predicted six miRNAs: sof-miR156, sof-miR159 (a, b, c, d), ssp-miR444c-3p (**Figure**

2D). No sof-miRNA/ssp-miRNA was predicted targeting the NIa-VPg region by the RNAhybrid algorithm (**Figure 2E**) (**File S1**) (**Table S2-S3**).

3.9. Sugarcane miRNAs Targeting NIa

The potyviruses NIa ORF (6110-6835 nt) encodes nuclear inclusion a (NIa) protein that is involved in RNA-binding and also interacts with NIb [85,86]. miRanda, RNA22 and RNAhybrid predicted binding of only sugarcane miRNA: ssp-miR528, sof-miR396 and ssp-miR827 at nucleotide positions 6376, 6821 and 6338, respectively (**Figure 2A-2B, 2E**). The psRNATarget identified three sugarcane miRNAs: sof-miR408e, ssp-miR444 (a, b) at nucleotide positions 6544 and 6641, respectively (**Figure 2D**). No miRNA-target pair was identified by TAPIR (**Figure 2C**) (**File S1**) (**Table S2-S3**).

3.10. Sugarcane miRNAs Targeting NIb

The potyviruses NIb ORF (6836-8398) encodes nuclear inclusion b (NIb) protein that is involved in translocation activity and also interacts with NIa [87]. It contains nuclear signals and also called as RdRp [88]. The miRanda algorithm detected binding of two sugarcane ssp-miRNAs: ssp-miR169 and ssp-miR1432 at nucleotide positions 7798 and 7523 respectively (**Figure 2A**). The psRNATarget algorithm predicted binding of two sugarcane ssp-miRNAs: sof-miR396 and ssp-miR444b at nucleotide positions 7798 and 7523 respectively (**Figure 2D**). No miRNA-target pairs were identified by the RNA22, TAPIR and RNAhybrid algorithms, (**Figure 2B-2C-2E**) (**File S1**) (**Table S2-S3**).

3.10.1. Sugarcane miRNAs Targeting CP

The potyviruses CP ORF (8399-9337) encodes multistaking protein, coat (CP) that is involved in the development of virion assembly. The CP is involved in all steps of potyviral life cycle [89-91]. The miRanda predicted binding of three sugarcane ssp-miRNAs (ssp-miR444 (a, b, c-3p) (start site 8501). ssp-miR444c-3p also targeted CP region at nt position 9268(**Figure 2A**). RNA22 predicted binding of ssp-miRNA444 family at nt positions 8502 and 9181(**Figure 2B**). The psRNATarget predicted bind of ssp-miR444c-3p at nt position 9282. Potential binding sites of sugarcane miRNAs: sof-miR159 (a, b, d, e), sof-miR408 (a, b, c, d), and ssp-miR169 were detected by RNAhybrid at nt positions 8953, 8355, and 8458 respectively (**Figure 2E**) (**File S1**) (**Table S2-S3**).

3.10.2. Sugarcane miRNAs Targeting UTR

The potyviruses 5' untranslated region (5' UTR) (1-148 nt) and 3' UTR (9341-9575 nt) are involved in replication and translocational activities of the ORFs [92,93]. The sof-miR408 (a, b, c, d) was predicted target the 5' UTR at nt positions 139 by miRanda (**Figure 2A**). Similarly, ssp-miR528 was identified to target the 5' UTR at nt position 122 by TAPIR and RNAhybrid (**Figure 2C-2E**). RNA22 predicted binding of sof-miR168 (a, b) at nt position 9520 in the 3' UTR (**Figure 2B**). RNAhybrid predicted binding of two sugarcane miRNAs in the 3'UTR: sof-miR156 and ssp-miR437c at nt positions 9402 and 9395 respectively (**Figure 2E**) (**File S1**) (**Table S2-S3**).

3.11. Identification of Consensual Sugarcane MicroRNAs

The current study was concluded on the basis of consensus genomic target binding sites of sugarcane miRNAs detected by different algorithms. Among them, we selected 9 sugarcane miRNAs (sof-miR159c, sof-miR168a, ssp-miR437a, ssp-miR528, ssp-miR444 (a, b), ssp-miR444c-3p), ssp-miR1128, and ssp-miR1432) that were detected on the basis of consensus genomic positions—3847 (target gene CI), 1296 (HC-Pro), 4869 (CI), 122 (5' UTR), 8502/1058 (CP/HC-Pro), 5583 (NIa-VPg), 4534 (CI) and 1316 (HC-Pro) respectively (**Table 2 and Table 3**). Of nine consensus sugarcane-encoded locus-derived sof-miRNAs/ ssp-miRNAs investigated in this study, only one sof-miRNA (sof-miR159c at nt position 3847 targeting CI) was detected by union of consensus genomic positions by at least three

algorithms (RNA22, TAPIR and psRNATarget) (**Figure 3, Table 2 and Table 3**) (**File S1**) (**Table S2-S3**).

Table 2. Predicted high-confidence binding sites of consensus sugarcane miRNAs targeting SCMV genome were detected by different computational algorithms.

Sugarcane miRNA	Position miRanda	Position RNA22	Position TAPIR	Position psRNATarget	Position RNAhybrid	MFE * miRanda	MFE ** RNA22	MFE Ratio TAPIR	Expectation psRNATarget	MFE* RNAhybrid
sof-miR159c		3847	3847	3847			-18.00	0.58	5.50	
sof-miR168a		1296			1296		-18.70			-25.80
ssp-miR437a			4869		4868			0.69		-21.20
ssp-miR528			122		121			0.60		-26.50
ssp-miR444a	8501	8502		1058	1057	-18.42	-18.00		7.00	-29.00
ssp-miR444b	8501	8502		1058	1057	-18.42	-18.00		7.00	
ssp-miR444c-3p			5583	5583				0.59	6.00	
ssp-miR1128			4534		4533			0.66		-27.30
ssp-miR1432		1315			1316		-15.40			-22.20




Figure 3. Intersection plot show consensus high-confidence binding sites of sugarcane mature miRNAs predicted by at least two computational tools. The colored dots represent sugarcane miRNA-binding sites targeting different genes of SCMV.

Table 3. Predicted consensus sugarcane-encoded miRNA-target sites localized in different target genes of SCMV-SO.

miRNA ID	Accession ID	Mature Sequence (5'-3')	Target Genes ORF(s)	Target Binding Locus Position
sof-miR159c	MIMAT0001662	CUUGGAUUGAAGGGAGCUCCU	CI	3847-3868
sof-miR168a	MIMAT0001665	UCGCUUGGUGCAGAU CGGGAC	HC-Pro	1296-1317
ssp-miR437a	MIMAT0020280	AAAGUUAGAGAAGUUUGACUU	CI	4869-4890
ssp-miR528	MIMAT0020288	UGGAAGGGGCAUGCAGAGGAG	5'UTR	122-143
ssp-miR444a	MIMAT0020284	UGCAGUUGUUGCCUCAAGCUU	CP	8501-8521
ssp-miR444a (1)	MIMAT0020284	UGCAGUUGUUGCCUCAAGCUU	HC-Pro	1058-1078
ssp-miR444b	MIMAT0020285	UGCAGUUGUUGCCUCAGGCUU	CP	8501-8521
ssp-miR444b (1)	MIMAT0020285	UGCAGUUGUUGCCUCAGGCUU	HC-Pro	1058-1079
ssp-miR444c-3p	MIMAT0020286	UGCAGUUGUUGUCUCAAGCUU	Nla-VPg	5583-5604
ssp-miR1128	MIMAT0020289	UACUACUCCCUCCGUCCCAA	CI	4534-4555
ssp-miR1432	MIMAT0020290	CUCAGGAAAGAUGACACCGAC	HC-Pro	1315-1336

3.12. Identification of miRNA-mRNA Regulatory Network

Circos plot represents predicted host–virus interactions of sugarcane miRNAs and SCMV target genes. The Circos plot was generated to visualize comprehensive master miRNA regulatory network with novel antiviral targets (**Figure 4**). Generation of miRNA-mRNA Regulatory Network was conducted using ‘Circos’ software [94]




Figure 4. Integrated Circos plot demonstrate multiple targets of sugarcane-encoded miRNAs. The colored connection lines are targeted genes (ORFs) in SCMV genome. Construction, exploration, target predictions and interactions between the sugarcane miRNAs and SCMV genes are mapped.

3.13. RNA Secondary Structures

The Computationally predicted consensual sugarcane mature miRNAs were analyzed by generating their secondary structures using original precursor sequences. The pre-miRNA hairpin sequences were used to perform manual curation. Salient parameters of predicted stable secondary structures were evaluated (**Table 6**). Stable secondary structures of potential consensual sugarcane precursors sequences were predicted by RNAfold algorithm [95].

Table 4. Features of predicted precursors of sugarcane were determined.

miRNA ID	Accession ID	MFE */Kcal/mol	AMFE **	MFEI ***	(G+C)%
sof-MIR159c	MI0001760	-110.60	-46.47	-0.87	53.36
sof-MIR168a	MI0001763	-66.20	-63.65	-0.83	75.96
ssp-MIR437a	MI0001763	-57.10	-32.62	-1.29	25.14
ssp-MIR528	MI0001763	-48.50	-52.71	-0.86	60.84
ssp-MIR444a	MI0001763	-57.70	-54.94	-1.28	42.86
ssp-MIR444b	MI0001763	-63.70	-60.09	-1.38	43.39
ssp-MIR444c	MI0001763	-61.80	-57.22	-1.31	43.52

ssp-MIR1128	MI0001763	-101.70	-36.98	-1.18	31.27
ssp-MIR1432	MI0001763	-57.10	-64.88	-1.14	56.82

4. Discussion

The SCMV is a monopartite potyvirus, suggested as etiological agent that spread to Pakistan and China due to highly transmissible pathogen, becoming an increasing potential, long lasting, threat to sugarcane and maize production over the past two decades [13,17,96]. In our previous studies, we have investigated experimentally validated sugarcane genome-encoded mature microRNAs that were predicted to target SCBGAV, SCYLV and SCBV based on in silico criteria [37-39]. Several studies have identified complex host-virus interactions and have investigated host-plant miRNAs targeting plant viruses using an in silico approach [97-103]. The miRNAs have evolved as novel endogenous targets for multiple layers of miRNA-gene level regulation [52,104,105]. In order to abate host plant-virus infection, several studies have indicated that efficacy of amiRNA-based RNA interference resulting specific gene silencing in transgenic crops [27,28,106-108]. In this computational research work, mature sugarcane sof-miRNAs were aligned with the target, SCMV genomic sequence to identify miRNA-mRNA binding sites hypothesized to understand complex host-virus specific interactions with the P1, HC-Pro, P3, 6K1, CI, 6K2, NIa-VPg, NIa-Pro, NIb, CP of SCMV.

Based on our reported finding, the SCMV genome (HC-Pro, CI, NIa-VPg and CP) is vulnerable to be targeted nine consensus sugarcane miRNAs. We revealed nine miRNAs could theoretically derive from sugarcane genome (Table 3 and Figure 4). In silico tools—RNA22, TAPIR and psRNATarget—screened a consensus genomic high-confidence binding of base-pairing complementarity sof-miR159c at nucleotide position 3847, (Figure 1 and Table 2). While all five algorithms identified ssp-miR444c-3p as the unique sugarcane miRNA (Figure 1 and Table 2). We identified the maximum folding energy of consensus functional miRNA-mRNA target pair which is -18.00 Kcal/mol using RNA22. RNA22 is a highly sensitive algorithm uses a pattern-based approach to screen target sites of miRNAs. While, we estimated expectation score 5.50 of consensual target pair by psRNATarget (Table 2). The [109]RNA22 and psRNATarget algorithms predict target sites using a non-seed-based approach. These results provide support for predicted consensus miRNA-mRNA duplex to represent a ‘true target’. Our findings demonstrated that sugarcane - miRNAs probably have putative role in host-virus pathogenesis interaction. Our results highlight the interaction of SCMV ss-RNA on the sugarcane miRNA: target interaction network.

The potyviral cylindrical inclusion helicase (CI) is required to initiate viral replication mechanism. It also controls cell-to-cell movement and plant-host protein-virus interaction [72,73]. Computational prediction and analysis implicated the sugarcane consensus sof-miR159c high-confidence target site potentially targeting the CI ORF. The conserved precursor MIR159 is reported to govern plant growth, and fertility [110]. The consensus sof-miR159e (Accession ID: MIMAT0001661), that have predicted effective target binding site at nucleotide position 5535 in SCBV genome, was identified as top effective miRNA by miRanda, RNA22 and RNAhybrid algorithms.

While miRNA-mRNA target pair interactions between sugarcane genome-encoded ghr-miRNAs and SCMV have been determined, development of amiRNA-based construct and further transformation in sugarcane to control SCMV is yet to fully understand. We reported first time a comprehensive analysis of SCMD-associated *Potyvirus* which is an initial step to construct miRNA-based anti-viral therapy. The amiRNA construct is based on highly specificity of nucleotide base-pairing to control detrimental off-target effects. The small size of amiRNA is a unique feature to develop a single gene expression vector to control multiple Potyviruses in transgenic sugarcane.

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Table S1: Sugarcane mature microRNA sequences used for prediction binding sites in the SCMV genome Table S2: Identification of high-confidence binding sites of

sugarcane miRNAs in the SCMV; Table S3: Gene wise prediction; File S1: Prediction results by computational tools.

Author Contributions: M.A.A. and S.Z. conceived the study. M.A.A., X.F and H.G analyzed data. M.A.A wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Central Public-Interest Scientific Institution Basal Research Fund (1630052023003).

Institutional Review Board Statement: The current study was approved by Board of Advanced Studies and Research (BASR).

Informed Consent Statement: Not applicable.

Data Availability Statement: All relevant data are available in the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Carvalho-Netto, O.V.; Bressiani, J.A.; Soriano, H.L.; Fiori, C.S.; Santos, J.M.; Barbosa, G.V.; Xavier, M.A.; Landell, M.G.; Pereira, G.A. The potential of the energy cane as the main biomass crop for the cellulosic industry. *Chemical and Biological Technologies in Agriculture* **2014**, *1*, 1-8.
2. Ahmed, A.; Dompok, E.; Gasparatos, A. Human wellbeing outcomes of involvement in industrial crop production: Evidence from sugarcane, oil palm and jatropha sites in Ghana. *PLoS One* **2019**, *14*, e0215433.
3. Zeng, F. Measuring convergence in the sugarcane industry in China's Guangxi province. *Plos one* **2021**, *16*, e0244617.
4. Piperidis, G.; Piperidis, N.; D'Hont, A. Molecular cytogenetic investigation of chromosome composition and transmission in sugarcane. *Molecular Genetics and Genomics* **2010**, *284*, 65-73.
5. Yu, F.; Wang, P.; Li, X.; Huang, Y.; Wang, Q.; Luo, L.; Jing, Y.; Liu, X.; Deng, Z.; Wu, J. Characterization of chromosome composition of sugarcane in nobilization by using genomic in situ hybridization. *Molecular cytogenetics* **2018**, *11*, 1-8.
6. Cuadrado, A.; Acevedo, R.; Moreno Díaz de la Espina, S.; Jouve, N.; De La Torre, C. Genome remodelling in three modern S. officinarum × S. spontaneum sugarcane cultivars. *Journal of experimental botany* **2004**, *55*, 847-854.
7. Zhang, J.; Zhang, X.; Tang, H.; Zhang, Q.; Hua, X.; Ma, X.; Zhu, F.; Jones, T.; Zhu, X.; Bowers, J. Allele-defined genome of the autopolyploid sugarcane Saccharum spontaneum L. *Nature genetics* **2018**, *50*, 1565-1573.
8. Shearman, J.R.; Pootakham, W.; Sonthirod, C.; Naktang, C.; Yoocha, T.; Sangsrakru, D.; Jomchai, N.; Tongsim, S.; Piriyaongsa, J.; Ngamphiw, C. A draft chromosome-scale genome assembly of a commercial sugarcane. *Scientific Reports* **2022**, *12*, 20474.
9. Yan, H.; Zhou, H.; Luo, H.; Fan, Y.; Zhou, Z.; Chen, R.; Luo, T.; Li, X.; Liu, X.; Li, Y. Characterization of full-length transcriptome in Saccharum officinarum and molecular insights into tiller development. *BMC Plant Biology* **2021**, *21*, 1-12.
10. Zhang, D.; Wang, R.; Han, S.; Li, Z.; Xiao, J.; Li, Y.; Wang, L.; Li, S. Transcriptome analysis of sugarcane young leaves and protoplasts after enzymatic digestion. *Life* **2022**, *12*, 1210.
11. Zhou, D.; Liu, Y.; Yao, J.; Yin, Z.; Wang, X.; Xu, L.; Que, Y.; Mo, P.; Liu, X. Characterization and Phylogenetic Analyses of the Complete Mitochondrial Genome of Sugarcane (Saccharum spp. Hybrids) Line A1. *Diversity* **2022**, *14*, 333.
12. Lu, G.; Wang, Z.; Xu, F.; Pan, Y.-B.; Grisham, M.P.; Xu, L. Sugarcane mosaic disease: Characteristics, identification and control. *Microorganisms* **2021**, *9*, 1984.
13. He, E.-Q.; Bao, W.-Q.; Sun, S.-R.; Hu, C.-Y.; Chen, J.-S.; Bi, Z.-W.; Xie, Y.; Lu, J.-J.; Gao, S.-J. Incidence and Distribution of Four Viruses Causing Diverse Mosaic Diseases of Sugarcane in China. *Agronomy* **2022**, *12*, 302.
14. Gadhav, K.R.; Gautam, S.; Rasmussen, D.A.; Srinivasan, R. Aphid transmission of Potyvirus: the largest plant-infecting RNA virus genus. *Viruses* **2020**, *12*, 773.
15. Diniz, A.L.; Ferreira, S.S.; Ten-Caten, F.; Margarido, G.R.; Dos Santos, J.M.; Barbosa, G.V.d.S.; Carneiro, M.S.; Souza, G.M. Genomic resources for energy cane breeding in the post genomics era. *Computational and Structural Biotechnology Journal* **2019**, *17*, 1404-1414.
16. Tang, W.; Yan, Z.; Zhu, T.; Xu, X.; Li, X.-D.; Tian, Y. The complete genomic sequence of Sugarcane mosaic virus from Canna spp. in China. *Virology journal* **2018**, *15*, 1-4.
17. Muhammad, K.; Herath, V.; Ahmed, K.; Tahir, M.; Verchot, J. Genetic diversity and molecular evolution of sugarcane mosaic virus, comparing whole genome and coat protein sequence phylogenies. *Archives of Virology* **2022**, *167*, 2239-2247.
18. He, Z.; Dong, Z.; Gan, H. Genetic changes and host adaptability in sugarcane mosaic virus based on complete genome sequences. *Molecular phylogenetics and evolution* **2020**, *149*, 106848.
19. Xie, X.; Chen, W.; Fu, Q.; Zhang, P.; An, T.; Cui, A.; An, D. Molecular variability and distribution of Sugarcane mosaic virus in Shanxi, China. *PLoS one* **2016**, *11*, e0151549.
20. Finnegan, E.J.; Matzke, M.A. The small RNA world. *Journal of cell science* **2003**, *116*, 4689-4693.
21. Millar, A.A. The function of miRNAs in plants. **2020**, *9*, 198.

22. Islam, W.; Waheed, A.; Idrees, A.; Rashid, J.; Zeng, F. Role of plant microRNAs and their corresponding pathways in fluctuating light conditions. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research* **2022**, 119304.
23. Jin, L.; Chen, M.; Xiang, M.; Guo, Z. RNAi-based antiviral innate immunity in plants. *Viruses* **2022**, *14*, 432.
24. Kuo, Y.W.; Falk, B.W. Artificial microRNA guide strand selection from duplexes with no mismatches shows a purine-rich preference for virus-and non-virus-based expression vectors in plants. *Plant Biotechnology Journal* **2022**, *20*, 1069.
25. Al-Roshdi, M.R.; Ammara, U.; Khan, J.; Al-Sadi, A.M.; Shahid, M.S. Artificial microRNA-mediated resistance against Oman strain of tomato yellow leaf curl virus. *Frontiers in Plant Science* **2023**, *14*, 1150.
26. Sharma, N.; Prasad, M. Silencing AC1 of Tomato leaf curl virus using artificial microRNA confers resistance to leaf curl disease in transgenic tomato. *Plant Cell Reports* **2020**, *39*, 1565-1579.
27. Miao, S.; Liang, C.; Li, J.; Baker, B.; Luo, L. Polycistronic artificial microRNA-mediated resistance to cucumber green mottle mosaic virus in cucumber. *International journal of molecular sciences* **2021**, *22*, 12237.
28. Zhou, L.; Yuan, Q.; Ai, X.; Chen, J.; Lu, Y.; Yan, F. Transgenic Rice Plants Expressing Artificial miRNA Targeting the Rice Stripe Virus MP Gene Are Highly Resistant to the Virus. *Biology* **2022**, *11*, 332.
29. Ali, I.; Amin, I.; Briddon, R.W.; Mansoor, S. Artificial microRNA-mediated resistance against the monopartite begomovirus Cotton leaf curl Burewala virus. *Virology journal* **2013**, *10*, 1-8.
30. Yang, Y.; Zhang, X.; Su, Y.; Zou, J.; Wang, Z.; Xu, L.; Que, Y. miRNA alteration is an important mechanism in sugarcane response to low-temperature environment. *BMC genomics* **2017**, *18*, 1-18.
31. Khan, M.S.; Khraiweh, B.; Pugalenth, G.; Gupta, R.S.; Singh, J.; Duttamajumder, S.K.; Kapur, R. Subtractive hybridization-mediated analysis of genes and in silico prediction of associated microRNAs under waterlogged conditions in sugarcane (*Saccharum* spp.). *FEBS Open Bio* **2014**, *4*, 533-541.
32. Ferreira, T.H.; Gentile, A.; Vilela, R.D.; Costa, G.G.L.; Dias, L.I.; Endres, L.; Menossi, M. microRNAs associated with drought response in the bioenergy crop sugarcane (*Saccharum* spp.). **2012**.
33. Swapna, M.; Kumar, S. MicroRNAs and their regulatory role in sugarcane. *Frontiers in Plant Science* **2017**, *8*, 997.
34. Meena, M.R.; Kumar, R.; Chinnaswamy, A.; Karupaiyan, R.; Kulshreshtha, N.; Ram, B. Current breeding and genomic approaches to enhance the cane and sugar productivity under abiotic stress conditions. *3 Biotech* **2020**, *10*, 1-18.
35. Silva, J.d.O.L.; Silva, R.G.d.; Nogueira, L.d.F.; Zingaretti, S.M. MicroRNAs regulate tolerance mechanisms in sugarcane (*Saccharum* spp.) under aluminum stress. *Crop Breeding and Applied Biotechnology* **2021**, *21*.
36. Wang, M.; Li, A.M.; Liao, F.; Qin, C.X.; Chen, Z.L.; Zhou, L.; Li, Y.R.; Li, X.F.; Lakshmanan, P.; Huang, D.L. Control of sucrose accumulation in sugarcane (*Saccharum* spp. hybrids) involves miRNA-mediated regulation of genes and transcription factors associated with sugar metabolism. *GCB Bioenergy* **2022**, *14*, 173-191.
37. Ashraf, F.; Ashraf, M.A.; Hu, X.; Zhang, S. A novel computational approach to the silencing of Sugarcane Bacilliform Guadeloupe A Virus determines potential host-derived MicroRNAs in sugarcane (*Saccharum officinarum* L.). *PeerJ* **2020**, *8*, e8359.
38. Ashraf, M.A.; Ashraf, F.; Feng, X.; Hu, X.; Shen, L.; Khan, J.; Zhang, S. Potential targets for evaluation of sugarcane yellow leaf virus resistance in sugarcane cultivars: in silico sugarcane miRNA and target network prediction. *Biotechnology & Biotechnological Equipment* **2021**, *35*, 1980-1991.
39. Ashraf, M.A.; Feng, X.; Hu, X.; Ashraf, F.; Shen, L.; Iqbal, M.S.; Zhang, S. In silico identification of sugarcane (*Saccharum officinarum* L.) genome encoded microRNAs targeting sugarcane bacilliform virus. *PloS one* **2022**, *17*, e0261807.
40. Zhang, N.; Feng, X.; Zeng, Q.; Lin, H.; Wu, Z.; Gao, X.; Huang, Y.; Wu, J.; Qi, Y. Integrated analysis of miRNAs associated with sugarcane responses to low-potassium stress. *Frontiers in Plant Science* **2022**, *12*, 3050.
41. Kozomara, A.; Birgaoanu, M.; Griffiths-Jones, S. miRBase: from microRNA sequences to function. *Nucleic acids research* **2019**, *47*, D155-D162.
42. Sayers, E.W.; Beck, J.; Bolton, E.E.; Bourex, D.; Brister, J.R.; Canese, K.; Comeau, D.C.; Funk, K.; Kim, S.; Klimke, W. Database resources of the national center for biotechnology information. *Nucleic acids research* **2021**, *49*, D10.
43. Enright, A.; John, B.; Gaul, U.; Tuschl, T.; Sander, C.; Marks, D. MicroRNA targets in *Drosophila*. *Genome biology* **2003**, *4*, 1-27.
44. John, B.; Enright, A.J.; Aravin, A.; Tuschl, T.; Sander, C.; Marks, D.S. Human microRNA targets. *PLoS biology* **2004**, *2*, e363.
45. Miranda, K.C.; Huynh, T.; Tay, Y.; Ang, Y.-S.; Tam, W.-L.; Thomson, A.M.; Lim, B.; Rigoutsos, I. A pattern-based method for the identification of MicroRNA binding sites and their corresponding heteroduplexes. *Cell* **2006**, *126*, 1203-1217.
46. Loher, P.; Rigoutsos, I. Interactive exploration of RNA22 microRNA target predictions. *Bioinformatics* **2012**, *28*, 3322-3323.
47. Bonnet, E.; He, Y.; Billiau, K.; Van de Peer, Y. TAPIR, a web server for the prediction of plant microRNA targets, including target mimics. *Bioinformatics* **2010**, *26*, 1566-1568.
48. Dai, X.; Zhao, P.X. psRNATarget: a plant small RNA target analysis server. *Nucleic acids research* **2011**, *39*, W155-W159.
49. Dai, X.; Zhuang, Z.; Zhao, P.X. psRNATarget: a plant small RNA target analysis server (2017 release). *Nucleic acids research* **2018**, *46*, W49-W54.
50. Krüger, J.; Rehmsmeier, M. RNAhybrid: microRNA target prediction easy, fast and flexible. *Nucleic acids research* **2006**, *34*, W451-W454.
51. Gandrud, C. *Reproducible research with R and RStudio*; Chapman and Hall/CRC: 2018.
52. Bajczyk, M.; Jarmolowski, A.; Jozwiak, M.; Pacak, A.; Pietrykowska, H.; Sierocka, I.; Swida-Barteczka, A.; Szewc, L.; Szweykowska-Kulinska, Z. Recent Insights into Plant miRNA Biogenesis: Multiple Layers of miRNA Level Regulation. *Plants* **2023**, *12*, 342.
53. Yu, Y.; Jia, T.; Chen, X. The 'how' and 'where' of plant micro RNA s. *New Phytologist* **2017**, *216*, 1002-1017.

54. Dong, Q.; Hu, B.; Zhang, C. microRNAs and their roles in plant development. *Frontiers in Plant Science* **2022**, *13*.
55. Gramzow, L.; Lobbes, D.; Innard, N.; Theißen, G. Independent origin of MIRNA genes controlling homologous target genes by partial inverted duplication of antisense-transcribed sequences. *The Plant Journal* **2020**, *101*, 401-419.
56. Valli, A.; Lopez-Moya, J.J.; Garcia, J.A. Recombination and gene duplication in the evolutionary diversification of P1 proteins in the family Potyviridae. *Journal of General Virology* **2007**, *88*, 1016-1028.
57. Pasin, F.; Simón-Mateo, C.; García, J.A. The hypervariable amino-terminus of P1 protease modulates potyviral replication and host defense responses. *PLoS Pathogens* **2014**, *10*, e1003985.
58. LaTourrette, K.; Garcia-Ruiz, H. Determinants of Virus Variation, Evolution, and Host Adaptation. *Pathogens* **2022**, *11*, 1039.
59. Elena, S.F. Local adaptation of plant viruses: lessons from experimental evolution. *Molecular Ecology* **2017**, *26*, 1711-1719.
60. Tatineni, S.; Qu, F.; Li, R.; Morris, T.J.; French, R. Triticum mosaic poacevirus enlists P1 rather than HC-Pro to suppress RNA silencing-mediated host defense. *Virology* **2012**, *433*, 104-115.
61. Kasschau, K.D.; Carrington, J.C. Long-distance movement and replication maintenance functions correlate with silencing suppression activity of potyviral HC-Pro. *Virology* **2001**, *285*, 71-81.
62. Atreya, C.; Atreya, P.; Thornbury, D.; Pirone, T. Site-directed mutations in the potyvirus HC-Pro gene affect helper component activity, virus accumulation, and symptom expression in infected tobacco plants. *Virology* **1992**, *191*, 106-111.
63. Sun, H.; Del Toro, F.; Makki, M.; Tenllado, F.; Canto, T. Adaptation of a Potyvirus Chimera Increases Its Virulence in a Compatible Host through Changes in HCPro. *Plants* **2022**, *11*, 2262.
64. Valli, A.A.; Gallo, A.; Rodamilans, B.; López-Moya, J.J.; García, J.A. The HCPro from the Potyviridae family: an enviable multitasking Helper Component that every virus would like to have. *Molecular plant pathology* **2018**, *19*, 744-763.
65. Sanobar, N.; Lin, P.-C.; Pan, Z.-J.; Fang, R.-Y.; Tjita, V.; Chen, F.-F.; Wang, H.-C.; Tsai, H.-L.; Wu, S.-H.; Shen, T.-L. Investigating the viral suppressor HC-pro inhibiting small rna methylation through functional comparison of HEN1 in angiosperm and bryophyte. *Viruses* **2021**, *13*, 1837.
66. De, S.; Pollari, M.; Varjosalo, M.; Mäkinen, K. Association of host protein VARICOSE with HCPro within a multiprotein complex is crucial for RNA silencing suppression, translation, encapsidation and systemic spread of potato virus A infection. *PLoS Pathogens* **2020**, *16*, e1008956.
67. Luan, H.; Shine, M.; Cui, X.; Chen, X.; Ma, N.; Kachroo, P.; Zhi, H.; Kachroo, A. The potyviral P3 protein targets eukaryotic elongation factor 1A to promote the unfolded protein response and viral pathogenesis. *Plant Physiology* **2016**, *172*, 221-234.
68. Jenner, C.E.; Wang, X.; Tomimura, K.; Ohshima, K.; Ponz, F.; Walsh, J.A. The dual role of the potyvirus P3 protein of Turnip mosaic virus as a symptom and avirulence determinant in brassicas. *Molecular plant-microbe interactions* **2003**, *16*, 777-784.
69. Luan, H.; Liao, W.; Niu, H.; Cui, X.; Chen, X.; Zhi, H. Comprehensive analysis of soybean mosaic virus P3 protein interactors and hypersensitive response-like lesion-inducing protein function. *International Journal of Molecular Sciences* **2019**, *20*, 3388.
70. Bera, S.; Arena, G.D.; Ray, S.; Flannigan, S.; Casteel, C.L. The Potyviral Protein 6K1 Reduces Plant Proteases Activity during Turnip mosaic virus Infection. *Viruses* **2022**, *14*, 1341.
71. Cui, H.; Wang, A. Plum pox virus 6K1 protein is required for viral replication and targets the viral replication complex at the early stage of infection. *Journal of virology* **2016**, *90*, 5119-5131.
72. Deng, P.; Wu, Z.; Wang, A. The multifunctional protein CI of potyviruses plays interlinked and distinct roles in viral genome replication and intercellular movement. *Virology journal* **2015**, *12*, 1-11.
73. Sorel, M.; García, J.A.; German-Retana, S. The Potyviridae cylindrical inclusion helicase: a key multipartner and multifunctional protein. *Molecular plant-microbe interactions* **2014**, *27*, 215-226.
74. Wei, T.; Zhang, C.; Hong, J.; Xiong, R.; Kasschau, K.D.; Zhou, X.; Carrington, J.C.; Wang, A. Formation of complexes at plasmodesmata for potyvirus intercellular movement is mediated by the viral protein P3N-PIPO. *PLoS pathogens* **2010**, *6*, e1000962.
75. Lohmus, A.; Varjosalo, M.; Mäkinen, K. Protein composition of 6K2-induced membrane structures formed during Potato virus A infection. *Molecular plant pathology* **2016**, *17*, 943-958.
76. Prakash, V.; Nihrenz, C.T.; Casteel, C.L. The potyviral protein 6K2 from Turnip mosaic virus increases plant resilience to drought. *Molecular Plant-Microbe Interactions* **2023**, *36*, 189-197.
77. Yang, Z.; Dong, M.; Cheng, G.; Liu, S.; Zhang, H.; Shang, H.; Zhou, Y.; Huang, G.; Zhang, M.; Wang, F. Selective interaction of sugarcane EIF4E with VPGS from sugarcane mosaic pathogens. *Viruses* **2021**, *13*, 518.
78. Goodfellow, I. The genome-linked protein VPg of vertebrate viruses—a multifaceted protein. *Current opinion in virology* **2011**, *1*, 355-362.
79. Wittmann, S.; Chatel, H.; Fortin, M.G.; Laliberté, J.-F. Interaction of the viral protein genome linked of turnip mosaic potyvirus with the translational Eukaryotic Initiation Factor (iso) 4E of Arabidopsis thaliana Using the Yeast two-hybrid system. *Virology* **1997**, *234*, 84-92.
80. Coutinho de Oliveira, L.; Volpon, L.; Rahardjo, A.K.; Osborne, M.J.; Culjkovic-Kraljacic, B.; Trahan, C.; Oeffinger, M.; Kwok, B.H.; Borden, K.L. Structural studies of the eIF4E–VPg complex reveal a direct competition for capped RNA: Implications for translation. *Proceedings of the National Academy of Sciences* **2019**, *116*, 24056-24065.
81. Płochocka, D.; Welnicki, M.; Zielenkiewicz, P.; Ostojka-Zagorski, W. Three-dimensional model of the potyviral genome-linked protein. *Proceedings of the National Academy of Sciences* **1996**, *93*, 12150-12154.
82. Puustinen, P.; Mäkinen, K. Uridylylation of the potyvirus VPg by viral replicase NIb correlates with the nucleotide binding capacity of VPg. *Journal of Biological Chemistry* **2004**, *279*, 38103-38110.

83. Eskelin, K.; Hafrén, A.; Rantalainen, K.I.; Mäkinen, K. Potyviral VPg enhances viral RNA translation and inhibits reporter mRNA translation in planta. *Journal of virology* **2011**, *85*, 9210-9221.
84. Schaad, M.C.; Lellis, A.D.; Carrington, J.C. VPg of tobacco etch potyvirus is a host genotype-specific determinant for long-distance movement. *Journal of virology* **1997**, *71*, 8624-8631.
85. Daròs, J.-A.; Carrington, J.C. RNA binding activity of NIa proteinase of tobacco etch potyvirus. *Virology* **1997**, *237*, 327-336.
86. Martínez, F.; Rodrigo, G.; Aragonés, V.; Ruiz, M.; Lodewijk, I.; Fernández, U.; Elena, S.F.; Daròs, J.-A. Interaction network of tobacco etch potyvirus NIa protein with the host proteome during infection. *BMC genomics* **2016**, *17*, 1-13.
87. Li, X.H.; Valdez, P.; Olvera, R.E.; Carrington, J.C. Functions of the tobacco etch virus RNA polymerase (NIb): subcellular transport and protein-protein interaction with VPg/proteinase (NIa). *Journal of virology* **1997**, *71*, 1598-1607.
88. Ivanov, K.; Eskelin, K.; Lohmus, A.; Mäkinen, K. Molecular and cellular mechanisms underlying potyvirus infection. *Journal of General Virology* **2014**, *95*, 1415-1429.
89. Saha, S.; Lohmus, A.; Dutta, P.; Pollari, M.; Mäkinen, K. Interplay of HCPro and CP in the Regulation of Potato Virus A RNA Expression and Encapsidation. *Viruses* **2022**, *14*, 1233.
90. Lindenau, S.; Winter, S.; Margaria, P. The amino-proximal region of the coat protein of cucumber vein yellowing virus (family Potyviridae) affects the infection process and whitefly transmission. *Plants* **2021**, *10*, 2771.
91. Martínez-Turiño, S.; García, J.A. Potyviral coat protein and genomic RNA: A striking partnership leading virion assembly and more. *Advances in Virus research* **2020**, *108*, 165-211.
92. Zhang, J.; Roberts, R.; Rakotondrafara, A.M. The role of the 5' untranslated regions of Potyviridae in translation. *Virus research* **2015**, *206*, 74-81.
93. Roberts, R.; Zhang, J.; Mayberry, L.K.; Tatineni, S.; Browning, K.S.; Rakotondrafara, A.M. A unique 5' translation element discovered in triticum mosaic virus. *Journal of virology* **2015**, *89*, 12427-12440.
94. Krzywinski, M.; Schein, J.; Birol, I.; Connors, J.; Gascoyne, R.; Horsman, D.; Jones, S.J.; Marra, M.A. Circos: an information aesthetic for comparative genomics. *Genome research* **2009**, *19*, 1639-1645.
95. Lorenz, R.; Bernhart, S.; Siederdisen, C.; Tafer, H.; Flamm, C.; Stadler, P.; Hofacker, I. ViennaRNA package 2.0. algorithms for molecular biology. *vol* **2013**, *6*, 26-26.
96. Li, Y.; Liu, R.; Zhou, T.; Fan, Z. Genetic diversity and population structure of Sugarcane mosaic virus. *Virus research* **2013**, *171*, 242-246.
97. Ashraf, M.A.; Tariq, H.K.; Hu, X.-W.; Khan, J.; Zou, Z. Computational Biology and Machine Learning Approaches Identify Rubber Tree (*Hevea brasiliensis* Muell. Arg.) Genome Encoded MicroRNAs Targeting Rubber Tree Virus 1. *Applied Sciences* **2022**, *12*, 12908.
98. Ashraf, M.A.; Ali, B.; Brown, J.K.; Shahid, I.; Yu, N. In Silico Identification of Cassava Genome-Encoded MicroRNAs with Predicted Potential for Targeting the ICMV-Kerala Begomoviral Pathogen of Cassava. *Viruses* **2023**, *15*, 486.
99. Shahid, M.N.; Rashid, S.; Iqbal, M.S.; Jamal, A.; Khalid, S. In Silico prediction of potential mirnas to target zymv in cucumis melo. *Pak. J. Bot* **2022**, *54*, 1319-1325.
100. Jabbar, B.; Iqbal, M.S.; Batcho, A.A.; Nasir, I.A.; Rashid, B.; Husnain, T.; Henry, R.J. Target prediction of candidate miRNAs from *Oryza sativa* for silencing the RYMV genome. *Computational biology and chemistry* **2019**, *83*, 107127.
101. Akhter, Y.; Khan, J.A. Genome wide identification of cotton (*Gossypium hirsutum*)-encoded microRNA targets against Cotton leaf curl Burewala virus. *Gene* **2018**, *638*, 60-65.
102. Iqbal, M.S.; Jabbar, B.; Sharif, M.N.; Ali, Q.; Husnain, T.; Nasir, I.A. In silico MCMV silencing concludes potential host-derived miRNAs in maize. *Frontiers in plant science* **2017**, *8*, 372.
103. Gaafar, Y.Z.A.; Ziebell, H. Novel targets for engineering Physostegia chlorotic mottle and tomato brown rugose fruit virus-resistant tomatoes: in silico prediction of tomato microRNA targets. *PeerJ* **2020**, *8*, e10096.
104. Yang, X.; Zhang, L.; Yang, Y.; Schmid, M.; Wang, Y. miRNA mediated regulation and interaction between plants and pathogens. *International Journal of Molecular Sciences* **2021**, *22*, 2913.
105. Mengistu, A.A.; Tenkegn, T.A. The role of miRNA in plant-virus interaction: a review. *Molecular Biology Reports* **2021**, *48*, 2853-2861.
106. Petchthai, U.; Yee, C.S.L.; Wong, S.-M. Resistance to CymMV and ORSV in artificial microRNA transgenic *Nicotiana benthamiana* plants. *Scientific reports* **2018**, *8*, 1-8.
107. Cui, C.; Wang, J.-J.; Zhao, J.-H.; Fang, Y.-Y.; He, X.-F.; Guo, H.-S.; Duan, C.-G. A Brassica miRNA regulates plant growth and immunity through distinct modes of action. *Molecular plant* **2020**, *13*, 231-245.
108. Yasir, M.; Motawaa, M.; Wang, Q.; Zhang, X.; Khalid, A.; Cai, X.; Li, F. Simple webserver-facilitated method to design and synthesize artificial miRNA gene and its application in engineering viral resistance. *Plants* **2022**, *11*, 2125.
109. Mortazavi, S.S.; Bahmanpour, Z.; Daneshmandpour, Y.; Roudbari, F.; Sheervalilou, R.; Kazeminasab, S.; Emamalizadeh, B. An updated overview and classification of bioinformatics tools for MicroRNA analysis, which one to choose? *Computers in Biology and Medicine* **2021**, *134*, 104544.
110. Li, Y.; Li, C.; Ding, G.; Jin, Y. Evolution of MIR159/319 microRNA genes and their post-transcriptional regulatory link to siRNA pathways. *BMC evolutionary biology* **2011**, *11*, 1-18.