

Brief Report

First report of the plastid genome of *Passiflora tripartita* var. *mollissima* (poro-poro) from Huánuco, Peru

Flavio Aliaga ^{1,2,3,*}, Mario Zapata-Cruz ² and Silvia A. Valverde-Zavaleta ^{2,*}

¹ Dirección de Investigación, Innovación y Responsabilidad Social, Universidad Privada del Norte (UPN), Trujillo 13011, Peru

² Plant Science Laboratory (PSL), Trujillo 13009, Peru

³ Capítulo de Ingeniería Agronómica, Consejo Departamental de La Libertad (CDLL), Colegio de Ingenieros del Perú (CIP), Trujillo 13008, Peru

* Correspondence: flavio.aliaga@upn.edu.pe

Abstract: Poro-poro is an important native fruit used in traditional Peruvian medicine with relevant agro-industrial and pharmaceutical potential for its antioxidant capacity for human health. However, to date, there have been few genetic studies. The lack of genomic exploration limits the possibility of expanding our knowledge of its molecular evolution, new molecular pathways, genetic traits, and evolutionary relationships. Here, we report the plastid sequence generated on the Illumina system platform using 26 *Passiflora* genomes and 1 outgroup from the NCBI database, and the reconstructed phylogenetic relationship. Our phylogenetic analysis showed that poro-poro is most closely related to *Passiflora menispermifolia* and *Passiflora oertedii*. In summary, our study provides the basis for developing new molecular markers that constitutes a valuable resource for studying molecular evolution and domestication. It also provides a powerful foundation for conservation genetics research and plant breeding programs. To our knowledge, this is the first report on the plastid genome of *Passiflora tripartita* var. *mollissima* from Peru.

Keywords: Chloroplast genome, *Passifloraceae*, *Passiflora tripartita*, poro-poro, Huánuco, Peru

1. Introduction

Passiflora tripartita var. *mollissima* (Kunth) Holms-Niels. & P.M. Jørg [1] previously known as *Passiflora mollissima* (Kunth) Bailey [2], is a semi-perennial fruit plant [3]. It is a diploid species with a small number of chromosomes ($2n = 18$) [4], which is placed in the section Elkea of supersection Tacsonia of subgenus *Passiflora* belonging to the *Passifloraceae* family [5, 6]. Poro-poro is a native fruit of the Andean region [5]. It grows in the Peruvian highlands in the departments of Ancash, Junín, Moquegua, Huancavelica, and Huánuco at altitudes of 1,000–4,000 m.a.s.l. [7, 8]. It is widely used in traditional medicine [8] and is considered one of the best *Passiflora* species based on its organoleptic characteristics [2]. This fruit provides a source of vitamins (A, B3, and C) and minerals (magnesium, potassium, phosphorus, sodium, chlorine, iron, calcium, sulfur, zinc, copper, selenium, cobalt, and nickel) [9, 10]. In addition, it has an elevated antioxidant activity and high content of carotenoids (118.8 mg β -carotene), phenols (460.1 mg gallic acid), and flavonoids (1907.6 mg catechin/100 g) [9, 10]. Specifically, the high concentration of flavan-3-ols (a group of bioactive compounds) has been associated with beneficial effects on human health, such as cardiovascular protection, neurodegenerative diseases, and as an anti-cancer, anti-microbial, and anti-parasitic agent [11, 12].

Plastome sequences of more than 800 sequenced genomes are small in size with high copy numbers and conserved sequences, enabling a significant understanding of plant molecular evolution, structural variations, and evolutionary relationships of plant diversity [13, 14]. The plastid genome has a quadripartite structure: a large single-copy (LSC) of 80–90 kilobase pairs (kb), a small single-copy (SSC) of 16–27 kb, and two sets of inverted

repeats (IRa and IRb) of 20–28 kb, with 110–130 unique genes, including protein-coding genes, transfer RNA (tRNA), and ribosomal RNA (rRNA) [15, 16].

In recent years, declining genome sequencing costs resulted in more than 780 complete plant genomes of different species becoming available [17, 18]. Recently, some *Passiflora* plastid genomes such as *Passiflora edulis* [19], *Passiflora xishuangbannaensis* [20], *Passiflora caerulea* [21], *Passiflora serrulata* [22], *Passiflora foetida* [23], and *Passiflora arbelaezii* [24], became publicly available. However, despite the scarcity of genomic information on underutilized crops [25], we have only begun to investigate the genomics of plants of great importance for plant breeding programs. The aim of the present study was to sequence, assemble, and annotate the plastid genome of poro-poro to contribute to plant breeding programs. In the present study, we report the first plastid genome sequence submitted for an isolate of *Passiflora tripartita* var. *mollissima* from Peru, a species with great agro-industrial and pharmaceutical potential because of its beneficial characteristics for human health.

2. Results

2.1. Plastid Genome Features

The plastid genome of *P. tripartita* var. *mollissima* (poro-poro) (Figure 1) was 163,451 bp in length, with a typical quadripartite structure consisting of a large single-copy (LSC) region of 85,525 bp (52.32% in total) and a small single-copy (SSC) region of 13,518 bp (8.27%), separated by a pair of inverted repeat regions (IRa and IRb) of 32,204 bp (19.70%). The overall AT content of the plastid genome was 63.13%, whereas the overall GC content was 36.87%. The plastome structure of the *P. tripartita* var. *mollissima* consisted of A = 30.79%, T(U) = 32.34%, C = 18.67% and G = 18.20%. Poro-poro plastid genome annotation identified 128 genes, of which 112 were unique, and 17 were duplicated in the inverted repeat (IR) region. The plastome contained 84 protein-coding genes, 37 transfer RNA (tRNA)-coding genes, 7 ribosomal RNA (rRNA)-coding genes, and 14 genes with introns (12 genes with one intron and 2 genes with two introns), as shown in Table 1.




Figure 1. Plastid genome of *Passiflora tripartita* var. *mollissima*. The thick lines indicate the IR1 and IR2 regions, which separate the SSC and LSC regions. Genes marked inside the circle are transcribed clockwise, and genes marked outside the circle are transcribed counterclockwise. Genes are color-coded based on their function, shown at the bottom left. The inner circle indicates the inverted boundaries and GC content.

Table 1. Plastid genome features of the *P. tripartita* var. *mollissima*

Features	Poro-poro ¹
Genome size (bp)	163,451
LSC length (bp)	85, 525
SSC length (bp)	13,518
IR length (bp)	32,204

Total GC content (%)	36.87
A content (%)	30.79
T(U) content (%)	32.34
G content (%)	18.20
C content (%)	18.67
Total number of genes	128
Protein-coding genes	84
rRNA coding genes	7
tRNA coding genes	37
Genes duplicated in IR regions	17
Total introns	14
Single introns (gene)	12
Double introns (gene)	2

¹ Poro-poro is the common name of *Passiflora tripartita* var. *mollissima* in Peru.

The poro-poro plastid genome contained 112 unique genes, of which there were 29 tRNA genes, 4 rRNA genes, and 79 protein-coding genes. The latter comprised 20 ribosomal subunit genes (9 large subunits and 11 small subunit), 4 DNA-directed RNA polymerase genes, 46 genes were involved in photosynthesis (11 encoded subunits of the NADH oxidoreductase, 7 for photosystem I, 15 for photosystem II, 6 for the cytochrome b6/f complex, 6 for different subunits of ATP synthase, and 1 for the large chain of ribulose biphosphate carboxylase), 8 genes were involved in different functions, and one gene was of unknown function (Table 2).

Table 2. Genes present in the plastid genome of *P. tripartita* var. *mollissima*

Group of genes	Gene names
Photosystem I	psaA, psaB, psaC, psaI, psaJ, ycf3 **, ycf4
Photosystem II	psbA, psbB, psbC, psbD, psbE, psbF, psbH, psbI, psbJ, psbK, psbL, psbM, psbN, psbT, psbZ
Cytochrome b/f complex	petA, petB, petD *, petG, petL, petN
ATP synthase	atpA, atpB, atpE, atpF, atpH, atpI
NADH dehydrogenase	ndhA*, ndhB * (X2), ndhC, ndhD, ndhE, ndhF, ndhG, ndhH, ndhI, ndhJ, ndhK
RubisCO large subunit	rbcL
DNA-dependent RNA polymerase	rpoA, rpoB, rpoC1 *, rpoC2
Ribosomal proteins (SSU)	rps2, rps3, rps4, rps8, rps11, rps12 ** (X2), rps14, rps15, rps16, rps18, rps19 (X2)
Ribosomal proteins (LSU)	rpl2 * (X2), rpl14, rpl16 *, rpl20, rpl22, rpl23 (X2), rpl32, rpl33, rpl36
Acetyl-CoA carboxylase	accD
C-type cytochrome synthesis	ccsA
Envelope membrane protein	cemA
Protease	clpP
Translational initiation factor IF-1	infA
Maturase	matK
Component of TIC complex	yct1, ycf2
Unknown function protein-coding	ycf15 (X2)
Ribosomal RNAs	rrn4.5, rrn5 (X2), rrn16 (X2), rrn23 (X2) trnA-UGC * (X2), trnC-GCA, trnD-GUC, trnE-UUC, trnF-GAA, trnG-GCC, trnG-UCC *, trnH-GUG, trnI-CAU (X2), trnI-GAU * (X2), trnK-UUU *, trnL-CAA (X2), trnL-UAA *, trnL-UAG, trnM-
Transfer RNAs	

CAU (X2), trnN-GUU (X2), trnP-UGG, trnQ-UUG, trnR-ACG (X2), trnR-UCU, trnS-GCU, trnS-GGA, trnS-UGA, trnT-GGU, trnT-UGU, trnV-GAC (X2), trnV-UAC *, trnW-CCA, trnY-GUA

* Gene contains one intron; ** gene contains two introns; (X2) indicates two gene copies in IRs.

In the plastid genome, 14 genes contained introns distributed as follows: the LSC, SSC, and IRs regions contained 8 genes (petD, rpl16, rpoC1, trnG-UCC, trnK-UUU, trnL-UAA, trnV-UAC, and ycf3), 1 gene (ndhA), and 5 genes (ndhB, rpl2, rps12, trnA-UGC, and trnI-GAU) respectively. Similarly, these genes included six protein-coding genes, each with a single intron (petD, ndhA, ndhB, rpoC1, rpl2, and rpl16); six tRNA genes, each with a single intron (trnA-UGC, trnG-UCC, trnI-GAU, trnK-UUU, trnL-UAA, and trnV-UAC); and two protein-coding genes with two introns (ycf3 and rps12). Except for 17 genes that were duplicated in the IR region (ndhB, rps19, rpl2, rpl23, rps12, ycf15, rrn5, rrn16, rrn23, trnA-UGC, trnI-CAU, trnI-GAU, trnL-CAA, trnM-CAU, trnN-GUU, trnR-ACG, and trnV-GAC) all genes contained a single copy, as shown in Table 2.

2.2. Phylogenetic Analysis

To identify the evolutionary position of *Passiflora tripartita* var. *mollissima* in the Passifloraceae family, phylogenetic relationships based on the OrthoFinder clustering method were used to perform orthologous group analyses with the complete annotated protein sequences of 27 plastid genomes, of which 26 were from *Passiflora* species. One species, *Vitis vinifera*, was chosen as the outgroup.

Maximum likelihood (ML) bootstrap values ranged from 38%–92% for 7 of the 25 nodes. All nodes except the indicated ones (seven nodes) exhibited bootstrap support (BS) values of 100%. These *Passiflora* species were divided into four groups: subgenus *Passiflora* (*P. nitida*, *P. quadrangularis*, *P. cincinnata*, *P. caerulea*, *P. edulis*, *P. laurifolia*, *P. vitifolia*, *P. serratifolia*, *P. serrulata*, *P. ligularis*, *P. serratodigilata*, *P. actinia*, *P. menispermifolia* and *P. oerstedii*), subgenus *Tetrapathea* (*P. tetrandra*), subgenus *Decaloba* (*P. microstipula*, *P. xishuangbannaensis*, *P. biflora*, *P. lutea*, *P. jatunsachensis*, *P. suberosa* and *P. tenuiloba*), and subgenus *Deidamoides* (*P. contracta* and *P. arbelaezii*). These results resolved *Passiflora tripartita* var. *mollissima* in the subgenus *Passiflora*, which was closely related to *P. menispermifolia* and *P. oerstedii* with 100% BS, as shown in the cladogram (Figure 2).




Figure 2. Phylogenetic tree of 27 plastid genomes using maximum likelihood analysis based on single-copy orthologous protein. Bootstrap values on the branches were calculated from 100 replicates.

3. Discussion

3.1. Plastome of *Passiflora tripartite* var. *mollissima*

The poro-poro plastome is 12,045 bp longer than that of one of the most economically important species, passion fruit (*P. edulis*) [19], and is only 7,117 bp longer than that of the longest *Passiflora* plastome reported, i.e., *P. arbelaezii* [24]. The plastome sequence of poro-poro has a quadripartite architecture consisting of an LSC and SSC separated by a pair of inverted repeat regions (IRa and IRb) as similar to other plants [26–28]. However, the LSC region is 4,150 bp longer than that of *P. xishuangbannaensis* but is 98bp, 195 bp, and 1,927 bp shorter than that of *P. caerulea*, *P. edulis*, and *P. arbelaezii*, respectively. The SSC region is 121 bp, 140 bp, 359 bp, and 754 bp longer than that of *P. caerulea*, *P. edulis*, *P. xishuangbannaensis*, and *P. arbelaezii*, respectively. The IRs regions are 6,024 bp, 6,050 bp, and 11,600 longer than that of *P. caerulea*, *P. edulis*, and *P. xishuangbannaensis*, respectively; however, it is 2,972 bp shorter than that of *P. arbelaezii* [19–21, 24, 29]. The overall GC content of the plastome (36.87%) was similar to that of other reported chloroplast genomes from the same family, such as 36.90% in *P. arbelaezii* [24], 37% in *P. edulis* and *P. serrulata* [19, 22], 37.03% in *P. caerulea* [21], and 37.1% in *P. xishuangbannaensis* [20].

The plastome of *P. tripartita* var. *mollissima* contained eight genes (*ycf1*, *ycf2*, *ycf15*, *rps16*, *rpl20*, *rpl22*, *accD*, *infA*) that were lost or non-functional genes in *P. edulis*; and compared to *P. edulis*, it has one absent gene (*trnfM-CAU*), as previously reported [19]. In this study, the *ycf1* sequence encodes a protein essential for plant viability and a vital component of the translocon on the inner chloroplast membrane (TIC) complex [30], and *ycf2* is a component of the ATPase motor protein associated with the TIC complex [31].

3.2. Phylogenetic Reconstruction of *P. tripartita* var. *mollissima*

Using concatenated single-copy orthologous proteins in phylogenetic relationships avoids erroneous rearrangements in phylogenetic tree reconstruction and provides a

more reliable evolutionary analysis [32, 33]. The phylogenetic tree was constructed based on single-copy orthologous genes [34] and maximum likelihood analysis. The relationships between the four subgenera of *Passiflora* species (*Passiflora*, *Tetrapathea*, *Decaloba*, and *Deidamoides*) were congruent and strongly supported by the same patterns as previously reported [35, 36]. *Passiflora tripartita* var. *mollissima* was placed in the subgenus *Passiflora* with *P. menispermifolia* and *P. oerstedii* in the same group and was sister to *P. tetrandra* (subgenus *Tetrapathea*), *P. biflora* (subgenus *Decaloba*), and *P. contracta* (subgenus *Deidamoides*).

4. Materials and Methods

4.1. Plant Material

In November 2022, the fresh leaves of *Passiflora tripartita* var. *mollissima* (Supplementary Figure S1) were collected from Raccha Cedrón locality of Quisqui District, Huánuco Province from Peru (9°53'37"S, 76°26'02"W, altitude 2,945 m.a.s.l.). A herbarium voucher specimen (USM<PER>:MHN331530) was deposited in the Herbario San Marcos (USM) of the Museo de Historia Natural (MHN) at the Universidad Nacional Mayor de San Marcos (UNMSM).

4.2. DNA Extraction

Total genomic DNA was extracted from approximately 100 mg fresh leaves (from voucher number USM<PER>:MHN331530) using a cetyl-trimethyl ammonium bromide (CTAB) protocol [37]. Genomic DNA quality was assessed using a fluorometry-based Qubit (Thermo Fisher Scientific, USA) coupled to a Broad Range Assay kit. High-quality DNA (230/260 and 260/280 ratios >1.8) was normalized (20 ng/μL) to examine its integrity using 1% (w/v) agarose gel electrophoresis.

4.3. Genome Sequencing, Assembly, and Annotation

4.3.1. Sequencing

Qualified DNA was fragmented, and the TruSeq Nano DNA kit (Illumina, San Diego, CA, USA) was used to construct an Illumina paired-end (PE) library. PE sequencing (2 × 150 bp) was performed using the Illumina NovaSeq 6000 platform [38] (Macrogen, Inc., Seoul, Republic of Korea).

4.3.2. Assembly

All adapters and low-quality reads were removed using the FastQC [39] and Cutadapt [40] programs. PE reads (2 × 150 bp) were evaluated for quality using QUAST [41] analysis, and subsequent steps used clean data. Then, clean reads obtained were assembled into a circular contig using NOVOPlasty v.4.3 [42], with *P. edulis* (NC_034285) as the reference.

4.3.3. Annotation

The plastid genome was annotated using the Dual Organellar GenoMe Annotator GeSeq [43] and CpGAVAS2 [44]. A circular genome map was constructed using OGDRAW v.1.3.1 [45]. Finally, the completed sequences were submitted to the NCBI GenBank under the accession number OQ910395.

4.4 Phylogenetic Analysis

We used 26 complete plastome sequences to infer the phylogenetic relationships among *Passiflora* species, and *Vitis vinifera* was used as an outgroup (Supplemental Table 1). Single-copy orthologous genes were identified using the Orthofinder version 2.2.6 pipeline [34]. For each gene family, the nucleotide sequences were aligned using the L-INS-i algorithm in MAFFT v7.453 [46]. A phylogenetic tree based on maximum likelihood

(ML) was constructed using RAxML v8.2.12 [47] with the GTRCAT model. A phylogenetic ML tree was reconstructed and edited using MEGA 11 [48] with 100 replicates.

5. Conclusions

This study provides information on the sequencing, assembly, and annotation of the plastid genome of *Passiflora tripartita* var. *mollissima* (poro-poro) from Huánuco, Peru. The plastome structure and gene content were relatively conserved, and the phylogenetic relationships illustrated that this species is positioned close to *Passiflora menispermifolia* and *Passiflora oerstedii*. This study will open up further avenues for research on plant molecular breeding, molecular markers, evolutionary studies, and conservation genetics of poro-poro.

Supplementary Materials:

The following supporting information can be downloaded from: www.mdpi.com/xxx/s1, Figure S1: Herbarium specimen voucher of *Passiflora tripartita* var. *mollissima* (Kunth) Holms-Niels. & P.M. Jørg (USM<PER>:MHN331530); Table S1: Details of the plastid genome sequences used for phylogenetic analysis.

Author Contributions: Conceptualization, F.A. M.Z.-C. and S.A.V.-Z.; methodology, F.A., M.Z.-C. and S.A.V.-Z.; formal analysis, F.A.; data curation, F.A.; writing—original draft preparation, F.A., M.Z.-C. and S.A.V.-Z.; writing—review and editing, F.A., M.Z.-C. and S.A.V.-Z.; visualization, M.Z.-C. and S.A.V.-Z.; supervision, F.A., M.Z.-C. and S.A.V.-Z.; project administration, M.Z.-C. and S.A.V.-Z.; funding acquisition, F.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Plant Science Laboratory E.I.R.L. (Sach'a Ruru grant: RIC-2022-101) and the APC was funded by Universidad Privada del Norte.

Data Availability Statement: The chloroplast genome sequence data have been deposited and openly available in GenBank of NCBI repository under the accession code: OQ910395.

Acknowledgments: We thank the Servicio Nacional Forestal y de Fauna Silvestre (SERFOR) for authorized this research project (AUT-IFL-2022-058). We thank Petr Sklenář (Charles University) and Filip Kolar (Charles University) for their help in the sample collection. We thank curator Julio C. Torres-Martínez (Museo de Historia Natural, Universidad Nacional Mayor de San Marcos) for the taxonomy identification and deposit of the plant specimen. We thank Mr. Julián Vasquez-Arriaga for administrative support (Plant Science Laboratory).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. ITIS. (2022) *Passiflora tripartita* var. *mollissima* (Kunth) Holms-Niels. & P.M. Jørg. Available online: <https://www.itis.gov/> (accessed on 01 Jun 2023).
2. Primot, S.; Coppens D'Eeckenbrugge, G.; Rioux, V.; Ocampo, J. A.; Garcin, F. Variación morfológica de tres especies de curubas (*Passiflora tripartita* var. *mollissima*, *P. tarminiana* y *P. mixta*) y sus híbridos en el Valle del Cauca (Colombia). *Rev. Bras. Frutic.* **2005**, *27*, 467–471, doi: [10.1590/S0100-29452005000300030](https://doi.org/10.1590/S0100-29452005000300030)
3. Mayorga, M.; Fischer, G.; Melgarejo, L. M.; Parra-Coronado, A. Growth, development and quality of *Passiflora tripartita* var. *mollissima* fruits under two environmental tropical conditions. *J. Appl. Bot. Food Qual.* **2020**, *93*, 66–75, doi: [10.5073/JABFQ.2020.093.009](https://doi.org/10.5073/JABFQ.2020.093.009)
4. Coppens D'Eeckenbrugge, G. Conservación y utilización de recursos genéticos de pasifloras. COLCIENCIAS Informe final 1999-2001. Available online: <https://agritrop.cirad.fr/576977> (accessed on 01 Jun 2023).
5. Ocampo J.; Coppens d'Eeckenbrugge, G.C. Morphological characterization in the genus *passiflora* l.: an approach to understanding its complex variability. *Plant Syst Evol.* **2017**, *303*, 531–558, doi: [10.1007/s00606-017-1390-2](https://doi.org/10.1007/s00606-017-1390-2)

6. Segura, S.D.; D'Eeckenbrugge, G.C.; Ocampo C.H.; Ollitrault, P. Isozyme variation in passiflora subgenus tacsonia: Geographic and interspecific differentiation among the three most common species. *Genet. Resour. Crop Evol.* **2005**, *52*, 455–463, doi: [10.1007/s10722-005-2255-z](https://doi.org/10.1007/s10722-005-2255-z)
7. Tapia, M. E; Fries, A. M. *Guía de campo de los cultivos andinos*, 1st ed.; FAO: Roma, Italy, 2007; pp. 22–114, ISBN 978-92-5-305682-8.
8. Ríos-García, G. Nivel de aceptabilidad del vino de tumbo serrano (*Passiflora mollissima*) elaborado con los parámetros tecnológicos óptimos en la ciudad de Huánuco 2015. Master thesis, Universidad Nacional Hermilio Valdizán de Huánuco, Huánuco, 06 November 2017.
9. Chaparro-Rojas, D.C.; Maldonado, M.E.; Franco-Londoño, M.C.; Urango-Marchena, L.A. Características nutricionales y antioxidantes de la fruta curuba larga (*Passiflora mollissima* Bailey). *Perspect. Nutr. Hum.* **2014**, *16*, 203–212, doi: [10.17533/udea.penh.v16n2a07](https://doi.org/10.17533/udea.penh.v16n2a07)
10. Leterme, P.; Buldgen, A.; Estrada, F.; Londoño, A.M. Mineral content of tropical fruits and unconventional foods of the Andes and the rain forest of Colombia. *Food. Chem.* **2006**, *95*, 644–652, doi: [10.1016/j.foodchem.2005.02.003](https://doi.org/10.1016/j.foodchem.2005.02.003)
11. Giambanelli, E.; Gómez-Caravaca, A.M.; Ruiz-Torralba, A.; Guerra-Hernández, E.J.; Figueroa-Hurtado, J.G.; García-Villanova, B.; Verardo, V. New advances in the determination of free and bound phenolic compounds of banana passion fruit pulp (*Passiflora tripartita*, var. *mollissima* (kunth) l.h. bailey) and their in vitro antioxidant and hypoglycemic capacities. *Antioxidants*, **2020**, *9*, 628, doi: [10.3390/antiox9070628](https://doi.org/10.3390/antiox9070628)
12. Luo, Y.; Jian, Y.; Liu, Y.; Jiang, S.; Muhammad, D.; Wang, W. Flavanols from nature: A phytochemistry and biological activity review. *Molecules*, **2022**, *27*, 719, doi: [10.3390/molecules27030719](https://doi.org/10.3390/molecules27030719)
13. Daniell, H.; Lin, C.S.; Yu, M.; Chang W.J. Chloroplast genomes: diversity, evolution, and applications in genetic engineering. *Genome Biol.*, **2016**, *17*, 134, doi: [10.1186/s13059-016-1004-2](https://doi.org/10.1186/s13059-016-1004-2)
14. Dobrogojski, J.; Adamiec, M.; Lucinski, R. The chloroplast genome: A review. *Acta Physiol. Plant.* **2020**, *42*, 98, doi: [10.1007/s11738-020-03089-x](https://doi.org/10.1007/s11738-020-03089-x)
15. Ozeki, H.; Umesono, K.; Inokuchi, H.; Kohchi, T.; Ohyama, K. The chloroplast genome of plants: a unique origin. *Genome*. **1989**, *31*, 169–174, doi: [10.1139/g89-029](https://doi.org/10.1139/g89-029)
16. Wang, W.; Lanfear, R. Long-reads reveal that the chloroplast genome exists in two distinct versions in most plants. *Genome Biol Evol.* **2019**, *11*, 3372–3381, doi: [10.1093/gbe/evz256](https://doi.org/10.1093/gbe/evz256)
17. Marks, R.A.; Hotaling, S.; Frandsen, P.B.; VanBuren, R. Representation and participation across 20 years of plant genome sequencing. *Nat Plants*. **2021**, *7*, 1571–1578, doi: [10.1038/s41477-021-01031-8](https://doi.org/10.1038/s41477-021-01031-8)
18. Sun, Y.; Shang, L.; Zhu, Q.H.; Fan, L.; Guo, L. Twenty years of plant genome sequencing: achievements and challenges. *Trends Plant Sci.* **2022**, *27*, 391–401, doi: [10.1016/j.tplants.2021.10.006](https://doi.org/10.1016/j.tplants.2021.10.006)
19. Cauz-Santos, L.A.; Munhoz, C.F.; Rodde, N.; Cauet, S.; Santos, A.A.; Penha, H.A.; Dornelas, M.C.; Varani, A.M.; Oliveira, G.C.X.; Gergès, H.; Vieira, M.L.C. The Chloroplast Genome of *Passiflora edulis* (Passifloraceae) Assembled from Long Sequence Reads: Structural Organization and Phylogenomic Studies in Malpighiales. *Front Plant Sci.* **2017**, *8*, 334, doi: [10.3389/fpls.2017.00334](https://doi.org/10.3389/fpls.2017.00334)
20. Hao, C.; Wu, F. The complete chloroplast genome sequence of *Passiflora xishuangbannaensis* (Passifloraceae), a vine endemic to Yunnan, China. *Mitochondrial DNA B Resour.* **2021**, *6*, 1763–1764, doi: [10.1080/23802359.2021.1928562](https://doi.org/10.1080/23802359.2021.1928562)
21. Niu, Y.F.; Ni, S.B.; Liu, S.H.; Liu, J. The complete chloroplast genome of *Passiflora caerulea*, a tropical fruit with a distinctive aroma. *Mitochondrial DNA B Resour.* **2021**, *6*, 488–490, doi: [10.1080/23802359.2021.1872442](https://doi.org/10.1080/23802359.2021.1872442)
22. Mou, H.F.; Huang, W.H.; Liu, J.Y.; Wen, F.; Tian, Q.L.; Wu, Y.Y.; Fu, L.F.; Zhang, Y.J.; Wei, Y.G. Complete chloroplast genome sequence of *Passiflora serrulata* Jacq. (Passifloraceae). *Mitochondrial DNA B Resour.* **2021**, *6*, 191–193, doi: [10.1080/23802359.2020.1860711](https://doi.org/10.1080/23802359.2020.1860711)
23. Hopley, T.; Webber, B.L.; Raghu, S.; Morin, L.; Byrne, M. Revealing the Introduction History and Phylogenetic Relationships of *Passiflora foetida* sensu lato in Australia. *Front Plant Sci.* **2021**, *12*, 651805, doi: [10.3389/fpls.2021.651805](https://doi.org/10.3389/fpls.2021.651805)
24. Shrestha, B.; Weng, M.L.; Theriot, E.C.; Gilbert, L.E.; Ruhlman, T.A.; Krosnick, S.E.; Jansen, R.K. Highly accelerated rates of genomic rearrangements and nucleotide substitutions in plastid genomes of *Passiflora* subgenus *Decaloba*. *Mol. Phylogenet. Evol.* **2019**, *138*, 53–64, doi: [10.1016/j.ympev.2019.05.030](https://doi.org/10.1016/j.ympev.2019.05.030)
25. Gioppato, H.A.; da Silva, M.B.; Carrara, S.; Palermo, B.R.Z.; de Souza Moraes, T.; Dornelas, M.C. Genomic and transcriptomic approaches to understand *Passiflora* physiology and to contribute to passionfruit breeding. *Theor. Exp. Plant Physiol.* **2019**, *31*, 173–181, doi: [10.1007/s40626-018-0134-1](https://doi.org/10.1007/s40626-018-0134-1)
26. Ohyama, K.; Fukuzawa, H.; Kohchi, T.; Shirai, H.; Sano, T.; Sano, S.; Umesono, K.; Shiki, Y.; Takeuchi, M.; Chang, Z.; Aota, S.I.; Inokuchi, H.; Ozeki, H. Chloroplast gene organization deduced from complete sequence of liverwort *Marchantia polymorpha* chloroplast DNA. *Nature*. **1986**, *322*, 572–574, doi: [10.1038/322572a0](https://doi.org/10.1038/322572a0)

27. Shinozaki, K.; Ohme, M.; Tanaka, M.; Wakasugi, T.; Hayashida, N.; Matsubayashi, T.; Zaita, N.; Chunwongse, J.; Obokata, J.; Yamaguchi-Shinozaki, K.; Ohto, C.; Torazawa, K.; Meng, B. Y.; Sugita, M.; Deno, H.; Kamogashira, T.; Yamada, K.; Kusuda, J.; Takaiwa, F.; Kato, A.; Tohdoh, N.; Shimada, H.; Sugiura, M. The complete nucleotide sequence of the tobacco chloroplast genome: its gene organization and expression. *EMBO J.*, **1986**, *5*, 2043–2049, doi: [10.1002/j.1460-2075.1986.tb04464.x](https://doi.org/10.1002/j.1460-2075.1986.tb04464.x)
28. Nguyen, H.Q.; Nguyen, T.N.L.; Doan, T.N.; Nguyen, T.T.N.; Pham, M.H.; Le, T.L.; Sy, D.T.; Chu, H.H.; Chu, H.M. Complete chloroplast genome of novel *Adrinandra megaphylla* Hu species: molecular structure, comparative and phylogenetic analysis. *Sci. Rep.* **2021**, *11*, 11731, doi: [10.1038/s41598-021-91071-z](https://doi.org/10.1038/s41598-021-91071-z)
29. Zheng, S.; Pocai, P.; Hyvönen, J.; Tang, J.; Amiroussi, A. Chloroplast: An Online Program for the Versatile Plotting of Organelle Genomes. *Front. Genet.* **2020**, *11*, 576124, doi: [10.3389/fgene.2020.576124](https://doi.org/10.3389/fgene.2020.576124)
30. Kikuchi, S.; Bédard, J.; Hirano, M.; Hirabayashi, Y.; Oishi, M.; Imai, M.; Takase, M.; Ide, T.; Nakai, M. Uncovering the protein translocon at the chloroplast inner envelope membrane. *Science*. **2013**, *339*, 571–574, doi: [10.1126/science.1229262](https://doi.org/10.1126/science.1229262)
31. Kikuchi, S.; Asakura, Y.; Imai, M.; Nakahira, Y.; Kotani, Y.; Hashiguchi, Y.; Nakai, Y.; Takafuji, K.; Bédard, J.; Hirabayashi-Ishioka, Y.; Mori, H.; Shiina, T.; Nakai, M. A Ycf2-FtsHi Heteromeric AAA-ATPase Complex Is Required for Chloroplast Protein Import. *Plant Cell*. **2018**, *30*, 2677–2703, doi: [10.1105/tpc.18.00357](https://doi.org/10.1105/tpc.18.00357)
32. Gabaldón T. Evolution of proteins and proteomes: a phylogenetics approach. *Evol. Bioinform. Online*. **2005**, *1*, 51–61, doi: [10.1177/117693430500100004](https://doi.org/10.1177/117693430500100004)
33. Zhang, N.; Zeng, L.; Shan, H.; Ma, H. Highly conserved low-copy nuclear genes as effective markers for phylogenetic analyses in angiosperms. *New Phytol.* **2012**, *195*, 923–937, doi: [10.1111/j.1469-8137.2012.04212.x](https://doi.org/10.1111/j.1469-8137.2012.04212.x)
34. Emms, D.M.; Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **2019**, *20*, 238, doi: [10.1186/s13059-019-1832-y](https://doi.org/10.1186/s13059-019-1832-y)
35. Pacheco, T.G.; Lopes, A.S.; Welter, J.F.; Yotoko, K.S.C.; Otoni, W.C.; Vieira, L.D.N.; Guerra, M.P.; Nodari, R.O.; Balsanelli, E.; Pedrosa, F.O.; de Souza, E.M.; Rogalski, M. Plastome sequences of the subgenus *Passiflora* reveal highly divergent genes and specific evolutionary features. *Plant Mol. Biol.* **2020**, *104*, 21–37, doi: [10.1007/s11103-020-01020-z](https://doi.org/10.1007/s11103-020-01020-z)
36. Cauz-Santos, L.A.; da Costa, Z.P.; Callot, C.; Cauet, S.; Zucchi, M.I.; Bergès, H.; van den Berg, C.; Vieira, M.L.C. A Repertory of Rearrangements and the Loss of an Inverted Repeat Region in *Passiflora* Chloroplast Genomes. *Genome Biol. Evol.* **2020**, *12*, 1841–1857, doi: [10.1093/gbe/evaa155](https://doi.org/10.1093/gbe/evaa155)
37. Doyle, J.J.; Doyle, J.L. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* **1987**, *19*, 11–15.
38. Modi, A.; Vai, S.; Caramelli, D.; Lari, M. The illumina sequencing protocol and the novaseq 6000 system. In: *Bacterial Pangenomics: Methods and Protocols*, 2nd edn; Mengoni, A.; Bacci, G.; Fondi, M., Eds.; Springer: New York, USA, 2021; Volume 2242, pp. 15–42, ISBN: 978-1-0716-1099-2.
39. Wingett, S.W.; Andrews, S. FastQ Screen: A tool for multi-genome mapping and quality control. *F1000Res*. **2018**, *7*, 1338, doi: [10.12688/f1000research.15931.2](https://doi.org/10.12688/f1000research.15931.2)
40. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* **2011**, *17*, 10–12, doi: [10.14806/ej.17.1.200](https://doi.org/10.14806/ej.17.1.200)
41. Gurevich, A.; Saveliev, V.; Vyahhi, N.; Tesler, G. QUAST: quality assessment tool for genome assemblies. *Bioinformatics*. **2013**, *29*, 1072–1075, doi: [10.1093/bioinformatics/btt086](https://doi.org/10.1093/bioinformatics/btt086)
42. Dierckxsens, N.; Mardulyn, P.; Smits, G. NOVOPlasty: de novo assembly of organelle genomes from whole genome data. *Nucleic Acids Res.* **2017**, *45*, e18, doi: [10.1093/nar/gkw955](https://doi.org/10.1093/nar/gkw955)
43. Tillich, M.; Lehwark, P.; Pellizzer, T.; Ulbricht-Jones, E.S.; Fischer, A.; Bock, R.; Greiner S. GeSeq - versatile and accurate annotation of organelle genomes. *Nucleic Acids Res.* **2017**, *45*, W6–W11, doi: [10.1093/nar/gkx391](https://doi.org/10.1093/nar/gkx391)
44. Shi, L.; Chen, H.; Jiang, M.; Wang, L.; Wu, X.; Huang, L.; Liu, C. CPGAVAS2, an integrated plastome sequence annotator and analyzer. *Nucleic Acids Res.* **2019**, *47*, W65–W73, doi: [10.1093/nar/gkz345](https://doi.org/10.1093/nar/gkz345)
45. Greiner, S.; Lehwark, P.; Bock, R. OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes. *Nucleic Acids Res.* **2019**, *47*, W59–W64, doi: [10.1093/nar/gkz238](https://doi.org/10.1093/nar/gkz238)
46. Katoh, K.; Standley, D.M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **2013**, *30*, 772–780, doi: [10.1093/molbev/mst010](https://doi.org/10.1093/molbev/mst010)
47. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. **2014**, *30*, 1312–1313, doi: [10.1093/bioinformatics/btu033](https://doi.org/10.1093/bioinformatics/btu033)
48. Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol. Biol. Evol.* **2021**, *38*, 3022–3027, doi: [10.1093/molbev/msab120](https://doi.org/10.1093/molbev/msab120)