

Article

An Integrated Multi-OMICS Approach Highlights Elevated Non-Esterified Fatty Acids Impact BeWo Trophoblast Metabolism and Lipid Processing

Zachary JW Easton ¹, Ousseynou Sarr ¹, Lin Zhao ¹, Adriana Zardini Buzatto ², Xian Luo ², Shuang Zhao ², Liang Li ^{2,3} and Timothy RH Regnault ^{1,4,5,6,*}

¹ Department of Physiology and Pharmacology, Western University, Medical Sciences Building Room 216, London, Ontario, Canada N6A 5C1

² The Metabolomics Innovation Centre (TMIC), University of Alberta, Edmonton, Alberta T6G 2G2, Canada

³ Department of Chemistry, University of Alberta, Edmonton, Alberta T6G 2G2, Canada

⁴ Department of Obstetrics and Gynaecology, London Health Science Centre-Victoria Hospital, B2-401, London, Ontario, Canada, N6H 5W9

⁵ Children's Health Research Institute, 800 Commissioners Road East, London, Ontario, Canada, N6C 2V5

⁶ Lawson Health Research Institute, 750 Base Line Rd E, London, Ontario, Canada, N6C 2R5

* Correspondence: tim.regnault@uwo.ca

Abstract: Maternal obesity and gestational diabetes mellitus (GDM) are linked with impaired placental function and early onset of non-communicable cardiometabolic diseases in offspring. Previous studies have highlighted that the dietary non-esterified fatty acids (NEFAs), palmitate (PA) and oleate (OA), key dietary metabolites associated with maternal obesity and GDM, are potential modulators of placental lipid processing. Using the BeWo cell line model, the current study integrated transcriptomic (mRNA microarray), metabolomic, and lipidomic readouts to characterize the underlying impacts of exogenous PA and OA on placental villous trophoblast cell metabolism. Targeted gas chromatography and thin layer chromatography highlighted that saturated and monounsaturated NEFAs differentially impact BeWo cell lipid profiles. Furthermore, cellular lipid profiles differed when exposed to single and multiple NEFA species. Additional multi-omic analyses suggested that PA exposure is associated with enrichment in β -oxidation pathways, while OA exposure is associated with enrichment in anti-inflammatory and antioxidant pathways. Overall, this study further demonstrated that dietary PA and OA are important regulators of placental lipid metabolism. Encouraging appropriate dietary advice and implementing dietary interventions that maintain appropriate placental function by limiting excessive exposure to saturated NEFAs such as PA, will continue to be crucial in the clinical management of at-risk obese and GDM pregnancies.

Keywords: BeWo cells; placental lipid metabolism; fatty acid desaturation; fatty acid elongation; transcriptome; metabolome; lipidome; OMIC integration

1. Introduction

The adverse intrauterine environments associated with maternal obesity and gestational diabetes mellitus (GDM) have been linked with an increased prevalence of pregnancy complications [1–3]. Offspring from these at-risk pregnancies are often at risk of developing non-communicable cardiometabolic health disorders including metabolic syndrome, obesity, and type 2 diabetes [4–8]. These children have been found to be impacted by these “adult-associated” metabolic diseases during early adolescence, highlighting the profound influence of the aberrant in utero programming resulting from maternal obesity and GDM [9–11]. Investigating the origins of these poor health outcomes in the exposed offspring is of great importance as the rates of obesity and GDM during pregnancy have been increasing globally over the past several decades [12,13]. It is currently estimated that one-third of pregnancies are impacted by maternal obesity, and one-sixth of pregnancies are impacted by GDM,

although these rates may be greater in some high-risk demographics [14–16]. The drastic increase in the prevalence of maternal obesity and GDM and subsequent increased rates of metabolic disease development in affected offspring will only create further strains on health care systems.

Aberrant placental metabolism in diabetic and obese pregnancies is an important factor in the in-utero programming of non-communicable cardiometabolic disorders [17–19]. These changes are thought to modulate placental nutrient metabolism and processing, ultimately leading to altered materno-fetal nutrient transport, and adverse fetal growth conditions [19–21]. Specifically, impairments in placental mitochondrial respiration have been identified in term villous trophoblast cells from obese and GDM pregnancies that leads to reduced placental energy production and cellular ATP content [22–26]. Furthermore, maternal obesity has been linked with altered placental lipid processing at term highlighted by increased fatty acid desaturation, lipid esterification, and ultimately increased placental triglyceride abundance and steatosis [20,27,28]. Placental lysates from obese pregnancies have also been found to have decreased abundance of short-chain acylcarnitine species (β -oxidation intermediates) and reduced carnitine Palmitoyltransferase 1b (CPT1b) expression that is indicative of reduced placental β -oxidative activity and suggests the increased placental steatosis in obese pregnancies may be facilitated by reduced placental lipid oxidation [20,29,30]. Likewise, maternal GDM has also been associated with increased placental steatosis at term that has been attributed to reduced placental β -oxidative activity, overall highlighting that adverse maternal environmental conditions can negatively regulate placental processing of lipid nutrient species [26,31–33].

Recently, studies have utilized “omic” based research approaches to identify and characterize the underlying biological mechanisms that facilitate placental dysfunction in at-risk pregnancies [34,35]. Metabolomic and lipidomic analyses have expanded our understanding of the mechanisms that facilitate impaired placental lipid processing. These omic-based readouts have been used to demonstrate altered neutral and polar lipid profiles (highlighted by changes in underlying fatty acid (FA) composition) in the placentae from obese and GDM pregnancies [36–40]. Pre-clinical mouse models of diet-induced maternal pre-gestational obesity have also described metabolomic and transcriptomic markers associated with altered placental lipid composition and metabolism [41,42]. Overall, these studies have suggested abnormal lipid processing occurs in the obese and diabetic placenta. Alterations in placental metabolic function and lipid handling likely lead to suboptimal transplacental nutrient transport that is associated with impacted fetal development and increased risk of metabolic disease in the exposed offspring [43,44].

Preceding the development of aberrant placental metabolic function in obese and GDM pregnancies is an increased fat supply to the placenta due to maternal dyslipidemia [45,46]. Analysis of the fasting serum of women during the third trimester of pregnancy has revealed that conditions of GDM and obesity are associated with elevated circulating levels of non-esterified fatty acids (NEFA) species [47]. These analyses also demonstrated that palmitate (C16:0; PA) and oleate (C18:1n9c; OA) are the most abundant NEFA species in serum during pregnancy [47,48]. Since PA and OA are the most abundantly consumed FA in the diets of Westernized populations, these fats themselves may be an important link between poor maternal diet and impaired placental function in obese and GDM pregnancies [49–52]. Notably, maternal dietary FA consumption has previously been shown to be an important independent regulator of placental lipid handling [27,53]. Thus, it is likely the increased circulating levels of NEFAs in obese and GDM pregnancies directly facilitates impairment in placental metabolism. However, the extent to which the PA and OA directly modulate important placental lipid metabolic functions such as lipid esterification, FA desaturation, and beta-oxidation remains ill defined.

The purpose of the current study was to elucidate the underlying mechanisms that regulate placental lipid metabolism in response to elevated dietary fat supply. Hence, we examined the impacts of a prolonged (72 hours) dietary-FA exposure using an in vitro cell culture-based system and BeWo trophoblast cells. First, we employed targeted lipidomic readouts to examine FA and neutral lipid profiles of BeWo progenitor cytotrophoblast (CT) and differentiated syncytiotrophoblast (SCT) cells. Our goals were to better understand how NEFA exposures impact placental lipid processing, and to

elucidate the impacts of NEFA exposures on placental FA desaturation and elongation processing. We postulated that exposure to elevated levels of dietary NEFA would alter BeWo trophoblast lipid profiles, which would be highlighted by increased TG abundance, and increased FA elongation and desaturation. Second, we used a transcriptomic mRNA microarray, in conjunction with untargeted metabolomic and lipidomic analyses to explore the underlying mechanisms that regulate BeWo CT metabolism in response to elevated dietary NEFA supply. Undifferentiated progenitor CT cells have previously been demonstrated to be the most metabolically active cells of the villous trophoblast layer as well as the primary site of nutrient localization and storage in the placenta [26,54–56]. Thus, the current study sought to examine “omic” profiles in these more metabolically active CT cells. We postulated that dietary-FA treated BeWo CT cells would display altered transcriptomic, metabolomic and lipidomic profiles indicative of altered lipid processing.

2. Materials and Methods

2.1. Materials

Materials for cell culture, and targeted lipidomic (gas chromatography and thin layer chromatography) procedures were purchased from Millipore Sigma (Oakville, Canada) unless otherwise indicated. The solvents and reagents for untargeted lipidomics and metabolomics were acquired from Fisher Scientific (Hampton, USA), Sigma-Aldrich (St. Louis, USA), and Honeywell (Charlotte, USA). Deuterated lipid standards for untargeted lipidomics were acquired from Avanti Polar Lipids (Alabaster, USA).

2.2. Cell culture protocol

BeWo trophoblast cells (CCL-98) were purchased from the American Type Culture Collection (ATCC; Cedarlane Labs, Burlington, Canada) and cultured in F12K media (Gibco, ThermoFisher Scientific, Mississauga, Canada) as per ATCC guidelines. Cell culture media was supplemented with 10% Fetal Bovine Serum (Gibco) and 1% Penicillin-Streptomycin (Invitrogen, ThermoFisher Scientific, Mississauga, Canada) for all experiments (complete F12K media). Cultures between passages 5–15 were utilized, and all cells were maintained at 5% CO₂/95% atmospheric air for all collections.

Cells were cultured with exogenous PA and OA as previously reported [56]. In brief, BeWo trophoblast cells were plated onto cell culture dishes at the specific densities stated for each experiment and allowed to adhere to cell culture plates overnight prior to NEFA treatment. At T0H, cells were treated with complete F12K media further supplemented with either 100 μ M PA, 100 μ M OA, or 50 μ M each PA and OA (P/O). All fatty acids were conjugated 2:1 (molar ratio) to Bovine Serum Albumin (BSA) to solubilize in cell culture media. Complete F12K media supplemented with BSA-alone (0.3%) was utilized as a control. Cell media was replenished every 24 hours, and subsets of cells were additionally treated with 250 μ M 8-Br-cAMP at T24H and T48H to induce CT-to-SCT differentiation. All cultures were collected for analysis after 72 hours of NEFA exposure.

The 100 μ M concentrations of NEFAs utilized, as used in this study, were reflective of physiological fasting levels of PA and OA during the third trimester of gestation [47], and more importantly has been previously demonstrated by our research group to not impact BeWo trophoblast viability or differentiation capacity at 72-hours [56]. Thus, the data presented in the current study highlight alterations in placental metabolic function independent from lipotoxic effects.

2.3. Fatty Acid and Neutral Lipid Profile Analysis of NEFA-treated BeWo trophoblasts

BeWo CT and SCT cells were plated at a density of 2×10^6 cells in 150 mm cell culture dishes and cultured with exogenous NEFA species as described. At T72H cells were collected via scrapping and washed twice with PBS, and the resulting cell pellet was flash frozen in liquid nitrogen and stored at -80°C until analyzed.

Total cellular lipids were then collected from cell pellets via chloroform/methanol extraction (Folch extraction method) as has previously been reported and adapted for use with cell pellets [57–59]. In brief, cells were lysed in 2:1 chloroform:methanol with 0.01% butylated hydroxytoluene (BHT). Subsequently, 0.25% KCl was added, and samples were heated at 70°C for 5 mins to separate organic and aqueous layers. The organic layer was then collected and dried under a gentle stream of N_2 gas. The collected lipids were resuspended in 2:1 chloroform:methanol (0.01% BHT) at 1 mg/mL concentrations and stored at -20°C until analysis.

To examine fatty acid profiles, 100 μg of lipids were methylated to create fatty acid methyl esters (FAMES), and subsequently separated and analyzed on an 6890 N gas chromatograph (Hewlett Packard, Palo Alto, CA, USA) with flame ionization detector (Agilent Technologies, Santa Clara, CA, USA) via a J&W Scientific High-Resolution Gas Chromatography Column (DB-23, Agilent Technologies; 30 m \times 250 μm ID \times 0.25 μm film thickness) as previously reported [58,59]. Separated FA species were then identified via comparison of retention times with known standards (Supelco 37 Component FAME Mix). The peak area of each fatty acid was then quantified and expressed as a percent of total fatty acids detected.

Ratios of FAs were then utilized to indirectly quantify the activities of fatty acid desaturase enzymes by calculating desaturation indices which reflect the ratio of enzyme product to substrate. Stearoyl-CoA Desaturase 1 (SCD1) activity was calculated as the ratio of 16:1n7/16:0 as well as 18:1n9/18:0; Fatty Acid Desaturase 1 (FADS1) activity was calculated as the ratio of 20:4n6/20:3n6; and Fatty Acid Desaturase 2 (FADS2) activity was calculated as the ratio of 18:3n6/18:2n6 as previously reported [60–63]. Additionally, FA elongation indices of 20:1n9/18:1n9; 18:0/16:0 and 22:5n3/20:5n3 were quantified to calculate the activity of FA elongase enzymes (reflective of enzyme product to substrate ratio) as previously reported in placental lysates [64].

Finally, the profiles of neutral lipid species in NEFA treated BeWo trophoblasts were analyzed via Thin Liquid Chromatography coupled Flame Ionization (TLC-FID) detection as previously described [65]. In brief, collected lipid samples were dried under a gentle stream of N_2 gas and resuspended in 100% chloroform. 1–4 μL of lipids were then spotted onto TLC rods (Chromarod Type S5; Shell-USA, Spotsylvania, VA, USA) and developed in a TLC chamber with benzene:chloroform:formic acid (70:30:0.5 v/v/v). Chromatography rods were then dried and analyzed using the Iatroscan MK-6 TLC-FID system (Shell-USA). Peaks were identified via comparison with known standards, and neutral lipid peak area was quantified and expressed as a percentage of all neutral lipid species identified.

Percent fatty acid composition, percent neutral lipid composition as well as FA desaturation and elongation index data were analyzed via Randomized Block Two-Way ANOVA (2WA) and Tukey's Multiple Comparisons Test. All statistical analysis was performed with GraphPad Prism v9.2.0 software (GraphPad Software, San Diego, CA, USA).

2.4. Transcriptomic Profiling of NEFA-treated BeWo Cytotrophoblast cells

BeWo CT cells were plated at a density of 3.5×10^5 cells/well in 60 mm cell culture dishes and cultured with exogenous NEFAs as described above. At T72H the CT cells were washed once with PBS, collected in 0.9 mL TRIzol reagent (Invitrogen), and stored at -80°C . Samples were then sent to the Genome Québec Innovation Centre for transcriptomic analysis using the Clariom S mRNA microarray (ThermoFisher Scientific, Mississauga, Canada), as previous detailed [66]. In brief, RNA was automatically extracted using the QIAcube Connect (Qiagen, Toronto, Canada). RNA concentration was determined using a Nanodrop 2000 (Nanodrop Technologies Inc.), and RNA integrity was confirmed using a Bioanalyzer 2100 (Agilent Technologies, Waldbronn, Germany). cDNA was created from 100 ng total extracted RNA, fragmented, and labelled using the GeneChip WT Terminal

Labeling Kit (ThermoFisher Scientific) following manufacturer's standard protocol. Labelled cDNA was hybridized to the Clariom S microarray chip using the GeneChip Hybridization oven 640 (Affymetrix, ThermoFisher Scientific) for 17 hours (60 rpm, 45 °C). The GeneChip Hybridization Wash and Stain Kit (ThermoFisher Scientific) was used to wash the microarray chips, and they were then scanned on a GeneChip scanner 3000 (ThermoFisher Scientific).

Microarray data was subsequently analyzed using Transcriptome Analysis Console v4.0 (ThermoFisher Scientific). Raw data was normalized using the Robust Multiple-Array Averaging (RMA) method. Each NEFA treatment condition was then compared to the BSA control group in binary, and samples collected from the same passage were paired for analysis. Transcripts with $\geq \pm 1.3$ fold-change (FC) vs BSA control, and raw p-value < 0.05 were determined to be differentially expressed.

To elucidate the impacts of gene expression changes in NEFA-treated BeWo CT cells, we next performed enrichment pathway analysis using both an over-representation analysis and Gene Set Enrichment Analysis (GSEA) approach. For over-representation enrichment analysis, each set of differentially expressed genes (DEGs) was imported into the WEB-based Gene SeT AnaLysis Toolkit (WebGestalt) for using the Wikipathways, Reactome, and Kyoto Encyclopedia of Genes and Genomes (KEGG) functional datasets, as well as the GeneOntology (GO) molecular function and biological processes datasets. Pathways with FDR-corrected p-value < 0.05 were considered significantly enriched. For the GSEA approach, all genes identified in the microarray panel were ranked according to signal-to-noise ratio using GSEA v4.2.3 software (Broad Institute Inc, Cambridge MA, USA) [67,68]. The ranked gene lists were then uploaded to WebGestalt for GSEA analysis using the Wikipathways, Reactome, and KEGG functional datasets, as well as the GO molecular function and biological pathways datasets [69]. GSEA pathways with an FDR-correct p-value < 0.25 were determined to be significantly enriched in the NEFA-treated BeWo CT cells, as has previously been reported [68,70].

2.5. RT-qPCR Validation of differentially expressed genes identified by mRNA microarray

Differentially expressed genes highlighted to be involved in lipid metabolic processes were selected for validation by RT-qPCR. RNA extracted by Genome Québec for mRNA microarray analysis was returned to our laboratory, and in conjunction with previously collected RNA (from an assessment BeWo trophoblast syncytialization following NEFA exposure [56]) was used for validation of DEGs. In brief, 2 μ g of RNA was reverse transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems; ThermoFisher Scientific). RT-qPCR was subsequently performed using sample cDNA and the CFX384 Real Time System (Bio-Rad, Mississauga, Canada). Relative expression of genes of interest was then determined using the $\Delta\Delta C_t$ method, with the geometric mean of PSMB6 and GAPDH utilized as a reference. The expression of genes of interest were analyzed in each NEFA treatment in binary versus BSA control samples via paired two-tailed T-Test. The sequences of RT-qPCR primers utilized for microarray validation are available in **Supplementary Table 1**.

2.6. Cell collections and lysis for Untargeted Metabolomic and Lipidomic Profiling of BeWo Cytotrophoblast cells.

BeWo trophoblasts were plated at a density of 2×10^6 cells/plate in 150 mm cell culture dishes and cultured with exogenous NEFAs as described above. AT T72H cell media was aspirated and cells were washed three times with cold PBS. Cellular metabolism was then quenched with ice-cold methanol, and cells were scrapped and collected in microcentrifuge tubes. Samples were then dried first under a gentle stream of N₂ gas and residual moisture was removed via lyophilization. Samples were then shipped to The Metabolomics Innovation Centre (TMIC, Edmonton, Canada) for untargeted metabolomic and lipidomic profiling of NEFA-treated BeWo CT cells. The dried cell pellets were reconstituted in 400 μ L 1:1 of (v/v) LC-MS grade methanol:water, and cells were lysed via five freeze-thaw cycles. Tubes were then centrifuged at 16,000g for 10 mins at 4°C. A 300 μ L aliquot of resulting supernatant was transferred to a fresh tube for metabolomic profiling, while the remaining supernatant and cell pellets were utilized for lipidomic profiling.

2.7. Untargeted Metabolomic Profiling of BeWo Cytotrophoblast Cells

Cellular metabolome profiles were analyzed using a High-Performance Chemical Isotope Labeling (HP CIL) liquid chromatography coupled mass spectrometry (LC-MS) approach, as previously described [66,71,72]. In brief, the 300 μ L aliquot of supernatant was dried down, and redissolved in water. A proprietary metabolome quantification kit from Nova Medical Testing Inc. (Product Number: NMT-6001-KT) was used to measure the total metabolite concentration in each sample. Different volumes of sample were taken and then dried down to adjust samples to the same concentration of 2 mM. The samples were then chemically labelled for amine-/phenol-, hydroxyl-, carboxyl, and carbonyl- submetabolomic profiling with dansylation reaction, base-activated dansylation reaction, DMPA bromide labeling, and dansylhydrazine labeling, respectively [73]. During labeling, individual samples were labeled with ^{12}C -reagents and a pooled sample, which was generated by mixing an aliquot from each individual sample, was labeled with ^{13}C -reagents. The ^{12}C -labelled samples were then mixed with a ^{13}C -labelled reference in a 1:1 ratio. LC-MS was subsequently utilized for metabolome analysis using the Agilent 1290 LC (with an Agilent eclipse plus reversed-phase C18 column (150 \times 2.1 mm, 1.8 μ m particle size) (Agilent Technologies) linked to the Bruker Impact II QTOF Mass Spectrometer (Bruker Corporation, Billerica, USA).

Metabolomics data was then imported into IsoMS pro v1.2.7 for data processing and analysis. Metabolites were identified in a three-tier approach with high confidence identifications in tiers 1 and 2, and putative matches in tier 3, as previously reported [73]. Metabolites with $\geq \pm 1.2$ FC and raw p-value < 0.05 in each NEFA treatment versus BSA control were determined to be differentially abundant. Metabolite peak-pairs identified in tiers 1 and 2 that had an associated KEGG pathway ID number were then imported into MetaboAnalyst v5.0 software for pathway enrichment analysis (using the homo sapiens KEGG Library) to determine biological impacts of differentially abundant metabolites. KEGG pathways with raw p-value < 0.05 were determined to be significantly enriched.

2.8. Integration of BeWo Cytotrophoblast Transcriptomic and Metabolomic Profiles

Peak Pair data of metabolites in tiers 1 and 2 that have an associated KEGG ID number were subsequently imported into MetaboAnalyst v5.0, and metabolites with a $\geq \pm 1.2$ FC and raw p-value < 0.05 in each NEFA treatment were identified. These differentially abundant metabolites were subsequently integrated with the list of DEGs identified in the Clariom S mRNA microarray using the Joint Pathway Analysis Tool in MetaboAnalyst v5.0 to determine enriched pathways containing both DEGs and differentially abundant metabolites.

2.9. Untargeted Lipidomic Profiling of BeWo Cytotrophoblast Cells

For lipidomic profiling, the remaining supernatant and cell pellets were processed using a modified Folch liquid-liquid lipid extraction protocol. In brief, samples were vortexed with 5 μ L Splash Lipidomix Mass Spec Standard (Avanti Polar Lipids) and methanol for 20s, and subsequently vortexed for an additionally 20s following the addition of dichloromethane. Samples were then incubated at room temperature for 10 mins before being centrifuged at 16,000g for 10 mins at 4°C. An aliquot of the resulting organic layer was then evaporated to dryness with a SpeedVac. The resulting dried material was resuspended in 15 μ L of mobile phase B (MPB, 10 mM ammonium formate in 95% isopropanol), vortexed for 1 min, and then diluted with 135 μ L of mobile phase A (MPA, methanol:acetonitrile:water 50:40:10 v/v/v). A pooled mixture of all extracts was then prepared for quality control samples that were injected into the mass spectrometer every 7 runs to monitor instrument performance. LC-MS analysis of the extracts was performed in both positive and negative ion modes using a ThermoFisher Dionex UltiMate 3000 UHPLC machine (ThermoFisher Scientific) with a Waters BEH C18 Column (Waters Corporation; 5 cm \times 2.1 mm with 1.7 μ m particles) coupled with a Bruker Impact II QTOF Mass Spectrometer (Bruker Corporation). Each sample was analyzed in technical duplicate. The chromatograms for the positive and negative ionization injections for each experimental replicate were aligned and were combined into a single feature-intensity table using Metaboscope v4.0 (Bruker Corporation) [74,75].

A three-tier approach was utilized to identify the detected MS/MS lipid-species. First, lipids were identified based on MS/MS spectral matches by searching the MS Dial LipidBlast database [76], the Human Metabolome database [77], and the MassBank of North America [78,79] LC-MS/MS libraries with Metaboscape v4.0, while lipid species in tier 3 were MS identified with the LipidMaps database. Compounds with MS/MS match score ≥ 500 ; precursor m/z error ≤ 5 mDa, and isotope pattern match (mSigma) score ≤ 50 were deemed Tier 1 (high-confidence) identifications, while MS/MS match scores between 100 and 500 were considered Tier 2 identifications. Second, the unidentified lipid species were mass-matched to the LipidMaps database (m/z error ≤ 5 mDa) for Tier 3. Since lipids can have many isomers and isobars within tight m/z tolerances, we employed a scoring approach to select the best identification candidate based on the expected performance of each lipid within our analytical platform (predicted retention time ranges, adduct form, predicted ionization efficiency and sensitivity, odd or even number of carbons in fatty acyl groups, and presence of unexpected modifications) [75].

The detected intensities for identified compounds were normalized to class-matches internal standards, i.e., a mixture of 14 deuterated lipids that represent the known lipidome (Splash Lipidomix Mass Spec Standard, Avanti Polar Lipids). Unidentified features could not be matched to the most similar internal standard; hence, they were not employed for statistics. The normalized feature intensities (detected peak intensity / intensity of the most similar internal standard) were then imported to MetaboAnalyst v5.0 for statistical analysis. Non-informative features (i.e., near-constant features in samples based on RSD calculation) and features with low repeatability (RSD $>30\%$ in QC samples) were filtered out. The dataset was further normalized by the median intensity ratio within each experiment and auto-scaled. For univariate statistics using Volcano plots, lipid species with $\geq \pm 1.5$ FC for NEFA treatment/BSA control, and raw p-value < 0.05 were considered differentially abundant.

The normalized peak intensities of lipid species from each lipid class identified in tiers 1 and 2 were subsequently added together to determine the total abundance of each lipid class in the NEFA treatments. These summed data were imported into MetaboAnalyst v5.0 and the total lipid class intensities of each NEFA condition were compared in to the BSA control in binary using a Wilcoxon rank-sum test to determine differentially abundant lipid classes in each NEFA treatment. Significance was determined using a Bonferroni-corrected cut-off of $p < 0.0167$.

3. Results

3.1. NEFA-treatments impact fatty acid profiles of BeWo trophoblasts

A total of 10 saturated fatty acids (SFA) species, 6 mono-unsaturated fatty acid (MUFA) species, and 9 polyunsaturated fatty acid (PUFA) species were detected in the BeWo trophoblast samples. Palmitic acid (16:0), stearic acid (18:0), Oleic acid (18:1n9c) and arachidonic acid (20:4n6) were highlighted as the most abundant fatty acid species in BeWo trophoblast cultures (**Table 1**, n=4-5/group).

Syncytialization of BeWo trophoblast cells was associated with significantly reduced relative abundances of 20:1n9 and 20:2, as well as significantly increased relative abundances of 18:2n6 and 20:4n6 compared to undifferentiated BeWo cells (**Table 1**; $p < 0.05$, n=4-5/group).

PA-treated BeWo cultures displayed significantly increased relative abundances of total SFA species, 16:0, 16:1n7, 22:2, and 20:5n3 as well as significantly decreased relative abundances of total MUFA species, 18:0, and 18:1n9c in both CT and SCT cultures relative respective differentiation state BSA-control (**Table 1**; $p < 0.05$, n=4-5/group). PA-treated BeWo CT cells further displayed a significantly increased relative abundance of 22:5n3, while PA-treated BeWo SCT cells displayed significantly reduced relative abundances of 18:1n7 and 18:2n6 versus respective BSA-control cultures (**Table 1**; $p < 0.05$, n=4-5/group).

Table 1. Fatty acid profiles (% mol) of NEFA-treated BeWo trophoblast cells.

Fatty Acid Species	CT				SCT				Differentiation State Difference
	BSA Ctrl	PA	OA	P/O	BSA Ctrl	PA	OA	P/O	
ΣSFA	39.64 ± 1.1 <i>a</i>	44.91 ± 1.63 <i>b</i>	20.73 ± 1.04 <i>c</i>	33.28 ± 2.99 <i>d</i>	37.20 ± 0.92 <i>a</i>	44.67 ± 0.85 <i>b</i>	24.32 ± 3.18 <i>c</i>	34.25 ± 3.29 <i>a</i>	NS
8:0	0.63 ± 0.23	0.32 ± 0.11	0.40 ± 0.16	0.54 ± 0.10	0.49 ± 0.14	0.34 ± 0.12	0.45 ± 0.11	0.45 ± 0.06	NS
10:0	0.40 ± 0.14	0.31 ± 0.09	0.37 ± 0.11	0.37 ± 0.11	0.21 ± 0.07	0.24 ± 0.05	0.30 ± 0.07	0.21 ± 0.05	NS
14:0	1.43 ± 0.35 <i>a</i>	0.85 ± 0.31 <i>ab</i>	0.52 ± 0.12 <i>b</i>	0.74 ± 0.25 <i>b</i>	1.03 ± 0.33	0.68 ± 0.24	0.40 ± 0.08	0.58 ± 0.18	NS
15:0	0.54 ± 0.16	0.51 ± 0.19	0.19 ± 0.03	0.34 ± 0.14	0.45 ± 0.10	0.59 ± 0.19	0.28 ± 0.02	0.48 ± 0.14	NS
16:0	19.73 ± 0.81 <i>a</i>	27.84 ± 2.12 <i>b</i>	8.27 ± 0.81 <i>c</i>	17.52 ± 2.87 <i>a</i>	16.32 ± 0.78 <i>ab</i>	28.07 ± 1.37 <i>c</i>	12.17 ± 3.01 <i>a</i>	19.74 ± 2.80 <i>b</i>	NS
18:0	14.98 ± 0.48 <i>a</i>	12.79 ± 0.76 <i>b</i>	9.23 ± 0.41 <i>c</i>	12.04 ± 0.79 <i>b</i>	16.20 ± 0.36 <i>a</i>	12.46 ± 0.89 <i>b</i>	9.26 ± 0.41 <i>c</i>	10.98 ± 0.43 <i>bc</i>	NS
20:0	0.78 ± 0.06	0.94 ± 0.09	0.79 ± 0.16	0.74 ± 0.07	1.21 ± 0.17	1.10 ± 0.24	0.73 ± 0.10	0.92 ± 0.14	NS
22:0	0.68 ± 0.11	0.80 ± 0.16	0.47 ± 0.10	0.53 ± 0.06	0.76 ± 0.11 <i>ab</i>	0.91 ± 0.08 <i>a</i>	0.47 ± 0.07 <i>b</i>	0.74 ± 0.08 <i>ab</i>	NS
23:0	0.49 ± 0.14	0.58 ± 0.11	0.52 ± 0.02	0.64 ± 0.05	0.23 ± 0.03	0.44 ± 0.12	0.27 ± 0.03	0.41 ± 0.09	NS
24:0	1.02 ± 0.05 <i>a</i>	1.08 ± 0.07 <i>a</i>	0.70 ± 0.06 <i>b</i>	0.79 ± 0.07 <i>b</i>	0.98 ± 0.05 <i>a</i>	0.89 ± 0.05 <i>a</i>	0.54 ± 0.02 <i>b</i>	0.63 ± 0.05 <i>b</i>	NS
ΣMUFA	42.82 ± 1.14 <i>a</i>	36.29 ± 1.21 <i>b</i>	68.11 ± 1.26 <i>c</i>	49.99 ± 4.19 <i>d</i>	42.91 ± 1.15 <i>a</i>	34.91 ± 0.43 <i>b</i>	63.40 ± 4.28 <i>c</i>	49.35 ± 4.35 <i>d</i>	NS
16:1n7	3.12 ± 0.18 <i>a</i>	5.23 ± 0.67 <i>b</i>	0.80 ± 0.12 <i>c</i>	1.87 ± 0.42 <i>c</i>	2.86 ± 0.33 <i>a</i>	6.20 ± 0.46 <i>b</i>	1.43 ± 0.41 <i>c</i>	2.42 ± 0.39 <i>ab</i>	NS
18:1n9c	33.09 ± 1.32 <i>a</i>	24.76 ± 1.59 <i>b</i>	63.73 ± 1.47 <i>c</i>	43.84 ± 4.64 <i>d</i>	32.65 ± 1.29 <i>a</i>	23.08 ± 0.54 <i>b</i>	62.64 ± 0.68 <i>c</i>	38.52 ± 0.83 <i>ab</i>	NS
18:1n9t	0.23 ± 0.02	0.34 ± 0.08	0.38 ± 0.07	0.23 ± 0.04	0.28 ± 0.03	0.46 ± 0.23	0.33 ± 0.07	0.41 ± 0.19	NS
18:1n7	5.99 ± 0.26 <i>a</i>	5.69 ± 0.23 <i>a</i>	2.91 ± 0.17 <i>b</i>	3.78 ± 0.31 <i>c</i>	6.86 ± 0.23 <i>a</i>	4.89 ± 0.20 <i>b</i>	3.03 ± 0.16 <i>c</i>	3.39 ± 0.18 <i>c</i>	NS
20:1n9	0.49 ± 0.02 <i>a</i>	0.45 ± 0.02 <i>a</i>	1.46 ± 0.08 <i>b</i>	0.92 ± 0.14 <i>c</i>	0.48 ± 0.04 <i>a</i>	0.37 ± 0.02 <i>a</i>	0.99 ± 0.10 <i>b</i>	0.77 ± 0.17 <i>c</i>	*
22:1n9	0.29 ± 0.13	0.21 ± 0.08	0.19 ± 0.01	0.20 ± 0.03	0.16 ± 0.03	0.24 ± 0.09	0.18 ± 0.02	0.20 ± 0.06	NS
ΣPUFA	16.30 ± 0.94 <i>ab</i>	17.47 ± 0.82 <i>a</i>	9.18 ± 0.44 <i>c</i>	15.02 ± 1.79 <i>b</i>	18.89 ± 0.78 <i>a</i>	18.96 ± 0.42 <i>a</i>	10.68 ± 1.24 <i>b</i>	14.84 ± 1.27 <i>b</i>	NS
16:3n4	0.34 ± 0.04 <i>ab</i>	0.23 ± 0.02 <i>ab</i>	0.16 ± 0.02 <i>b</i>	0.22 ± 0.03 <i>b</i>	0.37 ± 0.04 <i>a</i>	0.30 ± 0.05 <i>a</i>	0.19 ± 0.02 <i>b</i>	0.28 ± 0.03 <i>ab</i>	NS
18:2n6	2.52 ± 1.21 <i>ab</i>	1.36 ± 0.24 <i>a</i>	0.61 ± 0.04 <i>b</i>	0.82 ± 0.08 <i>b</i>	3.24 ± 0.98 <i>a</i>	1.45 ± 0.04 <i>b</i>	0.94 ± 0.04 <i>c</i>	1.11 ± 0.07 <i>c</i>	*
18:3n6	0.230 ± 0.005	0.234 ± 0.029	0.194 ± 0.006	0.226 ± 0.022	0.324 ± 0.016 <i>a</i>	0.265 ± 0.024 <i>ab</i>	0.201 ± 0.009 <i>c</i>	0.235 ± 0.011 <i>bc</i>	NS
20:2	1.48 ± 0.12 <i>ab</i>	3.03 ± 0.24 <i>b</i>	1.69 ± 0.11 <i>a</i>	3.23 ± 0.48 <i>b</i>	0.97 ± 0.15 <i>a</i>	2.00 ± 0.18 <i>b</i>	1.23 ± 0.27 <i>a</i>	1.83 ± 0.23 <i>b</i>	*
20:3n6	1.45 ± 0.07 <i>ab</i>	1.43 ± 0.08 <i>a</i>	0.72 ± 0.05 <i>b</i>	1.09 ± 0.12 <i>c</i>	1.66 ± 0.06 <i>a</i>	1.46 ± 0.05 <i>a</i>	0.76 ± 0.08 <i>b</i>	1.03 ± 0.08 <i>c</i>	NS
20:4n6	8.68 ± 0.39 <i>ab</i>	9.18 ± 0.50 <i>a</i>	5.08 ± 0.25 <i>b</i>	8.07 ± 0.94 <i>a</i>	11.12 ± 0.28 <i>a</i>	11.34 ± 0.38 <i>a</i>	6.52 ± 0.78 <i>b</i>	8.82 ± 0.85 <i>c</i>	*
20:4n3	0.15 ± 0.06	0.19 ± 0.10	0.07 ± 0.01	0.15 ± 0.04	0.14 ± 0.03	0.28 ± 0.11	0.08 ± 0.02	0.21 ± 0.09	NS
20:5n3	0.54 ± 0.14 <i>ac</i>	0.95 ± 0.14 <i>b</i>	0.16 ± 0.02 <i>a</i>	0.64 ± 0.15 <i>bc</i>	0.39 ± 0.05 <i>a</i>	0.92 ± 0.13 <i>b</i>	0.27 ± 0.06 <i>a</i>	0.63 ± 0.11 <i>ab</i>	NS
22:5n3	0.43 ± 0.05 <i>a</i>	0.52 ± 0.03 <i>b</i>	0.31 ± 0.01 <i>c</i>	0.39 ± 0.04 <i>ac</i>	0.53 ± 0.01 <i>a</i>	0.61 ± 0.02 <i>a</i>	0.33 ± 0.02 <i>b</i>	0.41 ± 0.04 <i>b</i>	NS

Data are mean ± SEM (n=4-5/group). Different lower-case letters denote statistical difference in FA abundance between NEFA treatments within a differentiation state. * denotes a differentiation state dependent difference in FA abundance between BeWo CT and SCT cells; NS indicates no statistical significance between differentiation states.

The OA-treatment in BeWo CT and SCT trophoblasts was associated with significantly increased relative abundance of total MUFA species, 18:1n9c, and 20:1n9 as well as significantly reduced relative abundance of total SFA and PUFA species, 18:0, 24:0, 16:1n7, 18:1n7, 16:3n4, 18:2n6, 20:3n6, 20:4n6, and 22:5n3 versus respective differentiation state BSA-control cultures (**Table 1**; $p < 0.05$, $n = 4-5/\text{group}$). Additionally, OA-treated BeWo CT cells displayed significantly decreased relative abundances of 14:0, and 16:0 while OA-treated BeWo SCT cells displayed a significantly reduced relative abundance of 18:3n6 versus respective differentiation state BSA-control (**Table 1**; $p < 0.05$, $n = 4-5/\text{group}$).

P/O-treated BeWo CT and SCT cells were found to have increased relative abundances of total MUFA species, 20:1n9, and 20:2 as well as decreased relative abundances of 18:0, 24:0, 18:1n7, 18:2n6, and 20:3n6 compared to BSA-control cultures (**Table 1**; $p < 0.05$). P/O-cultured BeWo CT cells further demonstrated an increased relative abundance of 18:1n9c as well as decreased relative abundance of total SFA species, 14:0, 16:1n7, and 16:3n4 versus BSA-control CT cultures (**Table 1**; $p < 0.05$, $n = 4-5/\text{group}$). Additionally, P/O-treated BeWo SCT cells displayed a decreased relative abundance of 18:3n6 compared to BSA-control SCT cultures (**Table 1**; $p < 0.05$, $n = 4-5/\text{group}$).

3.2. The impact of NEFA treatments on desaturation and elongation indices in BeWo trophoblasts

Fatty acid profile data was utilized to quantify the activity of fatty acid desaturase and elongase enzymes via calculation of the ratios of enzyme products to substrates (desaturation/elongation index). OA and P/O-treated BeWo SCT and SCT cells displayed significantly reduced SCD1 activity compared to respective BSA-control cultures when assessed by the 16:1n7/16:0 index (**Figure 1a**; $p < 0.05$, $n = 5/\text{group}$). However, when SCD1 activity was assessed by the ratio of 18:1n9c/18:0, OA and P/O-treated BeWo CT and SCT cells were found to have increased SCD1 activity compared to respective BSA-control cultures (**Figure 1b**; $p < 0.05$, $n = 5/\text{group}$). Furthermore, OA-treated BeWo CT cells displayed increased FADS2 activity (**Figure 1c**; $p < 0.05$, $n = 5/\text{group}$) while OA-treated SCT cells displayed increased FADS2 and FADS1 activity compared to BSA-control cultures (**Figure 1c-d**; $p < 0.05$, $n = 5/\text{group}$). P/O-treated BeWo CT and SCT also displayed increased FADS1 and FADS2 activity compared to BSA-controls (**Figure 1c-d**; $p < 0.05$, $n = 5/\text{group}$). Finally, PA-treated BeWo SCT cells displayed increased FADS2 activity compared to BSA-control SCT cells (**Figure 1c**; $p < 0.05$, $n = 5/\text{group}$). BeWo syncytialization was additionally linked with increased SCD1 activity (when assessed as 16:1n7/16:0), and FADS1 activity as well as decreased FADS2 activity (**Figure 1a,c-d**; $p < 0.05$, $n = 5/\text{group}$).

OA-treatment was additionally associated with increased elongation of 18:1n9, 16:0, and 20:5n3 in BeWo CT cells, while P/O-treated BeWo CT were only observed to have increased 18:1n9 elongation (**Figure 2a-c**). PA-treatment in BeWo SCT cells, however, was linked with reduced 16:0, and 20:5n3 elongation (**Figure 2b,c**). BeWo SCT also displayed an overall reduced elongation of 18:1n9 compared to undifferentiated BeWo CT cells (**Figure 2a**).




Figure 1. Fatty Acid Desaturase Enzyme Activity Indices in NEFA-treated BeWo CT and SCT cells. The abundance of desaturase enzyme substrate and product FAs were quantified via Gas Chromatography-coupled Flame Ionization Detection, and the activity of each desaturase enzyme was quantified via desaturation index, reflecting the ratio of enzyme product to substrate. The Stearoyl-CoA Desaturase-1 (SCD1) index was quantified via the ratios **(A)** 16:1n7/16:0 ratio and **(B)** 18:1n9/18:0; **(C)** the Fatty Acid Desaturase 1 index was quantified via the ratio of 18:3n6/18:2n6; and **(D)** the Fatty Acid Desaturase 2 index was quantified via the ratio of 20:4n6/20:3n6 (n=5/group; different upper-case letters denote statistical significance between NEFA treatments within each BeWo differentiation state, and different lower-case represents statistical significance between differentiation states; Two-Way Randomized Block ANOVA, Tukey’s multiple comparisons Test).




Figure 2. Fatty Acid Elongase Enzyme Activity Indices in NEFA-treated BeWo CT and SCT cells.

The abundance of products and substrates of FA elongase enzymes were quantified via gas chromatography coupled flame ionization detection. Subsequently the ratios of (A) 20:1n9/18:1n9, (B) 18:0/16:0, and (C) 22:5n3/20:5n3 (elongation indices) were quantified to determine elongase enzyme activities (n=5/group; different upper-case letters denote statistical significance between NEFA treatments within each BeWo differentiation state, and different lower-case represents statistical significance between differentiation states; Two-Way Randomized Block ANOVA, Tukey's multiple comparisons).

3.3. OA-treatment alters neutral lipid profiles of BeWo trophoblasts

Subsequent TLC-FID analysis of extracted lipids from BeWo cultures identified five neutral lipid species in all samples. Specifically, the analyses identified cholesterol ester, free fatty acids, triglycerides, free cholesterol, and diacylglycerols in BeWo trophoblast neutral lipid fraction. OA-treated BeWo CT and SCT displayed increased relative triglyceride abundance as well as decreased relative free cholesterol abundance relative to respective differentiation state BSA-control cultures (Table 2; 2WA: NEFA treatment $p < 0.05$, n=5/group).

Table 2. Neutral Lipid profiles of NEFA-treated BeWo trophoblast cells.

Neutral Lipid Species	CT				SCT			
	BSA Ctrl	PA	OA	P/O	BSA Ctrl	PA	OA	P/O
Cholesterol Esters	10.07 ± 2.66	25.46 ± 9.52	11.29 ± 2.28	15.44 ± 4.39	9.03 ± 1.69	17.56 ± 6.19	12.19 ± 4.33	19.18 ± 7.54
Free Fatty Acids	10.46 ± 3.67	11.91 ± 5.12	8.90 ± 3.36	10.05 ± 4.04	11.33 ± 3.69	9.93 ± 2.81	7.81 ± 2.89	8.72 ± 2.55
Triglycerides	15.99 ± 5.14 <i>a</i>	17.12 ± 2.63 <i>a</i>	39.47 ± 2.59 <i>b</i>	18.35 ± 2.88 <i>a</i>	14.34 ± 1.97 <i>a</i>	14.74 ± 1.51 <i>a</i>	30.86 ± 4.95 <i>b</i>	17.96 ± 1.19 <i>a</i>
Free Cholesterol	61.80 ± 6.21 <i>a</i>	43.6 ± 5.91 <i>bc</i>	37.84 ± 1.32 <i>b</i>	53.79 ± 3.88 <i>ac</i>	62.88 ± 3.19 <i>a</i>	56.39 ± 5.57 <i>ab</i>	46.40 ± 4.62 <i>b</i>	52.24 ± 6.66 <i>ab</i>
Diacylglycerols	1.68 ± 0.32	1.91 ± 0.77	2.51 ± 0.35	2.37 ± 0.42	2.43 ± 0.89	1.38 ± 0.24	2.75 ± 0.07	1.90 ± 0.26

Data are percent of neutral lipid fraction (mean ± SEM; n=5/group). Different lower-case letters denote statistical differences between NEFA-treatments within each differentiation state.

3.4. Transcriptomic profiles of BeWo Cytotrophoblasts are impacted by NEFA treatment.

PA-treated BeWo CT cultures displayed 340 differentially expressed genes (DEGs) (140 up-regulated, and 200 down-regulated); OA-treated CT cultures displayed 208 DEGs (88 up-regulated, and 120 down-regulated); and P/O-treated BeWo CT cultures displayed 221 DEGs (96 up-regulated, and 125 down-regulated) when compared to BSA-control CT cultures ($\geq \pm 1.3$ FC vs BSA control CT cultures; raw- $p < 0.05$, $n=5$ /group). A summary of the DEGs identified in each NEFA-treatment condition is available in **Supplementary Table 2**. Volcano plots were then constructed to visualize differentially expressed transcripts in each NEFA-treatment (**Figure 3a-c**).




Figure 3. Volcano plot visualization of differentially expressed genes in BeWo CT cells cultured with 100 μ M NEFAs for 72 hours. The transcriptome profiles of NEFA-treated BeWo cytotrophoblast cells was determined via Clariom S mRNA microarray, and differentially expressed genes were determined with ± 1.3 fold-change and raw $p < 0.05$ cut-offs. Volcano plots were constructed to visualize the differentially expressed genes (DEGs) in NEFA-treated BeWo CT cells. (A) PA-treated cells displayed 340 DEGs (140 up-regulated, and 200 down-regulated); (B) OA-treated cells displayed 208 DEGs (88 up-regulated, and 120 down-regulated); and (C) P/O-treated cells displayed 221 DEGs (96 up-regulated, and 125 down-regulated) compared to BSA-alone treated cells ($n=5$ /group). The x-axis of the volcano plots indicates fold-change vs BSA control, and the y-axis indicates p-value ($-\log_{10}$). Red dots represent up-regulated genes, and blue dots represent down-regulated genes. (D) A Venn-

diagram showing the number of differentially expressed genes that are unique to each NEFA-treatment or common to multiple NEFA-treatments.

A Venn-diagram was additionally constructed to visualize the DEGs that were unique to each NEFA-treatment, as well as those common in two or all three of the NEFA-treatment conditions (**Figure 3d**). Twenty-three differentially expressed transcripts were found to be common between PA and OA-treated CT cells; 32 DEGs were common between PA and P/O treatments; and 9 genes were found to be differentially expressed in both OA and PO-treated cells (**Figure 3d**). A further 9 genes were found to be differentially expressed in all NEFA-treated BeWo CT cells (**Figure 3d**). A summary list of unique and common DEGs is available in **Supplementary Table 3**.

Over-representation analysis revealed no significantly enriched pathways in the DEGs lists in PA, and OA treated BeWo CT cells (**Supplementary Table 4**). In the P/O-treated BeWo CT cells there was a significant enrichment in the Wikipathways “Fatty Acid Biosynthesis” dataset (associated with the DEGs: *ACACB* (+1.32 FC), *ACSL5* (+1.77 FC), *ACSL6* (-1.34 FC), and *SCD* (-1.24 FC)) as well as in the GO “CXCR3 Chemokine Receptor Binding” molecular function dataset (associated with the DEGs: *CXCL10* (+1.39 FC), *CXCL13* (+1.42 FC), and *CXCL11* (-1.37 FC)) (**Supplementary Table 4**, FDR- $p < 0.05$).

Subsequent enrichment analysis using a GSEA approach highlighted a significant downregulation of the Reactome “Adherens Junctions Interactions” pathway (associated with the DEGs: *CDH17* (-1.48 FC), and *CDH9* (-1.36 FC)) in PA-treated BeWo CT cells (**Supplementary Table 5**, FDR- $p < 0.25$). In PA-treated BeWo CT cells, a significant upregulation was also observed in the Reactome “Peptide Hormone Biosynthesis” pathway (no associated DEGs), as well as the KEGG datasets for: “Staphylococcus Aureus Infection” (associated with the DEG: *ITGB2* (+1.4 FC)); “PPAR Signaling Pathway” (associated with the DEGs: *ACSL5* (+1.77 FC), *APOA2* (+1.51 FC), *FABP2* (+1.31 FC), *FABP6* (+1.36 FC), *HMGCS2* (+1.34 FC), and *PLIN2* (+1.35 FC)); “Th1 and Th2 Cell Differentiation” (associated with the DEGs: *IKBK* (+1.32 FC), and *IL12A* (+1.35 FC)); “Fatty Acid Biosynthesis” (associated with the DEGs: *ACSL5* (+1.77 FC), and *ACACB* (+1.57 FC)); “Leishmaniasis” (associated with the DEGs: *ITGB2* (+1.4 FC), *NCF1* (+1.35 FC), and *IL12A* (+1.35 FC)); “Cortisol Synthesis and Secretion” (associated with the DEGs: *CREB3L3* (+2.31 FC), *CYP21A2* (+1.31 FC), and *HSD3B1* (+1.34 FC)); “Ovarian Steroidogenesis” (associated with the DEGs: *CYP11A1* (+1.37 FC), and *HSD3B1* (+1.34 FC)); “Adipocytokine Signaling Pathway” (associated with the DEGs: *ASCL5* (+1.77 FC), *ACACB* (+1.57 FC), and *IKBK* (+1.32 FC)); and “Primary Bile Acid Biosynthesis” (no associated DEGs) (**Supplementary Table 5**, FDR- $p < 0.25$).

In the OA-treated BeWo CT cells, there was a significant upregulation in the Wikipathways “PPAR Signaling Pathway” dataset (associated with the DEG: *ACSL5* (+1.87 FC)) and in the KEGG “PPAR Signaling Pathway” dataset (Associated with the DEGs: *ACSL5* (+1.87 FC), and *PLIN2* (+1.46 FC)) (**Supplementary Table 5**, FDR- $p < 0.25$). A significant downregulation was also observed in KEGG “Linoleic Acid Metabolism” dataset (no associated DEGs) in OA-treated BeWo CT cells (**Supplementary Table 5**, FDR- $p < 0.25$).

In the P/O-treated cells a significant upregulation was observed in the Reactome “Gene and protein expression by JAK-STAT signaling after Interleukin-12 stimulation” pathways (no associated DEGs), as well as in the GO “Amine Transport” biological processes dataset (no associated DEGs) (**Supplementary Table 5**, FDR- $p < 0.25$). Summary Ordered lists of the genes identified in the microarray panels, and their respective signal-to-noise ratio scores is available in **Supplementary Table 6**.

3.5. RT-qPCR Validation of Differentially Expressed Genes

In PA-treated BeWo CT cells, RT-qPCR analysis confirmed differential expression of *ACACB* (+1.57 FC microarray; +1.34 FC RT-qPCR); *ACADVL* (+1.32 FC microarray; +1.47 FC RT-qPCR); *ACSL5* (+1.77 FC microarray; +2.57 FC RT-qPCR); *CREB3L3* (+2.31 FC microarray; +5.29 FC RT-qPCR); *PLIN2* (+1.35 FC microarray; +1.83 FC RT-qPCR) (**Figure 4a**, $p < 0.05$, $n=9$). It is important to note that *SCD* mRNA abundance although statistically significant ($p < 0.05$) in the microarray panel did not meet the ± 1.3 FC cut-off to be considered differentially expressed in PA-treated CT cells (-1.16 FC).

Subsequent RT-qPCR analysis likewise highlighted a significant decrease in *SCD* mRNA abundance in PA-treated BeWo CT cells (-1.24 FC; Fig 4a, $p < 0.05$, $n = 9$). The microarray panel also highlighted a significant downregulation in a transcript best identified as *ACACA*, however RT-qPCR analysis demonstrated no significant alterations in *ACACA* mRNA abundance (Figure 4a, $p = 0.057$, $n = 9$). Finally, *AQP3* was highlighted as differentially expressed in the microarray panel, however there was no significant difference in *AQP3* mRNA abundance when assessed by RT-qPCR (Figure 4a, $p = 0.4287$, $n = 9$).




Figure 4. RT-qPCR validation of differentially expressed genes in NEFA-treated BeWo CT cells.

Differential expression of genes involved in lipid metabolic processes were validated in (A) PA, (B) OA, and (C) P/O-treated BeWo CT cells via RT-qPCR. RT-qPCR data were analyzed via paired two-tailed t-test ($n = 9$ /group), and the data expressed as fold-change vs BSA-control cultures (mean \pm SEM). *indicates $\geq \pm 1.3$ FC vs BSA-control, $p < 0.05$ for microarray data; and $p < 0.05$ for RT-qPCR data. #indicates < 1.3 FC vs BSA-control, $p < 0.05$ for microarray data.

In OA-treated BeWo CT cells, RT-qPCR analysis confirmed the differential expression of: *ACSL5* (+1.87 FC microarray; +2.60 FC RT-qPCR); *CREB3L3* (+2.48 FC microarray; +7.38 FC RT-qPCR); *PLIN2* (+1.46 FC microarray; +1.90 FC RT-qPCR); and *SCD* (-1.41 FC microarray; -1.98 FC RT-qPCR) (Figure 4b, $p < 0.05$, $n = 9$). The microarray panel also highlighted a significant increase in *ACADVL* mRNA abundance although this transcript did not meet the ± 1.3 FC cut-off to be considered differentially

expressed. However, it is important to note that RT-qPCR analysis also highlighted a significant increase in *ACADVL* mRNA abundance in OA-treated BeWo CT cells (+1.43 FC; **Figure 4b**, $p < 0.05$, $n = 9$).

In P/O-treated BeWo CT cells, RT-qPCR analysis confirmed the differential expression of *ACACB* (+1.32 FC microarray; +1.12 FC RT-qPCR); *ACSL5* (+1.77 FC microarray; +2.50 FC RT-qPCR); *CREB3L3* (+2.13 FC microarray; +6.23 FC RT-qPCR); *PLIN2* (+1.30 FC microarray; +1.79 FC RT-qPCR); and *SCD* (-1.34 FC microarray; -1.52 FC RT-qPCR) (**Figure 4c**; $p < 0.05$, $n = 9$ /group). The RT-qPCR analysis additionally highlighted a significant increase in the expression of *ACADVL* (+1.48 FC) and a decrease in the expression of *ACACA* (-1.32 FC) (**Figure 4c**, $p < 0.05$, $n = 9$ /group). Interestingly, the expression of both *ACADVL* (+1.22 FC, $p = 0.0609$) and *ACACA* (-1.11 FC, $p = 0.0862$) were trending towards significance in the microarray panel.

3.6. Metabolomic profiles of BeWo Cytotrophoblast in response to NEFA treatment

On average, 6569 ± 50 (mean \pm SD) metabolite features were identified in each NEFA-treated BeWo CT samples. A summary of the identified features, and metabolite peak-pair data is available in **Supplementary Table 7**. Of these features, 179 metabolites were identified with high confidence in tier 1, and a further 602 metabolites were identified with high confidence in tier 2. In PA-treated cultures, 2 metabolites in tier 1 (both decreased); 7 metabolites in tier 2 (4 increased, and 3 decreased); and 97 metabolites in tier 3 (28 increased, and 69 decreased) were found to be differentially abundant when compared to BSA-control samples ($\geq \pm 1.2$ FC, raw- $p < 0.05$, $n = 5$ /group). Furthermore, 5 metabolites identified in tier 1 (2 increased, and 3 decreased); 12 metabolites from tier 2 (2 increased, and 10 decreased); and 78 metabolites in tier 3 (51 increased, and 27 decreased) were found to be differentially abundant in OA-treated CT cells when compared to BSA-control samples ($\geq \pm 1.2$ FC, raw- $p < 0.05$, $n = 5$ /group). Finally, 7 metabolites identified in tier 1 (4 increased, and 3 decreased); 7 metabolites identified in tier 2 (4 increased, and 3 decreased); and 92 metabolites identified in tier 3 (48 increased, and 44 decreased) were found to be differentially abundant in P/O-treated BeWo CT cultures when compared to BSA-control samples ($\geq \pm 1.2$ FC, raw- $p < 0.05$, $n = 5$ /group). Volcano plots were constructed to visualize the differentially abundant metabolite species in each NEFA-treatment group versus BSA-control samples (**Figure 5a-c**). An unsupervised principal component analysis (PCA) 2D plot as well as supervised partial-least squares discriminant analysis (PLS-DA) 2D plot were constructed to visualize the degree of difference between metabolite profiles in all NEFA-treatment groups (**Supplementary Figure 1**). Summary lists of the differentially abundant metabolites identified in each NEFA condition are available in **Supplementary Table 8**.

A Venn-diagram was constructed to visualize the number of unique and shared differentially abundant metabolites in the NEFA treatments which highlighted that most differentially abundant metabolites were unique to one NEFA treatment (**Figure 5d**). Only 3 high confidence identified metabolites (from tiers 1 and 2) were found to be differentially abundant in at least 2 of the NEFA treatments. Specifically, the abundance of Kahweol was found to be significantly increased (+1.69 FC in OA; +1.64 FC in P/O); the abundance of N-acetyethanolamine was found to be significantly decreased in both the OA and P/O treatments (-1.66 FC in OA; -1.46 FC in P/O); and the abundance of 3-hydroxymethylglutaric acid was found to be significantly decreased in both the PA and P/O treatments (-2.89 FC in PA; -2.00 FC in P/O). Summary lists of the differentially abundant metabolites common between 2 or more NEFA-treatment groups is available in **Supplementary Table 9**.

Analysis of KEGG pathway enrichment of the metabolite sets from tiers 1 and 2 was subsequently performed using MetaboAnalyst software. Scatterplots were then constructed to visualize the KEGG pathway nodes highlighted by each metabolite set (**Figure 6a-c**). No KEGG pathways were identified as significantly enriched in the metabolite sets from PA and P/O-treated BeWo CT cells (**Figure 6a,c**). The KEGG pathways for i) Ether lipid metabolism (highlighted by increased sn-glycerol-3-phosphoethanolamine (+3.44 FC) levels), ii) Amino sugar and nucleotide sugar metabolism (highlighted by reduced 6-deoxy-L-galactose (-1.17 FC) levels); iii) Fructose and mannose metabolism (highlighted by reduced 6-deoxy-L-galactose (-1.17 FC) levels); and iv) Taurine and hypotaurine metabolism (highlighted by a trend towards reduced abundance of 5-L-Glutamyl taurine (-1.82 FC);

$p=0.062$) and 3-sulfinol-L-alanine (-1.47 FC; $p=0.054$) were determined to be significantly enriched in the OA-treated CT metabolite set (raw- $p<0.05$; **Figure 6b**).




Figure 5. Differentially abundant metabolites in NEFA-treated BeWo CT cultures. Volcano plots were constructed to visualize differentially abundant metabolites (DAMs) in NEFA treated BeWo CT cells. **(A)** PA-treated BeWo CT cells displayed 106 DAMs (32 increased, and 74 decreased); **(B)** OA-treated cells displayed 95 DAMs (55 increased, and 40 decreased); and **(C)** P/O-treated BeWo CT cultures displayed 106 DAMs (56 increased, and 50 decreased) compared to BSA-control cultures ($\geq \pm 1.2$ fold-change, raw- $p<0.05$). The x-axis of the volcano plots represents $\log_2(\text{fold-change})$ vs BSA-control and the y-axis represents p-value ($-\log_{10}$). Red dots represent individual metabolites with significantly increased abundance, and blue dots represent metabolites with significantly reduced abundance. **(D)** A Venn-diagram showing the number of differentially abundant metabolites that were unique to each NEFA-treatment or common to multiple NEFA-treatments.




Figure 6. Pathway Analysis of metabolite profiles of NEFA-treated BeWo CT cells. Metabolite peak pairs identified in tier 1 and tier 2 were imported into MetaboAnalyst v5.0 for analysis of enriched KEGG pathways. Pathways with raw- $p < 0.05$ were considered significantly enriched and scatterplots were created for (A) PA-treated; (B) OA-treated; and (C) P/O-treated cultures to visualize identified KEGG pathways. The x-axis of the scatterplot represents pathway impact and y-axis represents pathway enrichment p-value ($-\log_{10}$). Each individual node represents a unique KEGG pathway, and node size corresponds to pathway impact, while node color corresponds to the p-value of the pathway. The KEGG pathways for i) Ether lipid metabolism; ii) Amino sugar and nucleotide sugar metabolism; iii) Fructose and mannose metabolism; and iv) Taurine and hypotaurine metabolism were highlighted to be significantly enriched in the OA-treated BeWo CT cultures.

3.7. BeWo Cytotrophoblast transcriptome and metabolome integration

Integration of the transcriptomic and metabolomic datasets revealed no significantly enriched pathways in the KEGG database containing both differential genes and differential metabolites in the PA and P/O-treated BeWo CT cells (Figure 7a,c). The integration analysis, did however, highlight significantly enriched pathways only containing DEGs in the PA and P/O treatments. Specifically, there was a significant enrichment in the i) Inositol Phosphate Metabolism and ii) Phosphatidylinositol Signaling System KEGG pathways (both highlighted by downregulation of IMPA1 (-1.34 FC) and PIP5K1B (-1.37 FC) as well as upregulation of PLCE1 (1.50 FC) and INPP5D (1.447 FC)) in the PA-treated BeWo CT cells, as well as a significant enrichment in the i) Fatty Acid Biosynthesis KEGG Pathway (highlighted by upregulation of ACACB (+1.32 FC) and ACSL5 (+1.77 FC) and downregulation of ACSL6 -1.34 FC)) in the P/O-treated BeWo CT cells (raw- $p < 0.05$; Figure 7a,c).




Figure 7. Joint Pathway Analysis of Transcriptome and Metabolome Profiles of NEFA-treated BeWo CT cells. Differentially expressed gene and differentially abundant metabolite (from identification tiers 1 and 2) datasets were analyzed via the Joint Pathway Analysis module in MetaboAnalyst Software. Pathways with raw- $p < 0.05$ were considered significantly enriched, and scatterplots were constructed to visualize identified KEGG pathways. The x-axis of the scatterplot represents pathway impact and y-axis represents pathway enrichment p-value ($-\log_{10}$). Each individual node represents a unique KEGG pathway, and node size corresponds to pathway impact, while node color corresponds to the p-value of the pathway. **(A)** The i) inositol phosphate metabolism and ii) phosphatidylinositol signaling system KEGG pathways were highlighted as significantly enriched in PA-treated BeWo CT cells. **(B)** The i) ether lipid metabolism and ii) purine metabolism KEGG pathways were highlighted as significantly enriched in the OA-treated BeWo CT cells. **(C)** The i) fatty acid biosynthesis KEGG pathway was highlighted as significantly enriched in P/O-treated BeWo CT cells.

Integration analysis of the OA-treated BeWo CT cell transcriptome and metabolome profiles highlighted significantly enriched pathways containing both differential genes and metabolites. Specifically, the i) Ether Lipid Metabolism pathway (highlighted by increased accumulation of sn-glycero-3-phosphoethanolamine (+3.44 FC) in conjunction with upregulation of GPD1 (+1.34 FC) and downregulation of PLA2G12A (-1.35 FC) and ii) Purine Metabolism pathway (highlighted by increased levels of adenosine (+3.30 FC), inosine (+2.45 FC), and guanosine (+1.69 FC), in conjunction with upregulation of PRPS1 (+1.31 FC), and ADK (+1.31 FC), as well as downregulation of GMPR (-

1.76 FC) were found to be significantly enriched in OA-treated BeWo CT cells (raw- $p < 0.05$; **Figure 7b**).

3.8. The impact of NEFA-treatment of BeWo cytotrophoblast lipidome profiles

An average of 9199 ± 295 (mean \pm SD) lipid species were identified in each sample. Of these features, 950 were positively identified in tier 1, 161 were positively identified in tier 2, and a further 6808 features were putatively identified in tier 3. A summary of the identified features is available in **Supplementary Table 10**, and a list of lipid class abbreviations is available in **Supplementary Table 11**.

Non-parametric volcano plots were then constructed to visualize differentially abundant lipid species in each NEFA-treatment compared to BSA-control samples (**Figure 8**; FC (NEFA-treatment / BSA control $\geq \pm 1.5$; raw- $p < 0.05$). In PA-treated samples, 190 lipid species in tier 1 (80 decreased and 110 increased); 28 lipid species in tier 2 (12 increased, and 16 decreased); and 1137 lipid species in tier 3 (553 increased, and 584 decreased) were found to be significantly altered (**Figure 8a**). In OA-treated samples, 197 lipid species in tier 1 (73 increased, and 124 decreased); 30 lipid species in tier 2 (15 increased; and 15 decreased); and 1406 lipid species in tier 3 (711 increased and 695 decreased) were found to be significantly altered (**Figure 8b**). In P/O-treated BeWo CT cells, 163 lipid species in tier 1 (55 increased, and 108 decreased); 24 lipid species in tier 2 (9 increased, and 15 decreased); and 1092 lipid species in tier 3 (575 increased; and 517 decreased) were found to be significantly altered (**Figure 8c**). A summary of the altered lipid species in each NEFA condition is available in **Supplementary Table 12**.

A Venn-diagram was constructed to visualize the number of differentially abundant lipid species (DALs) that were unique to a single NEFA-treatment and shared between multiple NEFA treatments (**Figure 8d**). In total, 377 DALs were unique to PA-treated BeWo CT cells, 676 DALs were unique to OA-treated BeWo CT cells, and 100 DALs were unique to P/O-treated BeWo CT cells. Additionally, 86 lipid species were altered in both PA and OA-treated cells, 308 lipid species were altered in both PA and P/O-treated cells, and 287 lipid species were altered in both OA and P/O-treated BeWo CT cells. Finally, it was found that 584 lipid species were differentially abundant in all NEFA treatment groups. A summary of the lipid species unique to a single NEFA condition or shared between two or more groups is available in **Supplementary Table 13**.





Figure 8. Volcano Plot visualization of differentially abundant lipid species in NEFA-treated BeWo CT cells. Differentially abundant lipid species (DALs) were visualized via non-parametric volcano plot. The x-axis of the volcano plot represents $\log_2(\text{fold-change})$, and the y-axis represents p-value ($-\log_{10}$). Lipid species with significantly increased abundance vs BSA-control cultures are represented by red dots, and lipid species with significantly decreased abundance are represented by blue dots ($\geq \pm 1.5$ fold-change, raw p-value < 0.05). **(A)** PA-treated BeWo CT cells displayed 1355 differentially abundant lipids (676 increased, and 679 decreased); **(B)** OA-treated BeWo CT cells displayed 1633 differentially abundant lipids (799 increased, and 834 decreased); and **(C)** P/O-treated BeWo CT cells displayed 1279 differentially abundant lipids (639 increased, and 640 decreased). **(D)** A Venn-diagram showing the number of DALs unique to each NEFA-treatment and shared among two or more NEFA-treatments.

An unsupervised principal component analysis (PCA) 2D plot as well as supervised partial-least squares discriminant analysis (PLS-DA) 2D plot were constructed to visualize the degree of difference between lipidome profiles in all NEFA-treatment groups and highlighted a separation in lipid profiles between the different NEF treatments (**Supplementary Figure 2**).

Analysis of the totaled peak intensities of lipids from each independent class revealed 8 differentially abundant lipid classes in PA-treated BeWo CT cells; 6 differentially abundant lipid classes in OA-treated BeWo CT cell; and 7 differentially abundant lipid classes in P/O-treated BeWo CT cells (**Figure 9a-c**; raw- $p < 0.0167$). Specifically, the abundance of lipids in the Ceramide (CER), Diacylglycerophosphoethanolamine (PE), and Phosphatidylinositol (PI) classes were only significantly increased in PA-treated BeWo CT cells. The lipids in the Bis[monoacylglycerol]phosphate (BMP) class were only significantly decreased in OA-treated BeWo CT cells. No significantly altered lipid classes were unique to the P/O treatment group. The Fatty Acyl Carnitine (Car), Lysophosphatidylglycerol (LPG) and Triacylglycerol (TG) lipid classes were found to be significantly elevated in all NEFA treatment groups relative to BSA control. The Diacylglycerol (DG) class was significantly increased, and the cholesterol and derivatives (ST) lipid class was significantly decreased in both the PA and P/O treatments. Finally, the Glucosylceramide (HexCer) and Ceramide-1-phosphate (SM) lipid classes were found to be significantly elevated in both the OA and P/O treated BeWo CT cells.






Figure 9. Differentially Abundant Lipid Classes in NEFA-treated BeWo CT cells. Differentially abundant lipid species identified in tiers 1 and 2 were sorted by lipid sub-class and total peak intensity of lipids in each class was determined. The abundances of lipid class in each NEFA treatment were compared to the BSA-control samples in binary via Wilcoxon-rank sum test. Boxplot were constructed to visualize lipid classes with differential abundance versus BSA-control in (A) PA-treated, (B) OA-treated, and (C) P/O-treated BeWo CT cells (raw- $p < 0.0167$). The red boxplots in each figure represent the lipid-class abundance in the BSA-control BeWo CT cells, and the blue boxplots in each figure represent the lipid-class abundance in the respective NEFA-treated BeWo CT cells. .

4. Discussion

PA and OA are the most abundant circulating NEFA species in the serum of pregnant women, and the concentrations of these NEFA are reportedly elevated under conditions of maternal GDM and obesity [47]. Dietary fat consumption data has highlighted that PA and OA are the most abundantly consumed dietary fats in Westernized cultures [49–52], and as maternal diet has been demonstrated to be an important regulator of placental metabolic function [27,53], these fats themselves may be important in regulating placental function in obese and GDM pregnancies. The current study utilized a multi-omics research approach to characterize the independent and combined impacts of the dietary NEFA species PA and OA on placental lipid metabolic processing using BeWo villous trophoblast cells.

The first objective was to utilize targeted lipidomic analyses via GC-FID and TLC-FID to characterize cellular FA and neutral profiles, as well as FA desaturation and elongation indices to better understand lipid processing in BeWo CT and SCT cells cultured with dietary NEFA species. The current study subsequently utilized a multi-omics research approach (combining transcriptomic, with untargeted metabolomic and lipidomic analyses) to characterize the underlying biochemical impacts of increased NEFA supply on the metabolic function of BeWo CT cells, given that previous studies have highlighted the dominant metabolic profile of these progenitor placental trophoblast cells [26,54–56]. Overall, our results highlighted that the dietary NEFAs PA and OA have different metabolic fates in placental trophoblast cells, as well as that these FAs are processed differentially when exposed to trophoblasts individually (PA and OA treatments) and in combination (P/O treatments).

4.1. Impacts of dietary NEFA on BeWo FA and Neutral Lipid Profiles

The current study utilized GC-FID as a preliminary readout of the impacts of increased dietary NEFA supply on BeWo trophoblasts via analysis of cellular FA profiles. It is important to note that altered FA compositions have been described in placentae from obese and GDM pregnancies [38,43,80], as well as in the development of placental dysfunctions including preeclampsia [81]. Thus, specific dietary NEFA-induced alterations to FA profiles may be reflective of a transition towards aberrant placental function. Specifically, the current study highlighted that a 72-hour PA exposure was associated with increased SFA levels in both CT and SCT cultures, with specific elevations in the C16:0 content in both differentiation states. OA exposure however, resulted in a profound increase in cellular MUFA species, and strikingly C18:1n9 accounted for almost two-thirds of all FA content in both OA-treated CT and SCT cells. In contrast, the P/O treatments only moderately impacted BeWo SFA and PUFA compositions, and no specific alterations in C16:0 composition were observed in both the P/O-treated CT and SCT cells. However, the P/O-treated cells displayed altered MUFA profiles, and increased C18:1n9 levels similar to the trends in the OA- treatments although with lower magnitude of change. Interestingly, previous data from rodent models have highlighted that placental MUFA and not SFA content correlates with maternal plasma FA levels [82], similar to the findings in P/O-treated BeWo cells reported here. Overall, these data highlighted that altering the supply of dietary NEFA to placental trophoblasts independently impacts the trophoblast cell FA profile and suggests that increased circulating NEFA levels contribute to alterations to placental lipid compositions. Additionally, these data highlight that dietary NEFA differentially impact trophoblast FA profiles when elevated independently and in combination with other dietary fats, suggesting that both the quantity and composition of fats is important in regulating placental lipid processing.

The collected FA profile data also allowed for investigation into elongation and desaturation metabolic processing of dietary FA through the calculation of FA elongase and desaturase indices [60–64]. Of particular interest to the current study was FA desaturation mediated by Stearoyl-CoA Desaturase 1 (SCD1), which is responsible for the production of palmitoleate (POA; C16:1n7) and OA (C18:1n9) from PA (C16:0) and stearate (C18:0), respectively. Previously, obesity has been found to impact placental FA desaturation via SCD1, although current reports have been somewhat inconsistent. Specifically, placentae of obese pregnancies have been found to display increased placental abundance of SCD1 mRNA suggesting increased FA desaturation [20]. A subsequent report, however, described an overall decrease in SCD1 activity in placentae of obese pregnancies without alterations in SCD protein abundance, highlighting that changes in FA desaturation may also be regulated by post-translational modifications [83]. In the current study, we observed decreased SCD1 activity (as assessed by the POA/PA ratio) in OA and P/O-treated BeWo CT and SCT cells. These data could indicate that previously observed reductions in SCD1 activity in term placentae of obese pregnancies could be facilitated in part by increased placental OA or MUFA supply [83]. In contrast, PA-treated BeWo CT and SCT cells displayed an elevated SCD1 index (as assessed by POA/PA ratio). As POA has been linked with increased oxygen consumption and β -oxidation activity in adipocytes [84,85], we speculate that increased PA desaturation to POA in PA-treated BeWo CT cells may underlie the increased mitochondrial respiratory activity that our research group has previously observed in these cells [56].

Another measure of placental lipid processing of interest to the current study is the desaturation mediated through the enzyme Fatty Acid Desaturase 2 (FADS2). FADS2 is responsible for the synthesis of long-chain polyunsaturated FA species [86], and its expression is altered in the placentae of some pre-eclamptic pregnancies [87]. Our results demonstrated that OA-exposure both alone and in combination with PA resulted in increased FADS2 production of γ -Linoleic Acid (GLA; C18:2n6), as well as increased elongation of OA to Gondoic Acid (C20:1n9). Increased levels of these specific FA species have previously been linked to anti-inflammatory outcomes [88,89]. As OA has been demonstrated to attenuate the lipotoxic effects of increased PA levels in trophoblast cells [90,91], we speculate that OA partially exerts its protective effects on trophoblast cells via the increased production of anti-inflammatory lipid intermediaries. Overall, our data demonstrated that modulating dietary FA composition impacts placental FA elongation and desaturation, and in turn leads to altered

production of various FA species which themselves may be able to directly impact placental metabolic and inflammatory processes.

Using TLC-FID, the current study also demonstrated that the proportion of TG in the neutral lipid fractions of BeWo CT and SCT cells was elevated only in cells exposed to OA independently, likely reflecting that OA is highly lipogenic in trophoblasts. These results were consistent with previous reports in cultured placental explants and isolated trophoblasts [92,93] and NEFA-treated BeWo [94,95] cells that have demonstrated that imported OA is highly localized to TG lipid fractions. This may indicate that increased supply of OA to the placenta is an important factor that helps facilitate the increased TG synthesis that has previously been reported in obese and GDM placentae [20,26,28,31]. Interestingly, the P/O-treated BeWo CT and SCT cells were not observed to have altered neutral lipid and TG compositions when assessed by TLC-FID. These data aligned with reports from isolated primary human trophoblasts, demonstrating reduced lipid droplet formation with a combined OA and PA exposure compared to OA-alone exposures [93]. Subsequent analysis of lipid class abundance using untargeted LC-MS/MS also highlighted increased levels of PI, and PE lipid species in PA-treated BeWo CT cells. These results were also consistent with readouts in primary placental trophoblasts (PHT) that have highlighted PA is highly localized to trophoblast phospholipid fractions [83]. These data may indicate similarities in lipid processing in PHT and BeWo trophoblast cell culture systems. Overall, this suggests that saturated and monounsaturated dietary NEFA have different metabolic fates in placental trophoblasts as well as that the metabolic fates of these FA species are altered when present in combination with other dietary FAs.

The FA and neutral lipid profile data, however, highlighted limited alterations in trophoblast lipid profiles following syncytialization and, specifically, only four differentially abundant FA species were found in BeWo SCT cells. In contrast, previous work in cultured primary human trophoblasts (PHT) cells demonstrated profound reductions in the levels of many FA species in placental villous trophoblast cells following syncytialization [96]. The differences between the current study and previous reports are likely due to differences in the reporting of FA profile data and may highlight a limitation in the current study. The current study reported FA profiles as a percentage of total FA abundance, which contrasts with previous reports that highlighted protein normalized FA quantity. Thus, the current study was not able to report whether BeWo trophoblast cells demonstrate similar reductions in the absolute quantities of different FA species as was highlighted with PHT cultures [96]. Future works using quantitative readouts of FA abundances are therefore needed to establish if BeWo trophoblasts exhibit similar differences in lipid metabolism between differentiated and progenitor CT cells and differentiated SCT cells.

4.2. A multi-omics analysis of NEFA-treated BeWo CT metabolic function

The multi-omic analyses revealed that exposure to different dietary NEFA species extensively altered BeWo CT cellular transcriptome and lipidome profiles but led to limited alterations in cellular metabolome profiles. The mRNA microarray transcriptomic analyses highlighted 340 DEG in PA-treated BeWo CT cells; 308 DEGs in OA-treated BeWo CT cells; as well as 221 DEGs in P/O-treated BeWo CT cells. Subsequent over-representation enrichment analysis highlighted one significantly enriched functional pathway in the transcriptomic datasets. Specifically, the Wikipathways “Fatty Acid Biosynthesis Pathway” was significantly enriched in P/O-treated BeWo CT cells. Further enrichment analysis using a GSEA approach further highlighted alterations in lipid signaling and metabolic pathways in the PA, OA-treated BeWo CT cells. For example, PA-treated cells displayed a significant upregulation of genes involved in the Reactome “Fatty Acid Biosynthesis” and “PPAR Signaling Pathway” datasets, while OA-treated cells displayed a significant upregulation in the Wikipathways and KEGG “PPAR Signaling Pathway” datasets, as well as a significant downregulation in the KEGG “Linoleic Acid Metabolism” dataset. Overall, pathway enrichment analysis of NEFA-treated BeWo cells transcriptomic profiles demonstrated an altered regulation of key genes involved in lipid metabolic pathways. This data further indicates that placental trophoblasts modulate lipid processing functions in direct response to an increased supply of PA and OA.

Differentially expressed genes involved in lipid metabolism processes were identified, and their expression patterns were validated by RT-qPCR. The microarray readouts highlighted an upregulation of Acyl-CoA Synthetase Long Chain Family Member 5 (*ACSL5*), as well as perilipin 2 (*PLIN2*) in all NEFA-treated BeWo CT cells. Acyl-CoA synthetase (*ACSL*) enzymes have previously been highlighted to be directly involved in FA uptake and are responsible for conjugating FA species to Coenzyme A, an important first step in lipid metabolism that helps prevent FA efflux [83,97,98]. Interestingly, *ACSL5* expression has also been found to be induced in PUFA-exposed BeWo trophoblasts, leading to an overall increase in FA uptake [99]. The data from the current study therefore suggests that *ACSL5* is also involved in the processing of PA and OA in BeWo trophoblasts, and its increased expression likely facilitates increased cellular uptake of FA species. *PLIN2* on the other hand, has been shown to be required for lipid droplet synthesis in trophoblasts [100]. More importantly, placental expression of *PLIN2* has been demonstrated to be elevated in GDM pregnancies [101], and its expression has been found to be correlated with maternal pre-pregnancy Body-Mass-Index (BMI) [28]. Thus, the current study suggests that increased supply of dietary NEFAs to the placenta in obese and GDM pregnancies may directly facilitate this increased expression of *PLIN2*. In turn, this dietary NEFA-mediated modulation of *PLIN2* expression may promote TG accumulation in lipid droplets and subsequently placental steatosis in obese and GDM pregnancies.

Both *ACSL5* and *PLIN2* were DEGs involved in the enriched “PPAR signaling pathway” in both PA and OA-treated BeWo CT cells. These genes are known to be specifically induced by PPAR γ [102,103], suggesting that PPAR γ is activated in placental trophoblasts in response to elevated NEFA levels. However, all NEFA-treated BeWo CT cells also displayed reduced mRNA abundance of *SCD*, another lipid metabolism gene that has been shown in some tissue types to be induced by PPAR γ [104,105]. The reduced expression of *SCD* in the current study could indicate that certain lipid metabolism genes are differentially regulated by PPAR γ in placental trophoblast compared to what has previously been shown in other tissues. Notably, PPAR γ itself is regulated by multiple post-translational modifications (PTMs) including phosphorylation, SUMOylation, acetylation, and ubiquitination [106]. NEFA-mediated alterations in these PTMs could explain why some PPAR-associated genes were activated, and others were repressed in our BeWo trophoblast cells. Characterizing these modifications in future investigations will be important to fully understand the impacts of NEFAs on placental metabolism.

The microarray analysis further highlighted an increase in the mRNA abundance of cAMP responsive element-binding protein 3-like 3 (*CREB3L3*) in all NEFA treatment groups. *CREB3L3* is a transcription factor that has previously been demonstrated to regulate lipid metabolism via modulating the expression of genes involved in FA oxidation [107–109]. Interestingly, *CREB3L3* has also been implicated in the pathophysiology of inflammation and Endoplasmic Reticulum (ER) stress [108–110]. This could indicate that *CREB3L3* controls lipid processing functions in trophoblasts, as well as underlies the development of inflammation and ER stress that has previously been observed in some PA-treated placental trophoblasts [91,93,111]. The microarray analysis in the current study however, revealed that BeWo CT cells exposed to 100 μ M levels of PA for 72 hours do not have elevated expression of genes related to inflammation (such as *TNF α* , *IL6*, and *IL-32*), and ER stress (such as *BCL2*, *DDIT3*, and *XBP1*). In contrast, these inflammation and ER stress pathways have previously been shown to be induced in PA-treated PHT cultures [93,111]. These differential responses may arise from variations in PA dose utilized (100 μ M in the current study compared to 200–500 μ M in primary trophoblasts [93,111]). We speculate that the increased *CREB3L3* expression highlighted in the current study may reflect an early timepoint in transition towards ER stress and inflammation in PA-exposed BeWo trophoblasts.

Subsequently, the current study utilized an integrative approach combining readouts from transcriptomic, metabolomic, and lipidomic datasets to further describe the impacts of dietary NEFA exposure on BeWo CT metabolic function. Specifically, the transcriptome and metabolome datasets were integrated via the Joint Pathway Analysis module of the MetaboAnalyst software. This analysis highlighted a significant enrichment in the Purine Metabolism pathway in OA-treated BeWo CT cells that was associated with increased cellular accumulations of adenosine (+3.30 FC), inosine (+2.45 FC),

and guanosine (+1.69 FC) that may be suggestive of reduced purine breakdown. Increased degradation of purines has previously been shown to be involved in placental responses to increased inflammation [112]. The increased purine accumulation in OA-treated trophoblasts shown here may further suggest that anti-inflammatory processes are activated in dietary OA-treated trophoblasts. Alternatively, the increased purine abundances, in conjunction with the upregulation of phosphoribosyl pyrophosphate synthetase 1 (PRPS1) observed in the microarray panel, could reflect increased purine synthesis via the Pentose Phosphate Pathway (PPP), which has previously been shown to operate in placental tissues [113–115]. As the PPP increases generation of NADPH, a metabolite with antioxidant effects [116,117], these data may highlight that OA also acts to reduce oxidative stress in placental trophoblasts.

Combining readouts from the transcriptomic, metabolomic and lipidomic datasets also highlighted a potential shift in oxidative substrate selection in BeWo trophoblasts cultured with PA (both alone and in conjunction with OA) towards increased FA oxidation (β -oxidation). Examination of differentially abundant metabolites in PA and P/O-treated BeWo CT revealed a significant reduction in intracellular levels of 3-hydroxymethylglutaric acid (-2.89 FC in PA-treated CT cells; -2.00 FC in P/O-treated CT cells), a byproduct of leucine degradation [118]. We speculate that this may indicate a reduced breakdown of branched-chain amino acid species into byproducts that can enter The Tricarboxylic Acid Cycle (TCA Cycle) in BeWo trophoblasts cultured under increased PA levels. Additionally, we observed increased C16:0 carnitine (+13.92 FC in PA-treated CT cells; +3.79 FC in P/O-treated CT cells) levels in BeWo trophoblasts treated with PA. As FA species must first be conjugated to carnitine before being translocated to the mitochondrial matrix for oxidation [119], the increased acylcarnitine levels in the cells may reflect a metabolic shift towards using PA as an oxidative fuel.

Moreover, PA and P/O-treated BeWo CT cells were found to have elevated expression of genes involved in β -oxidative metabolism. Specifically, we observed increased expression of Very-Long Chain Acyl-CoA Dehydrogenase (*ACADVL*), the enzyme responsible for catalyzing the preliminary step in β -oxidation metabolism, and an increased expression of Acetyl-CoA Carboxylase Beta (*ACACB*), an enzyme thought to control β -oxidative activity by catalyzing the carboxylation of acetyl-CoA to malonyl-CoA. Notably, elevated β -oxidative activity has previously been linked with increased generation of Reactive Oxygen Species (ROS) [120,121], that ultimately promotes oxidative mitochondrial damage [122,123]. Therefore, the PA-mediated increases in β -oxidation associated genes and metabolites in BeWo trophoblasts could be indicative of an early transition towards placental mitochondrial dysfunction.

Interestingly, malonic acid, an end-point metabolic byproduct of β -oxidation that is synthesized by *ACACB*, was only found to be increased in P/O-cultured BeWo CT cells (+1.85 FC), and not in PA-cultured cells. As malonate has been found to limit β -oxidation activity by inhibiting acylcarnitine transport into the mitochondrial matrix [124], these data may highlight that BeWo trophoblasts exposed to PA in combination with OA, but not PA-alone, are able to restrict excessive β -oxidation activity. Malonate-mediated inhibition of β -oxidation in the P/O-cultured BeWo CT cells may act to suppress excessive mitochondrial ROS production and could therefore limit mitochondrial membrane damage. We additionally speculate that these data may highlight a shift towards incomplete β -oxidation in BeWo trophoblasts cultured with PA alone, as these cells (but not P/O-treated CT cells) also displayed increased intracellular levels of the shortened-chain C14:0 acylcarnitine (+3.99 FC). Increased production of short-chain acylcarnitine species has previously been linked with pro-inflammatory processes [125], and mitochondrial overload [126], and thus these data may further highlight a transition towards inflammation and mitochondrial dysfunction in PA-exposed trophoblasts. However, the insights of the current study into incomplete β -oxidation are minimized due to the limited identifications of short chain acylcarnitine species in our untargeted lipidomic readouts. Future studies may need to utilize targeted metabolomic approaches or utilize radio-labelled FA species to better identify the production of short-chain acylcarnitine species and incomplete β -oxidation in PA-treated BeWo trophoblasts.

4.3. Conclusion

The results of the current study highlight the continued utility of the BeWo preparation as a pre-clinical placental model, and that the dietary FAs PA and OA are important regulators of placental trophoblast lipid metabolism. Moreover, it is shown that excessive availability of these FAs is associated with changes in FA desaturation, elongation, esterification, and oxidation processes in placental trophoblast cells. These results further support the concept that increased levels of certain dietary FAs in maternal circulation are involved in facilitating the aberrant placental function in obese and GDM pregnancies. In turn these data suggest that monitoring circulating FA levels in obese and GDM pregnancies, and subsequently implementing dietary interventions through modulating dietary fat content and composition continues to be needed in the clinical management of these at-risk pregnancies. Implementation of healthy diet and lifestyle choices in these “at-risk” pregnancies could act to preserve appropriate placental metabolic function throughout gestation and in turn reduce risk of future metabolic health complication in the offspring.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org., S1 file: Supplementary data tables, S2 file: Supplementary figures.

Author Contributions: Conceptualization: Z.E. and T.R.; Investigation: Z.E., O.S., L.Z., A.Z.B., X.L., and S.Z.; Formal Analysis: Z.E., A.Z.B., X.L., and S.Z.; Visualization: Z.E., A.Z.B., X.L., and S.Z.; Writing – original draft: Z.E., and T.R.; Writing – Review & Editing: Z.E., A.Z.B., S.Z., and T.R.; Funding Acquisition: L.L., and T.R.

Funding: This project was supported by a National Institutes of Health (NIH) Human Placenta Project Grant (grant No. U01 HD087181-01). The metabolomics work done at the Metabolomics Innovation Centre (TMIC) of Canada was also supported by Genome Canada and Canada Foundation for Innovation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data Availability Statement: Microarray data is available on the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (GEO) database (GSE197385). A full list of metabolome peak pairs, and their high confidence/putative identifications is available in **Supplementary Table 7**. A full list of lipidome peak intensities, and their high confidence/putative identifications is available in **Supplementary Table 10**.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Zhuang, W.; Lv, J.; Liang, Q.; Chen, W.; Zhang, S.; Sun, X. Adverse Effects of Gestational Diabetes-Related Risk Factors on Pregnancy Outcomes and Intervention Measures. *Exp Ther Med* **2020**, doi:10.3892/etm.2020.9050.
2. Yang, Z.; Phung, H.; Freebairn, L.; Sexton, R.; Raulli, A.; Kelly, P. Contribution of Maternal Overweight and Obesity to the Occurrence of Adverse Pregnancy Outcomes. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **2019**, *59*, 367–374, doi:10.1111/ajo.12866.
3. Yegorov, Y.; Visser, G.H.A. Obesity, Gestational Diabetes and Pregnancy Outcome. *Semin Fetal Neonatal Med* **2009**, *14*, 77–84, doi:10.1016/j.siny.2008.09.002.
4. Armitage, J.A.; Poston, L.; Taylor, P.D. Developmental Origins of Obesity and the Metabolic Syndrome: The Role of Maternal Obesity. *Front Horm Res* **2008**, *36*, 73–84, doi:10.1159/000115355.
5. Manderson, J.; Mullan, B.; Patterson, C.; Hadden, D.; Traub, A.; McCance, D. Cardiovascular and Metabolic Abnormalities in the Offspring of Diabetic Pregnancy. *Diabetologia* **2002**, *45*, 991–996, doi:10.1007/s00125-002-0865-y.
6. Agarwal, P.; Morriveau, T.S.; Kereliuk, S.M.; Doucette, C.A.; Wicklow, B.A.; Dolinsky, V.W. Maternal Obesity, Diabetes during Pregnancy and Epigenetic Mechanisms That Influence the Developmental Origins of Cardiometabolic Disease in the Offspring. *Crit Rev Clin Lab Sci* **2018**, *55*, 71–101, doi:10.1080/10408363.2017.1422109.
7. Catalano, P.M. The Impact of Gestational Diabetes and Maternal Obesity on the Mother and Her Offspring. *J Dev Orig Health Dis* **2010**, *1*, 208–215, doi:10.1017/S2040174410000115.
8. Shrestha, N.; Ezechukwu, H.C.; Holland, O.J.; Hryciw, D.H. Developmental Programming of Peripheral Diseases in Offspring Exposed to Maternal Obesity during Pregnancy. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **2020**, *319*, R507–R516, doi:10.1152/ajpregu.00214.2020.
9. Whitaker, R.C. Predicting Preschooler Obesity at Birth: The Role of Maternal Obesity in Early Pregnancy. *Pediatrics* **2004**, *114*, e29–e36, doi:10.1542/peds.114.1.e29.

10. Boney, C.M.; Verma, A.; Tucker, R.; Vohr, B.R. Metabolic Syndrome in Childhood: Association with Birth Weight, Maternal Obesity, and Gestational Diabetes Mellitus. *Pediatrics* **2005**, *115*, e290-6, doi:10.1542/peds.2004-1808.
11. Vääräsmäki, M.; Pouta, A.; Elliot, P.; Tapanainen, P.; Sovio, U.; Ruokonen, A.; Hartikainen, A.-L.; McCarthy, M.; Järvelin, M.-R. Adolescent Manifestations of Metabolic Syndrome Among Children Born to Women With Gestational Diabetes in a General-Population Birth Cohort. *Am J Epidemiol* **2009**, *169*, 1209–1215, doi:10.1093/aje/kwp020.
12. Ferrara, A. Increasing Prevalence of Gestational Diabetes Mellitus: A Public Health Perspective. *Diabetes Care* **2007**, *30*, S141–S146, doi:10.2337/dc07-s206.
13. Kim, S.Y.; Dietz, P.M.; England, L.; Morrow, B.; Callaghan, W.M. Trends in Pre-Pregnancy Obesity in Nine States, 1993–2003*. *Obesity* **2007**, *15*, 986–993, doi:10.1038/oby.2007.621.
14. Li, G.; Wei, T.; Ni, W.; Zhang, A.; Zhang, J.; Xing, Y.; Xing, Q. Incidence and Risk Factors of Gestational Diabetes Mellitus: A Prospective Cohort Study in Qingdao, China. *Front Endocrinol (Lausanne)* **2020**, *11*, doi:10.3389/fendo.2020.00636.
15. Hod, M.; Kapur, A.; Sacks, D.A.; Hadar, E.; Agarwal, M.; Di Renzo, G.C.; Roura, L.C.; McIntyre, H.D.; Morris, J.L.; Divakar, H. The International Federation of Gynecology and Obstetrics (FIGO) Initiative on Gestational Diabetes Mellitus: A Pragmatic Guide for Diagnosis, Management, and Care #. *International Journal of Gynecology & Obstetrics* **2015**, *131*, S173–S211, doi:10.1016/S0020-7292(15)30033-3.
16. Driscoll, A.K.; Gregory, E.C.W. Increases in Prepregnancy Obesity: United States, 2016–2019. *NCHS Data Brief* **2020**, 1–8.
17. Howell, K.R.; Powell, T.L. Effects of Maternal Obesity on Placental Function and Fetal Development. *Reproduction* **2017**, *153*, R97–R108, doi:10.1530/REP-16-0495.
18. Hebert, J.F.; Myatt, L. Placental Mitochondrial Dysfunction with Metabolic Diseases: Therapeutic Approaches. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **2021**, *1867*, 165967, doi:10.1016/j.bbdis.2020.165967.
19. Kelly, A.C.; Powell, T.L.; Jansson, T. Placental Function in Maternal Obesity. *Clin Sci* **2020**, *134*, 961–984, doi:10.1042/CS20190266.
20. Calabuig-Navarro, V.; Haghiac, M.; Minium, J.; Glazebrook, P.; Ranasinghe, G.C.; Hoppel, C.; Hauguel de Mouzon, S.; Catalano, P.; O'Tierney-Ginn, P. Effect of Maternal Obesity on Placental Lipid Metabolism. *Endocrinology* **2017**, *158*, 2543–2555, doi:10.1210/en.2017-00152.
21. Nogues, P.; Dos Santos, E.; Couturier-Tarrade, A.; Berveiller, P.; Arnould, L.; Lamy, E.; Grassin-Delyle, S.; Vialard, F.; Dieudonne, M.-N. Maternal Obesity Influences Placental Nutrient Transport, Inflammatory Status, and Morphology in Human Term Placenta. *J Clin Endocrinol Metab* **2021**, *106*, 1880–1896, doi:10.1210/clinem/dgaa660.
22. Hastie, R.; Lappas, M. The Effect of Pre-Existing Maternal Obesity and Diabetes on Placental Mitochondrial Content and Electron Transport Chain Activity. *Placenta* **2014**, *35*, 673–683, doi:10.1016/j.placenta.2014.06.368.
23. Mandò, C.; Anelli, G.M.; Novielli, C.; Panina-Bordignon, P.; Massari, M.; Mazzocco, M.I.; Cetin, I. Impact of Obesity and Hyperglycemia on Placental Mitochondria. *Oxid Med Cell Longev* **2018**, *2018*, 1–10, doi:10.1155/2018/2378189.
24. Maloyan, A.; Mele, J.; Muralimanoharan, S.; Myatt, L. Placental Metabolic Flexibility Is Affected by Maternal Obesity. *Placenta* **2016**, *45*, 69, doi:10.1016/j.placenta.2016.06.031.
25. Mele, J.; Muralimanoharan, S.; Maloyan, A.; Myatt, L. Impaired Mitochondrial Function in Human Placenta with Increased Maternal Adiposity. *Am J Physiol Endocrinol Metab* **2014**, *307*, E419-25, doi:10.1152/ajpendo.00025.2014.
26. Valent, A.M.; Choi, H.; Kolahi, K.S.; Thornburg, K.L. Hyperglycemia and Gestational Diabetes Suppress Placental Glycolysis and Mitochondrial Function and Alter Lipid Processing. *The FASEB Journal* **2021**, *35*, doi:10.1096/fj.202000326RR.
27. Calabuig-Navarro, V.; Puchowicz, M.; Glazebrook, P.; Haghiac, M.; Minium, J.; Catalano, P.; Hauguel de Mouzon, S.; O'Tierney-Ginn, P. Effect of ω -3 Supplementation on Placental Lipid Metabolism in Overweight and Obese Women. *Am J Clin Nutr* **2016**, *103*, 1064–1072, doi:10.3945/ajcn.115.124651.
28. Hirschmugl, B.; Desoye, G.; Catalano, P.; Klymiuk, I.; Scharnagl, H.; Payr, S.; Kitzinger, E.; Schliefssteiner, C.; Lang, U.; Wadsack, C.; et al. Maternal Obesity Modulates Intracellular Lipid Turnover in the Human Term Placenta. *Int J Obes* **2017**, *41*, 317–323, doi:10.1038/ijo.2016.188.
29. Bucher, M.; Montani, K.R.C.; Myatt, L.; Weintraub, S.; Tavori, H.; Maloyan, A. Dyslipidemia, Insulin Resistance, and Impairment of Placental Metabolism in the Offspring of Obese Mothers. *J Dev Orig Health Dis* **2021**, *12*, 738–747, doi:10.1017/S2040174420001026.
30. Powell, T.L.; Barner, K.; Madi, L.; Armstrong, M.; Manke, J.; Uhlson, C.; Jansson, T.; Ferchaud-Roucher, V. Sex-Specific Responses in Placental Fatty Acid Oxidation, Esterification and Transfer Capacity to Maternal Obesity. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **2021**, *1866*, 158861, doi:10.1016/j.bbalip.2020.158861.

31. Hulme, C.H.; Nicolaou, A.; Murphy, S.A.; Heazell, A.E.P.; Myers, J.E.; Westwood, M. The Effect of High Glucose on Lipid Metabolism in the Human Placenta. *Sci Rep* **2019**, *9*, 14114, doi:10.1038/s41598-019-50626-x.
32. Visiedo, F.; Bugatto, F.; Quintero-Prado, R.; Cózar-Castellano, I.; Bartha, J.L.; Perdomo, G. Glucose and Fatty Acid Metabolism in Placental Explants From Pregnancies Complicated With Gestational Diabetes Mellitus. *Reproductive Sciences* **2015**, *22*, 798–801, doi:10.1177/1933719114561558.
33. Visiedo, F.; Bugatto, F.; Sánchez, V.; Cózar-Castellano, I.; Bartha, J.L.; Perdomo, G. High Glucose Levels Reduce Fatty Acid Oxidation and Increase Triglyceride Accumulation in Human Placenta. *American Journal of Physiology-Endocrinology and Metabolism* **2013**, *305*, E205–E212, doi:10.1152/ajpendo.00032.2013.
34. Ortega, M.A.; Saez, M.A.; Sainz, F.; Fraile-Martínez, O.; García-Gallego, S.; Pekarek, L.; Bravo, C.; Coca, S.; Mon, M.Á.-; Buján, J.; et al. Lipidomic Profiling of Chorionic Villi in the Placentas of Women with Chronic Venous Disease. *Int J Med Sci* **2020**, *17*, 2790–2798, doi:10.7150/ijms.49236.
35. Mohammad, S.; Bhattacharjee, J.; Vasanthan, T.; Harris, C.S.; Bainbridge, S.A.; Adamo, K.B. Metabolomics to Understand Placental Biology: Where Are We Now? *Tissue Cell* **2021**, *73*, 101663, doi:10.1016/j.tice.2021.101663.
36. Fattuoni, C.; Mandò, C.; Palmas, F.; Anelli, G.M.; Novielli, C.; Parejo Laudicina, E.; Savasi, V.M.; Barberini, L.; Dessì, A.; Pintus, R.; et al. Preliminary Metabolomics Analysis of Placenta in Maternal Obesity. *Placenta* **2018**, *61*, 89–95, doi:10.1016/j.placenta.2017.11.014.
37. Segura, M.T.; Demmelmair, H.; Krauss-Etschmann, S.; Nathan, P.; Dehmel, S.; Padilla, M.C.; Rueda, R.; Koletzko, B.; Campoy, C. Maternal BMI and Gestational Diabetes Alter Placental Lipid Transporters and Fatty Acid Composition. *Placenta* **2017**, *57*, 144–151, doi:10.1016/j.placenta.2017.07.001.
38. Yang, Y.; Pan, Z.; Guo, F.; Wang, H.; Long, W.; Wang, H.; Yu, B. Placental Metabolic Profiling in Gestational Diabetes Mellitus: An Important Role of Fatty Acids. *J Clin Lab Anal* **2021**, doi:10.1002/jcla.24096.
39. Furse, S.; Fernandez-Twinn, D.S.; Jenkins, B.; Meek, C.L.; Williams, H.E.L.; Smith, G.C.S.; Charnock-Jones, D.S.; Ozanne, S.E.; Koulman, A. A High-Throughput Platform for Detailed Lipidomic Analysis of a Range of Mouse and Human Tissues. *Anal Bioanal Chem* **2020**, *412*, 2851–2862, doi:10.1007/s00216-020-02511-0.
40. Uhl, O.; Demmelmair, H.; Segura, M.T.; Florido, J.; Rueda, R.; Campoy, C.; Koletzko, B. Effects of Obesity and Gestational Diabetes Mellitus on Placental Phospholipids. *Diabetes Res Clin Pract* **2015**, *109*, 364–371, doi:10.1016/j.diabres.2015.05.032.
41. Bidne, K.L.; Rister, A.L.; McCain, A.R.; Hitt, B.D.; Dodds, E.D.; Wood, J.R. Maternal Obesity Alters Placental Lysophosphatidylcholines, Lipid Storage, and the Expression of Genes Associated with Lipid Metabolism. *Biol Reprod* **2021**, *104*, 197–210, doi:10.1093/biolre/iaaa191.
42. Stuart, T.J.; O'Neill, K.; Condon, D.; Sasson, I.; Sen, P.; Xia, Y.; Simmons, R.A. Diet-Induced Obesity Alters the Maternal Metabolome and Early Placenta Transcriptome and Decreases Placenta Vascularity in the Mouse†. *Biol Reprod* **2018**, *98*, 795–809, doi:10.1093/biolre/iory010.
43. Segura, M.T.; Demmelmair, H.; Krauss-Etschmann, S.; Nathan, P.; Dehmel, S.; Padilla, M.C.; Rueda, R.; Koletzko, B.; Campoy, C. Maternal BMI and Gestational Diabetes Alter Placental Lipid Transporters and Fatty Acid Composition. *Placenta* **2017**, *57*, 144–151, doi:10.1016/j.placenta.2017.07.001.
44. Delhaes, F.; Giza, S.A.; Koreman, T.; Eastabrook, G.; McKenzie, C.A.; Bedell, S.; Regnault, T.R.H.; de Vrijer, B. Altered Maternal and Placental Lipid Metabolism and Fetal Fat Development in Obesity: Current Knowledge and Advances in Non-Invasive Assessment. *Placenta* **2018**, *69*, 118–124, doi:10.1016/j.placenta.2018.05.011.
45. Herrera Martínez, A. Hyperlipidemia during Gestational Diabetes and Its Relation with Maternal and Offspring Complications. *Nutr Hosp* **2018**, doi:10.20960/nh.1539.
46. Scifres, C.M.; Catov, J.M.; Simhan, H.N. The Impact of Maternal Obesity and Gestational Weight Gain on Early and Mid-Pregnancy Lipid Profiles. *Obesity* **2014**, *22*, 932–938, doi:10.1002/oby.20576.
47. Chen, X.; Scholl, T.O.; Leskiw, M.; Savaille, J.; Stein, T.P. Differences in Maternal Circulating Fatty Acid Composition and Dietary Fat Intake in Women with Gestational Diabetes Mellitus or Mild Gestational Hyperglycemia. *Diabetes Care* **2010**, *33*, 2049–2054, doi:10.2337/dc10-0693.
48. Villa, P.M.; Laivuori, H.; Kajantie, E.; Kaaja, R. Free Fatty Acid Profiles in Preeclampsia. *Prostaglandins Leukot Essent Fatty Acids* **2009**, *81*, 17–21, doi:10.1016/j.plefa.2009.05.002.
49. Denomme, J.; Stark, K.D.; Holub, B.J. Directly Quantitated Dietary (n-3) Fatty Acid Intakes of Pregnant Canadian Women Are Lower than Current Dietary Recommendations. *J Nutr* **2005**, *135*, 206–211, doi:10.1093/ajcn.135.2.206 [pii].
50. Savard, C.; Lemieux, S.; Weisnagel, S.; Fontaine-Bisson, B.; Gagnon, C.; Robitaille, J.; Morisset, A.-S. Trimester-Specific Dietary Intakes in a Sample of French-Canadian Pregnant Women in Comparison with National Nutritional Guidelines. *Nutrients* **2018**, *10*, 768, doi:10.3390/nu10060768.
51. Iggman, D.; Risérus, U. Role of Different Dietary Saturated Fatty Acids for Cardiometabolic Risk. *Clin Lipidol* **2011**, *6*, 209–223, doi:10.2217/clp.11.7.

52. Kien, C.L.; Bunn, J.Y.; Stevens, R.; Bain, J.; Ikayeva, O.; Crain, K.; Koves, T.R.; Muoio, D.M. Dietary Intake of Palmitate and Oleate Has Broad Impact on Systemic and Tissue Lipid Profiles in Humans. *Am J Clin Nutr* **2014**, *99*, 436–445, doi:10.3945/ajcn.113.070557.
53. Alvarado, F.L.; Calabuig-Navarro, V.; Haghiac, M.; Puchowicz, M.; Tsai, P.-J.S.; O'Tierney-Ginn, P. Maternal Obesity Is Not Associated with Placental Lipid Accumulation in Women with High Omega-3 Fatty Acid Levels. *Placenta* **2018**, *69*, 96–101, doi:10.1016/j.placenta.2018.07.016.
54. Kolahi, K.; Valent, A.; Thornburg, K.L. Cytotrophoblast, Not Syncytiotrophoblast, Dominates Glycolysis and Oxidative Phosphorylation in Human Term Placenta. *Sci Rep* **2017**, 1–12, doi:10.1038/srep42941.
55. Kolahi, K.; Louey, S.; Varlamov, O.; Thornburg, K. Real-Time Tracking of BODIPY-C12 Long-Chain Fatty Acid in Human Term Placenta Reveals Unique Lipid Dynamics in Cytotrophoblast Cells. *PLoS One* **2016**, *11*, 1–23, doi:10.1371/journal.pone.0153522.
56. Easton, Z.J.W.; Delhaes, F.; Mathers, K.; Zhao, L.; Vanderboor, C.M.G.; Regnault, T.R.H. Syncytialization and Prolonged Exposure to Palmitate Impacts BeWo Respiration. *Reproduction* **2021**, *161*, 73–88, doi:10.1530/REP-19-0433.
57. Folch, J.; Lees, M.; Stanley, G.H.S. A SIMPLE METHOD FOR THE ISOLATION AND PURIFICATION OF TOTAL LIPIDES FROM ANIMAL TISSUES. *Journal of Biological Chemistry* **1957**, *226*, 497–509, doi:10.1016/S0021-9258(18)64849-5.
58. Klaiman, J.M.; Price, E.R.; Guglielmo, C.G. Fatty Acid Composition of Pectoralis Muscle Membrane, Intramuscular Fat Stores and Adipose Tissue of Migrant and Wintering White-Throated Sparrows (*Zonotrichia Albicollis*). *Journal of Experimental Biology* **2009**, *212*, 3865–3872, doi:10.1242/jeb.034967.
59. Sarr, O.; Mathers, K.E.; Zhao, L.; Dunlop, K.; Chiu, J.; Guglielmo, C.G.; Bureau, Y.; Cheung, A.; Raha, S.; Lee, T.-Y.; et al. Western Diet Consumption through Early Life Induces Microvesicular Hepatic Steatosis in Association with an Altered Metabolome in Low Birth Weight Guinea Pigs. *J Nutr Biochem* **2019**, *67*, 219–233, doi:10.1016/j.jnutbio.2019.02.009.
60. Merino, D.M.; Johnston, H.; Clarke, S.; Roke, K.; Nielsen, D.; Badawi, A.; El-Soheemy, A.; Ma, D.W.L.; Mutch, D.M. Polymorphisms in FADS1 and FADS2 Alter Desaturase Activity in Young Caucasian and Asian Adults. *Mol Genet Metab* **2011**, *103*, 171–178, doi:10.1016/j.ymgme.2011.02.012.
61. Chajès, V.; Joulin, V.; Clavel-Chapelon, F. The Fatty Acid Desaturation Index of Blood Lipids, as a Biomarker of Hepatic Stearoyl-CoA Desaturase Expression, Is a Predictive Factor of Breast Cancer Risk. *Curr Opin Lipidol* **2011**, *22*, 6–10, doi:10.1097/MOL.0b013e3283404552.
62. Jeyakumar, S.M.; Lopamudra, P.; Padmini, S.; Balakrishna, N.; Giridharan, N. V.; Vajreswari, A. Fatty Acid Desaturation Index Correlates with Body Mass and Adiposity Indices of Obesity in Wistar NIN Obese Mutant Rat Strains WNIN/Ob and WNIN/GR-Ob. *Nutr Metab (Lond)* **2009**, *6*, 27, doi:10.1186/1743-7075-6-27.
63. Massih, Y.N.; Hall, A.G.; Suh, J.; King, J.C. Zinc Supplements Taken with Food Increase Essential Fatty Acid Desaturation Indices in Adult Men Compared with Zinc Taken in the Fasted State. *J Nutr* **2021**, *151*, 2583–2589, doi:10.1093/jn/nxab149.
64. Gómez-Vilarrubla, A.; Mas-Parés, B.; Díaz, M.; Xargay-Torrent, S.; Carreras-Badosa, G.; Jové, M.; Martin-Gari, M.; Bonmatí-Santané, A.; de Zegher, F.; Ibañez, L.; et al. Fatty Acids in the Placenta of Appropriate-versus Small-for-Gestational-Age Infants at Term Birth. *Placenta* **2021**, *109*, 4–10, doi:10.1016/j.placenta.2021.04.009.
65. Dunlop, K.P. Skeletal Muscle Lipid Metabolism and Markers of Insulin Resistance in Young Male Low Birth Weight Offspring in Combination With a Postnatal Western Diet. *Electronic Thesis and Dissertation Respository* **2015**, 2946.
66. Easton, Z.J.W.; Luo, X.; Li, L.; Regnault, T.R.H. The Impact of Hyperglycemia upon BeWo Trophoblast Cell Metabolic Function: A Multi-OMICS and Functional Metabolic Analysis. *PLoS One* **2023**, *18*, doi:10.1371/journal.pone.0283118.
67. Mootha, V.K.; Lindgren, C.M.; Eriksson, K.-F.; Subramanian, A.; Sihag, S.; Lehar, J.; Puigserver, P.; Carlsson, E.; Ridderstråle, M.; Laurila, E.; et al. PGC-1 α -Responsive Genes Involved in Oxidative Phosphorylation Are Coordinately Downregulated in Human Diabetes. *Nat Genet* **2003**, *34*, 267–273, doi:10.1038/ng1180.
68. Subramanian, A.; Tamayo, P.; Mootha, V.K.; Mukherjee, S.; Ebert, B.L.; Gillette, M.A.; Paulovich, A.; Pomeroy, S.L.; Golub, T.R.; Lander, E.S.; et al. Gene Set Enrichment Analysis: A Knowledge-Based Approach for Interpreting Genome-Wide Expression Profiles. *Proceedings of the National Academy of Sciences* **2005**, *102*, 15545–15550, doi:10.1073/pnas.0506580102.
69. Liao, Y.; Wang, J.; Jaehnig, E.J.; Shi, Z.; Zhang, B. WebGestalt 2019: Gene Set Analysis Toolkit with Revamped UIs and APIs. *Nucleic Acids Res* **2019**, *47*, W199–W205, doi:10.1093/nar/gkz401.
70. Khan, M.I.; Dębski, K.J.; Dabrowski, M.; Czarnecka, A.M.; Szczylik, C. Gene Set Enrichment Analysis and Ingenuity Pathway Analysis of Metastatic Clear Cell Renal Cell Carcinoma Cell Line. *American Journal of Physiology-Renal Physiology* **2016**, *311*, F424–F436, doi:10.1152/ajprenal.00138.2016.

71. Guo, K.; Li, L. Differential ^{12}C -/ ^{13}C -Isotope Dansylation Labeling and Fast Liquid Chromatography/Mass Spectrometry for Absolute and Relative Quantification of the Metabolome. *Anal Chem* **2009**, *81*, 3919–3932, doi:10.1021/ac900166a.
72. Zhao, S.; Li, L. Chemical Derivatization in LC-MS-Based Metabolomics Study. *TrAC Trends in Analytical Chemistry* **2020**, *131*, 115988, doi:10.1016/j.trac.2020.115988.
73. Zhao, S.; Li, H.; Han, W.; Chan, W.; Li, L. Metabolomic Coverage of Chemical-Group-Submetabolome Analysis: Group Classification and Four-Channel Chemical Isotope Labeling LC-MS. *Anal Chem* **2019**, *91*, 12108–12115, doi:10.1021/acs.analchem.9b03431.
74. Zardini Buzatto, A.; Abdel Jabar, M.; Nizami, I.; Dasouki, M.; Li, L.; Abdel Rahman, A.M. Lipidome Alterations Induced by Cystic Fibrosis, CFTR Mutation, and Lung Function. *J Proteome Res* **2021**, *20*, 549–564, doi:10.1021/acs.jproteome.0c00556.
75. Zardini Buzatto, A.; Tatlay, J.; Bajwa, B.; Mung, D.; Camicioli, R.; Dixon, R.A.; Li, L. Comprehensive Serum Lipidomics for Detecting Incipient Dementia in Parkinson's Disease. *J Proteome Res* **2021**, *20*, 4053–4067, doi:10.1021/acs.jproteome.1c00374.
76. Kind, T.; Liu, K.-H.; Lee, D.Y.; DeFelice, B.; Meissen, J.K.; Fiehn, O. LipidBlast in Silico Tandem Mass Spectrometry Database for Lipid Identification. *Nat Methods* **2013**, *10*, 755–758, doi:10.1038/nmeth.2551.
77. Wishart, D.S.; Guo, A.; Oler, E.; Wang, F.; Anjum, A.; Peters, H.; Dizon, R.; Sayeeda, Z.; Tian, S.; Lee, B.L.; et al. HMDB 5.0: The Human Metabolome Database for 2022. *Nucleic Acids Res* **2022**, *50*, D622–D631, doi:10.1093/nar/gkab1062.
78. Aoyagi, R.; Ikeda, K.; Isobe, Y.; Arita, M. Comprehensive Analyses of Oxidized Phospholipids Using a Measured MS/MS Spectra Library. *J Lipid Res* **2017**, *58*, 2229–2237, doi:10.1194/jlr.D077123.
79. Ma, Y.; Kind, T.; Vaniya, A.; Gennity, I.; Fahrman, J.F.; Fiehn, O. An in Silico MS/MS Library for Automatic Annotation of Novel FAHFA Lipids. *J Cheminform* **2015**, *7*, 53, doi:10.1186/s13321-015-0104-4.
80. Fattuoni, C.; Mandò, C.; Palmas, F.; Anelli, G.M.; Novielli, C.; Parejo Laudicina, E.; Savasi, V.M.; Barberini, L.; Dessì, A.; Pintus, R.; et al. Preliminary Metabolomics Analysis of Placenta in Maternal Obesity. *Placenta* **2018**, *61*, 89–95, doi:10.1016/j.placenta.2017.11.014.
81. Brown, S.H.J.; Eather, S.R.; Freeman, D.J.; Meyer, B.J.; Mitchell, T.W. A Lipidomic Analysis of Placenta in Preeclampsia: Evidence for Lipid Storage. *PLoS One* **2016**, *11*, e0163972, doi:10.1371/journal.pone.0163972.
82. Amusquivar, E.; Herrera, E. Influence of Changes in Dietary Fatty Acids during Pregnancy on Placental and Fetal Fatty Acid Profile in the Rat. *Neonatology* **2003**, *83*, 136–145, doi:10.1159/000067963.
83. Ferchaud-Roucher, V.; Barner, K.; Jansson, T.; Powell, T.L. Maternal Obesity Results in Decreased Syncytiotrophoblast Synthesis of Palmitoleic Acid, a Fatty Acid with Anti-inflammatory and Insulin-sensitizing Properties. *The FASEB Journal* **2019**, *33*, 6643–6654, doi:10.1096/fj.201802444R.
84. Cruz, M.M.; Simão, J.J.; Sá, R.D.C.C. de; Farias, T.S.M.; Silva, V.S. da; Abdala, F.; Antraco, V.J.; Armelin-Correa, L.; Alonso-Vale, M.I.C. Palmitoleic Acid Decreases Non-Alcoholic Hepatic Steatosis and Increases Lipogenesis and Fatty Acid Oxidation in Adipose Tissue From Obese Mice. *Front Endocrinol (Lausanne)* **2020**, *11*, doi:10.3389/fendo.2020.537061.
85. Cruz, M.M.; Lopes, A.B.; Crisma, A.R.; de Sá, R.C.C.; Kuwabara, W.M.T.; Curi, R.; de Andrade, P.B.M.; Alonso-Vale, M.I.C. Palmitoleic Acid (16:1n7) Increases Oxygen Consumption, Fatty Acid Oxidation and ATP Content in White Adipocytes. *Lipids Health Dis* **2018**, *17*, 55, doi:10.1186/s12944-018-0710-z.
86. Bláhová; Harvey; Pšenička; Mráz Assessment of Fatty Acid Desaturase (Fads2) Structure-Function Properties in Fish in the Context of Environmental Adaptations and as a Target for Genetic Engineering. *Biomolecules* **2020**, *10*, 206, doi:10.3390/biom10020206.
87. Dave, K.; Kaur, L.; Sundrani, D.; Sharma, P.; Bayyana, S.; Mehendale, S.; Randhir, K.; Chandak, G.R.; Joshi, S. Association of Placental Fatty Acid Desaturase 2 (FADS2) Methylation with Maternal Fatty Acid Levels in Women with Preeclampsia. *Prostaglandins Leukot Essent Fatty Acids* **2022**, *184*, 102472, doi:10.1016/j.plefa.2022.102472.
88. Muralidharan, J.; Papandreou, C.; Sala-Vila, A.; Rosique-Esteban, N.; Fitó, M.; Estruch, R.; Angel Martínez-González, M.; Corella, D.; Ros, E.; Razquín, C.; et al. Fatty Acids Composition of Blood Cell Membranes and Peripheral Inflammation in the PREDIMED Study: A Cross-Sectional Analysis. *Nutrients* **2019**, *11*, 576, doi:10.3390/nu11030576.
89. Kapoor, R.; Huang, Y.-S. Gamma Linolenic Acid: An Antiinflammatory Omega-6 Fatty Acid. *Curr Pharm Biotechnol* **2006**, *7*, 531–534, doi:10.2174/138920106779116874.
90. Colvin, B.N.; Longtine, M.S.; Chen, B.; Costa, M.L.; Nelson, D.M. Oleate Attenuates Palmitate-Induced Endoplasmic Reticulum Stress and Apoptosis in Placental Trophoblasts. *Reproduction* **2017**, *153*, 369–380, doi:10.1530/REP-16-0576.
91. Natarajan, S.K.; Bruett, T.; Muthuraj, P.G.; Sahoo, P.K.; Power, J.; Mott, J.L.; Hanson, C.; Anderson-Berry, A. Saturated Free Fatty Acids Induce Placental Trophoblast Lipoapoptosis. *PLoS One* **2021**, *16*, e0249907, doi:10.1371/journal.pone.0249907.
92. Watkins, O.C.; Islam, M.O.; Selvam, P.; Pillai, R.A.; Cazenave-Gassiot, A.; Bendt, A.K.; Karnani, N.; Godfrey, K.M.; Lewis, R.M.; Wenk, M.R.; et al. Metabolism of ^{13}C -Labeled Fatty Acids in Term Human

- Placental Explants by Liquid Chromatography–Mass Spectrometry. *Endocrinology* **2019**, *160*, 1394–1408, doi:10.1210/en.2018-01020.
93. Colvin, B.N.; Longtine, M.S.; Chen, B.; Costa, M.L.; Nelson, D.M. Oleate Attenuates Palmitate-Induced Endoplasmic Reticulum Stress and Apoptosis in Placental Trophoblasts. *Reproduction* **2017**, *153*, 369–380, doi:10.1530/REP-16-0576.
 94. Campbell, F.M.; Clohessy, A.M.; Gordon, M.J.; Page, K.R.; Dutta-Roy, A.K. Uptake of Long Chain Fatty Acids by Human Placental Choriocarcinoma (BeWo) Cells: Role of Plasma Membrane Fatty Acid-Binding Protein. *J Lipid Res* **1997**, *38*, 2558–2568.
 95. Tobin, K.A.R.; Johnsen, G.M.; Staff, A.C.; Duttaroy, A.K. Long-Chain Polyunsaturated Fatty Acid Transport across Human Placental Choriocarcinoma (BeWo) Cells. *Placenta* **2009**, *30*, 41–47, doi:10.1016/j.placenta.2008.10.007.
 96. Ferchaud-Roucher, V.; Rudolph, M.C.; Jansson, T.; Powell, T.L. Fatty Acid and Lipid Profiles in Primary Human Trophoblast over 90 h in Culture. *Prostaglandins Leukot Essent Fatty Acids* **2017**, *121*, 14–20, doi:10.1016/j.plefa.2017.06.001.
 97. Mashek, D.G.; McKenzie, M.A.; Van Horn, C.G.; Coleman, R.A. Rat Long Chain Acyl-CoA Synthetase 5 Increases Fatty Acid Uptake and Partitioning to Cellular Triacylglycerol in McArdle-RH7777 Cells. *Journal of Biological Chemistry* **2006**, *281*, 945–950, doi:10.1074/jbc.M507646200.
 98. Dutta-Roy, A.K. Cellular Uptake of Long-Chain Fatty Acids: Role of Membrane-Associated Fatty-Acid-Binding/Transport Proteins. *Cellular and Molecular Life Sciences* **2000**, *57*, 1360–1372, doi:10.1007/PL00000621.
 99. Johnsen, G.M.; Weedon-Fekjær, M.S.; Tobin, K.A.R.; Staff, A.C.; Duttaroy, A.K. Long-Chain Polyunsaturated Fatty Acids Stimulate Cellular Fatty Acid Uptake in Human Placental Choriocarcinoma (BeWo) Cells. *Placenta* **2009**, *30*, 1037–1044, doi:10.1016/j.placenta.2009.10.004.
 100. Bildirici, I.; Schaiff, W.T.; Chen, B.; Morizane, M.; Oh, S.-Y.; O'Brien, M.; Sonnenberg-Hirche, C.; Chu, T.; Barak, Y.; Nelson, D.M.; et al. PLIN2 Is Essential for Trophoblastic Lipid Droplet Accumulation and Cell Survival During Hypoxia. *Endocrinology* **2018**, *159*, 3937–3949, doi:10.1210/en.2018-00752.
 101. Stirm, L.; Kovářová, M.; Perschbacher, S.; Michlmaier, R.; Fritsche, L.; Siegel-Axel, D.; Schleicher, E.; Peter, A.; Pauluschke-Fröhlich, J.; Brucker, S.; et al. BMI-Independent Effects of Gestational Diabetes on Human Placenta. *J Clin Endocrinol Metab* **2018**, *103*, 3299–3309, doi:10.1210/jc.2018-00397.
 102. Adamo, K.B.; Dent, R.; Langefeld, C.D.; Cox, M.; Williams, K.; Carrick, K.M.; Stuart, J.S.; Sundseth, S.S.; Harper, M.-E.; McPherson, R.; et al. Peroxisome Proliferator-Activated Receptor γ 2 and Acyl-CoA Synthetase 5 Polymorphisms Influence Diet Response*. *Obesity* **2007**, *15*, 1068–1075, doi:10.1038/oby.2007.630.
 103. Schubert, M.; Becher, S.; Wallert, M.; Maeß, M.B.; Abhari, M.; Rennert, K.; Mosig, A.S.; Große, S.; Heller, R.; Grün, M.; et al. The Peroxisome Proliferator-Activated Receptor (PPAR)- γ Antagonist 2-Chloro-5-Nitro-N-Phenylbenzamide (GW9662) Triggers Perilipin 2 Expression via PPAR δ and Induces Lipogenesis and Triglyceride Accumulation in Human THP-1 Macrophages. *Mol Pharmacol* **2020**, *97*, 212–225, doi:10.1124/mol.119.117887.
 104. Ikeda, J.; Ichiki, T.; Takahara, Y.; Kojima, H.; Sankoda, C.; Kitamoto, S.; Tokunou, T.; Sunagawa, K. PPAR γ Agonists Attenuate Palmitate-Induced ER Stress through Up-Regulation of SCD-1 in Macrophages. *PLoS One* **2015**, *10*, e0128546, doi:10.1371/journal.pone.0128546.
 105. Yao-Borengasser, A.; Rassouli, N.; Varma, V.; Bodles, A.M.; Rasouli, N.; Unal, R.; Phanavanh, B.; Ranganathan, G.; McGehee, R.E.; Kern, P.A. Stearoyl-Coenzyme A Desaturase 1 Gene Expression Increases after Pioglitazone Treatment and Is Associated with Peroxisomal Proliferator-Activated Receptor- γ Responsiveness. *J Clin Endocrinol Metab* **2008**, *93*, 4431–4439, doi:10.1210/jc.2008-0782.
 106. Brunmeir, R.; Xu, F. Functional Regulation of PPARs through Post-Translational Modifications. *Int J Mol Sci* **2018**, *19*, 1738, doi:10.3390/ijms19061738.
 107. Nakagawa, Y.; Satoh, A.; Tezuka, H.; Han, S.; Takei, K.; Iwasaki, H.; Yato, S.; Yahagi, N.; Suzuki, H.; Iwasaki, Y.; et al. CREB3L3 Controls Fatty Acid Oxidation and Ketogenesis in Synergy with PPAR α . *Sci Rep* **2016**, *6*, 39182, doi:10.1038/srep39182.
 108. Wade, H.; Pan, K.; Su, Q. CREBH: A Complex Array of Regulatory Mechanisms in Nutritional Signaling, Metabolic Inflammation, and Metabolic Disease. *Mol Nutr Food Res* **2021**, *65*, 2000771, doi:10.1002/mnfr.202000771.
 109. Sampieri, L.; Di Giusto, P.; Alvarez, C. CREB3 Transcription Factors: ER-Golgi Stress Transducers as Hubs for Cellular Homeostasis. *Front Cell Dev Biol* **2019**, *7*, doi:10.3389/fcell.2019.00123.
 110. Zhang, K.; Shen, X.; Wu, J.; Sakaki, K.; Saunders, T.; Rutkowski, D.T.; Back, S.H.; Kaufman, R.J. Endoplasmic Reticulum Stress Activates Cleavage of CREBH to Induce a Systemic Inflammatory Response. *Cell* **2006**, *124*, 587–599, doi:10.1016/j.cell.2005.11.040.
 111. Saben, J.; Zhong, Y.; Gomez-Acevedo, H.; Thakali, K.M.; Borengasser, S.J.; Andres, A.; Shankar, K. Early Growth Response Protein-1 Mediates Lipotoxicity-Associated Placental Inflammation: Role in Maternal Obesity. *American Journal of Physiology-Endocrinology and Metabolism* **2013**, *305*, E1–E14, doi:10.1152/ajpendo.00076.2013.

112. Lien, Y.-C.; Zhang, Z.; Barila, G.; Green-Brown, A.; Elovitz, M.A.; Simmons, R.A. Intrauterine Inflammation Alters the Transcriptome and Metabolome in Placenta. *Front Physiol* **2020**, *11*, doi:10.3389/fphys.2020.592689.
113. Stincone, A.; Prigione, A.; Cramer, T.; Wamelink, M.M.C.; Campbell, K.; Cheung, E.; Olin-Sandoval, V.; Grüning, N.; Krüger, A.; Tauqeer Alam, M.; et al. The Return of Metabolism: Biochemistry and Physiology of the Pentose Phosphate Pathway. *Biological Reviews* **2015**, *90*, 927–963, doi:10.1111/brv.12140.
114. Moe, A.J.; Farmer, D.R.; Nelson, D.M.; Smith, C.H. Pentose Phosphate Pathway in Cellular Trophoblasts from Full-Term Human Placentas. *American Journal of Physiology-Cell Physiology* **1991**, *261*, C1042–C1047, doi:10.1152/ajpcell.1991.261.6.C1042.
115. Beaconsfield, P.; Ginsburg, J.; Jeacock, M.K. Glucose Metabolism via the Pentose Phosphate Pathway Relative to Nucleic Acid and Protein Synthesis in the Human Placenta. *Dev Med Child Neurol* **2008**, *6*, 469–474, doi:10.1111/j.1469-8749.1964.tb10811.x.
116. Riganti, C.; Gazzano, E.; Polimeni, M.; Aldieri, E.; Ghigo, D. The Pentose Phosphate Pathway: An Antioxidant Defense and a Crossroad in Tumor Cell Fate. *Free Radic Biol Med* **2012**, *53*, 421–436, doi:10.1016/j.freeradbiomed.2012.05.006.
117. Grant, C.M. Metabolic Reconfiguration Is a Regulated Response to Oxidative Stress. *J Biol* **2008**, *7*, 1, doi:10.1186/jbiol63.
118. Duan, Y.; Li, F.; Li, Y.; Tang, Y.; Kong, X.; Feng, Z.; Anthony, T.G.; Watford, M.; Hou, Y.; Wu, G.; et al. The Role of Leucine and Its Metabolites in Protein and Energy Metabolism. *Amino Acids* **2016**, *48*, 41–51, doi:10.1007/s00726-015-2067-1.
119. Longo, N.; Frigeni, M.; Pasquali, M. Carnitine Transport and Fatty Acid Oxidation. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **2016**, *1863*, 2422–2435, doi:10.1016/j.bbamcr.2016.01.023.
120. Lambertucci, R.H.; Hirabara, S.M.; Silveira, L. dos R.; Levada-Pires, A.C.; Curi, R.; Pithon-Curi, T.C. Palmitate Increases Superoxide Production through Mitochondrial Electron Transport Chain and NADPH Oxidase Activity in Skeletal Muscle Cells. *J Cell Physiol* **2008**, *216*, 796–804, doi:10.1002/jcp.21463.
121. Rosca, M.G.; Vazquez, E.J.; Chen, Q.; Kerner, J.; Kern, T.S.; Hoppel, C.L. Oxidation of Fatty Acids Is the Source of Increased Mitochondrial Reactive Oxygen Species Production in Kidney Cortical Tubules in Early Diabetes. *Diabetes* **2012**, *61*, 2074–2083, doi:10.2337/db11-1437.
122. Zhao, M.; Wang, Y.; Li, L.; Liu, S.; Wang, C.; Yuan, Y.; Yang, G.; Chen, Y.; Cheng, J.; Lu, Y.; et al. Mitochondrial ROS Promote Mitochondrial Dysfunction and Inflammation in Ischemic Acute Kidney Injury by Disrupting TFAM-Mediated MtDNA Maintenance. *Theranostics* **2021**, *11*, 1845–1863, doi:10.7150/thno.50905.
123. Rizwan, H.; Pal, S.; Sabnam, S.; Pal, A. High Glucose Augments ROS Generation Regulates Mitochondrial Dysfunction and Apoptosis via Stress Signalling Cascades in Keratinocytes. *Life Sci* **2020**, *241*, 117148, doi:10.1016/j.lfs.2019.117148.
124. Honda, A.; Yamashita, K.; Ikegami, T.; Hara, T.; Miyazaki, T.; Hirayama, T.; Numazawa, M.; Matsuzaki, Y. Highly Sensitive Quantification of Serum Malonate, a Possible Marker for de Novo Lipogenesis, by LC-ESI-MS/MS. *J Lipid Res* **2009**, *50*, 2124–2130, doi:10.1194/jlr.D800054-JLR200.
125. Rutkowski, J.M.; Knotts, T.A.; Ono-Moore, K.D.; McCoin, C.S.; Huang, S.; Schneider, D.; Singh, S.; Adams, S.H.; Hwang, D.H. Acylcarnitines Activate Proinflammatory Signaling Pathways. *Am J Physiol Endocrinol Metab* **2014**, *306*, E1378–87, doi:10.1152/ajpendo.00656.2013.
126. Koves, T.R.; Ussher, J.R.; Noland, R.C.; Slentz, D.; Mosedale, M.; Ilkayeva, O.; Bain, J.; Stevens, R.; Dyck, J.R.B.; Newgard, C.B.; et al. Mitochondrial Overload and Incomplete Fatty Acid Oxidation Contribute to Skeletal Muscle Insulin Resistance. *Cell Metab* **2008**, *7*, 45–56, doi:10.1016/j.cmet.2007.10.013.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.