

Case Report

Not peer-reviewed version

# Can Hepatitis B Virus (Hbv) Reactivation Be the Result of a Mild COVID-19 Infection With No Need of Systemic Immunosuppressive Therapy?

[Ioannis Braimakis](#) , Sofia Vasileiadi , [Eleni-Myrto Trifylli](#) , [Nikolaos Papadopoulos](#) <sup>\*</sup> , [Melanie Deutsch](#)

Posted Date: 19 June 2023

doi: 10.20944/preprints202306.1305.v1

Keywords: HBV reactivation; Covid-19; immunosuppressive therapy



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Case report*

# Can Hepatitis B Virus (HBV) Reactivation Be the Result of a Mild Covid-19 Infection with No Need of Systemic Immunosuppressive Therapy?

Ioannis Braimakis <sup>1,\*</sup>, Sofia Vasileiadi <sup>2</sup>, Eleni-Myrto Trifylli <sup>3</sup>, Nikolaos Papadopoulos <sup>4</sup> and Melanie Deutsch <sup>2</sup>

<sup>1</sup> Gastroenterology Department, 417 Army Share Fund Hospital of Athens, 11521 Athens, Greece

<sup>2</sup> 2nd Department of Internal Medicine, Hippokration General Hospital of Athens, Medical School of National & Kapodistrian University of Athens, 11527 Athens, Greece

<sup>3</sup> 1st Department of Internal Medicine, 417 Army Share Fund Hospital of Athens, 11521 Athens, Greece

<sup>4</sup> 2nd Department of Internal Medicine, 401 General Army Hospital of Athens, 11525 Athens, Greece

\* Correspondence: giannis\_br@hotmail.com; Tel.: +306984168389

**Abstract:** Hepatitis B virus reactivation (HBVr) is a well described result of immunosuppressive therapy initiation in a variety of diseases with dose and duration of treatment being the main factors determining the probability for reactivation. Such cases have been described also in Covid-19 patients treated with immunosuppressive therapies. Nevertheless, there have also been reported cases of Covid-19 infection that led to HBVr with no concurrent immunosuppressive therapy or any other described cause. In accordance with that observation, we present a patient followed for a period spanning 20 years with HBeAg negative chronic HBV infection and non-detectable HBV DNA who after a mild Covid-19 infection treated only with low dose and short duration inhaled corticosteroids (ICS), developed elevated AST and ALT as well as elevated HBV DNA levels. During the diagnostic workout other etiologies of abnormal liver biochemistries were excluded and thus the diagnosis of HBV reactivation was established and treated with entecavir. Since other causes of reactivation were excluded, and ICS dose and duration was found baring only a very low risk (<1%) for HBVr, Covid-19 infection could be considered the most probable cause of reactivation.

**Keywords:** HBV reactivation; Covid-19; immunosuppressive therapy

## 1. Introduction

Hepatitis B virus (HBV), part of the Hepadnaviridae family [3], is responsible for about 296 million chronic hepatitis infections according to latest World Health Organization (WHO) data [1]. Following HBV infection and based on the interaction between HBV and the host's immune system response, the acute infection can either resolve or progress to chronic hepatitis B [3]. Current nomenclature for HBV chronic infection marks the presence of five separate phases, not always as an infection sequence, but based on HBeAg seropositivity as well as serum levels of HBV DNA and alanine aminotransferase (ALT) [3]. Thus, the following phases arise: HBeAg (+) chronic HBV infection, HBeAg (+) chronic hepatitis B, HBeAg (-) chronic HBV infection, HBeAg (-) chronic hepatitis B and HBsAg-negative phase [2,3]. Common ground in chronic Hepatitis B or infection and regardless of the phase, is the persistent presence of the covalently closed circular DNA (cccDNA) inside hepatocytes nucleus [5]. It is cccDNA that acts as the template for HBV transcripts that are subsequently translated into the main proteins of the virus [3,4], remaining also the most important cause of HBV reactivation (HBVr) at any time and any phase especially under conditions of immunosuppression [6-9].

As of late 2019, Covid-19 pandemic has proved a serious global public health issue. The responsible virus itself -SARS-CoV-2 not only affects the respiratory system but as already reported in several studies it also causes liver injury with abnormal liver biochemistry [10,11,12,16]. Multiple possible liver injury mechanisms have been proposed, such as direct cytopathic effect of SARS-CoV-

2 through angiotensin-converting enzyme 2 (ACE2) and transmembrane serine protease 2 (TMPRSS2) receptors of cholangiocytes and hepatocytes [13]. Other mechanisms include liver injury caused by severe inflammation, hypoxia, toxicity of the drugs used for Covid-19 treatment and vascular changes attributed to coagulopathy [14,15].

Taking into consideration that both HBV and SARS-CoV-2 can potentially cause liver injury, several studies attempted to investigate coinfection cases and highlight the effect of the latter on chronic HBV natural history on one hand, and the effect of HBV seropositivity on the severity of Covid-19 infection on the other hand [17]. Attempting to help address this controversial matter we report a case of a patient presenting with HBV reactivation soon after Covid-19 infection without administration of systemic immunosuppressive medication.

3. Case report

A 49-year-old female presented in the outpatient Department with elevated serum AST and ALT level. The patient has been followed for HBeAg negative chronic HBV infection every six months for twenty years presenting always normal liver enzymes and non-detectable HBV DNA Apart from that, patient’s prior medical history did not include any other health problem in need of long-term medical treatment, while no alcohol consumption, no medications over the counter and no smoking was mentioned.

The patient reported symptoms indicative of respiratory tract infection 3 weeks prior to her presentation. She was tested positive for Covid-19 infection and due to mild symptoms only a combination of budesonide 160 µg plus formoterol fumarate dehydrate 4.5 µg, for a duration of five days were prescribed. Over the course of SARS-CoV-2 infection, the patient experienced no complications and had no need of hospitalization or further treatment.

At her first hepatology consultation appointment, apart from abnormal liver chemistries, the patient experienced no other symptoms. More specifically, in the blood tests presented by the patient, aminotransferases were mildly elevated with AST level double the upper limit of normal (ULN) and ALT level triple the ULN (Table 1). No other important deviations from normal were noted in the blood tests presented (Table 1). Liver and hepatitis panel tests as well as upper abdominal ultrasound were prescribed and at her second appointment, one week later, the patient presented with the results. Persisting elevated AST and ALT levels double and triple the ULN respectively were observed (Table 1). As for her Hepatitis B known status the patient was HBsAg-positive, HBeAg-negative, anti-HBe-positive and anti-HBc-positive with HBV-DNA at 9.350.000 iu/ml.

Table 1. Blood test results.

	1 <sup>st</sup> consultation	2 <sup>nd</sup> consultation	2 months after entecavir
AST (U/L)	82	71	28
ALT (U/L)	106	90	30
ALP (U/L)	106	80	84
γGT (U/L)	16	19	20

Moreover, the patient was tested negative for anti-HCV antibodies, anti-HDV antibodies, antinuclear antibodies (ANA), antimitochondrial antibodies (AMA), anti-smooth muscle antibodies (ASMA) and anti-dsDNA antibodies (Table 2). Immunoglobulin levels showed no deviation from normal (Table 2). The upper abdominal ultrasound revealed no abnormalities and alpha-fetoprotein (a-FP) levels were normal. Abnormal liver chemistries were attributed to HBV reactivation and thus entecavir was prescribed.

Table 2. Trend over time of HBV serologic markers, anti-HCV and anti-HDV.

	Prior six months	2 <sup>nd</sup> consultation	2 months after entecavir
--	------------------	------------------------------	--------------------------

HBsAg	+	+	
HBeAg	-	-	
Anti-HBc	+	+	
Anti-HBe	+	+	
Anti-HCV	-	-	
Anti-HDV	-	-	
HBV DNA (IU/ml)	Non-Detectable	9.350.000	54.000
IgG (mg/dl)		1555	

Two months after the initiation of entecavir, AST and ALT levels had normalized (Table 1) whereas HBV DNA levels demonstrated a decreasing trend (Table 2). The patient remains asymptomatic under close surveillance.

#### 4. Discussion

We present a case of HBV reactivation (HBVr) in a female patient with a 20-year history of chronic HBeAg negative HBV infection occurring after a mild Covid-19 infection. Usually, HBVr occurs when the host immune system loses the ability to suppress the virus replication [24]. Both HBsAg-positive/anti-HBc-positive and HBsAg-negative/anti-HBc-positive patients preserve the potential of HBVr [25] under certain circumstances. These comprise either concomitant use of immunosuppressive therapies or variable viral interactions. Such interaction is well described in HBV/HCV coinfection cases, where HCV usually suppresses HBV replication leading to low levels of HBV DNA [39-42]. Consequently, a Direct-Acting antiviral (DAA) therapy aiming for sustained virologic response (SVR) of HCV infection may lead to HBV reactivation in cases where no concurrent nucleoside/nucleotide analogue (NA) HBV therapy is applied [39,40].

Regarding the definition of HBVr a variety of variants have been proposed and used in different studies [23]. We used the latest proposed nomenclature presented by Papatheodoridis, et al. (2022), at their latest systematic review, meta-analysis, and expert opinion [6]. Accordingly, our patient with a known chronic HBV infection (HBsAg-positive) diagnosed 20 years ago that was currently found with elevated levels of HBV DNA  $\geq 10.000$  IU/ml, met the proposed criteria for HBVr [6].

In our patient, to ensure the accuracy of HBVr diagnosis, other causes of viral hepatitis such as HAV, HEV, HCV and HDV were excluded. Moreover, a diagnostic workup for other etiologies of abnormal liver biochemistry such as autoimmune hepatitis (AIH) and Primary Biliary Cholangitis (PBC) using non-organ specific autoantibodies such as ANA, ASMA and AMA was carried out [18-20]. Imaging with upper abdominal ultrasound was also used to rule out focal liver lesions as a cause for elevated AST and ALT in combination with normal levels of a-FP [21,22].

Following the establishment of the diagnosis of HBVr, the main question arising concerned the pathophysiology of HBVr 20 years after the initial diagnosis. In general, the main reason leading to impaired ability to suppress HBV replication and thus to a possible reactivation is the initiation of immunosuppressive treatment [6,23]. A broad spectrum of immunosuppressants has been incriminated for HBVr [23]. Among others, corticosteroid treatment for any indication including Covid-19 has been in the spotlight of studies investigating its ability to induce HBVr [26-28]. This ability is brought about by the activation of a DNA sequence known as Glucocorticoid Response Element (GRE) enclosed in HBV genome on one hand [29], and by immunosuppressive properties of corticosteroids on the other hand [23]. Nevertheless, reactivation probability depends on the dose and treatment duration of corticosteroids [6,23]. The American Gastroenterological Association addressed this matter [23] by categorizing the HBVr risk induced by immunosuppressants as high when the risk of reactivation is expected to be more than 10%, moderate when frequency of reactivation is between 1% and 10% and low when reactivation risk is  $< 1\%$ . Thus, for HBsAg-positive patients a moderate (10–20 mg) or high ( $>20$  mg) dose of prednisone (or equivalent) for  $\geq 4$  weeks results in a high risk for HBVr, a low dose ( $<10$  mg) of prednisone (or equivalent) for  $\geq 4$  weeks results in a moderate risk for HBVr while corticosteroid therapy for  $\leq 1$  week results in low risk for HBVr

[23]. Moreover, a dose >20mg for a treatment duration >2 weeks results in significant immunosuppression [30].

In our patient, as already mentioned, the course of Covid-19 infection was mild and as a result only inhaled budesonide 160 µg plus formoterol fumarate dehydrate 4.5 µg, for a five-day duration was prescribed. Could this possibly be the reason of HBVr in our case? Data regarding inhaled corticosteroids (ICSs) derive from studies addressing their efficacy and safety profile especially in asthma and chronic obstructive pulmonary disease. In general, ICS's action is mainly applied locally at the airway level and is effective in low doses [31]. Apart from that, according to a study from Majers et.al. (2020) an increase of 1000 µg of fluticasone propionate is commensurate with 5mg of prednisone, while 1000 µg of budesonide is commensurate with 2mg of prednisone [32]. Thus, even the high doses used in severe respiratory disorders such as budesonide dose of up to 1600 µg/day or fluticasone propionate dose of up to 1000 µg/day [32] correspond to low dose (<10mg) of prednisone. As a result, HBVr risk with ICS courses administered for less than a 1week, as in our patient, is lower than 1%.

As ICS administration did not seem to explain the sudden HBVr of our patient, a review of the literature was carried out in order to investigate the possibility of HBVr induced by Covid-19. Most cases of HBVr after SARS-CoV-2 infection were attributed to immunosuppressive treatment such as dexamethasone 6mg per day [5], other systemic corticosteroids like methylprednisolone 40mg per day [17,34] or tocilizumab which was used in severe cases of Covid-19 [35]. The aforementioned systemic corticosteroid doses correspond to a high dose of prednisone (40mg of prednisone equivalent for dexamethasone and > 40mg of prednisone equivalent for methylprednisolone) [36] that even when administered for <7 days, can result in increasing risk for hepatitis flare [27,33].

Interestingly enough, we also came across patient cases where Covid-19 was considered to be the trigger for HBVr with other causes of reactivation excluded [14,17,37,38]. The pathogenesis pathway for this reactivation in patients with HBV and SARS-CoV-2 coinfection remains to be elucidated, but the disruption of the balance of immune system activity on one hand and HBV replication on the other has been proposed as a possible mechanism [14].

## 5. Conclusions

In conclusion, due to the low risk for HBVr provoked by ICS, and in accordance with other cases presented, our patient's HBVr was attributed to the recent Covid-19 infection, underlining the need for close monitoring of patients who are HBsAg -positive in case of infection with SARS-CoV-2, even when no systemic immunosuppressive treatment is administered. In order to answer the question if HBVr may be the result of viral interaction with a non-hepatotropic virus like SARS-CoV-2 more research is needed.

**Author Contributions:** Authors contributed equally to this manuscript. Conceptualization, I.B., N.P. and M.D.; validation, I.B., N.P. and M.D.; investigation, I.B., S.V. and E-M.T.; resources, I.B.; data curation, I.B.; writing—original draft preparation, I.B., S.V. and E-M.T.; writing—review and editing, N.P. and M.D.; visualization, I.B.; supervision, M.D. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding

**Institutional Review Board Statement:** Not applicable

**Informed Consent Statement:** Written informed consent has been obtained from the patient to publish this paper.

**Data Availability Statement:** Data is contained within the article.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. World Health Organization. (2022, June 24). Hepatitis B. Retrieved from: <https://www.who.int/news-room/fact-sheets/detail/hepatitis-b>



2. Spera AM. Hepatitis B virus infection reactivation in patients under immunosuppressive therapies: Pathogenesis, screening, prevention and treatment. *World J Virol.* 2022;11(5):275-282. doi:10.5501/wjv.v11.i5.275
3. European Association for the Study of the Liver. Electronic address: easloffice@easloffice.eu; European Association for the Study of the Liver. EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol.* 2017;67(2):370-398. doi:10.1016/j.jhep.2017.03.021
4. Lucifora J, Protzer U. Attacking hepatitis B virus cccDNA--The holy grail to hepatitis B cure. *J Hepatol.* 2016;64(1 Suppl):S41-S48. doi:10.1016/j.jhep.2016.02.009
5. Sagnelli C, Montella L, Grimaldi P, et al. COVID-19 as Another Trigger for HBV Reactivation: Clinical Case and Review of Literature. *Pathogens.* 2022;11(7):816. Published 2022 Jul 21. doi:10.3390/pathogens11070816
6. Papatheodoridis GV, Lekakis V, Voulgaris T, et al. Hepatitis B virus reactivation associated with new classes of immunosuppressants and immunomodulators: A systematic review, meta-analysis, and expert opinion. *J Hepatol.* 2022;77(6):1670-1689. doi:10.1016/j.jhep.2022.07.003
7. Moghooei M, Mostafaei S, Ashraf-Ganjouei A, Kavosi H, Mahmoudi M. HBV reactivation in rheumatic diseases patients under therapy: A meta-analysis. *Microb Pathog.* 2018;114:436-443. doi:10.1016/j.micpath.2017.12.014
8. Sagnelli C, Pisaturo M, Calò F, Martini S, Sagnelli E, Coppola N. Reactivation of hepatitis B virus infection in patients with hemo-lymphoproliferative diseases, and its prevention. *World J Gastroenterol.* 2019;25(26):3299-3312. doi:10.3748/wjg.v25.i26.3299
9. Mori S, Fujiyama S. Hepatitis B virus reactivation associated with antirheumatic therapy: Risk and prophylaxis recommendations. *World J Gastroenterol.* 2015;21(36):10274-10289. doi:10.3748/wjg.v21.i36.10274
10. Zhang C, Shi L, Wang FS. Liver injury in COVID-19: management and challenges. *Lancet Gastroenterol Hepatol.* 2020;5(5):428-430. doi:10.1016/S2468-1253(20)30057-1
11. Sultan S, Altayar O, Siddique SM, et al. AGA Institute Rapid Review of the Gastrointestinal and Liver Manifestations of COVID-19, Meta-Analysis of International Data, and Recommendations for the Consultative Management of Patients with COVID-19. *Gastroenterology.* 2020;159(1):320-334.e27. doi:10.1053/j.gastro.2020.05.001
12. Youssef M, H Hussein M, Attia AS, et al. COVID-19 and liver dysfunction: A systematic review and meta-analysis of retrospective studies. *J Med Virol.* 2020;92(10):1825-1833. doi:10.1002/jmv.26055
13. Pirola CJ, Sookoian S. SARS-CoV-2 virus and liver expression of host receptors: Putative mechanisms of liver involvement in COVID-19. *Liver Int.* 2020;40(8):2038-2040. doi:10.1111/liv.14500
14. Giugliano L, Pinon M, Calvo PL. COVID-19 as a Trigger of Acute-on-Chronic Hepatitis B Presenting With Undetectable INR Due to Hypercoagulability in a 16-Year-Old Girl. *Pediatr Infect Dis J.* 2023;42(2):143-145. doi:10.1097/INF.0000000000003771
15. Baroiu L, Anghel L, Laurențiu Tatu A, et al. Risk of hepatitis B reactivation: From biologic therapies for psoriasis to immunosuppressive therapies for COVID-19 (Review). *Exp Ther Med.* 2022;23(6):385. doi:10.3892/etm.2022.11312
16. Marjot T, Webb GJ, Barritt AS 4th, et al. COVID-19 and liver disease: mechanistic and clinical perspectives. *Nat Rev Gastroenterol Hepatol.* 2021;18(5):348-364. doi:10.1038/s41575-021-00426-4
17. Liu J, Wang T, Cai Q, et al. Longitudinal changes of liver function and hepatitis B reactivation in COVID-19 patients with pre-existing chronic hepatitis B virus infection. *Hepatol Res.* 2020;50(11):1211-1221. doi:10.1111/hepr.13553
18. Carey EJ, Ali AH, Lindor KD. Primary biliary cirrhosis [published correction appears in Lancet. 2015 Oct 17;386(10003):1536]. *Lancet.* 2015;386(10003):1565-1575. doi:10.1016/S0140-6736(15)00154-3
19. Selmi C, Bowlus CL, Gershwin ME, Coppel RL. Primary biliary cirrhosis. *Lancet.* 2011;377(9777):1600-1609. doi:10.1016/S0140-6736(10)61965-4
20. Hennes EM, Zeniya M, Czaja AJ, et al. Simplified criteria for the diagnosis of autoimmune hepatitis. *Hepatology.* 2008;48(1):169-176. doi:10.1002/hep.22322
21. European Association for the Study of the Liver. Electronic address: easloffice@easloffice.eu; European Association for the Study of the Liver. EASL Clinical Practice Guidelines: Management of hepatocellular carcinoma [published correction appears in J Hepatol. 2019 Apr;70(4):817]. *J Hepatol.* 2018;69(1):182-236. doi:10.1016/j.jhep.2018.03.019
22. Marrero JA, Kulik LM, Sirlin CB, et al. Diagnosis, Staging, and Management of Hepatocellular Carcinoma: 2018 Practice Guidance by the American Association for the Study of Liver Diseases. *Hepatology.* 2018;68(2):723-750. doi:10.1002/hep.29913
23. Perrillo RP, Gish R, Falck-Ytter YT. American Gastroenterological Association Institute technical review on prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology.* 2015;148(1):221-244.e3. doi:10.1053/j.gastro.2014.10.038
24. Perrillo RP. Acute flares in chronic hepatitis B: the natural and unnatural history of an immunologically mediated liver disease. *Gastroenterology.* 2001;120(4):1009-1022. doi:10.1053/gast.2001.22461

25. Hoofnagle JH. Reactivation of hepatitis B. *Hepatology*. 2009;49(5 Suppl):S156-S165. doi:10.1002/hep.22945
26. Zhong Z, Liao W, Dai L, et al. Average corticosteroid dose and risk for HBV reactivation and hepatitis flare in patients with resolved hepatitis B infection. *Ann Rheum Dis*. 2022;81(4):584-591. doi:10.1136/annrheumdis-2021-221650
27. Wong GL, Wong VW, Yuen BW, et al. Risk of hepatitis B surface antigen seroreversion after corticosteroid treatment in patients with previous hepatitis B virus exposure. *J Hepatol*. 2020;72(1):57-66. doi:10.1016/j.jhep.2019.08.023
28. Kim TW, Kim MN, Kwon JW, et al. Risk of hepatitis B virus reactivation in patients with asthma or chronic obstructive pulmonary disease treated with corticosteroids. *Respirology*. 2010;15(7):1092-1097. doi:10.1111/j.1440-1843.2010.01798.x
29. Tur-Kaspa R, Shaul Y, Moore DD, et al. The glucocorticoid receptor recognizes a specific nucleotide sequence in hepatitis B virus DNA causing increased activity of the HBV enhancer. *Virology*. 1988;167(2):630-633.
30. Stuck AE, Minder CE, Frey FJ. Risk of infectious complications in patients taking glucocorticosteroids. *Rev Infect Dis*. 1989;11(6):954-963. doi:10.1093/clinids/11.6.954
31. Brattsand R, Selroos O. May a different kinetic mode explain the high efficacy/safety profile of inhaled budesonide?. *Pulm Pharmacol Ther*. 2022;77:102167. doi:10.1016/j.pupt.2022.102167
32. Majiers I, Kearns N, Harper J, Weatherall M, Beasley R. Oral steroid-sparing effect of high-dose inhaled corticosteroids in asthma. *Eur Respir J*. 2020;55(1):1901147. Published 2020 Jan 2. doi:10.1183/13993003.01147-2019
33. Yip TC, Gill M, Wong GL, Liu K. Management of hepatitis B virus reactivation due to treatment of COVID-19. *Hepatol Int*. 2022;16(2):257-268. doi:10.1007/s12072-022-10306-x
34. Wu YF, Yu WJ, Jiang YH, et al. COVID-19 or treatment associated immunosuppression may trigger hepatitis B virus reactivation: A case report. *World J Clin Cases*. 2021;9(19):5266-5269. doi:10.12998/wjcc.v9.i19.5266
35. Campbell C, Andersson MI, Ansari MA, et al. Risk of Reactivation of Hepatitis B Virus (HBV) and Tuberculosis (TB) and Complications of Hepatitis C Virus (HCV) Following Tocilizumab Therapy: A Systematic Review to Inform Risk Assessment in the COVID-19 Era. *Front Med (Lausanne)*. 2021;8:706482. Published 2021 Aug 20. doi:10.3389/fmed.2021.706482
36. Mager DE, Lin SX, Blum RA, Lates CD, Jusko WJ. Dose equivalency evaluation of major corticosteroids: pharmacokinetics and cell trafficking and cortisol dynamics. *J Clin Pharmacol*. 2003;43(11):1216-1227. doi:10.1177/0091270003258651
37. Aldhalei WA, Alnuaimi A, Bhagavathula AS. COVID-19 Induced Hepatitis B Virus Reactivation: A Novel Case From the United Arab Emirates. *Cureus*. 2020;12(6):e8645. Published 2020 Jun 15. doi:10.7759/cureus.8645
38. Libroero Jiménez M, López Garrido MÁ, Fernández Cano MC. Letter to the editor: Reactivation of HBV triggered by SARS-CoV-2 in a patient with cirrhosis. *Hepatology*. 2022;75(3):765-766. doi:10.1002/hep.32271
39. European Association for the Study of the Liver. Electronic address: easloffice@easloffice.eu; Clinical Practice Guidelines Panel: Chair;; EASL Governing Board representative;; Panel members:. EASL recommendations on treatment of hepatitis C: Final update of the series\* [published correction appears in *J Hepatol*. 2023 Feb;78(2):452]. *J Hepatol*. 2020;73(5):1170-1218. doi:10.1016/j.jhep.2020.08.018
40. Wang C, Ji D, Chen J, et al. Hepatitis due to Reactivation of Hepatitis B Virus in Endemic Areas Among Patients With Hepatitis C Treated With Direct-acting Antiviral Agents. *Clin Gastroenterol Hepatol*. 2017;15(1):132-136. doi:10.1016/j.cgh.2016.06.023
41. Papadopoulos N, Papavdi M, Pavlidou A, et al. Hepatitis B and C coinfection in a real-life setting: viral interactions and treatment issues. *Ann Gastroenterol*. 2018;31(3):365-370. doi:10.20524/aog.2018.0255
42. Konstantinou D, Deutsch M. The spectrum of HBV/HCV coinfection: epidemiology, clinical characteristics, viral interactions and management. *Ann Gastroenterol*. 2015;28(2):221-228.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.