

Article

Not peer-reviewed version

Does Omeprazole and Melatonin Enhance Photosynthetic Efficiency in Bean Plants?

[Esteban Sánchez](#)*, Carlos Abel Ramírez-Estrada, [María Antonia Flores-Córdova](#), [Sandra Pérez-Álvarez](#), Linda Citlalli Noperi-Mosqueda, [Celia Chávez-Mendoza](#)

Posted Date: 30 June 2023

doi: 10.20944/preprints202306.2189.v1

Keywords: Phaseolus vulgaris L.; photochemical quenching; benzimidazole; photosystem II; CO₂



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Does Omeprazole and Melatonin Enhance Photosynthetic Efficiency in Bean Plants?

Carlos Abel Ramírez-Estrada ¹, Esteban Sánchez ^{1,*}, María Antonia Flores-Córdova ², Sandra Pérez-Álvarez ³, Linda Citlalli Noperi-Mosqueda ² and Celia Chavez-Mendoza ¹

¹ Food and Development Research Center, A.C. (CIAD) Delicias Unit, 33089 Delicias, Chihuahua, México

² Faculty of Agrotechnological Sciences, Autonomous University of Chihuahua (UACH), 31310 Chihuahua, Chihuahua, México

³ Faculty of Agricultural and Forestry Sciences, Autonomous University of Chihuahua (UACH), 33000 Delicias, Chihuahua, México

* Correspondence: esteban@ciad.mx

Abstract: Current environmental conditions have prompted agri-food systems to explore new strategies for improving the production of crops in a sustainable manner. The use of omeprazole (OMP) and melatonin (MEL) have been proposed as innovative strategy for crop improvement when applied as biostimulants. However, although their role in the growth of several species has been studied, the results in photosynthetic efficiency parameters are still scarce. Therefore, the objective of the present study was to evaluate the effect of foliar application of OMP and MEL on biomass, yield, SPAD values, leaf chlorophyll fluorescence (Fv/Fm), photochemical quenching (qP), Non-photochemical quenching (NPQ), quantum yield of photosystem II (PhiPSII) and electron transport rate (ETR) in bean plants. Treatments were applied separately at doses of 1, 10, and 100 µM, plus a control without application. The results obtained indicate that OMP and MEL were able to increase biomass, yield, SPAD values, as well as photochemical quenching, fluorescence, and PhiPSII coefficients. Finally, it is concluded that foliar application of OMP and MEL at a dose of 1 and 10 µM can increase photosynthetic efficiency and decrease photoinhibition, which is reflected in higher biomass accumulation and yield in green bean plants cv. Strike.

Keywords: *Phaseolus vulgaris* L.; photochemical quenching; benzimidazole; photosystem II; CO₂

1. Introduction

Currently, the global demand for food is increasing at the rate of the world population [1]. Because of this, technological innovations in agriculture are striving to increase yields of highly important crops in a sustainable manner [2]. Some of these innovations include genetic improvement, rethinking fertilization, and the use of new generation fertilizers as well as various biostimulants [3]. However, due to its great potential, the use of biostimulant products has gained great relevance as a tool for crop development improvement and efficient use of inputs.

One of the most innovative strategies that has shown favorable results in the last decade in terms of growth, production and efficient use of nutrients is the use of low weight molecules as biostimulants, such as omeprazole (OMP) and melatonin (MEL) [4,5]. Omeprazole, 5-metoxi-2-[(4-metoxi3,5-dimetil-2-piridinil)metil]sulfinil]-1H-benzimidazole, is a substituted benzimidazole widely used as an anti-ulcer drug [6]. Melatonin (N-acetil-5-metoxitriptamine) is a multifunctional molecule with physiological effects in various organisms. In humans, it is related to the enhancement of the immune system, circadian rhythms, sleep, mood, body temperature and appetite [7].

The biological function of both molecules is well identified in humans. However, recent research has shown that micromolar doses of both OMP and MEL in plants improved yield, root structure, nitrogen use efficiency, tolerance to different types of stress, photosynthesis parameters and the expression of genes that regulate these mechanisms [8,9]. In the same way, it has been reported that

one of the key mechanisms affecting the application of OMP and MEL to increase yield and biomass are related to carbon metabolism [10].

Carbon metabolism through the process of photosynthesis is a fundamental process for plants to carry out their routine and primary cellular activities during growth and development [11]. Furthermore, the study of chlorophyll fluorescence parameters, as a method of studying photosynthesis, is a valuable technique, which can serve as an indicator of the efficiency of the thylakoid membranes and the functional changes of the photosynthetic apparatus [12]. There is interaction between this process and key areas of plant function. Therefore, an improvement in photosynthesis would lead to an improvement in crop yields [13].

Although the responses of OMP and MEL in major metabolic processes, such as growth and physiological responses under stress have been previously studied, information about the specific function of these molecules related to light harvesting and transformation processes, as well as the efficiency of the photosynthetic machinery is limited. Therefore, the present research aims to evaluate the effect of foliar application of OMP and MEL on biomass, yield, SPAD values, leaf chlorophyll fluorescence (Fv/Fm), photochemical quenching (qP), photosystem II quantum yield (PhiPSII) and electron transport rate (ETR) in green bean plants cv. Strike.

2. Materials and Methods

2.1. Crop management

The experiment was carried out at the Food and Development Research Center, A.C. Delicias unit (CIAD), located in Cd. Delicias, Chihuahua, Mexico, during the months of September and October 2022. The experiment was established under shade netting at a mean temperature of 20.2 ± 7.8°C. Seeds of green bean (*Phaseolus vulgaris* L.) cv. Strike were grown in 13.4 L plastic pots (two plants per pot) in a substrate mixture composed of vermiculite and perlite in a 2:1 ratio. During development, plants were irrigated with a standard nutrient solution composed of 6 mM NH₄NO₃, 1.6 mM K₂HPO₄, 0.3 mM K₂SO₄, 4 mM CaCl₂, 1.4 mM MgSO₄, 5 µM Fe-EDDHA, 2 µM MnSO₄, 0.25 µM CuSO₄ y 0.5 µM H₃BO₃ y 0.3 µM Na₂MoO₄, at 6.0 ± 0.1 pH. A 500 mL nutrient solution was applied every third day and once the flowering stage (38 days after sowing (DAS)) was initiated, 1 L was applied. The treatments used were applied by foliar application every week from 15 DAS to 43 DAS.

2.2. Experimental design and description of treatments

A completely randomized design (CRD) was used with 7 treatments; the foliar application of omeprazole and melatonin at 3 doses each (1, 10 and 100 µM) and a control without application (Table 1). Each treatment had a total of 6 replicates (pots) with two plants each.

Table 1. Description of treatments applied.

Applied molecule	Dose (µM)	Code
Control	0	CONTROL
Omeprazole	1	OMP 1
Melatonin	1	MEL 1
Omeprazole	10	OMP 10
Melatonin	10	MEL 10
Omeprazole	100	OMP 100
Melatonin	100	MEL 100

2.3. Plant sampling

Once the plants reached physiological maturity, at 53 DAS, the plants were harvested for sampling. Fresh weight was recorded, as well as pod production. The harvested material was washed and dried in a drying oven (Shel Lab Forced Air Laboratory Oven SMO14-108 2, Baltimore, USA) at

70 °C for 48 h. Finally, dry weight was recorded. Fresh weight was expressed as grams of fresh weight per plant ($\text{g} \cdot \text{plant}^{-1}$ FW) and dry weight was expressed as grams of dry weight per plant ($\text{g} \cdot \text{plant}^{-1}$ DW).

2.4. Plant analysis

al biomass was evaluated considering all the organs of the plant. An analytical balance (AND HR-120, San Jose, California, USA) was used to quantify the weight. Biomass was expressed as grams of dry weight per plant ($\text{g} \cdot \text{plant}^{-1}$ DW).

2.4.1. Yield

The total fresh weight of pods per plant was quantified and expressed as grams of fresh weight per plant ($\text{g} \cdot \text{plant}^{-1}$ FW).

2.4.2. SPAD values

The quantification of the SPAD (Soil-Plant Analysis Development) values was obtained through a Minolta SPAD 502 chlorophyll reader (Konica Minolta Sensing, Inc., Osaka, Japan. The reading was achieved by projecting light through a leaf. Measurement was made on rib-free parts at 32 DAS. The results were expressed as SPAD units [37].

2.4.3. Measurements of chlorophyll fluorescence parameters

Chlorophyll fluorescence parameters were calculated at 32 DAS to evaluate the transfer, dissipation, and distribution of light in the photosystem of green bean plants grown under different biostimulant molecules, at different doses. Measurement of the variables was performed with a gas exchange system (Li-Cor 6400, Li-Cor, Inc., Lincoln, NE, USA) and an integrated fluorescence camera head (Li-Cor-6400, Li-Cor, Inc. Lincoln, NE, USA). The PPFD (photosynthetic photon flux density) value was $200 \mu\text{M photons m}^{-2} \text{ s}^{-1}$ and the CO_2 concentration was $400 \pm 2 \mu\text{mol} \cdot \mu\text{mol}^{-1}$ for the reference cell. The vapor pressure deficit of the air vapor pressure in the sample chamber was less than 1.5 and the leaf block temperature was 25 °C.

Table 2. Description of chlorophyll fluorescence parameters.

Variable	Parameter	Formula	Description
Fv/Fm	Maximum quantum yield of PSII	$\frac{F_m - F_o}{F_m}$	Maximum variable fluorescence in the state in which all non-photochemical processes are at a minimum. That is, in the dark-adapted state.
Fv'/Fm'	Maximum variable quantum yield of PSII	$\frac{F_m' - F_o'}{F_m'}$	Maximum effective variable fluorescence of the open centers of PSII. That is, in any light-adapted state (actinic radiation).
qP	Photochemical quenching of chlorophyll fluorescence	$\frac{F_m' - F_s'}{F_m' - F_o'}$	Estimation of the dissipation of light energy, transformed into chemical energy, which is used to carry out the reactions that drive photosynthesis.
NPQ	Non-photochemical quenching of chlorophyll fluorescence	$\frac{F_m - F_m'}{F_m'}$	Estimation of the attenuation of light energy, dissipated as heat during adaptation to actinic radiation. Acts as a protective mechanism for photoinhibition under conditions of excess energy.

PhiPSII	Quantum yield of PSII	$\frac{F_m' - F_s}{F_m'}$	Also represented as Φ PSII or ϕ PSII (Derived from the greek letter "Phi"). Estimate of the ratio of energy used by the PSII centers for transport to the absorbed light.
ETR	Electron transport rate	$\text{PhiPSII} \times \text{PPDF} \times f \times a$	Estimation of the actual electron transport rate of PSII. PPDF: Photosynthetic photon flux density. f: Factor of the energy partition between PSI and PSII, assuming it is equal between the two systems, 0.5. a: Common ratio of light absorption by photosynthetic tissue of C3 plants, 0.84.




Figure 1. Schematic of the different parameters of chlorophyll fluorescence measured in dark-adapted and light-adapted plants.

3.5. Statistical analysis

Once the measurements were obtained, a one-way analysis of variance (ANNOVA) was performed, with a separation of means test using the LSD method. For all the analyzes, 6 replicates were used. In addition, a Pearson correlation test between all the variables was performed, using the SAS 9.0 statistical package at ($p < 0.05$) level of significance.

3. Results and Discussion

3.1. Total Biomass

A key variable to determine the effectiveness of the treatments applied is the quantification of accumulated dry biomass [14]. In the present study, significant differences ($p < 0.05$) were found in total biomass accumulation by effect of OMP and MEL application at different doses (Figure 2). The treatment that was most favored was MEL 100 with a significant increase of 70% over the control, followed by MEL 1 with a significant increase of 55.4% over the control (Figure 3). However, although the increase in biomass accumulation was higher when the dose was increased from 1 to 100 μ M melatonin, there was no significant difference between these treatments (9.3%), so the most efficient

treatment was MEL 1. The data obtained in the present work agree with those obtained by [15], who reported a 71.1% increase in total dry biomass of maize seedlings when a 100 μM dose of melatonin was applied to plants fertilized with an optimum dose of nitrogen. Other authors such as [16], indicated that maize seedlings treated with melatonin at 1 μM increased their dry biomass by 50.6% under nitrogen-sufficient conditions at 28 days after planting compared to untreated plants.

According to the foliar application of omeprazole, the highest biomass accumulation was recorded in the OMP 100 treatment, with an increase of 32.4% with respect to the control (Figure 3). However, no significant differences were obtained between the different doses of this molecule and the control. Investigations such as the one conducted by [17] obtained significant increases of 14.3%, 21.9% and 25.7% in dry weight in lettuce plants, after adding 10, 50 and 100 μM of omeprazole respectively.

Models have previously been proposed to explain biomass distribution and relate it to carbon supply from leaf photosynthesis in the aerial part [18]. In addition, it has been indicated that the use of low weight molecules such as omeprazole and melatonin can intervene in growth processes, mainly in photosynthesis, through which the constituent sugars of biomass are obtained [19].

Similarly, the interaction of membrane ATPases and OMP has been described as responsible for the increase in vegetative growth in treated plants. ATPases are regulators of the polar transport of auxin from the source organs (leaves) to the root and fruit. It is possible that this mechanism promotes an increase in the number and size of pods in green bean, ultimately resulting in an increase in yield [20].

Also, multiple studies have shown the protective role of MEL in the photosynthetic apparatus of plants of different species, especially under stress conditions [21,22]. One of the processes that is disrupted under stress conditions is photosynthesis. However, the application of MEL favors that this process is carried on continuously, through different mechanisms such as maintaining the integrity of chloroplasts, improving water uptake to boost energy transfer between PSI and PSII, and improving the nitrogen assimilation process, allowing NO_3^- reduction and subsequently the utilization of carbon skeletons from the glutamine-glutamate pathway coming from the photosynthesis process [10,23]. The result in the enhancement of all these mechanisms may be the possible cause of increased biomass accumulation in bean plants.




Figure 2. Effect of foliar application of omeprazole (OMP) and melatonin (MEL) at doses of 1, 10 and 100 μM in green bean plants cv. Strike.

2.2. Yield

Yield is linked to biomass production and the physiological state of the plants, which makes it an indicator of the functionality of the treatments applied [24]. The data obtained in this research work showed significant differences ($p < 0.05$) in yield by effect of the application of OMP and MEL at different doses (Figure 3). The treatment that obtained the highest pod production was MEL1, which increased by 222.7% with respect to the control, with no significant difference when increasing the dose 10 and 100 times (Figure 4). The dose of omeprazole that obtained the best result was OMP 1, with a significant increase of 182% with respect to the control (Figure 3). In contrast to the application of melatonin, when the dose of omeprazole was increased by 10 and 100 times, the yield

decreased, which could indicate that this molecule modifies the metabolism of the plant according to the dose.

Similarly, MEL has been linked to the regulation of endogenous growth promoters such as auxin. This regulation has described accelerated growth in the root zone. When root architecture is modified by MEL action, nutrient and water uptake is promoted [25]. Adequate mineral supply and proper water uptake ultimately leads to increased plant capacity to produce biomass and yield. As shown in Figure 4, plants with the highest biomass accumulation were able to produce the highest amount of yield ($r = 0.6859^{**}$). In addition, application at the lowest doses (1 μM) of both OMP and MEL were the most effective in increasing yield by 1.8 and 2.2 times respectively with respect to the control, with a 100-fold lower dose.




Figure 3. Effect of foliar application of omeprazole (OMP) and melatonin (MEL) at doses of 1, 10 and 100 μM on total biomass accumulation (a) and yield (b) in green bean plants cv. Strike. Means with equal letters do not differ according to LSD test ($p < 0.05$).

2.3. SPAD values

Measurement of pigment concentration through SPAD values can be a good indicator of leaf chlorophyll composition and photosynthetic process, as well as plant health status and proper plant functioning [26]. In the present study, significant differences ($p < 0.05$) were found in photosynthetic activity expressed through SPAD units by effect of OMP and MEL application at different doses (Figure 4). The most favored treatment and, therefore, with the highest photosynthetic activity was OMP 1, with a significant increase over the control of 9.3%, being even higher than the photosynthetic activity recorded when the foliar dose of OMP was increased to 10 and 100 μM , making it the most efficient dose. Regarding the application of melatonin, the treatment that showed the highest photosynthetic activity was MEL10, with a significant increase of 6.5% over the control, being significantly equal to the doses of 1 and 100 μM .

The data obtained agree with those reported by [17], who found significant increases of 5.9% in basil cultivation under greenhouse conditions after the application of omeprazole at a dose of 10 μM . Similar results are reported by [27], who observed an increase in photosynthetic activity with exogenous addition of melatonin in maize seedlings.

The role of omeprazole in promoting the photosynthetic process is mainly related to the enhancement of nitrate and potassium uptake, due to the regulation of ATPases, and to the protection of the photosynthetic apparatus and regulation that these nutrients provide to the transpiration process [8]. In addition, a protective role against harmful ions such as chlorine has been described, facilitating their compartmentalization in vacuoles and excluding them from active sites during photosynthesis. Thus, with adequate nitrate uptake, leaves avoid the senescence state, allowing chloroplasts to maintain integrity and photosynthetic activity to be favored [28].

As for the role of melatonin, it is believed that one of the mechanisms driving the photosynthetic process is ionic balance in the cytosol, especially in the vegetative growth stage, by increasing the

uptake of potassium and calcium ions [29]. Also, according to [30], the presence of melatonin affects the expression of the genes responsible for photosynthetic pigment biosynthesis. More importantly, it has been reported to down-regulate the production of chlorophyllase, the enzyme responsible for chlorophyll degradation [31]. Finally, mention that the improvement in photosynthetic activity of the treatments compared to the control is due to different protective mechanisms that OMP and MEL molecules trigger inside the plant and lead to an improvement in the efficiency in light utilization for transformation into energy.




Figure 4. Effect of foliar application of omeprazole (OMP) and melatonin (MEL) at doses of 1, 10 and 100 μM on SPAD values in green bean plants cv. Strike. Means with equal letters do not differ according to LSD test ($p < 0.05$).

3.4. Chlorophyll Fluorescence Parameters

3.4.1. Chlorophyll fluorescence

Quantification of chlorophyll fluorescence is one of the most important parameters when considering photosynthetic efficiency. The energy absorbed by photosynthetic pigments is sent to the reaction centers of photosystems I (PSI) and II (PSII) for transformation and remission as fluorescence, photochemical energy for photosynthesis or dissipated as heat [32] (Figure 5).




Figure 5. Changes in the electronic state of chlorophyll (Chl) when absorbing a photon ($h\nu$). When chlorophyll absorbs energy, a transition to the excited state (Chl^*) occurs. In the excited state, chlorophyll can dissipate the energy in different ways: 1) The energy can be used to drive chemical reactions in the subsequent process of photosynthesis. 2) Chlorophyll can re-emit a photon, to return to the ground state, a process known as fluorescence. 3) Energy can be dissipated directly as heat.

In the present study, significant differences ($p < 0.05$) were found in chlorophyll fluorescence as a result of the application of OMP and MEL at different doses (Figure 6). All applied treatments showed increases with respect to the control, except for MEL 1. The treatment that expressed the highest fluorescence was MEL 10, with a significant increase over the control of 2.6%; however, it did not show a significant difference against MEL 100, whose increase was 2.5% over the control (Figure 6).

The omeprazole application that obtained the best fluorescence values was MEL 100, with a significant increase of 2.4% with respect to the control. However, the MEL 1 treatment obtained a similar increase of 2.2%.

The range of fluorescence values for bean plants from non-stress plants is between 0.7 and 0.8 [33]. As indicated by numerous authors, a decrease in fluorescence expressed as (Fv/Fm) is indicative of a decrease in PSII [34]. The results found in this study agree with those reported by [35], who found increases in fluorescence, as well as an increase in photosynthesis driven by the production of protective molecules such as catalase, peroxidase and superoxide dismutase in tomato plants treated with melatonin via foliar at 50 and 100 μM . Other authors have described that the increase in PSII efficiency related to the application of melatonin is due to a decrease in PSII electron flow [36]. This decrease in flux allows the captured energy to be used to drive energetic reactions for photosynthesis and less energy is dissipated by other pathways (Figure 7).

Another mechanism described for the increase in photosynthesis efficiency caused by melatonin is protection against reactive oxygen species. When the energy stored in chlorophyll is not dissipated by any of the possible pathways (Figure 5 and figure 7), the excited state of chlorophyll (Chl *) can react with molecular oxygen to form singlet oxygen (O_2^*), which is highly reactive and can degrade cellular components [37]. Thanks to the high stability of the central ring of the molecule, and its two side chains, once a functional group interacts with ROS to stabilize them, melatonin retains its stability and its ability to scavenge free radicals [38].




Figure 6. Effect of foliar application of omeprazole (OMP) and melatonin (MEL) at doses of 1, 10 and 100 μM on chlorophyll fluorescence (Fv/Fm) in green bean plants cv. Strike. Means with equal letters do not differ according to LSD test ($p < 0.05$).




Figure 7. Diagram of the energy flow of photosystems I and II. The blue arrows indicate the energy outputs of PSII. Black arrows indicate energy inputs. Under conditions of low radiation or optimal light, most of the energy is transferred to the reaction center for photochemistry. Under conditions of excessive radiation, energy dissipation takes place by exchange with PSI, heat emission, fluorescence or in extreme cases, photoinhibition.

3.4.2. Photochemical and non-photochemical quenching

Photochemical quenching (qP) refers to the set of processes of photosynthetic electron consumption in excited state through metabolic sinks, i.e., a different pathway from non-photosynthetic quenching (NPQ) processes, which returns electrons in excited state to their normal state through their remission by heat [39]. In the present research work, significant differences ($p < 0.05$) were found in the photochemical quenching coefficient due to the effect of applying OMP and MEL at different doses (Figure 8). The treatment with the highest rate was MEL 10, with a significant increase of 54.8%, followed by the OMP 1 treatment, with an increase of 15.5% both with respect to the control (Figure 8).

The trend observed in the decrease in yield as the dose of omeprazole increased (Figure 4), is repeated in this variable ($r = 0.4821$), so it is possible that the lower dose is the one that gives the plant the greatest capacity to transfer the "extra" energy to sinks, avoiding photoinhibition. The enhancement of photosynthetic quenching in omeprazole-treated plants has already been reported. Authors explain that there is a group of genes related to the integral maintenance of PSII and protein synthesis of the thylakoids that are overexpressed in the presence of omeprazole, as well as photosynthetic genes that protect against ROS, such as SIPSII and SIFTSH. One of the enzymes that are biosynthesized with the application of OMP is catalase, related to H_2O_2 scavenging [40].

Similarly, a higher qP value is observed in MEL 10 than in MEL 1, both increased with respect to the control, and their difference in yield is only 9%, indicating that the use of melatonin at low doses (between 1 to 10 μM), may confer the ability to protect the PSII reaction centers. Finally, indicate that both treatments at low doses obtained the highest operational efficiency by being able to achieve the highest electron transport rate ($r = 0.9554$) to possible acceptors, avoiding photoinhibition by excess energy [41].

Through the thermal dissipation contained in PSI and PSII, the plant avoids inactivation of the photosynthesis process or damage caused by excess energy. This thermal dissipation is known as non-photosynthetic quenching (NPQ) and its quantification can be an important index to measure the health of the photosynthetic system in plants [42]. Significant differences ($p < 0.05$) in the present work were found in the non-photochemical quenching coefficient due to the effect of applying OMP and MEL at different doses (Figure 8). The treatment with the lowest value was MEL 10, with a

significant decrease of 7.4% with respect to the control. On the omeprazole side, the lowest index was obtained in the OMP 100 treatment, with a significant decrease with respect to the control of 5.6%.

Although an increase in NPQ has been reported as a protective mechanism against photoinhibition, it may also indicate a loss of energy that can be used for photosynthesis, leading to a decrease in growth and production [43]. As demonstrated above, the increase in photochemical quenching (qP) caused by the application of omeprazole and melatonin may induce an improvement in the conversion of energy used for photosynthesis, but in turn, a decrease in the NPQ index because the protection mechanism against excess light is not as necessary in plants with high photosynthetic efficiency. The increase in NPQ, found in CONTROL treatment may be due to the fact that plants without OMP and MEL-driven production of antioxidant molecules, require the activation of alternative mechanisms such as the induction of the xanthophyll cycle and carotenoid biosynthesis [44]. However, these mechanisms, being energy-dependent, may have an energetic cost to the plant, which will ultimately lead to a reduction in growth and yield [45].




Figure 8. Effect of foliar application of omeprazole (OMP) and melatonin (MEL) at doses of 1, 10 and 100 μM on photochemical quenching (a) and non-photochemical quenching (b) in green bean plants cv. Strike. Means with equal letters do not differ according to LSD test ($p < 0.05$).

3.4.3. Electron transport quantum yield of PSII and electron transport rate

To complement the photochemical quenching results, an important indicator can be quantified: the electron transport quantum yield of PSII (PhiPSII). This measurement allows an estimation of the overall process of photosynthesis, as it quantifies the linear transport of electrons [46].

In the present study, significant differences ($p < 0.05$) were found in the coefficients of PhiPSII and ETR by the effect of OMP and MEL application at different doses (Table 3). With respect to PhiPSII, the highest yield was recorded in the MEL 10 treatment, with a significant increase of 71.4% over the control, followed by the OMP 1 treatment, with a significant increase of 52.8% over the control (Table 3). These are similar to those obtained by [47], who reported a greater increase in PhiPSII in tomato seedlings to which exogenous melatonin was foliar applied compared to solution application and the control. It is possible that the increased electron transport quantum yield of PSII is related to a higher amount of open PSII centers, which maintain their activity due to the protective role of melatonin as a redox regulator and promoter of D1 protein synthesis, the central subunit of PSII [48].

Table 3. Effect of foliar application of omeprazole (OMP) and melatonin (MEL) at doses of 1, 10 and 100 μ M on electron transport quantum yield of PSII in green bean plants cv. Strike. Means with equal letters do not differ according to LSD test ($p < 0.05$).

Treatment	PhiPSII	ETR (μ mol e $^{-}$ m $^{-2}$ s $^{-1}$)
CONTROL	0.1492 $\pm$ 0.030 e	76.42 $\pm$ 15.7 e
OMP 1	0.2278 $\pm$ 0.013 d	116.21 $\pm$ 6.46 d
MEL 1	0.1873 $\pm$ 0.017 b	95.58 $\pm$ 8.85 b
OMP 10	0.2070 $\pm$ 0.005 cd	105.47 $\pm$ 2.66 cd
MEL 10	0.1492 $\pm$ 0.030 a	130.34 $\pm$ 9.54 a
OMP 100	0.2212 $\pm$ 0.013 bc	112.85 $\pm$ 6.37 bc
MEL 100	0.2177 $\pm$ 0.014 bc	110.80 $\pm$ 7.18 bc

Authors such as [49], have reported an increase in electron transport rate caused by higher photochemical quenching. The results of the present experiment showed the treatment with higher qP (Figure 8) as the most favored treatment in terms of PhiPSII (Table 3). Similarly, the treatment with lower qP and higher NPQ, recorded the lowest PhiPSII.

The improvement in electron transport and PSII quantum yield could indicate a positive effect of melatonin and omeprazole on the reaction centers. In addition, these molecules could enhance quinone activation in PSII and PSI, which is primarily responsible for electron uptake and transfer between photosystems [50].

When quinones (Q_A) are in the oxidized state, PSII reaction centers are open and drained (F_o), ready to initiate light harvesting, which would enhance both electron reception and electron transport of photosystems (Figure 7) [51]. Finally, the role of melatonin in the antioxidant protection of the photosynthetic apparatus has been highlighted. Possibly, the reduction of ROS stress was reduced in the treatments to which MEL was applied. Photoinhibition and blockage of electron transport was lower in the applied treatments, thus improving the capacity of photosynthetic pigments to transform energy used for photosynthesis [52].

Table 4. Pearson correlation analysis for 8 green bean cv. Strike variables.

	TB	YD	SPAD	Fv/Fm	qP	NPQ	PhiPSII	ETR
TB	1	0.6859 **	0.1671	0.2420	0.2971	-0.1399	0.3173 *	0.3140 *
YD	0.6859 **	1	0.3345 *	0.1961	0.4821 *	-0.0118	0.4684 *	0.4618 *
SPAD	0.1671	0.3345 *	1	0.2640	0.3908 *	-0.0615	0.4302 *	0.4259 *
Fv/Fm	0.2420	0.1961	0.2640	1	0.7812 **	-0.1638	0.7721 **	0.7688 **
qP	0.2971	0.4821 *	0.3908 *	0.7812 **	1	-0.1396	0.9572 **	0.9554 **
NPQ	-0.1399	-0.0118	-0.0615	-0.1638	-0.1396	1	-0.1435	-0.1435
PhiPSII	0.3173 *	0.4684 *	0.4302 *	0.7721 **	0.9572 **	-0.1435	1	0.9997 **
ETR	0.3140 *	0.4618 *	0.4259 *	0.7688 **	0.9554 **	-0.1435	0.9997 **	1

** The correlation is significant ($p < 0.01$, bilateral) *. The correlation is significant ($p < 0.05$, bilateral). Abbreviations: TB= Total biomass, YD= Yield, SPAD= SPAD values, Fv/Fm= Fluorescence, qP= Photochemical quenching of chlorophyll fluorescence, NPQ= Non-photochemical quenching, PhiPSII= Quantum yield of electron transport of PSII, ETR= Electron transport rate.




Figure 9. Summary of key mechanisms affected by foliar application of omeprazole (OMP) and melatonin (MEL) in green bean plants cv. Strike.

In general, the application at the lower doses stimulated growth, expressed in a higher biomass recorded in treatments MEL 1 and OMP 1 (Figure 3). Similarly, greater leaf greenness expressed in SPAD units was found in the OMP 1 treatment (Figure 4). The treatments that benefited the most in photosynthetic efficiency parameters were generally MEL 10 and OMP 1 (Figures 6 and 8, Table 3). Finally, this resulted in higher fruit yield in treatments MEL 1 and OMP 1. As indicated by numerous authors, bioactive agents such as salicylic acid, brassinosteroids, calcium ion, polyamine, abscisic acid, and now melatonin and omeprazole, achieve remarkable effects on plant metabolism after a millimolar dose [53].

4. Conclusions

Foliar application of omeprazole and melatonin, both at 1 μ M doses, were the treatments that most benefited biomass accumulation and yield in plants of green bean cv. Strike. When the dose of these molecules was increased to 10 and 100 μ M, improvements were obtained with respect to the control, but not with respect to the lower doses. The treatment at a dose of 1 μ M omeprazole (OMP 1) was the treatment that increased SPAD units and obtained the best results in terms of photosynthesis fluorescence and photochemical quenching. On the other hand, the treatment at a dose of 10 μ M (MEL 10) of melatonin obtained the highest PhiPSII coefficient. These results could indicate that the use of both omeprazole and melatonin at doses of 1 μ M applied via foliar in green bean plants can increase photosynthetic efficiency and decrease photoinhibition, which is reflected in higher biomass accumulation and yield. Finally, it should be noted that further studies on the use of low weight molecules in agriculture are required to elucidate the mechanisms of action at the physiological, biochemical and molecular levels that are involved in the processes of photosynthesis, nitrogen assimilation, carbon metabolism and crop growth.

Author Contributions: To carry out the present study, E.S. and C.R. designed the study, E.M., C.R. and E.S. conducted the experiment and collected the samples, C.C. conducted the statistical analysis, and C.R., E.S., E. M., M.F., L.N., S.P. and C.C., contributed to the writing, revision and editing of the article. All authors have read and agreed to the published version of the manuscript.

Funding: "This work was supported by the National Science and Technology Council (CONACYT, México) under the project of the Call to Address National Problems: Project #1529 "Biofortification of basic agricultural crops representing the key to combat malnutrition and ensure food security in Mexico".

Acknowledgments: We are grateful to the National Science and Technology Council (CONACYT, México) for the financial support provided through the national program of scholarships for postgraduate studies under CVU No. 1011744. We also thank the Food and Development Research Center, A.C. (CIAD) Delicias unit, the Faculty of Agrotechnological Sciences and the Autonomous University of Chihuahua (UACH), for providing the facilities and laboratory equipment necessary to carry out the study.

Conflicts of Interest: “The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.”

References

1. Bajželj, B.; Richards, K. S.; Allwood, J. M.; Smith, P.; Dennis, J. S.; Curmi, E.; Gilligan, C. A. Importance of food-demand management for climate mitigation. *Nat. Clim. Change* **2014**, *4*(10), 924-929. <https://doi.org/10.1038/nclimate2353>
2. Simkin, A. J.; López-Calcagno, P. E.; Raines, C. A. Feeding the world: improving photosynthetic efficiency for sustainable crop production. *J. Exp. Bot.* **2019**, *70*(4), 1119-1140. <https://doi.org/10.1093/jxb/ery445>
3. Rouphael, Y.; Colla, G. Biostimulants in agriculture. *Front. Plant Sci.* **2020**, *11*, 40. <https://doi.org/10.3389/fpls.2020.00040>
4. Hardeland, R. Melatonin in plants—diversity of levels and multiplicity of functions. *Front. Plant Sci.* **2016**, *7*, 198. <https://doi.org/10.3389/fpls.2016.00198>
5. Van Oosten, M. J.; Dell'Aversana, E.; Ruggiero, A.; Cirillo, V.; Gibon, Y.; Woodrow, P.; Maggio, A.; Carillo, P. Omeprazole treatment enhances nitrogen use efficiency through increased nitrogen uptake and assimilation in corn. *Front. Plant Sci.* **2019**, *10*, 1507. <https://doi.org/10.3389/fpls.2019.01507>
6. Shah, N.; Gossman, W. Omeprazole. *StatPearls Publishing*. **2019**. PMID: 30969608.
7. Arnao, M. B.; Hernández-Ruiz, J. Functions of melatonin in plants: a review. *J. Pineal Res.*, **2015**, *59*(2), 133-150. <https://doi.org/10.1111/jpi.12253>
8. Cirillo, V.; Van Oosten, M. J.; Izzo, M.; Maggio, A. Omeprazole treatment elicits contrasting responses to salt stress in two basil genotypes. *Ann. Appl. Biol.* **2019**, *174*(3), 329-338. <https://doi.org/10.1111/aab.12496>
9. Elansary, H. O.; El-Abedin, T. K. Z. Omeprazole alleviates water stress in peppermint and modulates the expression of menthol biosynthesis genes. *Plant Physiol. Biochem.* **2019**, *139*, 578-586. <https://doi.org/10.1016/j.plaphy.2019.04.018>
10. Erdal, S. Melatonin promotes plant growth by maintaining integration and coordination between carbon and nitrogen metabolisms. *Plant Cell Rep.* **2019**, *38*(8), 1001-1012. <https://doi.org/10.1007/s00299-019-02423-z>
11. Sun, J.; Lu, N.; Xu, H.; Maruo, T.; Guo, S. Root zone cooling and exogenous spermidine root-pretreatment promoting *Lactuca sativa* L. growth and photosynthesis in the high-temperature season. *Front. Plant Sci.* **2016**, *7*, 368. <https://doi.org/10.3389/fpls.2016.00368>
12. Yamori, W.; Kondo, E.; Sugiura, D.; Terashima, I.; Suzuki, Y.; Makino, A. Enhanced leaf photosynthesis as a target to increase grain yield: insights from transgenic rice lines with variable Rieske FeS protein content in the cytochrome b6/f complex. *Plant Cell Environ.* **2016**, *39*(1), 80-87. <https://doi.org/10.1111/pce.12594>
13. Wang, L. Y.; Liu, J. L.; Wang, W. X.; Sun, Y. Exogenous melatonin improves growth and photosynthetic capacity of cucumber under salinity-induced stress. *Photosynthetica* **2016**, *54*(1), 19-27. <https://doi.org/10.1007/s11099-015-0140-3>
14. Sánchez, E.; Ruiz, J. M.; Romero, L. Compuestos nitrogenados indicadores de estrés en respuesta a las dosis tóxicas y deficientes de Nitrógeno en frijol ejotero. *Nova scientia* **2016**, *8*(16), 228-244.
15. Ahmad, S.; Wang, G. Y.; Muhammad, I.; Chi, Y. X.; Zeeshan, M.; Nasar, J.; Zhou, X. B. Interactive effects of melatonin and nitrogen improve drought tolerance of maize seedlings by regulating growth and physiochemical attributes. *Antioxidants* **2022**, *11*(2), 359. <https://doi.org/10.3390/antiox11020359>
16. Qiao, Y.; Yin, L.; Wang, B.; Ke, Q.; Deng, X.; Wang, S. Melatonin promotes plant growth by increasing nitrogen uptake and assimilation under nitrogen deficient condition in winter wheat. *Plant Physiol. Biochem.* **2019**, *139*, 342-349. <https://doi.org/10.1016/j.plaphy.2019.03.037>
17. Carillo, P.; Raimondi, G.; Kyriacou, M. C.; Pannico, A.; El-Nakhel, C.; Colla, G.; De Pascale, S.; Cirillo, V.; Rouphael, Y. Morpho-physiological and homeostatic adaptive responses triggered by omeprazole enhance lettuce tolerance to salt stress. *Sci. Hortic.* **2019**, *249*, 22-30. <https://doi.org/10.1016/j.scienta.2019.01.038>
18. Turner, M. F.; Heuberger, A. L.; Kirkwood, J. S.; Collins, C. C.; Wolfrum, E. J.; Broeckling, C. D.; Prenni, J. E.; Jahn, C. E. Non-targeted metabolomics in diverse sorghum breeding lines indicates primary and secondary metabolite profiles are associated with plant biomass accumulation and photosynthesis. *Front. Plant Sci.* **2016**, *7*, 953. <https://doi.org/10.3389/fpls.2016.00953>
19. Liang, D.; Ni, Z.; Xia, H.; Xie, Y.; Lv, X.; Wang, J.; Ling, L.; Deng, Q.; Luo, X. Exogenous melatonin promotes biomass accumulation and photosynthesis of kiwifruit seedlings under drought stress. *Sci. Hortic.* **2019**, *246*, 34-43. <https://doi.org/10.1016/j.scienta.2018.10.058>

20. Li, L.; Gallei, M.; Friml, J. Bending to auxin: fast acid growth for tropisms. *Trends Plant Sci.* **2021**, 27(5), 440-449. <https://doi.org/10.1016/j.tplants.2021.11.006>
21. Jahan, M. S.; Guo, S.; Sun, J.; Shu, S.; Wang, Y.; El-Yazied, A. A.; Alabdallah, N. M.; Hikal, M.; Mostafa H. M.; Ibrahim, F. M.; Hasan, M. M. Melatonin-mediated photosynthetic performance of tomato seedlings under high-temperature stress. *Plant Physiol. Biochem.* **2021**, 167, 309-320. <https://doi.org/10.1016/j.plaphy.2021.08.002>
22. Zhang, R.; Yue, Z.; Chen, X.; Wang, Y.; Zhou, Y.; Xu, W.; Huang, R. Foliar applications of urea and melatonin to alleviate water-logging stress on photosynthesis and antioxidant metabolism in sorghum seedlings. *Plant Growth Regul.* **2021**, 1-10. <https://doi.org/10.1007/s10725-021-00705-9>
23. Yan, F.; Zhang, J.; Li, W.; Ding, Y.; Zhong, Q.; Xu, X.; Wei, H.; Li, G. Exogenous melatonin alleviates salt stress by improving leaf photosynthesis in rice seedlings. *Plant Physiol. Biochem.* **2021**, 163, 367-375. <https://doi.org/10.1016/j.plaphy.2021.03.058>
24. Salcido-Martínez, A.; Sanchez, E.; Licon-Trillo, L. P.; Perez-Alvarez, S.; Palacio-Márquez, A.; Amaya-Olivas, N. I.; Preciado-Rangel, P. Impact of the foliar application of magnesium nanofertilizer on physiological and biochemical parameters and yield in green beans. *Not. Bot. Horti Agrobot. Cluj-Napoca.* **2020**, 48(4), 2167. <https://doi.org/10.15835/nbha48412090>
25. Sukumar, P.; Maloney, G. S.; Muday, G. K. Localized induction of the ATP-binding cassette B19 auxin transporter enhances adventitious root formation in *Arabidopsis*. *Plant Physiol.* **2013**, 162(3), 1392-1405. <https://doi.org/10.1104/pp.113.217174>
26. Jiang, C.; Johkan, M.; Hohjo, M.; Tsukagoshi, S.; Maruo, T. A correlation analysis on chlorophyll content and SPAD value in tomato leaves. *Hort. Res.* **2017**, 71, 37-42. <https://doi.org/10.20776/s18808824-71-p37>
27. Siddiqui, M. H.; Alamri, S.; Alsubaie, Q. D.; Ali, H. M. Melatonin and gibberellic acid promote growth and chlorophyll biosynthesis by regulating antioxidant and methylglyoxal detoxification system in tomato seedlings under salinity. *J. Plant Growth Regul.* **2020**, 39, 1488-1502. <https://doi.org/10.1007/s00344-020-10122-3>
28. Soltabayeva, A.; Srivastava, S.; Kurmanbayeva, A.; Bekturova, A.; Fluhr, R.; Sagi, M. Early senescence in older leaves of low nitrate-grown *Atxhd1* uncovers a role for purine catabolism in N supply. *Plant Physiol.* **2018**, 178(3), 1027-1044. <https://doi.org/10.1104/pp.18.00795>
29. Hussain, S.; Khalid, M. F.; Hussain, M.; Ali, M. A.; Nawaz, A.; Zakir, I.; Fatima, Z.; Ahmad, S. Role of micronutrients in salt stress tolerance to plants. In *Plant nutrients and abiotic stress tolerance*, (eds) Springer, Singapore, **2018**, 363-376. https://doi.org/10.1007/978-981-10-9044-8_15
30. Mir, A. R.; Siddiqui, H.; Alam, P.; Hayat, S. Melatonin modulates photosynthesis, redox status, and elemental composition to promote growth of *Brassica juncea*—a dose-dependent effect. *Protoplasma.* **2020**, 257, 1685-1700. <https://doi.org/10.1007/s00709-020-01537-6>
31. Weeda, S.; Zhang, N.; Zhao, X.; Ndip, G.; Guo, Y.; Buck, G. A.; Fu, C.; Ren, S. *Arabidopsis* transcriptome analysis reveals key roles of melatonin in plant defense systems. *PloS one.* **2014**, 9(3), e93462. <https://doi.org/10.1371/journal.pone.0093462>
32. Zhuang, J.; Wang, Y.; Chi, Y.; Zhou, L.; Chen, J.; Zhou, W.; Song, J. Zhao, N.; Ding, J. Drought stress strengthens the link between chlorophyll fluorescence parameters and photosynthetic traits. *PeerJ.* **2020**, 8, e10046. <https://doi.org/10.7717/peerj.10046>
33. Sánchez-Reinoso, A. D.; Ligarreto-Moreno, G. A.; Restrepo-Díaz, H. Physiological and biochemical expressions of a determinated growth common bean genotype (*Phaseolus vulgaris* L.) to water deficit stress periods. *The J. Anim. Plant Sci.* **2018**, 28(2).
34. Guidi, L.; Lo Piccolo, E.; Landi, M. Chlorophyll fluorescence, photoinhibition and abiotic stress: does it make any difference the fact to be a C3 or C4 species?. *Front. Plant Sci.* **2019**, 10, 174. <https://doi.org/10.3389/fpls.2019.00174>
35. Khan, T. A.; Saleem, M.; Fariduddin, Q. Melatonin influences stomatal behavior, root morphology, cell viability, photosynthetic responses, fruit yield, and fruit quality of tomato plants exposed to salt stress. *J. Plant Growth Regul.* **2022**, 1-25. <https://doi.org/10.1007/s00344-022-10713-2>
36. Zhang, Y. P.; Yang, S. J.; Chen, Y. Y. Effects of melatonin on photosynthetic performance and antioxidants in melon during cold and recovery. *Biol. Plant.* **2017**, 61(3), 571-578. <https://doi.org/10.1007/s10535-017-0717-8>
37. Dumanović, J.; Nepovimova, E.; Natić, M.; Kuča, K.; Jaćević, V. The significance of reactive oxygen species and antioxidant defense system in plants: A concise overview. *Front. Plant Sci.* **2021**, 11, 552969. <https://doi.org/10.3389/fpls.2020.552969>
38. Tan, D. X.; Manchester, L. C.; Esteban-Zubero, E.; Zhou, Z.; Reiter, R. J. Melatonin as a potent and inducible endogenous antioxidant: synthesis and metabolism. *Molecules.* **2015**, 20(10), 18886-18906. <https://doi.org/10.3390/molecules201018886>
39. Papageorgiou, G.C.; Govindjee. The Non-Photochemical Quenching of the Electronically Excited State of Chlorophyll a in Plants: Definitions, Timelines, Viewpoints, Open Questions. In *Non-Photochemical Quenching and Energy Dissipation in Plants, Algae and Cyanobacteria*. Demmig-Adams, B., Garab, G., Adams

- III, W., Govindjee (eds). Springer: Dordrecht, 2014; volume 40, 1-44. https://doi.org/10.1007/978-94-017-9032-1_1
40. Van Oosten, M. J.; Silletti, S.; Guida, G.; Cirillo, V.; Di Stasio, E.; Carillo, P.; Woodrow, P.; Maggio, A.; Raimondi, G. A benzimidazole proton pump inhibitor increases growth and tolerance to salt stress in tomato. *Front. Plant Sci.* **2017**, 8, 1220. <https://doi.org/10.3389/fpls.2017.01220>
 41. Tyystjärvi, E. Photoinhibition of photosystem II. In *International review of cell and molecular biology*; Elsevier, 2013; volume 300, 243-303. <https://doi.org/10.1016/B978-0-12-405210-9.00007-2>
 42. Murchie, E. H.; Ruban, A. V. Dynamic non-photochemical quenching in plants: from molecular mechanism to productivity. *The Plant J.* **2020**, 101(4), 885-896. <https://doi.org/10.1111/tpj.14601>
 43. Berteotti, S.; Ballottari, M.; Bassi, R. Increased biomass productivity in green algae by tuning non-photochemical quenching. *Sci. Rep.* **2016**, 6(1), 21339. <https://doi.org/10.1038/srep21339>
 44. Quaas, T.; Berteotti, S.; Ballottari, M.; Flieger, K.; Bassi, R.; Wilhelm, C.; Goss, R. Non-photochemical quenching and xanthophyll cycle activities in six green algal species suggest mechanistic differences in the process of excess energy dissipation. *J. Plant Physiol.* **2015**, 172, 92-103. <https://doi.org/10.1016/j.jplph.2014.07.023>
 45. Kromdijk, J.; Walter, J. Relaxing non-photochemical quenching (NPQ) to improve photosynthesis in crops. Burleigh Dodds Science Publishing: Cambridge, UK, 2023; pp. 1-19. <https://doi.org/10.19103/AS.2022.0119.09>
 46. Wang, P.; Sun, X.; Li, C.; Wei, Z.; Liang, D.; Ma, F. Long-term exogenous application of melatonin delays drought-induced leaf senescence in apple. *J. Pineal Res.* **2013**, 54(3), 292-302. <https://doi.org/10.1111/jpi.12017>
 47. Yang, X. L.; Xu, H.; Li, D.; Gao, X.; Li, T. L.; Wang, R. Effect of melatonin priming on photosynthetic capacity of tomato leaves under low-temperature stress. *Photosynthetica* **2018**, 56(3), 884-892. <https://doi.org/10.1007/s11099-017-0748-6>
 48. Zhou, X.; Zhao, H.; Cao, K.; Hu, L.; Du, T.; Baluška, F.; Zou, Z. Beneficial roles of melatonin on redox regulation of photosynthetic electron transport and synthesis of D1 protein in tomato seedlings under salt stress. *Front. Plant Sci.* **2016**, 7, 1823. <https://doi.org/10.3389/fpls.2016.01823>
 49. Perez, C. E. A.; Rodrigues, F. A.; Moreira, W. R.; DaMatta, F. M. Leaf gas exchange and chlorophyll a fluorescence in wheat plants supplied with silicon and infected with *Pyricularia oryzae*. *Phytopathology*. **2014**, 104(2), 143-149. <https://doi.org/10.1094/PHYTO-06-13-0163-R>
 50. Pereira, Y. C.; Rodrigues, W. S.; Lima, E. J. A.; Santos, L. R.; Silva, M. H. L.; Lobato, A. K. S. Brassinosteroids increase electron transport and photosynthesis in soybean plants under water deficit. *Photosynthetica*. **2019**, 57(1), 181-191. <https://doi.org/10.32615/ps.2019.029>
 51. Lu, X. F.; Zhang, H.; Lyu, S. S.; Du, G. D.; Wang, X. Q.; Wu, C. H.; Lyu, D. G. Effects of exogenous phenolic acids on photosystem functions and photosynthetic electron transport rate in strawberry leaves. *Photosynthetica*. **2018**, 56, 616-622. <https://doi.org/10.1007/s11099-017-0702-7>
 52. Wang, D. Y.; Wang, J.; Shi, S. H.; Huang, L. X.; Zhu, M.; Li, F. H. Exogenous melatonin ameliorates salinity-induced oxidative stress and improves photosynthetic capacity in sweet corn seedlings. *Photosynthetica*. **2021**, 59(2), 327-336. <https://doi.org/10.32615/ps.2021.031>
 53. Borges, A. A.; Jiménez-Arias, D.; Expósito-Rodríguez, M.; Sandalio, L. M.; Pérez, J. A. Priming crops against biotic and abiotic stresses: MSB as a tool for studying mechanisms. *Front. Plant Sci.* **2014**, 5, 642. <https://doi.org/10.3389/fpls.2014.00642>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.