

Article

Not peer-reviewed version

A Current Approach to Non-Alcoholic Steatohepatitis in Type 2 Diabetes Mellitus Patients

Ionelia Sorina Stan , Teodora Biciusca , [Biciusca Viorel](#) * , [Ramona Cioboata](#) , [Adina Mitrea](#) , Diana Clenciu , [Alice Elena Ghenea](#) , Suzana Danoiu

Posted Date: 3 July 2023

doi: [10.20944/preprints202307.0120.v1](https://doi.org/10.20944/preprints202307.0120.v1)

Keywords: non-alcoholic fatty liver disease; type 2 diabetes mellitus; inflammatory process; biological markers



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

A Current Approach to Non-Alcoholic Steatohepatitis in Type 2 Diabetes Mellitus Patients

Sorina I. Stan ^{1,†}, Teodora Biciusca ^{2,†}, Viorel Biciusca ^{3,*}, Ramona Cioboata ³, Diana Clenciu ⁴, Adina Mitrea ⁴, Alice Elena Ghenea ⁵ and Suzana Danoiu ⁶

¹ Doctoral School, University of Medicine and Pharmacy of Craiova, 200349, Romania

² Institute of Diagnostic and Interventional Radiology, Frankfurt, Germany; teodora.biciusca@kgu.de

³ Department of Internal Medicine, Faculty of Medicine, University of Medicine and Pharmacy of Craiova, 200349, Romania

⁴ Department of Bacteriology-Virology-Parasitology, University of Medicine and Pharmacy of Craiova, 200349 Craiova, Romania

⁵ Department of Pathophysiology, Faculty of Medicine, University of Medicine and Pharmacy of Craiova, 200349, Romania

* Correspondence: biciuscaviorel@gmail.com

† These authors contributed equally to this work.

Abstract: (1) **Background:** The relationship between non-alcoholic fatty liver disease (NAFLD) and type 2 diabetes (T2DM) is bidirectional: NAFLD increases the risk of T2DM, and T2DM promotes the progression of the disease to non-alcoholic steatohepatitis (NASH). (2) **Material and methods:** We performed a retrospective, open study that included 59 patients with NAFLD and T2DM who were distributed into two groups: 44 (74.57%) patients were diagnosed with HS and 15 (25.42%) patients were diagnosed with NASH. (3) **Results:** Among the non-specific inflammatory biomarkers statistically significant ($p=0.003$ respectively $p=0.03$) higher mean values were recorded in patients with NASH and T2DM in the case of ferritin and the neutrophil percentage-to-albumin ratio (NPATR). (4) **Conclusions:** Consequently, it is essential to identify alternative "surrogate" markers for the inflammatory process, particularly in individuals with diabetes, as it is a key characteristic of NASH. This need arises from the desire to avoid the risks associated with liver biopsy procedures and to prevent the unpredictable and unfavorable progression of NAFLD in patients with T2DM.

Keywords: non-alcoholic fatty liver disease; type 2 diabetes mellitus; inflammatory process; biological markers

1. Introduction

The term NAFLD dates back more than 40 years and encompasses a broad spectrum of diseases ranging from simple HS with low mortality risk to NASH with progression to cirrhosis and hepatocellular carcinoma (HCC) [1]. Worldwide, NAFLD represents the most common liver disease with a prevalence of 25% and is closely related to T2DM and obesity. The association of this disease with T2DM increases the likelihood of developing steatohepatitis, which is characterized by the presence of lobular inflammation and hepatocellular ballooning. This combination poses an increased risk of progression to advanced stages of the disease [2,3]. Current evidence suggests that the overall burden of NAFLD is closely related to the occurrence of extrahepatic complications, such as cardio-metabolic pathologies including diabetes, dyslipidemia, metabolic syndrome (MS), and cardiovascular disease (CVD) which are the leading cause of death among NAFLD patients [4-6].

The relationship between T2DM and NAFLD is supported by numerous studies, and there is a complex bidirectional relationship that pejoratively affects the prognosis and outcome of both pathologies [7-9]. Patients with NAFLD and diabetes develop advanced forms of the disease, even in the absence of other risk factors for advanced fibrosis such as age, dyslipidemia, hypertension, and an inherited history of NAFLD-associated cirrhosis [10].



The diagnosis of NAFLD is typically confirmed through liver biopsy puncture (LBP), an invasive procedure associated with various complications, including bleeding, pain, and abdominal discomfort, which limit its practical utility. Non-invasive techniques such as computed tomography (CT), nuclear magnetic resonance (NMR), and liver ultrasonography, are commonly included in the diagnosis of NAFLD. Among these techniques, liver ultrasonography is the most frequently used method in medical practice due to its simplicity and non-invasive nature. It can detect steatosis when it affects around 20-30% of hepatocytes [11]. To facilitate the diagnosis of the disease, experts have introduced a range of blood biomarkers into medical practice. Currently, the fatty liver index (FLI), hepatic steatosis index (HSI), and liver fat score (LFS) have been validated for diagnosing HS [12]. HS is generally considered to have a benign clinical course while NASH has the potential to contribute to progressive liver fibrosis, eventually leading to cirrhosis, liver failure, and HCC [13]. Thus, it is necessary to distinguish between fatty liver and NASH (inflammation with or without fibrosis). Given the prevalence of hepatic steatosis, a liver biopsy cannot be considered for all patients. Indeed, in a recent study by an Italian group, fatty liver regressed in 50% of all examined cases and had a benign evolution [12,14]. Therefore, there have been efforts to determine predictive markers of fibrosis and/or steatohepatitis to avoid biopsies [15]. For the diagnosis of NASH, the American Gastroenterological Association (AGA) recommends considering an initial stage of biochemical suspicion, characterized by a moderate increase in transaminase levels, reaching up to five times the upper limit of normal [16]. Additionally, elevated levels of gamma-glutamyltransferase (GGT) have been observed in these patients and are associated with advanced fibrosis and higher mortality rates. GGT is included as a biomarker in several non-invasive tests used to assess liver fibrosis, such as FibroTest and Hepascore [17].

Chronic inflammation plays a crucial role in the development of the disease, involving various cell types such as neutrophils, lymphocytes, monocytes, and platelets. The leukocytes are assessed through a range of biological markers that evaluate the inflammatory status, including the neutrophil-to-lymphocyte ratio (NLR), monocyte-to-lymphocyte ratio (MLR), and systemic inflammatory index (SII) [18]. Platelets also actively contribute to the inflammatory process by facilitating the recruitment of leukocytes and their morphology changes in patients with T2DM and MS. Therefore, mean platelet volume (MPV) and platelet distribution width (PDW) serve as biological markers for platelet activation and are associated with the presence of insulin resistance (IR) and the severity of its complications [19]. Other inflammatory markers used in current medical practice are serum C-reactive protein (CRP), C-reactive protein/albumin ratio (CAR), and neutrophil percentage-to-albumin ratio (NPAR). NPAR is a cost-effective, easy-to-calculate biomarker initially used to calculate the prognosis of cancer patients and for cardiovascular disease prognosis, but it has been reported to have predictive value in NAFLD as well [20,21]. The present study aimed to determine which biological markers ("surrogate" biomarkers), other than aminotransferases (alanine aminotransferase (ALT) and aspartate aminotransferase (AST)), differentiate the two forms of liver disease in NAFLD (HS from NASH), with emphasis on inflammatory factors.

2. Materials and Methods

2.1. Selection of patients

We conducted a retrospective, open study at the Diabetes and Nutritional Diseases Clinic of the Filantropia Municipal Clinical Hospital, Craiova, in the period 2020-2023. This study involved 59 patients diagnosed with NAFLD and T2DM who met the specified inclusion and exclusion criteria and were closely monitored at the Filantropia Municipal Clinical Hospital (Figure 1).




Figure 1. Study Flow Chart depicts the patient selection process for our study. Out of the initial 120 patients, after applying the exclusion criteria, a total of 59 patients remained. The diagnosis of non-alcoholic fatty liver disease (NAFLD) in these individuals was established based on the results of the fatty liver index (FLI), hepatic steatosis index (HSI), and abdominal ultrasonography.

To facilitate the comparative analysis of the outcomes, we divided the patients into two groups: one comprising 44 individuals diagnosed with HS, and the other consisting of 15 patients diagnosed with NASH. The composition of the patient groups under study was determined according to specific criteria. To be included in the study, patients needed to meet certain inclusion criteria, including having a diagnosis of T2DM, exhibiting ultrasonographic evidence of HS, and demonstrating elevated values of the two diagnostic indicators, namely HSI above 30 and FLI above 60 [22]. The diagnosis of T2DM was established on diabetes symptoms (polyuria, polydipsia, and polyphagia) in combination with a random venous plasma glucose concentration ≥ 11.1 mmol/l (200 mg/dL), a fasting plasma glucose concentration ≥ 7.0 mmol/l (126 mg/dL), two-hour plasma glucose concentration ≥ 11.1 mmol/l (200 mg/dL) two hours after an oral glucose tolerance test (OGTT) with 75g anhydrous glucose, glycated hemoglobin (HbA1c) greater than 6.4% or ongoing therapy for T2DM. The diagnosis of NAFLD was established based on abdominal ultrasound (AU), FLI, and HSI values [22]. AU was performed by specialist doctors in module B, which allowed the subjective estimation of steatosis. HS was defined according to the criteria of the AGA, such as the increase of hepatic echogenicity compared to renal echogenicity, the presence of intensity as well as the lack of differentiation of periportal and vascular wall intensity as a result of hepatic hyperechogenicity and eventually the presence of hepatomegaly. According to EASL, AU is the first screening imaging investigation performed in patients with NAFLD due to its profitability; however, it does have several limitations [23]. AU exhibits a sensitivity of 90% for detecting liver fat content above 30%, but this sensitivity decreases as the liver fat content decreases as well [24]. FLI was calculated based on the formula:

$$ey/(1+ey) \times 100, \quad (1)$$

where:

$$y = 0.953 \times TG \text{ (mg/dL)} + 0.139 \times BMI \text{ (kg/m}^2\text{)} + 0.718 \times GGT \text{ (U/L)} + 0.053 \times WC \text{ (cm)} - 15,745. \quad (2)$$

In this formula TG stands for triglycerides (mg/dL), GGT for γ -glutamyl transferase (U/L), and WC represents the waist circumference (cm) [25]. FLI ranges from 0 to 100. In a study conducted by Bedogni et al., a diagnosis of NAFLD was confirmed when the FLI value reached or exceeded 60, while a value below 30 excluded the diagnosis, demonstrating excellent diagnostic accuracy [26].

HSI serves as a screening tool to optimize the management of NAFLD. HSI is calculated based on the formula:

$$8 \times \text{ALT/AST} + \text{BMI} \quad (3)$$

where 2 points more will be added if the patient has diabetes and 2 points more if the individual is a female [27].

The validation of HSI enables clinicians to effectively identify patients who require dietary and lifestyle modifications, as well as those who may need further assessment through AU. A value of HIS <30 excludes the diagnosis of NAFLD (negative probability of 0.186), while a value >36 supports the diagnosis. The AUROC value is 0.182 (95% CI, 0.801-0.24) and the sensitivity of AU for a value below 30 was 93.1%, effectively excluding the diagnosis, while a value above 36 demonstrated a specificity of 92.4% to confirm the diagnosis [28]. The diagnostic criteria used to distinguish HS from NASH in diabetic patients with NAFLD was the increased value of ALT and AST with values for ALT>55 U/L and for AST>34 U/L.

The exclusion criteria were excessive alcohol consumption (for men more than 30ml/day and women more than 20ml/day), the presence of chronic viral hepatitis C, especially genotype 3, chronic viral hepatitis B, autoimmune hepatitis, the utilization of medications known for their hepatotoxic properties (amiodarone, methotrexate, glucocorticoids), type 1 diabetes, severe infections, and neoplasias.

2.2. Clinical parameters

Detailed clinical data were collected from all patients included in the study. The anthropometric data obtained from the patients included measurements of height and body weight, which were utilized to evaluate or calculate the body mass index (BMI) and waist circumference (WC). The BMI was calculated based on the formula weight/height². Based on the resulting value, the degree of obesity was determined. Patients with a BMI between 24 and 29.9 kg/m² were classified as overweight, while those with a BMI exceeding 30 kg/m² were categorized as having obesity. WC was measured halfway between the lowest rib and the iliac crest to assess the degree of abdominal adiposity. In women, values greater than 80 cm were associated with abdominal adiposity, while men with abdominal adiposity presented values greater than 94 cm. Comorbidities such as hypertension (systolic blood pressure \geq 130 mmHg or diastolic blood pressure \geq 85 mm Hg, or those on treatment) and dyslipidemia (serum triglyceride concentration \geq 150 mg/dl or fasting HDL-C <40 mg/dl in men and <50 mg/dl in women, or patients on treatment) were also recorded. Based on these comorbidities, the diagnosis of MS was established, according to the already known diagnostic criteria.

Measurements of hematological (complete blood count), biochemical parameters (glycemia, total cholesterol, triglycerides, albumin), liver tissue function (ALT, AST, GGT), and inflammatory parameters (erythrocyte sedimentation rate: ESR, C-reactive protein: CRP, fibrinogen: Fg, and serum ferritin: SF) were performed by standard laboratory methods in the Clinical Laboratory of the "Filantropia" Municipal Clinical Hospital, Craiova, with the help of analyzers using the Mindray BC Auto Hematology and Architect Clinical Chemistry Systems. To assess the subclinical systemic inflammation commonly observed in patients with NASH, we conducted a comprehensive analysis of various biological markers, apart from traditional inflammatory markers like WBC (white blood cell count), ESR, CRP, Fg, SF levels, and other hematological markers and biochemical parameters described in the existing literature to evaluate the inflammatory status. In this context, we calculated the ratio neutrophils%/lymphocytes% (NLR), the ratio monocytes%/lymphocytes% (MLR), the systemic inflammatory index (SII) based on the formula: neutrophils (/mm³) x platelets (/mm³)/lymphocytes (no /mm³), the CAR as well as the NPAR, which was calculated based on the formula

neutrophil count (%) $\times 100$ /albumin (d/dL). We also used MPV (mean platelet volume) and PDW (platelet distribution width), which are closely correlated with IR, considering that platelets are active participants in the liver inflammatory process, as a result of their role in recruiting leukocytes through the liver sinusoids.

2.4. Statistical analysis

Patient data and data processing were possible using Microsoft Excel together with IBM SPSS Statistics 20.0. Since the numerical variables studied presented a normal data distribution in each studied group, we were allowed to use parametric statistical tests, and the results were summarized as mean value \pm standard deviation. P values less than 0.05 were considered statistically significant for all statistical tests.

2.5. Ethical considerations

The present study was carried out after obtaining the approval of the Scientific Academic Ethics and Deontology Committee of the University of Medicine and Pharmacy in Craiova, number 94/27.04.2023. Each patient signed an informed consent for inclusion in the study.

3. Results

3.1. Study composition

The study population comprised 59 individuals diagnosed with both NAFLD and T2DM, with 44 patients (74.57%) having HS and T2DM, while 15 patients (25.42%) had NASH and T2DM. Demographic and epidemiological studies followed the distribution of the studied patients and highlighted the main risk factors that were associated with NAFLD (Table 1).

Table 1. The incidence and the mean value of the demographic and diagnostic parameters of patients with NAFLD and T2DM, and the p-value of the comparative analysis of the two groups studied.

Parameter	Incidence and Mean Value in Patients with NAFLD and T2DM	Incidence and Mean Value in Patients with HS and T2DM	Incidence and Mean Value in Patients with NASH and T2DM	Statistical Significance p-Value
Women (number)	25/59 (42.37%)	21/44 (47.27%)	4/15 (26.66%)	0.07
Men (number)	34/59 (57.62%)	23/44 (52.27%)	11/15 (73.33%)	0.15
Age (years)	61.52 \pm 10.07	62.20 \pm 0.70	59.53 \pm 10.06	0.38
The environment of urban origin (number)	36/59 (61.01%)	25/44 (56.81%)	11/15 (73.33%)	0.26
The rural environment of origin (number)	23/59 (38.89%)	19/44 (43.18%)	4/15 (26.66%)	0.29
T2DM	59/59 (100%)	44/44 (100%)	15/15 (100%)	-
Hypertension	53/59 (89.83%)	39/44 (88.63%)	14/15 (93.33%)	0.61
Obesity	54/59 (91.52%)	41/44 (93.18%)	13/15 (86.66%)	0.44
Dyslipidemia	47/59 (79.66%)	36/44 (81.81%)	11/15 (73.33%)	0.48
MS	56/59 (94.91%)	42/44 (95.45%)	14/15 (93.33 %)	0.75
FLI	77.76 \pm 22.42	74.40 \pm 22.62	88.93 \pm 10.59	0.04*
HSI	46.58 \pm 8,71	45.31 \pm 7.91	50.39 \pm 10.17	0.05*
US-LLL_diameter (cm)	7.54 \pm 1.14	7.38 \pm 1.83	8 \pm 1.37	0.07
US-RHL_diameter (cm)	14.3 \pm 1.95	14.05 \pm 2.03	15.02 \pm 1.48	0.09

US-PV_diameter (cm)	1.084±0.108	1.072±0.07	1.12±0.104	0.15
US-Spleen_diameter (cm)	10.35±1.09	10.21±1	10.7±1.22	0.09

Abbreviations: NAFLD: non-alcoholic fatty liver disease; HS: hepatic steatosis; NASH: non-alcoholic steatohepatitis; T2DM: type 2 diabetes mellitus; MS: metabolic syndrome; FLI: fatty liver index; HSI: hepatic steatosis index; US: ultrasonography, US-LLL: left liver lobe measured US; US-RHL: right hepatic lobe evaluated US, US-PV: portal vein assessed US.

Among the patients, there were 25 women (42.37%) and 34 men (57.62%). In both the HS and T2DM subgroup (52.27%) and the NASH and T2DM subgroup (73.33%), men constituted the majority. The mean age of patients with NAFLD was 61.52 ± 10.07 years, with small differences between the values of the mean age of patients with SH and T2DM (62.20 ± 0.70 years) compared to those with NASH and T2DM (59.53 ± 10.06 years). Regarding the patients' geographical background, 36 (61.01%) patients came from urban areas, while 23 (38.98%) patients were from rural areas. In the studied patients, T2DM presented an equal incidence in both groups of patients. This equality was due to the inclusion criteria of the study, which required patients to have diabetes mellitus. Patients with HS and DM2 presented a higher incidence of obesity (93.18% vs 86.66%), dyslipidemia (81.81% vs 86.66%), and MS (95.45% vs 93.33%), while patients with NASH and T2DM showed a higher incidence of hypertension (93.33% vs 88.63%).

In these patients, the diagnosis of NAFLD was established based on AU and the calculation of the FLI and HSI diagnostic index values. The imaging study showed that hyperechoic appearance with inhomogeneous structure was the echographic appearance encountered in all selected patients. The sagittal diameter of the left lobe recorded a mean value of 7.54 ± 1.14 cm in patients with NAFLD. NASH patients presented a higher mean value (8 ± 1.37 cm) than HS patients (7.38 ± 1.83 cm). The longitudinal diameter of the right liver lobe, measured on the midclavicular line, showed a mean value of 14.3 ± 1.95 cm in patients with NAFLD. Patients with NASH presented a higher mean value of this diameter (15.02 ± 1.48 cm) compared to patients with HS (14.05 ± 2.03 cm). The diameter of the portal vein assessed by ultrasound exhibited a mean value of 1.084 ± 0.108 cm in patients with NAFLD. This value was higher in patients with NASH (1.12 ± 0.104 cm) compared to patients with HS (1.072 ± 0.07 cm). The longitudinal diameter of the spleen measured by echography registered a mean value of 10.35 ± 1.09 cm, a higher mean value being observed in patients with NASH (10.7 ± 1.22 cm) compared to patients with HS (10.21 ± 1 cm). Patients with NAFLD recorded a mean value of FLI of 77.76 ± 22.42 , while patients with NASH (88.93 ± 10.59) presented a statistically significant higher mean value ($p=0.04$), compared to patients with HS (74.40 ± 22.62). The mean value of HSI in patients with NAFLD was 46.58 ± 8.71 , with patients with NASH (50.39 ± 10.17) showing a statistically significant higher mean value ($p=0.05$), compared to patients with HS (45.31 ± 7.91).

3.2. The biochemical study

The biochemical study analyzed the main metabolic and enzymatic parameters used to diagnose HS and NASH in patients with T2DM (Table 2).

Table 2. The mean values of the metabolic and enzymatic parameters found in patients with NAFLD and T2DM, as well as the values of the statistical test of the comparative analysis of the two studied groups.

Parameter	Incidence and mean value in patients with NAFLD and T2DM	Incidence and mean value in patients with HS and T2DM	Incidence and mean value in patients with NASH and T2DM	Statistical significance p-value
Glucose (mg/dL)	220.74±103.61	216.11±107.59	234.33±89.56	0.56
HbA1c (%)	8.34±1.97	8.26±2.74	8.57±1.67	0.60

Cholesterol (mg/dL)	204.40±88.62	193.59±16.26	235.73±145.13	0.10
Triglycerides (mg/dL)	271.30±290.061	232.79±14.14	384.26±450.90	0.05*
ALT (U/L)	43.89±44.39	23.61±9.19	103.4±54.26	0.0001*
AST (U/L)	28.33±23.64	18.18±5.11	56.26±32.80	0.005*
GGT (U/L)	47.74±47.58	30.15±21.21	99.33±70.17	0.002*
Albumin (g/dL)	4.48±0.60	4.66±0.44	4.42±0.35	0.19

Abbreviations: NAFLD: non-alcoholic fatty liver disease; HS: hepatic steatosis; NASH: non-alcoholic steatohepatitis; T2DM: type 2 diabetes mellitus; HbA1c: glycosylated hemoglobin; ALT: alanine transferase; AST: aspartate transferase; GGT: gamma-glutamyl transferase.

"Fasting" blood glucose showed a higher mean value in patients with NASH and T2DM (234.33±89.56 g/dL) compared to patients with SH and T2DM (216.11±107.59 g/dL). Regarding the mean value of HbA1c, patients with NASH and T2DM presented higher mean values (8.57±1.67%), compared to the group of patients with HS and T2DM (8.26±2.74 %). Cholesterolemia showed higher mean values in patients with NASH and T2DM (235.73±145.13 mg/dL) compared to patients with HS and T2DM (193.59±16.26 mg/dL). Triglyceridaemia also exhibited statistically significant (p=0.05) higher mean values in patients with NASH and T2DM (384.26±45.90 mg/dL) compared to patients with HS and T2DM (232.79±14.14 mg/dL) (Figure 2).




Figure 2. Mean values and statistical significance of biochemical parameters in the studied patients.
Abbreviations: NAFLD: non-alcoholic fatty liver disease; NASH: non-alcoholic steatohepatitis.

The enzymes of the liver cytolysis syndrome (ALT and AST) recorded statistically significant higher mean values (p=0.0001 respectively p=0.005) in patients with NASH and T2DM (103.4±54.26 U/L and 56.26±32.80 U/L) compared to the values in patients with SH and T2DM (23.61±9.19 U/L and 18.18±5.11 U/L). Among the enzymes of hepatic cholestasis syndrome, GGT recorded statistically significant higher values (p=0.002) in patients with NASH and T2DM (99.33±70.17 U/L) in comparison to patients with HS and T2DM (30.15±21.21 U/L) (Figure 3). Regarding the liver synthesis function,

the mean value of albumin was lower in patients with NASH and T2DM (4.42 ± 0.35 g/dL) compared to the group of patients with HS and T2DM (4.66 ± 0.44 g/dL).




Figure 3. Mean values and statistical significance of cytolysis enzymes and hepatic cholestasis. Abbreviations: NAFLD: non-alcoholic fatty liver disease; NASH: non-alcoholic steatohepatitis; ALT: alanine aminotransferase; AST: aspartate aminotransferase; GGT: gamma-glutamyl transpeptidase.

3.3. The hematological study

The hematological study analyzed the behavior of the hematological parameters in the studied patients (Table 3).

Table 3. The mean values of the hematological parameters found in patients with NAFLD and T2DM, as well as the value of the statistical test of the comparative analysis of the two studied groups.

Parameter	Incidence and Mean Value in Patients with NAFLD and T2DM	Incidence and Mean Value in Patients with HS and T2DM	Incidence and Mean Value in Patients with NASH and T2DM	Statistical Significance
				p-Value
Hb (g/dL)	13.35 \pm 1.58	13.09 \pm 1.64	14.1 \pm 1.27	0.03*
WBC (no/mm ³)	7901 \pm 1893	8066 \pm 1945	7415 \pm 1635	0.25
Neutrophils (%)	59.37 \pm 7.88	59.67 \pm 0.94	58.46 \pm 6.13	0.62
Lymphocytes (%)	29.16 \pm 8.13	28.65 \pm 4.10	30.65 \pm 6.96	0.42
Monocytes (%)	6.17 \pm 1.38	5.99 \pm 1.72	6.72 \pm 1.47	0.07
Platelets (no/mm ³)	245502 \pm 62489	251992 \pm 91923	226666 \pm 77524	0.45
MLR	0.22 \pm 0.07	0.21 \pm 0.07	0.23 \pm 0.07	0.62
NLR	2.37 \pm 0.93	2.31 \pm 0.97	2.53 \pm 0.80	0.45
PDW (%)	41.19 \pm 14.37	41.98 \pm 12.62	38.87 \pm 18.36	0.47
MPV (fL)	9.86 \pm 1.03	9.92 \pm 0.92	9.68 \pm 1.28	0.44

Abbreviations: NAFLD: non-alcoholic fatty liver disease; HS: hepatic steatosis; NASH: non-alcoholic steatohepatitis; T2DM: type 2 diabetes mellitus; Hb: hemoglobin; WBC: white blood cells; NLR: neutrophil/lymphocyte ratio; MLR: monocyte/lymphocyte ratio; PDW: platelet distribution width; MPV: mean platelet volume.

Hemoglobin (Hb) recorded a statistically significant lower mean value ($p=0.03$) in patients with HS and T2DM (13.09 ± 1.64 g/dL) in comparison with patients with NASH and T2DM (14.1 ± 1.27 g/dL). Regarding the leukocyte series, the patients with HS and T2DM showed higher mean leukocyte (WBC) values (8066 ± 1945 /mmc vs 7415 ± 1635 /mmc) compared to patients with NASH and T2DM. The behavior of leukocyte subpopulations was different in the two groups of patients. Thus, the percentage of neutrophils (N) registered a slightly higher but statistically insignificant increased mean value in patients with HS and T2DM ($59.67\pm0.94\%$) compared to patients with NASH and T2DM ($58.46\pm6.13\%$), while the mean value of the percentage of lymphocytes (L) was higher in NASH and T2DM patients ($30.65\pm6.96\%$) in comparison to the mean value in HS and T2DM patients ($28.65\pm4.10\%$). Moreover, the mean value of the percentage of monocytes (M) was higher in patients with NASH and T2DM ($6.72\pm1.47\%$) compared to the mean percentage value recorded in patients with HS and T2DM ($5.99\pm1.72\%$). Inflammatory biomarkers, NLR and MLR showed higher mean values in patients with NASH and T2DM (2.53 ± 0.80 respectively 0.23 ± 0.07) compared to patients with HS and T2DM (2.31 ± 0.97 respectively 0.21 ± 0.07). In the case of platelets, patients with HS and T2DM presented a higher mean value compared to patients with NASH and T2DM (251992 ± 91923 /mmc vs 226666 ± 77524 /mmc). Regarding platelet indices, the mean value of PDW in patients with HS and T2DM ($41.98\pm12.62\%$) was higher than in patients with NASH and T2DM ($38.87\pm18.36\%$). The same behavior was found in the case of the MPV, which presented a higher value in patients with HS and T2DM (9.92 ± 0.92 fl) compared to patients with NASH and T2DM (9.68 ± 1.28 fl).

3.5. Study of inflammatory parameters

The study of systemic inflammation parameters that evaluate the subclinical inflammatory process from T2DM, allowed us to identify the parameters that are associated with inflammation in NASH, apart from ALT and AST (Table 4).

Table 4. The mean values of the inflammatory parameters found in patients with NAFLD and T2DM as well as the values of the statistical test of the comparative analysis of the two studied groups.

Parameter	Incidence and mean value in patients with NAFLD and T2DM	Incidence and mean value in patients with HS and T2DM	Incidence and mean value in patients with NASH and T2DM	Statistical significance p-value
ESR (mm)	38.42 ± 27.08	37.54 ± 26.16	41 ± 25.29	0.67
CRP (mg/dL)	0.81 ± 1.55	0.60 ± 0.21	1.44 ± 2.87	0.07
Fg (mg/dL)	361.84 ± 103.65	351.76 ± 195.16	391.4 ± 111.27	0.20
SF (ng/ml)	206.93 ± 92.97	186.70 ± 86.26	266.26 ± 96.74	0.003*
SII	487.59 ± 190.19	476.20 ± 36.75	520.97 ± 238.10	0.43
CAR	0.39 ± 1.56	0.38 ± 1.77	0.40 ± 0.67	0.97
NPAR	1386.9 ± 429.88	1317.79 ± 235.88	1589.6 ± 713.15	0.03*

Abbreviations: NAFLD: non-alcoholic fatty liver disease; HS: hepatic steatosis; NASH: non-alcoholic steatohepatitis; T2DM: type 2 diabetes mellitus; ESR: erythrocyte sedimentation rate; CRP: C-reactive protein; Fb: fibrinogen; SF: serum ferritin; SII: systemic inflammatory index; CAR: C-reactive protein/albumin ratio; NPAR: ratio of neutrophils x100/albumin.

The mean value of the ESR in one hour (ESR/1h) was higher in patients with NASH and T2DM (41 ± 25.29 mm/1h) compared to that recorded in HS and T2DM patients (37.54 ± 26.16 mm/1h). Additionally, the CRP recorded a higher mean value in patients with NASH and T2DM (1.44 ± 2.87

mg/dL) compared to the values found in those with HS and T2DM (0.60 ± 0.21 mg/dL). The same behavior was found in the case of Fg, which showed a higher mean value in patients with NASH and T2DM (391.4 ± 111.27 mg/dL) compared to patients with HS and T2DM (351.76 ± 195.16 mg/dL). Moreover, there was a statistically significant difference ($p=0.003$) in the mean serum ferritin levels between patients with NASH and T2DM (266.26 ± 96.74 ng/mL) compared to patients with hepatic HS and T2DM (186.70 ± 86.26 ng/mL) (Figure 4).

The mean value of the SII was higher in patients with NASH and T2DM (520.97 ± 238.10) compared to patients with HS and T2DM (476.20 ± 36.75). The mean CAR ratio in patients with NASH and T2DM (0.40 ± 0.67) was slightly higher compared to patients with HS and T2DM (0.38 ± 1.77), although this difference was not statistically significant. Furthermore, the mean value of the NPAR exhibited a statistically significant increase ($p=0.03$) in patients with NASH and T2DM (1589.6 ± 713.15) when compared to patients with HS and T2DM (1317.79 ± 235.88) (Figure 5).




Figure 4. The mean values and statistical significance of serum ferritin in the two groups of patients. Abbreviations: NAFLD: non-alcoholic fatty liver disease; NASH: non-alcoholic steatohepatitis.




Figure 5. The mean value and the statistical significance of NPAR in the two groups studied.

Abbreviations: NPAR: neutrophil percentage to albumin ratio; NAFLD: non-alcoholic fatty liver disease; NASH: non-alcoholic steatohepatitis.

4. Discussion

The increase in aminotransferases in patients with NAFLD is the result of the interaction between IR, obesity and MS components [29]. In our study, the factors that were associated with the progression of NAFLD in T2DM patients were: gender, advanced age, and MS components. Diabetes is a well-known factor in the progression of the disease, thus identifying biomarkers other than ALT and AST that differentiate the two forms of the disease is an important step for stratifying the risk of the disease. In both groups of patients, the male gender was predominant; in the group of patients with HS and T2DM, the incidence was 52.27%, while in the group of patients with NASH and T2DM, the incidence was higher, 73.33%. NAFLD is known as a sex-dysmorphic pathology, with a higher incidence for the male sex compared to the female sex, at least during the fertile period. After the onset of menopause, the incidence is comparable. The same higher incidence for the male gender, but lower compared to our study (51.8%) was present in a Chinese study that investigated the factors of progression to fibrosis in patients with NAFLD and T2DM. [30]. Regarding the mean age of the patients, it was higher in patients with HS and T2DM compared to the group of patients with NASH and T2DM. Age is recognized as a factor that contributes to the progression of the disease, an aspect that was also present in our study. Also, in the study by Lamonaco et al., it was observed that the mean age of patients with HS and T2DM was higher compared to the mean age of patients with NASH and T2DM [31]. Both patients with HS and those with NASH and T2DM mostly came from the urban environment. The incidence of urban patients in the literature is lower (34.1%) compared to our study [32].

The main risk factors of the progression of NAFLD to NASH and subsequently to advanced stages of the disease are represented by MS components. T2DM is one of these factors through IR and compensatory hyperinsulinemia that favors the deposition of lipids in the liver [32]. Lonardo et al [33] and Popescu et al [34] also stated that hepatic steatosis and fibrosis are the consequences of IR. In our study, the inclusion criteria required the presence of diabetes, resulting in a 100% incidence of diabetes in both patient groups. Regarding the comorbidities of the patients included in the study,

patients with NASH and T2DM had a higher incidence of arterial hypertension (93.33%), while patients with HS and T2DM presented a higher incidence of obesity (93.18%), dyslipidemia (81.81%) and MS (94.45%). The incidence of arterial hypertension in the study by Bazick et al. [35] was lower (40.7%) compared to our study, while Chen's study only reported a higher incidence of dyslipidemia (84.44%) in patients with HS and T2DM, the incidence of obesity and MS being lower (68.89% respectively 63.63%) compared to the results of our study [36].

The analysis of carbohydrate metabolism parameters revealed a higher value of both blood glucose and HbA1c in patients with NASH and T2DM compared to the group of patients with HS and T2DM. IR is known as the most important pathogenic stage of MS but is also the main factor in the development and progression of non-alcoholic liver disease. Currently, the risk factors associated with the development of fibrosis in patients with NAFLD and T2DM are not fully known. To emphasize these factors and their prevalence in patients with T2DM, Julian's study which included 930 patients with NAFLD and T2DM, showed lower mean blood glucose values in patients with HS and T2DM (148 ± 45 mg/dL) compared to patients with NASH and T2DM (151 ± 50 mg/dL). Regarding the mean value of HbA1c, patients with HS ($7.0 \pm 1.2\%$) had lower values in comparison to patients with NASH ($7.1 \pm 1.3\%$), without statistically significant differences [37].

The analysis of lipid metabolism parameters demonstrated the presence of dyslipidemia with statistically significant higher values (in the case of triglycerides) in patients with NASH and T2DM. The mechanisms of dyslipidemia are related to the appearance of hepatic and peripheral insulin resistance associated with disorders of hepatic glucose metabolism and lipoproteins [38]. In our study, both the mean value of cholesterol and triglycerides were higher and statistically significant in the case of triglycerides ($p=0.05$) in patients with NASH and T2DM. The mean values of the two components of lipid metabolism were documented in a study made up of 204 patients with NAFLD and T2DM that examined the prevalence of risk factors in these patients. Both the mean values of cholesterol and triglycerides were higher but statistically insignificant in patients with NASH and T2DM (190.56 ± 44.02 mg/dL respectively 173.51 ± 68.84 mg/dL) compared to patients with HS and T2DM (183.97 ± 48.84 mg/dL respectively 147.37 ± 74.86 mg/dL) [39]. The analysis of protein metabolism demonstrated a lower mean value of albumin in patients with NASH and T2DM compared to patients with HS and T2DM. Albumin levels can fall in patients with inflammatory disorders and other conditions. This may be due to downregulated production of albumin mRNA by the liver, leading to reduced synthesis, increased albumin catabolism, and increased vascular permeability [40]. In a study made up of 1249 patients that explored the determinants of fibrosis in patients with NAFLD and T2DM, the mean value of albumin was also lower in patients with NASH and T2DM (4.15 ± 0.41 g/dL) compared to patients with HS and T2DM (4.21 ± 0.43 g/dL) [32].

The analysis of the enzymatic parameters of the liver functions highlighted significantly higher mean values of liver cytolysis enzymes, ALT and AST ($p=0.0001$ respectively $p=0.005$) in patients with NASH and T2DM, compared to patients with HS and T2DM. Most patients with NAFLD are asymptomatic and are typically identified when abnormal liver studies are noted on routine laboratory assessment. In particular, the liver enzymes, ALT, and AST levels are elevated. However, these enzymes may not be elevated in all cases of NAFLD, and the level of aminotransferases may not reliably predict the extent of inflammation and cirrhosis [41]. In a study by Mandal et al., an independent association between liver enzymes, particularly ALT, and NAFLD in a population of individuals with T2DM was observed [42]. Mandal et al. also found that 40.4% of the diabetic population had elevated ALT levels, while increases in AST and ALP levels were observed in only 17% and 16% of diabetic individuals, respectively [42]. Similar findings were reported by Shrestha and Bhatt in Nepal, where they also observed a statistically significant elevation in ALT levels among diabetic patients compared to the control groups [44]. Furthermore, a higher prevalence of abnormalities in liver function tests has been linked to individuals with T2DM compared to those without T2DM [45]. Also, the enzyme of hepatic cholestasis syndrome, GGT presented a higher and statistically significant mean value ($p=0.002$) in patients with NASH and T2DM compared to patients with HS and T2DM. GGT plays a crucial role in the breakdown of glutathione, facilitating the transfer of the glutamyl component and preserving intracellular balance against oxidative stress [45].

Previous research has established connections between serum GGT levels and various chronic conditions, including NAFLD, cardiovascular diseases, diabetes, and MS [46]. The mechanisms underlying elevated GGT levels in NAFLD patients involve oxidative stress, inflammatory response, and excessive hepatic fat accumulation, all of which contribute to impaired insulin signaling and IR [47].

The analysis of the mean values of the hematological parameters highlighted a statistically significant higher mean hemoglobin value ($p=0.03$) in patients with NASH and T2DM. In the literature, the mean value of Hb in a study that investigated the association between glycated albumin (GA) and glycated hemoglobin (GA/HbA1c) ratio and the grade of NAFLD in patients with NAFLD and T2DM, who were ultrasonographical evaluated, was also statistically significant lower ($p<0.001$) in patients with HS and T2DM (12.5 ± 2.2 g/dL) compared to patients with NASH and T2DM (13.8 ± 1.6 g/dL) [48]. The mean value of the parameters of the leukocyte series was different in the two groups of patients, without statistically significant differences being evident. Thus, the mean value of WBC was higher in patients with HS and T2DM compared to patients with NASH and T2DM. In Bazick's study, which investigated the determinants of fibrosis in 1249 patients with NAFLD, of which 435 (34.8%) had been diagnosed with diabetes, the mean value of leukocytes in patients with HS and T2DM (7180 ± 230 mm 3) was statistically insignificant lower, compared to the group of patients with NASH and T2DM (7230 ± 210 mm 3) [35]. Regarding the percentage of neutrophils (NP), their mean value was higher in patients with HS and T2DM compared to patients with NASH and T2DM. Instead, the mean value of the percentage of lymphocytes (LP) was lower in patients with HS and T2DM compared to patients with NASH and T2DM. The mean values of the percentage of neutrophils and lymphocytes have been studied by many authors, as well as the relationship between the value of the NLR ratio and the severity of the degree of steatosis in patients with NAFLD and T2DM. The mean value of the NP in Kahraman's study, which included 143 patients with NAFLD and T2DM, evaluated by liver ultrasonography, reported a lower value in patients with NAFLD and mild fatty liver ($59.2\pm8.2\%$) compared to patients with NAFLD and severe fattening ($68.6\pm7.9\%$). The mean LP varied significantly, with patients diagnosed with NAFLD and mild fatty liver ($30.8\pm8.6\%$) exhibiting a higher mean value compared to patients with NAFLD and severe fatty liver ($22.3\pm6.4\%$) [49].

The mean value of the inflammatory indices derived from the leukocyte series in our study was lower in patients with HS and T2DM compared to patients with NASH and T2DM. NLR is known as an easily obtained blood biomarker that evaluates both the inflammatory state and the severity of liver fibrosis in patients with NAFLD. This aspect was suggested in Kahraman's study which concluded that NLR increased with the degree of obesity in patients with T2DM [49]. To determine the severity of fibrosis in patients with NAFLD and T2DM, a Chinese study investigated the association between NLR, NPAR, and the presence of NAFLD or advanced liver fibrosis in a group of 3991 patients with T2DM, of which 1355 (33.95%) had NAFLD and T2DM, while 162 (4.05%) patients had T2DM and advanced liver fibrosis. In this study, it was concluded that the NLR is an easily measurable inflammatory marker, that correlated significantly and positively with the degree of steatosis and fibrosis [20]. Another blood biomarker used to assess the inflammatory state in patients with NAFLD is MLR, which has not yet been studied in patients with T2DM. It has been used as a blood biomarker in several studies in patients with different degrees of steatosis. In this regard, a recent study that consisted of 81 patients with NAFLD and associated comorbidities, including diabetes and obesity, reported a higher mean value of MLR in patients with mild steatosis (0.25) compared to patients with severe steatosis (0.19), results differing compared to our study [50]. The platelets count and their characteristics were also investigated as another hematological parameter in the study. In our study, the average value of the PLT was higher in patients with HS and T2DM compared to the group of patients with NASH and T2DM. The role of platelets in the inflammatory process in NAFLD is well known, as they are included in numerous scores that evaluate the process of non-invasive fibrosis. Thus, a study that included 565 patients with NAFLD and diabetes, and aimed to highlight the factors that are associated with liver fibrosis in these patients, showed a statistically significant ($p<0.01$) higher average value of platelets in NAFLD

without liver fibrosis patients (a mean value of 226.000/mmc, with limits of 211.000-279.000/mmc³) compared to those with NAFLD and liver fibrosis (a mean value of 201.000/mmc, with limits of 173.500-238.000/mmc) [51]. Related to the characteristics of platelets, in our study, the average value of PDW was higher in the group of patients with HS and T2DM, compared to the group of patients with NASH and T2DM. The same behavior was found in the case of MPV, where the average value in the group of patients with HS and T2DM was higher compared to the group of patients with NASH and T2DM. To demonstrate the diagnostic accuracy of platelets and platelet indices, a study by Alempijevic et al. consisting of 98 patients with NAFLD and associated risk factors (diabetes, MS), divided according to the presence or absence of liver fibrosis proven using non-invasive tests, showed a statistically significantly ($p<0.01$) higher average PDW distribution in patients with fibrosis (16.91 ± 0.35) compared to patients without fibrosis (16.55 ± 0.77). In another study, a different behavior was found in the case of MPV, where the average value in patients with liver fibrosis (9.21 ± 1.05 fL) was higher but statistically insignificant compared to the group of patients without liver fibrosis (9.04 ± 1.46 fL) [52].

In our study, we observed that patients diagnosed with both NASH and T2DM had higher levels of all inflammatory markers compared to those with HS and T2DM. Particularly noteworthy was the mean NPAR, which exhibited significantly elevated values ($p=0.03$) in patients with NASH and T2DM, in contrast to patients with HS and T2DM. Chi-Feng Liu et al. also conducted a study that revealed significant connections between the NPAR, NLR, and the presence of NAFLD or advanced liver fibrosis. They found that incremental increases in both the NLR and NPAR were linked to a higher risk of developing NAFLD. Furthermore, among individuals with T2DM, a notable and independent correlation existed between higher NPAR levels and the likelihood of having NAFLD. Based on the findings, the NPAR demonstrates promise as a reliable predictive biomarker for NAFLD. As far as we know this is the only study besides ours that has looked at the association between NPAR and NAFLD [20]. The NPAR has also been employed as a predictive tool for systemic inflammation in various medical conditions, including unresectable pancreatic cancer [53]. Furthermore, NPAR has been linked to all-cause mortality in critically ill patients with acute kidney injury [54] and septic shock [55]. It is well known that chronic inflammation plays a crucial role in the progression of NAFLD, which encompasses a spectrum ranging from simple steatosis to NASH, advanced liver fibrosis, cirrhosis, and ultimately end-stage liver disease and hepatocellular carcinoma [56]. Consequently, NAFLD represents a significant contributor to chronic and progressive liver damage, emphasizing the importance of noninvasive diagnostic and monitoring methods for progressive liver disease. The findings of Chi-Feng Liu et al. indicate that an elevated NPAR level is significantly linked to an increased risk of NAFLD, surpassing the effectiveness of albumin, neutrophil percentage, and NLR as biomarkers for predicting NAFLD [20]. Additionally, the relationship between the NPAR and NAFLD was influenced by the presence of T2DM, a recognized risk factor for NAFLD development and progression [57]. Furthermore, T2DM itself is associated with systemic inflammation [58]. Hence, it is plausible that the presence of concurrent T2DM might conceal the correlation between the NPAR and NAFLD [20].

In our study, we also observed a statistically significant rise ($p=0.003$) in the average serum ferritin (SF) level among patients diagnosed with NASH and T2DM compared with patients with HS and T2DM. Previous studies have shown that the level of SF, which is the main storage form of iron in the body, is correlated with T2DM and NAFLD [59,60,61]. A study by Yan et al. revealed that individuals with T2DM and NAFLD displayed notably higher levels of serum ferritin when compared to T2DM patients without NAFLD. Moreover, as SF levels increased, the proportions of NAFLD cases also showed a significant rise. Thus, SF was identified as an independent risk factor for NAFLD among patients with T2DM [62]. Research has indicated that elevated levels of SF without corresponding high transferrin saturation can serve as an indicator of glucose or lipid metabolism dysfunction [63]. Increased iron levels may contribute to the development of diabetes by triggering oxidative stress, inflammation, insulin resistance, insulin insufficiency, liver impairment, and other factors [64]. Conversely, diabetes itself can disrupt the balance of iron metabolism and lead to iron overload, thereby establishing a detrimental cycle [65]. IR stands as a prominent characteristic of

NAFLD [66]. Under conditions of IR, the β -oxidation process of free fatty acids becomes impeded, which subsequently fosters the accumulation of fat within the liver [67] Mayneris-Perxachs et al. also documented a positive association between SF levels and the extent of hepatic fat accumulation [68]

Furthermore, we assessed the clinical utility of C-reactive protein (CRP) as a diagnostic tool for NAFLD. CRP displayed a higher mean value in our study in patients with NASH and DM2, though statistically insignificant, compared to patients with HS and DM2. Several studies have suggested a potential association between elevated serum CRP levels and NAFLD [69,70,71]. Since CRP has a short half-life of approximately 18 hours, an increase in its serum levels typically indicates its synthesis in response to a pathological process. As a result, CRP is regarded as a valuable non-specific biochemical marker for chronic inflammation [72]. Oruc et al. further validated that elevated circulating CRP levels can serve as an indicative marker for the presence of NAFLD. However, it is important to note that due to its limited specificity, interpreting CRP values accurately requires consideration of all other clinical and laboratory information. Consequently, it was proposed that serial CRP measurements could prove beneficial in the clinical management and ongoing monitoring of NAFLD patients [73]. Ndumele et al. conducted a study involving 2388 individuals, both diabetic and non-diabetic, without known coronary heart disease, which revealed a significant correlation between hepatic steatosis, as detected by ultrasound, and elevated levels of high-sensitivity C-reactive protein (hs-CRP). HS was associated with higher hs-CRP levels, irrespective of obesity status, presence of metabolic syndrome, or absence. Notably, obesity and MS were independently linked to increased hs-CRP levels, as expected [74]. It is important to mention that the liver plays a crucial role as the main producer of CRP. Previous research suggests that the level of hepatic steatosis and inflammation, as determined by histology, is linked to the systemic levels of inflammatory biomarkers. In a study involving 85 patients, it was observed that higher grades of hepatic steatosis, necroinflammation, and fibrosis in biopsy samples were associated with progressively elevated hs-CRP levels, reaching the "high risk" range [75]. Additional studies have discovered a direct correlation between the severity of NAFLD and the expression of inflammatory mediators in hepatocytes [74,76].

The results of our study provide evidence supporting an independent association between NASH and systemic inflammation in patients with T2DM. Nevertheless, further research is required in the future to explore the link between inflammatory factors and NAFLD in patients with diabetes. This is crucial since the existing studies on this subject are limited in number.

5. Conclusions

NAFLD, the most prevalent liver disease globally, is characterized by its tendency to advance to more severe stages, although the specific factors influencing this progression remain uncertain. Type 2 diabetes mellitus is recognized as one of these factors, as it may contribute to the unfavorable development of NAFLD through insulin resistance. The subclinical, systemic inflammatory process seen in diabetic patients, triggered by lipid accumulation at the hepatocellular level, can be evidenced both histologically and biologically.

Therefore, finding "surrogate" biomarkers of the inflammatory process, which is a defining feature of NASH in diabetic individuals, is an imperative current need to avoid performing liver biopsy puncture, which is fraught with risks, and avoiding the unpredictable, unfavorable evolution of NAFLD in patients with T2DM. The inflammatory biomarkers that recorded statistically significantly higher serum values in NASH patients compared to HS patients were serum ferritin and the neutrophil percentage-to-albumin ratio (NPAR).

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used "Conceptualization: I.S.S., T.B., and V.B.; Methodology: I.S.S.; Software: T.B.; Validation: V.B., I.S.S., T.B., and S.D.; formal analysis: I.S.S., T.B., and V.B.; investigation: I.S.S., T.B., and V.B.; resources: I.S.S., T.B., A.M., D.C., R.C., A.E.G., and V.B.; data curation: I.S.S., T.B.; writing—original draft preparation: I.S.S., T.B., and V.B.; writing—review and editing: I.S.S., T.B., and V.B.; visualization: I.S.S., T.B., S.D.; and V.B.; supervision: B.V.; project administration: I.S.S., T.B., V.B., and S.D.;

funding acquisition: I.S.S.; T.B.; V.B.; and S.D.; All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki and approved by the Scientific Academic Ethics and Deontology Committee of the University of Medicine and Pharmacy in Craiova, number 94/27.04.2023.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Not applicable.

Acknowledgments: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kosmalski M.; Ziolkowska S.; Czarny P.; Szemraj J.; Pietras T. The Coexistence of Nonalcoholic Fatty Liver Disease and Type 2 Diabetes Mellitus. *Journal of Clinical Medicine* 2022, 11, 1375. <https://doi.org/10.3390/jcm11051375>
2. Spiers J.; Brindley J.H.; Li W; Alazawi Q. What's new in non-alcoholic fatty liver disease? *Frontline Gastroenterology* 2022, 13, e102-e108. doi:10.1136/flgastro-2022-102122
3. Estes C.; Anstee Q.M.; Arias-Loste M.T.; et al. Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, the United Kingdom, and the United States for the period 2016-2030. *J Hepatol* 2018, 69, 896–904. doi: 10.1016/j.jhep.2018.05.036
4. Chalasani N.; Younossi Z.; Lavine J.E.; Charlton M; Cusi K; Rinella M; Harrison SA., Brunt EM; Sanyal AJ. The diagnosis and management of nonalcoholic fatty liver disease: Practice guidance from the American Association for the Study of Liver Diseases. *Hepatology*. 2017, 67, 328–357. <https://doi.org/10.1002/hep.29367>
5. Loomba R.; Wong R.; Fraysse J.; Shreay S.; Li S.; Harrison S.; Gordon S.C. Nonalcoholic fatty liver disease progression rates to cirrhosis and progression of cirrhosis to decompensation and mortality: a real-world analysis of Medicare data. *Aliment Pharmacol Ther* 2020, 51, 1149–1159. <https://doi.org/10.1111/apt.15679>
6. Long M.T.; Zhang X.; Xu H.; Liu C.; Corey K.E.; Chung R.T; Loomba R; Benjamin E.J. Hepatic Fibrosis Associates With Multiple Cardiometabolic Disease Risk Factors: The Framingham Heart Study. *Hepatology* 2021, 73, 548–559. <https://doi.org/10.1002/hep.31608>
7. Watt G.P.; De La Cerdá I.; Pan J.; Fallon M.B.; Beretta L; Loomba R; Lee M; McCormick J.B; Fisher-Hoch S.P. Elevated Glycated Hemoglobin Is Associated With Liver Fibrosis, as Assessed by Elastography, in a Population-Based Study of Mexican Americans. *Hepatology Communications* 2020, 4, 1793–1801. <https://doi.org/10.1002/hep4.1603>
8. Caussy C.; Aubin A.; Loomba R. The Relationship Between Type 2 Diabetes, NAFLD, and Cardiovascular Risk. *Current Diabetes Reports* 2021, 21. <https://doi.org/10.1007/s11892-021-01383-7>
9. Zarghamravanbakhsh P.; Frenkel M.; Poretsky L. Metabolic causes and consequences of nonalcoholic fatty liver disease (NAFLD). *Metabolism Open* 2021, 12, 100149. <https://doi.org/10.1016/j.metop.2021.100149>
10. Caussy C.; Soni M.; Cui J.; Bettencourt R.; Schork N.; Chen C.H; Ikhwan M.A; Bassirian S.; Cepin S.; et al. Nonalcoholic fatty liver disease with cirrhosis increases the familial risk for advanced fibrosis. *Journal of Clinical Investigation* 2017, 127, 2697–2704. <https://doi.org/10.1172/jci93465>
11. Papatheodoridi M.; Cholongitas E. Diagnosis of Non-alcoholic Fatty Liver Disease (NAFLD): Current Concepts 2019, 24, 4574–4586. <https://doi.org/10.2174/1381612825666190117102111>
12. Manousou P.; Kalambokis G.; Grillo F.; Watkins J.; Xirouchakis E.; Pleguezuelo M.; Leandro G.; Arvaniti V.; Germani G.; Patch D.; Calvaruso V.; Mikhailidis D.P.; Dhillon A.P.; Burroughs A.K. Serum ferritin is a discriminant marker for both fibrosis and inflammation in histologically proven non-alcoholic fatty liver disease patients. *Liver Int* 2011, 31, 730–739. doi: 10.1111/j.1478-3231.2011.02488.x.
13. Bedogni G.; Miglioli L.; Masutti F.; et al. Incidence and natural course of fatty liver in the general population: the Dionysos study. *Hepatology* 2007, 46, 1387–1391. doi: 10.1002/hep.21827.
14. Wieckowska A.; McCullough A.J.; Feldstein A.E. Noninvasive diagnosis and monitoring of nonalcoholic steatohepatitis: present and future. *Hepatology* 2007, 46, 582–589. doi: 10.1002/hep.21768.
15. Eslam M.; George J. Reply to: correspondence regarding "A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement." *Journal of Hepatology* 2020, 73, 1575. <https://doi.org/10.1016/j.jhep.2020.07.045>
16. Kanwal F.; Shubrook J.H.; Adams L.A.; Pfotenhauer K.; Wai-Sun Wong V.; Wright E.; Abdelmalek M.F.; Harrison S.A.; Loomba R.; Mantzoros C.S.; et al. Clinical Care Pathway for the Risk Stratification and Management of Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2021, 161, 1657–1669. doi: 10.1053/j.gastro.2021.07.049.

17. Martinou E.; Pericleous M.; Stefanova I.; Kaur V.; Angelidi A.M. Diagnostic Modalities of Non-Alcoholic Fatty Liver Disease: From Biochemical Biomarkers to Multi-Omics Non-Invasive Approaches. *Diagnostics* 2022, 12, 407. doi: 10.3390/diagnostics12020407.
18. Schuster S.; Cabrera D.; Arrese M.; Feldstein A.E. Triggering and resolution of inflammation in NASH. *Nat Rev Gastroenterol Hepatol* 2018, 15, 349–64. doi: 10.1038/s41575-018-0009-6
19. Milovanovic Alempijevic T.; Stojkovic Lalosevic M.; Dumić I.; Jocić N.; Pavlović Marković A.; Dragasević S.; Jović I.; Lukić S.; Popović D.; Milošavljević T. Diagnostic Accuracy of Platelet Count and Platelet Indices in Noninvasive Assessment of Fibrosis in Nonalcoholic Fatty Liver Disease Patients. *Canadian Journal of Gastroenterology and Hepatology* 2017, 1–5. <https://doi.org/10.1155/2017/6070135>
20. Liu C.F.; Chien L.W. Predictive Role of Neutrophil-Percentage-to-Albumin Ratio (NPAR) in Nonalcoholic Fatty Liver Disease and Advanced Liver Fibrosis in Nondiabetic US Adults: Evidence from NHANES 2017–2018. *Nutrients* 2023, 15, 1892. <https://doi.org/10.3390/nu15081892>
21. Zhu C.; Huang D.; Ma H.; Qian C.; You H.; Bu L.; Qu S. High-Sensitive CRP Correlates With the Severity of Liver Steatosis and Fibrosis in Obese Patients With Metabolic Dysfunction Associated Fatty Liver Disease. *Frontiers in Endocrinology* 2022, 13. <https://doi.org/10.3389/fendo.2022.848937>.
22. Zhang J.Z.; Cai J.J.; Yu Y.; She Z.G.; Li H. Nonalcoholic Fatty Liver Disease: An Update on the Diagnosis. *Gene Expr* 2019, 19, 187–198. doi: 10.3727/105221619X15553433838609.
23. European Association for the Study of the Liver. EASL Clinical Practice Guidelines on non-invasive tests for evaluation of liver disease severity and prognosis - 2021 update. *J Hepatol* 2021, 75, 659–689. doi: 10.1016/j.jhep.2021.05.025.
24. Zhou J.H.; Cai J.J.; She Z.G.; Li H.L. Noninvasive evaluation of nonalcoholic fatty liver disease: Current evidence and practice. *World J Gastroenterol* 2019, 25, 1307–1326. doi: 10.3748/wjg.v25.i11.1307.
25. Fatty Liver Index (FLI) MD App Available on <https://www.mdcalc.com/calc/10001/fatty-liver-index#next-steps>
26. Bedogni G.; Bellentani S.; Miglioli L.; Masutti F.; Passalacqua M.; Castiglione A.; Tiribelli C. The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterology* 2006, 6, 33. <https://doi.org/10.1186/1471-230x-6-33>
27. Hepatic Steatosis Index (HSI) MD App Available on <https://www.mdapp.co/hepatic-steatosis-index-hsi-calculator-357> (Assessed on 23.06.2023)
28. Lee J.H.; Kim D.; Kim H.J.; Lee C.H.; Yang J.I.; Kim W.; Kim Y.J.; Yoon J.H.; Cho S.H.; Sung M.W.; Lee H.S. Hepatic steatosis index: a simple screening tool reflecting nonalcoholic fatty liver disease. *Dig Liver Dis* 2010, 42, 503–508. doi: 10.1016/j.dld.2009.08.002.
29. Cordeiro A.; Pereira S.E.; Saboya C.J.; Ramalho A. Nonalcoholic Fatty Liver Disease Relationship with Metabolic Syndrome in Class III Obesity Individuals. *Biomed Res Int* 2015, 2015, 839253. doi: 10.1155/2015/839253.
30. Dai C.Y.; Fang T.J.; Hung W.W.; Tsai H.J.; Tsai Y.C. The Determinants of Liver Fibrosis in Patients with Nonalcoholic Fatty Liver Disease and Type 2 Diabetes Mellitus. *Biomedicines* 2022, 10, 1487. <https://doi.org/10.3390/biomedicines10071487>.
31. Lomonaco R.; Bril F.; Portillo-Sánchez P.; Ortiz-López C.; Orsak B.; Biernacki D.; Lo M.; Suman A.; Weber M.; Cusi K. Metabolic Impact of Nonalcoholic Steatohepatitis in Obese Patients With Type 2 Diabetes. *Diabetes Care* 2016, 39, 632–638. <https://doi.org/10.2337/dc15-1876>
32. Yu Y.; Yu Y.; Wang Y.; Chen Y.; Wang N.; Wang B.; Lu Y. Nonalcoholic fatty liver disease and type 2 diabetes: an observational and Mendelian randomization study. *Frontiers in Endocrinology* 2023, 14. <https://doi.org/10.3389/fendo.2023.1156381>
33. Lonardo, A.; Ballestri, S.; Guaraldi, G.; Nascimbeni, F.; Romagnoli, D.; Zona, S.; Targher, G. Fatty liver is associated with an increased risk of diabetes and cardiovascular disease - Evidence from three different disease models: NAFLD, HCV and HIV. *World J. Gastroenterol.* 2016, 22, 9674–9693.
34. Popescu, M.S.; Drocas, A.I.; Osman, A.; Firu, D.M.; Padureanu, V.; Marginean, C.M.; Pirvu, D.C.; Mitruț, R.; Margaritescu, D.N.; Radu, A.; et al. The Extent of Insulin Resistance in Patients That Cleared Viral Hepatitis C Infection and the Role of Pre-Existent Type 2 Diabetes Mellitus: A Meta-Analysis. *Reports* 2022, 5, 42. <https://doi.org/10.3390/reports5040042>.
35. Bazick J.; Donithan M.; Neuschwander-Tetri B.A.; Kleiner D.; Brunt E.M.; Wilson L.; Doo E.; Lavine J.; Tonascia J.; Loomba R. Clinical Model for NASH and Advanced Fibrosis in Adult Patients With Diabetes and NAFLD: Guidelines for Referral in NAFLD. *Diabetes Care* 2015, 38, 1347–1355. <https://doi.org/10.2337/dc14-1239>.
36. Chen K.; Sng W.K.; Hui-Min Quah J.; Liu J.; Chong B.Y.; et al. Clinical spectrum of non-alcoholic fatty liver disease in patients with diabetes mellitus. *PLoS One* 2020, 15, e0236977. doi: 10.1371/journal.pone.0236977
37. Julián MT.; Pera G.; Soldevila B.; Caballería L.; Julve J.; Puig-Jové C.; Morillas R.; Torán P.; Expósito C.; Puig-Domingo M.; Castelblanco E.; Franch-Nadal J.; Cusi K.; Mauricio D.; Alonso N. Atherogenic dyslipidemia, but not hyperglycemia, is an independent factor associated with liver fibrosis in subjects

with type 2 diabetes and NAFLD: a population-based study. *European Journal of Endocrinology* 2021, **184**, 587–596. <https://doi.org/10.1530/eje-20-1240>

- 38. Martin A.; Lang S.; Goeser T.; Demir M.; Steffen H.M.; Kasper F. Management of Dyslipidemia in Patients with Non-Alcoholic Fatty Liver Disease. *Current Atherosclerosis Reports* 2022, **24**, 533–546. doi: 10.1007/s11883-022-01028-4
- 39. Prashanth M.; Ganesh H.K.; Vimal M.V.; John M.; Bandgar T.; et al. Prevalence of Nonalcoholic fatty Liver Disease in Patients with Type 2 Diabetes Mellitus. *Japi* 2009, **57**.
- 40. Friedman A.N.; Fadem S.Z. Reassessment of albumin as a nutritional marker in kidney disease. *J Am Soc Nephrol*, 2010, **21**, 223–230. doi: 10.1681/ASN.2009020213.
- 41. Mofrad P.; Contos M.J.; Haque M.; et al. Clinical and histologic spectrum of nonalcoholic fatty liver disease associated with normal ALT values. *Hepatolog* 2003, **37**, 1286–1292. doi: 10.1053/jhep.2003.50229.
- 42. Mandal A.; Bhattacharai B.; Kafle P.; Khalid M.; Jonnadula S.K.; Lamicchane J.; Kanth R.; Gayam V. Elevated Liver Enzymes in Patients with Type 2 Diabetes Mellitus and Non-alcoholic Fatty Liver Disease. *Cureus* 2018, **10**, e3626. doi: 10.7759/cureus.3626.
- 43. Harris HE. Elevated Liver Function Tests in Type 2 Diabetes. *Clin Diabetes* 2005, **23**, 115–119. <https://doi.org/10.2337/diaclin.23.3.115>
- 44. Shrestha N.; Bhatt N.P.; Neopane P.; Dahal S.; Regmi P.; Khanal M.; Shrestha R. Hepatic involvement with elevated liver enzymes in Nepalese subjects with type 2 diabetes mellitus. *Int J Biochem Res Rev* 2017, **16**, 1–8. doi: 10.7759/cureus.3626
- 45. Dominici S.; Paolicchi A.; Corti A.; Maellaro E.; Pompella A. Prooxidant reactions promoted by soluble and cell-bound γ -glutamyltransferase activity. *Methods Enzymol* 2005, **401**, 484–501. doi: 10.1016/S0076-6879(05)01029-3.
- 46. Kunutsor S.K. Gamma-glutamyltransferase-friend or foe within? *Liver Int* 2016, **36**, 1723–1734. doi: 10.1111/liv.13221.
- 47. Chen L.W.; Huang M.S.; Shyu Y.C.; Chien R.N. Gamma-glutamyl transpeptidase elevation is associated with metabolic syndrome, hepatic steatosis, and fibrosis in patients with nonalcoholic fatty liver disease: A community-based cross-sectional study. *Kaohsiung J Med Sci* 2021, **37**, 819–827. doi: 10.1002/kjm2.12395.
- 48. Jung C.H.; Lee B.; Choi D.H.; Jung S.H.; Kim B.Y.; Kim C.H.; Kang S.K.; Mok J.O. Association of grade of non-alcoholic fatty liver disease and glycated albumin to glycated hemoglobin ratio in patients with type 2 diabetes mellitus. *Diabetes Research and Clinical Practice* 2017, **125**, 53–61. <https://doi.org/10.1016/j.diabres.2016.12.017>
- 49. Kahraman N.K.; Kahraman C.; Kocak F.E.; Cosgun S.; Sanal B.; Korkmaz M.; Bayham Z.; Zeren S. Predictive value of neutrophil-to-lymphocyte ratio in the severity of non-alcoholic fatty liver disease among type 2 diabetes patients. *Acta Gastroenterol Belg* 2016, **79**, 295–300
- 50. Kapici O.B.; Abus S.; Ayhan S.; Karaagac M.; Sirik M. Investigation of hemogram and biochemistry parameters in non-alcoholic fatty liver disease. *Medicine Science | International Medical Journal* 2023, **12**, 63. <https://doi.org/10.5455/medscience.2022.12.258>
- 51. Luo Y.; Wang C.; Zhang T.; He X.; Hao J.; Shen A.; Zhao H.; Chen S.; Ren L. Factors Associated with Liver Fibrosis in Chinese Patients with Type 2 Diabetes Mellitus and Non-Alcoholic Fatty Liver Disease. *International Journal of General Medicine* 2023, **16**, 293–302. <https://doi.org/10.2147/ijgm.s395948>
- 52. Milovanovic A.T.; Stojkovic L.M.; Dumić I.; Jocić N.; Pavlović Marković A.; Dragasević S.; Jovićić I.; Lukić S.; Popović D.; Milošavljević T. Diagnostic Accuracy of Platelet Count and Platelet Indices in Noninvasive Assessment of Fibrosis in Nonalcoholic Fatty Liver Disease Patients. *Canadian Journal of Gastroenterology and Hepatology* 2017, 1–5. <https://doi.org/10.1155/2017/6070135>
- 53. Tingle S.J.; Ma G.R.S.; Goodfellow M.; Moir J.A.; White S.A. NARCA: A novel prognostic scoring system using neutrophil albumin ratio and Ca19-9 to predict overall survival in palliative pancreatic cancer. *J. Surg. Oncol* 2018, **118**, 680–686. doi: 10.1002/jso.25209.
- 54. Wang B.; Li D.; Cheng B.; Ying B.; Gong Y. The Neutrophil Percentage-to-Albumin Ratio Is Associated with All-Cause Mortality in Critically Ill Patients with Acute Kidney Injury. *Biomed Res Int* 2020, **18**, 2020, 5687672. doi: 10.1155/2020/5687672.
- 55. Gong Y.; Li D.; Cheng B.; Ying B.; Wang B. Increased neutrophil percentage-to-albumin ratio is associated with all-cause mortality in patients with severe sepsis or septic shock. *Epidemiol Infect* 2020, **148**, e87. doi: 10.1017/S0950268820000771.
- 56. Marjot T.; Moola A.; Cobbold J.F.; Hodson L.; Tomlinson J.W. Nonalcoholic Fatty Liver Disease in Adults: Current Concepts in Etiology, Outcomes, and Management. *Endocr. Rev* 2020, **41**, 66–117. doi: 10.1210/endrev/bnz009.
- 57. Loosen SH.; Demir M.; Kunstein A.; Jördens M.; Qvarskhava N.; Luedde M.; Luedde T.; Roderburg C.; Kostev K. Variables associated with increased incidence of non-alcoholic fatty liver disease (NAFLD) in patients with type 2 diabetes. *BMJ Open Diabetes Res. Care* 2021, **9**, e002243. doi: 10.1136/bmjdrc-2021-002243.

58. Okdahl T.; Wegeberg AM.; Pociot F.; Brock B.; Størling J.; Brock C. Low-grade inflammation in type 2 diabetes: A crosssectional study from a Danish diabetes outpatient clinic. *BMJ Open* 2022, 12, e062188. doi: 10.1136/bmjopen-2022-062188.

59. You G.; Ding J.; Shen J.; Wang Y.; Sun Y. Association between serum ferritin and non-alcoholic fatty liver disease among middle-aged and elderly Chinese with normal weight. *Asia Pacific Journal of Clinical Nutrition* 2019, 28, 747–753. [https://doi.org/10.6133/apjcn.201912_28\(4\)0011](https://doi.org/10.6133/apjcn.201912_28(4)0011)

60. Kunutsor S.K.; Apekey T.A.; Walley J.; Kain K. Ferritin levels and risk of type 2 diabetes mellitus: an updated systematic review and meta-analysis of prospective evidence. *Diabetes/Metabolism Research and Reviews* 2013, 29, 308–318. (<https://doi.org/10.1002/dmrr.2394>).

61. Amin R.F.; El Bendary A.S.; Ezzat S.E.; Mohamed W.S. Serum ferritin level, microalbuminuria and non-alcoholic fatty liver disease in type 2diabetic patients. *Diabetes and Metabolic Syndrome* 2019, 13, 2226–2229. <https://doi.org/10.1016/j.dsx.2019.05.030>.

62. Yan J.X.; Pan B.J.; Zhao P.P.; Wang L.T.; Liu J.F.; Fu S.B. Serum ferritin is crrelated with non-alcoholic fatty liver disease in middle-aged and older patients with type 2 diabetes. *Endocr Connect* 2021, 29, 10, 1560-1569. doi: 10.1530/EC-21-0367.

63. Lombardi R.; Pisano G.; Fargion S. Role of serum uric acid and ferritin in the development and progression of NAFLD. *International Journal of Molecular Sciences* 2016, 17, 548. 10.3390/ijms17040548

64. Sachinidis A.; Doumas M.; Imprialos K.; Stavropoulos K.; Katsimardou A.; Athyros V.G. Dysmetabolic iron overload in metabolic syndrome. *Current Pharmaceutical Design* 2020, 26, 1019–1024. doi: 10.2174/1381612826666200130090703.

65. Lee H.J.; Choi J.S.; Lee H.J.; Kim W.H.; Park S.I.; Song J. Effect of excess iron on oxidative stress and gluconeogenesis through hepcidin during mitochondrial dysfunction. *Journal of Nutritional Biochemistry* 2015, 26, 1414–1423. doi: 10.1016/j.jnubio.2015.07.008.

66. Park S.H.; Kim B.I.; Yun J.W.; Kim J.W.; Park D.I.; Cho Y.K.; Sung I.K.; Park C.Y.; Sohn CI.; Jeon W.K.; et al. Insulin resistance and C-reactive protein as independent risk factors for non-alcoholic fatty liver disease in non-obese Asian men. *Journal of Gastroenterology and Hepatology* 2004, 19, 694–698. doi: 10.1111/j.1440-1746.2004.03362.x.

67. Wong V.W.; Hui A.Y.; Tsang S.W.; Chan J.L.; Tse A.M.; Chan K.F.; So W.Y.; Cheng A.Y.; Ng W.F.; Wong G.L.; et al. Metabolic and adipokine profile of Chinese patients with nonalcoholic fatty liver disease. *Clinical Gastroenterology and Hepatology* 2006, 4, 1154–1161. doi: 10.1111/j.1365-2036.2005.02334.x.

68. Mayneris-Perxachs J.; Cardellini M.; Hoyles L.; Latorre J.; Davato F.; Moreno-Navarrete J.M.; Arnoriaga-Rodríguez M.; Serino M.; Abbott J.; Barton R. H.; et al. Iron status influences non-alcoholic fatty liver disease in obesity through the gut microbiome. *Microbiome* 2021, 104. doi: 10.1186/s40168-021-01052-7.

69. Haukeland J.W.; Damås J.K.; Konopski Z.; Løberg E.M.; Haaland T.; Goverud I.; Torjesen P.A.; Birkeland K.; Bjøro K.; Aukrust P: Systemic inflammation in nonalcoholic fatty liver disease is characterized by elevated levels of CCL2. *J Hepatol* 2006, 44, 1167-1174.

70. Yoneda M.; Mawatari H.; Fujita K.; Iida H.; Yonemitsu K.; Kato S.; Takahashi H.; Kirikoshi H.; Inamori M.; Nozaki Y.; Abe Y.; Kubota K.; Saito S.; Iwasaki T.; Terauchi Y.; Togo S.; Maeyama S.; Nakajima A: High-sensitivity C-reactive protein is an independent clinical feature of nonalcoholic steatohepatitis (NASH) and also of the severity of fibrosis in NASH. *J Gastroenterol* 2007, 42, 573-582.

71. Riquelme A.; Arrese M.; Soza A.; Morales A.; Baudrand R.; Pérez-Ayuso RM.; González R.; Alvarez M.; Hernández V.; García-Zattera MJ.; Otarola F.; Medina B.; Rigotti A.; Miquel JF.; Marshall G.; Nervi F. Non-alcoholic fatty liver disease and its association with obesity, insulin resistance and increased serum levels of C-reactive protein in Hispanics. *Liver Int* 2009, 29, 82-88.

72. Kao PC.; Shiesh SC.; Wu TJ. Serum C-reactive protein as a marker for wellness assessment. *Ann Clin Lab Sci* 2006, 36, 163-169.

73. Oruc N.; Ozutemiz O.; Yuce G.; et al. Serum procalcitonin and CRP levels in non-alcoholic fatty liver disease: a case control study. *BMC Gastroenterol* 2009, 9, 16. <https://doi.org/10.1186/1471-230X-9-16>

74. Ndumele C.E.; Nasir K.; Conceição R.D.; Carvalho J.A.; Blumenthal R.S.; Santos R.D. Hepatic steatosis, obesity, and the metabolic syndrome are independently and additively associated with increased systemic inflammation. *Arterioscler Thromb Vasc Biol*, 2011 31, 1927-1932. doi: 10.1161/ATVBAHA.111.228262.

75. Viallon A.; Zeni F.; Pouzet V.; Lambert C.; Quenet S.; Aubert G.; Guyomarch S.; Tardy B.; Bertrand J.C. Serum and ascitic procalcitonin levels in cirrhotic patients with spontaneous bacterial peritonitis: diagnostic value and relationship to pro-inflammatory cytokines. *Intensive Care Med* 2000, 26, 1082-1088. doi: 10.1007/s001340051321.

76. Dandona P.; Nix D.; Wilson MF.; Aljada A.; Love J.; Assicot M.; Bohuon C. Procalcitonin increase after endotoxin injection in normal subjects. *J Clin Endocrinol Metab* 1994, 79:1605-1608.

disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.