

Article

Not peer-reviewed version

Using Low-Cost Sensing Technology to Assess Ambient and Indoor Air Quality in New York during and after the COVID-19 Pandemic

Justin Holder , Jamelia Jordan , [Kera Johnson](#) , [Ayodele Akinremi](#) , [Dawn Roberts-Semple](#) *

Posted Date: 5 July 2023

doi: 10.20944/preprints202307.0321.v1

Keywords: AirBeam2; indoor air; fine particulate matter; seasonal variations; sensor; COVID-19



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Using Low-Cost Sensing Technology to Assess Ambient and Indoor Air Quality in New York during and after the COVID-19 Pandemic

Justin Holder ¹, Jamelia Jordan ¹, Kera Johnson ¹, Ayodele Akinremi ² and Dawn Roberts-Semple ^{1,*}

¹ Department of Earth and Physical Sciences, City University of New York – York College, 94-20 Guy R. Brewer Blvd. Jamaica, NY 11451; justin.holder@yorkmail.cuny.edu; jamelia.jordan@yorkmail.cuny.edu; kera.johnson@yorkmail.cuny.edu

² Department of Physical Therapy, School of Sciences, Hampton University, 100 E Queen St., Hampton, VA 23669 United States; ayodele.akinremi@hamptonu.edu

* Correspondence: drobertssemple@york.cuny.edu; Tel.: +(718)-262-2775

Abstract: Air pollution is a leading cause of death in the United States, and is associated with adverse health outcomes, including increased vulnerability to coronavirus disease 2019 (COVID-19). The AirBeam2 was used to measure particulate matter with a diameter of 2.5 micrometers or smaller (PM_{2.5}), to investigate differences between indoor and ambient levels at seven private homes in New York during and after the COVID-19 lockdown. Measurements taken in 2020 winter, spring and fall and in fall 2022 showed that 90% of the time, indoor PM_{2.5} levels exceeded outdoor levels both during and after the COVID-19 lockdown, $p = 0.03$, and exceeded safety levels. Higher indoor PM_{2.5} levels attributed to little or no ventilation from cooking and smoke in kitchens and basements, were significantly greater in fall than in winter. Higher ambient PM_{2.5} levels were attributed to vehicular traffic at a street-facing sampling site. PM_{2.5} sources identified in this study may help in devising control strategies to improve indoor air quality (IAQ), and consequently alleviate respiratory health effects. These findings may be used as a basis for in-house modifications including natural ventilation and use of air purifiers to reduce exposures, mitigate future risks, and prevent potential harm to vulnerable residents.

Keywords: AirBeam2; indoor air; fine particulate matter; seasonal variations; sensor; COVID-19

1. Introduction

Poor air quality is associated with adverse health outcomes and is a leading cause of death in the U.S. [1–3]. Since air pollutants are ubiquitous, many studies have investigated indoor and outdoor air quality, as human activities and emission sources vary both in outdoor and indoor environments, mainly vehicular emission, thermal power plants, combustion, food grilling, and frying [4]. Particulate matter (PM) is a major pollutant known to degrade air quality and is categorized by size fractions. Coarse particles referred to as PM₁₀ have a diameter of 2.5 – 10 microns; they are produced mainly from construction sites, mines, farms, and dirt roads, and can irritate the eyes, nose, and throat. Fine particles that measure 2.5 micrometers or less in diameter, referred to as PM_{2.5} includes a subcategory, or ultrafine particles less than 1 micron in diameter [4]; these pose a greater health risk because they can reach deep into the lungs and enter the bloodstream. PM_{2.5} is usually found in smoke from fires, power plants and motor vehicles and have been linked to chronic and acute respiratory, cardiovascular, and nervous system illnesses that can result in mortality, even at concentrations below the 24-hour national ambient air quality standard (NAAQS) of 35 $\mu\text{g}/\text{m}^3$ [4,5].

The association between PM_{2.5} and respiratory infections is a serious concern that heightened after COVID-19 was declared by the World Health Organization (WHO) as a public health emergency [6]. Indoor air pollution (IAP) remains a concern, yet data are seriously lacking. IAP has been classified as one of the top five environmental health hazards [7], and recent studies have found that

it could exceed ambient levels [8]. Since most Americans spend more than 90% of their time indoors, a healthy indoor environment is very important [9]. While compulsory lockdown during the COVID-19 pandemic apparently contributed to improved ambient air quality in most cities, exposure to IAP was significant [10]. IAQ issues are generally caused by household energy conservation, a lack of adequate ventilation and the use of cleaning products and disinfectants which intensified during the COVID-19 pandemic. Higher IAP is also attributed to the absence of meteorological influence particularly wind speed, known to dilute air pollutant concentrations outdoor [11]. Recent scientific evidence has shown that indoor air can be more seriously polluted than outdoor air in urban areas including the largest and most industrialized cities in the world [9]. With no IAQ health standards, high IAP levels imply greater health risks among vulnerable residents especially children, the elderly and individuals suffering chronic respiratory and cardiovascular diseases [9].

During the COVID-19 pandemic lockdown, restrictions on business and transportation resulted in a dramatic reduction in air pollution levels in most cities around the world; however, the apparent improvement in air quality did not necessarily reduce exposure to IAP and presumably, vulnerability to respiratory health effects. As most people utilized online sources for work and shopping, this change in reality raised the question of whether the risk of exposure to PM_{2.5} in homes increased with more time spent indoors. Air quality in homes could be worse because pollutants in such environments are restricted by walls and ceilings which are absent in outdoor spaces. Exposure to PM_{2.5} could result in an elevated risk for acute or chronic respiratory infections and disease, susceptibility, and exacerbation of inflammatory stimulus in young and even healthy individuals [12]. PM exposure has also been associated with increased medical visits and hospitalizations [13]. Wu et al. (2020) found that an increase of only 1 µg/m³ in long-term average PM_{2.5} levels was associated with a significant increase of 15% in the COVID-19 death rate [14].

Cooking is a major contributor of indoor air pollution because the different types of foods prepared may produce a variety of volatile organic compounds (VOCs) and PM, that may cause detrimental health effects [15]. Approximately 3.8 million people worldwide die every year from illnesses attributable to the harmful indoor air from cookstoves and fuel [16]. Additionally, kitchens are regularly characterized by a micro-climate due to food preparation processes with higher temperatures and humidity [17]. Generally, most adults spend over 10% of their time in kitchen areas to prepare meals, eat, or clean up after meal preparation [18]. Harmful compounds and particles are produced from incomplete combustion of biomass at fireplaces, open fires and stoves due to an insufficient supply of oxygen [18]. IAQ problems also result from the use of solid fuels in stoves for household heating and cooking, especially from stir-frying and deep-frying methods, and smoking in homes [19]. The basement is also a microenvironment where dust, mold and high relative humidity may persist due to limited ventilation and stored items such as unused furniture, lacquers, paints, and gasoline contribute to an odor called the "basement smell" mainly due to putrefaction processes and the growth of microorganisms [18]. Such biological air pollution may cause serious health effects. Even under normal conditions, infiltration of PM_{2.5} through ventilation systems could further degrade IAQ. Outdoor air tends to enter and leave buildings by natural or mechanical ventilation through open windows or unsealed cracks in doors; IAQ can also be influenced by ambient temperature and humidity [17,20]. Depending on the weather conditions, people tend to open or close their windows, operate air conditioning systems, humidifiers, and heaters. Through these processes, outdoor air moves indoor at a rate of replacement termed the air exchange rate which when low, contributes to an increase in air pollutant levels [4].

This pilot study aimed to provide a simple uncomplicated way to determine PM_{2.5} levels indoors for comparison with ambient levels, and to identify the potential risks of exposure to indoor emission sources that can be targeted for reduction or removal through increased ventilation or supplemental filtration for improved IAQ. The study does not intend to discuss or draw conclusions about health impacts of measured PM_{2.5} concentrations. The complexity of identifying health risks, personal health status, level of exposure, and other environmental factors; however, is noteworthy, as it is not possible to fully understand potential health impacts or risks solely by detecting PM_{2.5} levels using low-cost technology such as the AirBeam2.

2. Materials and Methods

2.1. Study Sites

This study was conducted at seven (7) residential homes in New York (Figure 1) using two AirBeam2 concurrently: one for indoor and another for outdoor sampling at each location. Measurements were taken in the winter, spring and fall in the months of February, October and November of 2020, April of 2021 and September through November of 2022. PM_{2.5} samples were taken indoors (kitchen, basement, bedroom) and outdoors for approximately 2-6 hours during and after the COVID-19 pandemic lockdown. Indoor samples were collected in the living room and basement at Home 1 in Queens Village; kitchen of Home 2 in Cambria Heights; living room of Home 3 in Flushing; basement of Home 4 in Hollis; and kitchens of Home 5 in Garden City, Home 6 in Springfield Gardens and Home 7 in St. Albans. After each sampling session, the data were downloaded and STATA version 15 was used to perform statistical analysis. Daily average ambient PM_{2.5} levels in this study were compared to those obtained from the EPA regulatory/central monitoring station in Queens, NY during the same period [21].




Figure 1. Map of Residential sites in New York.

2.2. Suitability of AirBeam2 for PM_{2.5} Air Quality Monitoring

The AirBeam2 used in this study is a portable palm-sized device that collects PM_{2.5}, while paired with a BLU android cellular phone. It uses a light scattering method to measure PM₁, PM_{2.5} and PM₁₀, as air is drawn through a sensing chamber and light from a laser scatters particles in the airstream [22]. A record of the measurement is sent each second to the Aircasting mobile app to permit real-time visualization, and pooled data can be crowdsourced on an AirCasting air quality map. The AirBeam2 is among low-cost sensors approved by the United States Environmental Protection Agency (USEPA) [23], to measure indoor and outdoor PM concentrations [17,24–26]. It provides a simple and quick way to determine spatial and temporal PM_{2.5} levels, informing possible actions that may be taken to reduce unacceptable levels of exposure. Therefore, data quality and performance of the air sensor can help to advance IAQ management and improve our understanding of total exposure to particle pollutants in the home environment. The device can also provide greater insights into potential benefits and limitations of applying the sampled data to improve IAQ, including the

need to monitor other pollutants besides PM and assess their impacts on human health. For future studies that include larger sample sizes would be considered for correlation between PM levels and health risks in homes.

3. Results

3.1. Comparison of PM_{2.5} Levels in 2020, 2021 and 2022

There was no significant difference in indoor, outdoor, and ambient PM_{2.5} measurements across the three years (Fall 2020, Winter 2021, Winter 2022). Data for the same study periods obtained from the nearest central monitoring station were compared to the sampled outdoor PM_{2.5} concentrations. In 2020, 2021 and 2022, average indoor PM_{2.5} levels were higher than average outdoor levels at 13.33 $\mu\text{g}/\text{m}^3$, 9.0 $\mu\text{g}/\text{m}^3$ and 16.5 $\mu\text{g}/\text{m}^3$, respectively. The city ambient PM_{2.5} levels, on the other hand, were found to be the lowest across the years. In 2020 and 2021, there was an increase in indoor PM_{2.5} levels, whereas there was a decrease in outdoor and city ambient levels of PM_{2.5}. In 2022, indoor concentrations of PM_{2.5} increased whereas the concentrations for outdoor and city ambient decreased.

As shown in Figure 2, there were no significant changes in PM_{2.5} measurements indoor versus outdoor in the fall season, and sampled outdoor PM_{2.5} levels in the fall versus City ambient levels recorded at the regulatory monitoring station.




Figure 2. Comparison of PM_{2.5} levels by year.

3.2. Comparison of PM_{2.5} Levels by Season

Indoor PM_{2.5} levels were consistently high in both winter and fall, even though they were not statistically significant ($p=0.42$); indoor PM_{2.5} levels were higher than sampled outdoor and city ambient levels in the winter. Additionally, indoor PM_{2.5} levels in the fall were higher than outdoor and city ambient levels with values of 14.6 $\mu\text{g}/\text{m}^3$, 6.2 $\mu\text{g}/\text{m}^3$ and 6.16 $\mu\text{g}/\text{m}^3$, respectively. Overall, indoor PM_{2.5} levels during the fall were significantly greater than indoor winter levels. Indoor PM_{2.5} levels in the fall were also higher than the measured outdoor and City ambient levels.




Figure 3. Comparison of PM_{2.5} levels by seasons.

The mean PM_{2.5} levels for indoor and outdoor were the same ($p = 0.03$), and the sampled outdoor PM_{2.5} levels were comparable to the City ambient levels. Conversely, 90% (1 out of 7) of the time indoor PM_{2.5} levels exceeded outdoor levels. Additionally, approximately 50% of the time, indoor concentrations of PM_{2.5} exceeded the safety level ($8.75 \mu\text{g}/\text{m}^3$ for 6-hr exposure) and 100% of outdoor levels were within the safety levels ($8.75 \mu\text{g}/\text{m}^3$ for 6-hr exposure). The average indoor PM_{2.5} level for Home 1 was $4 \mu\text{g}/\text{m}^3$ and sampling was conducted in the living room in February 2021 for four weeks (27 days). Higher PM_{2.5} levels were recorded at Homes 2 and 3, averaging $14 \mu\text{g}/\text{m}^3$. At Home 2, samples were measured in the kitchen in February 2021 for two weeks (14 days) while at home 3, samples were measured for one week (7 days) in the living room in October 2020. At Home 4, the average PM_{2.5} indoor level was $5 \mu\text{g}/\text{m}^3$ and the samples were collected in the basement at the end of October 2020 into November 2020 for one week (7 days).

The average indoor PM_{2.5} level for Homes 5 and 6 were $21 \mu\text{g}/\text{m}^3$ and $24 \mu\text{g}/\text{m}^3$, respectively and samples were collected in the kitchen at both homes. PM_{2.5} was measured for one week (7 days) at Home 5 in November 2020, and at Home 6, samples were collected for almost two weeks (12 days) from late September 2022 into early November 2022. Finally, the average PM_{2.5} level at Home 7 was $9 \mu\text{g}/\text{m}^3$ and samples were collected in the kitchen for almost three weeks (18 days) in late October 2022 and early November 2022. The outdoor PM_{2.5} levels for Home 1, Home 3 and Home 6 were $7 \mu\text{g}/\text{m}^3$, respectively. Home 4 had the lowest PM_{2.5} levels, while Home 2 had the highest level of $9 \mu\text{g}/\text{m}^3$. Overall, the data obtained showed that outdoor PM_{2.5} concentrations were 1.8 times greater in the winter than in the fall.

4. Discussion

4.1. Temporal Variations of Indoor PM_{2.5} Levels

In this study, higher indoor PM_{2.5} levels both during and after the COVID-19 lockdown suggested a combination of emissions from indoor activities and outside sources [27,28]. Indoor PM_{2.5} levels were mostly influenced by human activity and smoke [29]; especially from cooking, as meal preparation by frying, toasting and boiling may release PM_{2.5} with various compositions including saturated and unsaturated fatty acids [18,30]. Extended exposure times in confined spaces of residential homes substantially affect air quality because they often lack proper ventilation; hence, PM_{2.5} levels tend to be more concentrated than outdoor levels [31]. Fuel oil and natural gas was used in most residential homes in this study, and only a few stoves were furnished with vents to extract fumes while cooking during sampling, potentially increasing PM_{2.5} levels [17]. Indoor gas stove use

for cooking is associated with an increased risk of asthma among children and is prevalent in 35% of households in the US. Pollutants released by combustion in homes with range hoods may not vent outdoors or be removed effectively [32]. Source reduction and control is the most effective solution to indoor air pollution, although ventilation, supplemental filtration and air cleaning can help to remove PM_{2.5} and other pollutants.

4.2. Temporal Variations of Outdoor PM_{2.5}

Previous studies have reported greater variability in ambient PM levels than indoor levels due to local meteorological conditions, road traffic, power generation plants, industries, etcetera [33]. Restrictions in human activities and the consequent dramatic decrease in ambient air pollution occurred during the COVID-19 lockdown, as most people stayed at home at the onset of the pandemic [34]. The lowest ambient PM_{2.5} levels were measured in 2020 but increased in 2021 and 2022, and City ambient measurements from the central monitoring station in NYC also reflected this decline (Table 1). Similarly, ground-based observations of PM_{2.5} and other air pollutants showed reductions in metropolitan cities in South and East Asia, Europe and North America during COVID 19 lockdown, that accounted for 67% of anthropogenic emissions and 5% from the transportation sector [35]. Higher ambient PM_{2.5} levels were only recorded at one sampling site facing a street with moderate vehicular traffic. Similarly, a study conducted at 39 schools in Barcelona, Spain, reported that school playgrounds in close proximity to road traffic exhibited higher levels of PM_{2.5} than other areas [22]. In Po Valley, Italy, poor air quality pre-lockdown was a result of large population, industrial activities, geographical factors and weather conditions but although most anthropogenic activities halted during the lockdown, there was no significant decrease in PM during the lockdown compared to 2019 [11]. This suggested that metrological factors could have a greater influence on PM levels than emissions from traffic [11].

Table 1. Comparison of PM_{2.5} levels by year.

	2020 (Fall) Mean (SE)	2021 (Winter) Mean (SE)	2022 (Fall) Mean (SE)	p-Value
Indoor	13.33 (4.63)	9.0 (5.0)	16.5 (7.5)	0.93
Outdoor	5.33 (1.20)	8.0 (1.0)	7.5 (0.5)	0.65
City Ambient	6.03 (1.24)	6.5 (0.69)	6.35 (0.05)	0.11

4.3. Seasonal Variations of Indoor PM_{2.5}

PM_{2.5} was measured up to 6 hours with an ambient standard of 8.75 µg/m³ (35 µg/m³ for 24-hr period); but most indoor levels exceeded this standard (Table 2). During the winter and fall seasons, people typically keep their windows and doors closed to conserve energy; however, this could result in less air circulation, gas exchange, and increased indoor PM_{2.5} levels. During the winter, particles are usually resuspended in the air from daily indoor activities because of poor ventilation but reduce in the summer season when air exchange rates are higher as people open their windows and doors [36]. A major contributing factor to higher PM_{2.5} levels could be cooking, as Begum et al. (2009) observed that food preparation in the kitchen could create a specific micro-climate because of high temperatures and humidity [18]. Although the increase in temperature is expected to cause PM to decrease as the particles gain kinetic energy, move faster, expand and are dispersed [37] and relative humidity could lead to the deposition of larger particles as they become moisture laden [38], smaller particles remain suspended in indoor air for longer periods. With no meteorological influence indoors to disperse particles, PM_{2.5} increased; however, there is no national standard for indoor air quality to which the measurements could appropriately be compared.

Table 2. Comparison of PM_{2.5} levels by season.

	Winter (n = 2) Mean (SE)	Fall (n = 5) Mean (SE)	p-Value
Indoor	9.0 (5)	14.6 (3.5)	0.42
Outdoor	8.0 (1.0)	6.2 (0.69)	0.29
City Ambient	6.5 (0.98)	6.16 (0.68)	0.78

Table 3. Seasonal comparison between indoor and outdoor PM_{2.5} levels.

Homes	Season	InAv PM _{2.5} ($\mu\text{g}/\text{m}^3$)	OutAv PM _{2.5} ($\mu\text{g}/\text{m}^3$)	City Ambient
1	2021 (W)	4	7	5.8
2	2021 (W)	14	9	7.2
3	2020 (F)	14	7	3.9
4	2020 (F)	5	3	8.2
5	2020 (F)	21	6	6.0
6	2022 (F)	24	7	6.4
7	2022 (F)	9	8	6.3
Mean (Std Error)		13.0 (2.87)	6.71 (0.71)	6.25 (0.50)

4.4. Seasonal Variations of Outdoor PM_{2.5}

The samples obtained in this study showed that outdoor PM_{2.5} levels were 1.8 times greater than the fall as seen in Table 2. Previous studies have shown that during the winter season, anthropogenic emissions can influence outdoor PM_{2.5} levels via multiple sources and processes including the increased use of passenger vehicles and meteorological effects [39]. Seasonal variations affect outdoor air quality when combined with factors such as fossil fuel combustion for the heating of buildings and automobile operation which contribute to higher PM_{2.5} during the winter season [40]. The phenomenon of temperature inversion can also affect densely populated industrialized areas and prevent the dispersion of pollutants in cold weather and cloudless skies, thus increasing the concentration of pollutants in countries such as China, with health implications from PM_{2.5} pollution in the winter [41]. Jiang et al. (2022) observed two air pollution episodes, one mainly affected by air masses in which increased levels of water-soluble ions led to increased PM_{2.5} [42].

4.5. Health Implications of Higher Indoor PM_{2.5} Levels during COVID-19

Before the COVID-19 pandemic, ambient PM_{2.5} levels were much higher during the same months that the COVID-19 lockdown occurred [8]; however, being indoors during the COVID-19 pandemic lockdown seemed not to reduce exposure to PM_{2.5}. Previous studies have shown that inhalation of particle pollution was related to the prevalence of, and mortality due to COVID-19 [43]. Many pre-existing conditions that increased the morbidity of COVID-19 were the same as those that affected individuals due to long-term exposure of air pollution. Wu et al. (2020) found that 1 $\mu\text{g}/\text{m}^3$ of PM_{2.5} was linked to 15% in COVID-19 deaths with a magnitude that was 20 times greater than what was observed for PM_{2.5} and other mortalities [14]. Biological particles that can suspend in the air have the ability to attach themselves usually to coarser PM; however, they can also attach to PM_{2.5}, allowing them to directly transport into the respirable regions causing serious respiratory and circulation illnesses [44,45]. Further PM_{2.5} can upregulate ACE-2 receptor of SARS-CoV-2 virus, increasing the chance of the viral RNA entering the lungs and potentiating the risk of COVID-19 symptoms [46]. Although there is no robust evidence that short-term PM_{2.5} exposure could increase the chance of dying from COVID-19, the risk increases with long-term exposure [45]. Additionally, the use of ethanol-based and other disinfectants increased during the COVID-19 pandemic to eliminate SARS-CoV-2 on surfaces and indoor spaces could have changed the chemical composition of the air and increase PM concentrations after initial use [30]. A pilot study utilized the Foobot®, a low-cost

consumer air monitor to assess air quality in offices and households in South Texas before the COVID-19 pandemic (May – July 2019) and during the pandemic (June – September 2020) while employees worked from home. PM_{2.5}, air temperature and relative humidity were collected at 5-minute intervals to assess the health outcomes of participants before and during the COVID-19 pandemic. The study found that for all participants, PM_{2.5} levels in households were significantly higher than those measured in offices ($p < 0.05$), and the PM_{2.5} levels in all households also exceeded the health-based annual mean standard of 12 $\mu\text{g}/\text{m}^3$. Conversely, 90% of the offices studied were in compliance with the health standard [7]. This finding suggests that there could be a higher health risk for residents in their homes than at their places of work.

5. Conclusions

This study provided insights into the potential increased risk to PM_{2.5} exposure in micro-environments and micro-climates of residential homes in urban environments. We found that sampled PM_{2.5} levels indoors significantly exceeded outdoor levels at most private homes due to insufficient ventilation and absence of meteorological influence. Sampled PM_{2.5} levels outdoor were comparable to City ambient PM_{2.5} levels and were below the acceptable ambient standard unlike indoor PM_{2.5} levels. The link between PM_{2.5} and COVID-19 in previous studies suggests that high indoor PM_{2.5} in this study could potentially have posed greater respiratory health effects on residents than ambient PM_{2.5} concentrations. Although there are currently no widely accepted performance criteria used to standardize how measurements are made by low-cost air pollutant monitors or what thresholds limits should be used to determine health effects indoors, this study may initiate an alert to such cues. The findings in this study are valuable in the absence of widely accepted indoor PM_{2.5} health-based standards or thresholds for comparison. To ensure that indoor environments are safe and healthy in the meantime; however, air purifiers with filtration technologies, particularly high efficiency particulate air (HEPA) filters and systems based on the principle of electrostatic precipitation, are two of the most common indoor air-purification technologies that can reduce PM_{2.5} concentrations indoors. Additionally, natural ventilation with open windows is a common and economical approach that can help to dilute indoor PM. Future studies, should consider establishing a network of two or more monitors to compare pollutant levels in different spaces and at different times: before, during, and after activities such as cooking. Additionally, stricter limits than the current national ambient air quality standards should be considered for indoor environments; larger sample sets would better assess seasonal variations of PM_{2.5} and the health status of residents should be surveyed to explain the potential health impacts of indoor PM_{2.5} exposure.

Author Contributions: Investigation, literature review, methodology, writing—original draft preparation, and editing to completion, J.H.; Conceptualization, writing—original draft preparation, methodology, review and editing to completion, D.R.-S.; Literature review, writing, GIS mapping and visualization, methodology, and formatting, J.J.; Methodology, formal analysis, validation and review, A.A.; Writing, literature review, methodology, review and editing, K.J. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: This study neither involved human subjects nor required identifiable information to be used, studied, analyzed, or generated.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data discussed in this study are contained within the article.

Acknowledgments: We thank the Environmental Health Science (EHS) students who assisted with data collection including data for the EHS 351: Fundamentals of Air Pollution course project.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Caiazzo F, Ashok A, Waitz IA, Dong GH, Barrett SF. Air pollution and early deaths in the United States. Part I: Quantifying the impact of major sectors in 2005. *Atmospheric Environment*. 2013;79:198-208. doi:10.1016/j.atmosenv.2013.05.081
2. NIEHS. Air Pollution. National Institute of Environmental Health Science. Published 2018. <https://www.niehs.nih.gov/health/topics/agents/air-pollution/index.cfm>
3. Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E. Environmental and Health Impacts of Air Pollution: A Review. *Frontiers in Public Health*. 2020;8. doi:10.3389/fpubh.2020.00014
4. US EPA. Particulate Matter (PM). US EPA. Published 2022. <https://www.epa.gov/pm-pollution/particulate-matter-pm-basics>
5. Jones E, Laurent JGC, Young AM, et al. The effects of ventilation and filtration on indoor PM2.5 in office buildings in four countries. *Building and Environment*. 2021;200:107975. doi:10.1016/j.buildenv.2021.107975
6. Sohrabi C, Alsafi Z, O'Neill N, et al. World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). *International Journal of Surgery*. 2020;76:71-76. doi:10.1016/j.ijsu.2020.02.034
7. Roh T, Moreno-Rangel A, Baek J, Obeng AB, Hasan NT, Carrillo G. Indoor Air Quality and Health Outcomes in Employees Working from Home during the COVID-19 Pandemic: A Pilot Study. *Atmosphere*. 2021;12(12):1665. doi:10.3390/atmos12121665
8. Ferreira A, Barros N. COVID-19 and Lockdown: The Potential Impact of Residential Indoor Air Quality on the Health of Teleworkers. *International Journal of Environmental Research and Public Health*. 2022;19(10):6079. doi:10.3390/ijerph19106079
9. Indoor Air Quality. US EPA. Published 2021. <https://www.epa.gov/report-environment/indoor-air-quality>
10. Domínguez-Amarillo S, Fernández-Agüera J, García SC, González-Lezcano RA. Bad Air Can Also Kill: Residential Indoor Air Quality and Pollutant Exposure Risk during the COVID-19 Crisis. *International Journal of Environmental Research and Public Health*. 2020;17(19):7183. doi:10.3390/ijerph17197183
11. Pala D, Casella V, Larizza C, Malovini A, Bellazzi R. Impact of COVID-19 lockdown on PM concentrations in an Italian Northern City: A year-by-year assessment. *PLOS ONE*. 2022;17(3):e0263265. doi:10.1371/journal.pone.0263265
12. Kyung SY, Kim HS. Particulate-Matter Related Respiratory Diseases. *Tuberculosis and Respiratory Diseases*. 2020;83(2):116. doi:10.4046/trd.2019.0025
13. Roberts-Semple D, Gao Y. Evaluation of air pollution, local meteorology and urban public health. *International Journal of Environmental Technology and Management*. 2013;16(1/2):160. doi:10.1504/ijetm.2013.050656
14. Wu X, Nethery RC, Sabath MB, Braun D, Dominici F. Exposure to air pollution and COVID-19 mortality in the United States: A nationwide cross-sectional study. *medRxiv (Cold Spring Harbor Laboratory)*. Published online April 7, 2020. doi:10.1101/2020.04.05.20054502
15. Akteruzzaman M, Rahman MdA, Rabbi FM, et al. The impacts of cooking and indoor air quality assessment in the southwestern region of Bangladesh. *Heliyon*. 2023;9(1):e12852. doi:10.1016/j.heliyon.2023.e12852
16. World Health Organization: WHO. Household air pollution. *www.who.int*. Published online November 28, 2022. <https://www.who.int/news-room/fact-sheets/detail/household-air-pollution-and-health>
17. Bergmans B, Cattaneo A, Duarte RMBO, et al. Particulate matter indoors: a strategy to sample and monitor size-selective fractions. *Applied Spectroscopy Reviews*. 2022;57(8):675-704. doi:10.1080/05704928.2022.2088554
18. Begum BA, Paul SK, Hossain MS, Biswas SK, Hopke PK. Indoor air pollution from particulate matter emissions in different households in rural areas of Bangladesh. *Building and Environment*. 2009;44(5):898-903. doi:10.1016/j.buildenv.2008.06.005
19. Hao J, Li X, Zhao YP. Indoor Air Pollution and Its Control in China. In: *Springer EBooks*. ; 2014:145-170. doi:10.1007/978-1-4939-2572-3_7
20. Mansor AA, Hisham ANB, Abdullah S, Napi NNLM, Ahmed A, Ismail M. Indoor-Outdoor Air Quality Assessment in Nurseries. *IOP Conference Series: Earth and Environmental Science*. 2020;616(1):012001. doi:10.1088/1755-1315/616/1/012001
21. Us Epa O. Download daily data. Published online 2016. <https://www.epa.gov/outdoor-air-quality-data/download-daily-data>
22. HabitatMap. <https://www.habitatmap.org/airbeam/how-it-works>
23. Ryder O, Minor H, Brown S, Duvall R, Clements A, Freed R. AirBeam2 Quick Start Guide. Air Sensor Loan Programs. Published 2020.
24. Gao SS, Hiltz RW, Ross MK, Styler SA. Particulate matters: student-led air quality research in the third-year environmental chemistry classroom and the field. *Analytical and Bioanalytical Chemistry*. 2018;410(14):3223-3229. doi:10.1007/s00216-018-0972-3
25. Kortoçi P, Motlagh NH, Zaidan MA, et al. Air pollution exposure monitoring using portable low-cost air quality sensors. *Smart Health*. 2022;23:100241. doi:10.1016/j.smhl.2021.100241

26. Hegde S, Min KT, Moore JE, et al. Indoor Household Particulate Matter Measurements Using a Network of Low-cost Sensors. *Aerosol and Air Quality Research*. 2020;20(2):381-394. doi:10.4209/aaqr.2019.01.0046
27. Weschler CJ. Changes in indoor pollutants since the 1950s. *Atmospheric Environment*. 2009;43(1):153-169. doi:10.1016/j.atmosenv.2008.09.044
28. Kaunelienė V, Prasauskas T, Krugly E, et al. Indoor air quality in low energy residential buildings in Lithuania. *Building and Environment*. 2016;108:63-72. doi:10.1016/j.buildenv.2016.08.018
29. Xu R, Qi X, Dai G, et al. A Comparison Study of Indoor and Outdoor Air Quality in Nanjing, China. *Aerosol and Air Quality Research*. 2020;20(10):2128-2141. doi:10.4209/aaqr.2019.10.0496
30. Jiang J, Ding X, Isaacson KP, et al. Ethanol-based disinfectant sprays drive rapid changes in the chemical composition of indoor air in residential buildings. *Journal of Hazardous Materials Letters*. 2021;2:100042. doi:10.1016/j.hazl.2021.100042
31. Cao Z, Wang Y, Wang M. A comparison of concentrated contaminant removal in enclosure by using mixing ventilation and vortex ventilation. In: *Proceedings of the International Conference on Building Energy & Environment*, 2018:139-143.
32. Gruenwald T, Seals BA, Knibbs LD, Hosgood HD. Population Attributable Fraction of Gas Stoves and Childhood Asthma in the United States. *International Journal of Environmental Research and Public Health*. 2022;20(1):75. doi:10.3390/ijerph20010075
33. Bo M, Salizzoni P, Clerico M, Buccolieri R. Assessment of Indoor-Outdoor Particulate Matter Air Pollution: A Review. *Atmosphere*. 2017;8(8):136. doi:10.3390/atmos8080136
34. Li J, Tartarini F. Changes in Air Quality during the COVID-19 Lockdown in Singapore and Associations with Human Mobility Trends. *Aerosol and Air Quality Research*. 2020;20(8):1748-1758. doi:10.4209/aaqr.2020.06.0303
35. Zhang Z, Arshad A, Zhang C, Hussain S, Li W. Unprecedented Temporary Reduction in Global Air Pollution Associated with COVID-19 Forced Confinement: A Continental and City Scale Analysis. *Remote Sensing*. 2020;12(15):2420. doi:10.3390/rs12152420
36. Massey DB, Kulshrestha A, Masih J, Taneja A. Seasonal trends of PM₁₀, PM_{5.0}, PM_{2.5} & PM_{1.0} in indoor and outdoor environments of residential homes located in North-Central India. *Building and Environment*. 2012;47:223-231. doi:10.1016/j.buildenv.2011.07.018
37. Bagley M, Dobrijevic D. Properties of Matter: Gases. *livescience.com*. Published online February 16, 2022. <https://www.livescience.com/53304-gases.html>
38. Hernandez G. Temperature and humidity effects on particulate matter concentrations in a sub-tropical climate during winter. Published 2017. <https://www.semanticscholar.org/paper/Temperature-and-humidity-effects-on-particulate-in-Hernandez-Berry/e2d8748fa838f69fa8554ea6f3e0fcc3fbd13c9b>
39. Li L, Li QX, Huang L, et al. Air quality changes during the COVID-19 lockdown over the Yangtze River Delta Region: An insight into the impact of human activity pattern changes on air pollution variation. *Science of the Total Environment*. 2020;732:139282. doi:10.1016/j.scitotenv.2020.139282
40. Clements N, Keady P, Emerson J, Fierier N, Miller S. Seasonal variability of airborne particulate matter and bacterial concentrations in Colorado homes. *Atmosphere (Basel)*. 2018;9(4):133. doi:10.3390/atmos9040133
41. Sarmadi M, Rahimi S, Rezaei M, Sanaei D, Dianatinasab M. Air quality index variation before and after the onset of COVID-19 pandemic: a comprehensive study on 87 capital, industrial and polluted cities of the world. *Environmental Sciences Europe*. 2021;33(1). doi:10.1186/s12302-021-00575-y
42. Jiang S, Kong S, Zheng H, Zeng X, Chen N, Qi S. [Real-time Source Apportionment of PM_{2.5} and Potential Geographic Origins of Each Source During Winter in Wuhan]. *PubMed*. 2022;43(1):61-73. doi:10.13227/j.hjx.202105171
43. Comunian SO, Dongo D, Milani C, Palestini P. Air Pollution and COVID-19: The Role of Particulate Matter in the Spread and Increase of COVID-19's Morbidity and Mortality. *International Journal of Environmental Research and Public Health*. 2020;17(12):4487. doi:10.3390/ijerph17124487
44. Sharma AK, Balyan P. Air pollution and COVID-19: Is the connect worth its weight? *Indian J Public Health*. 2020;64(Supplement):S132-S134. doi:10.4103/ijph.IJPH_466_20
45. Shao L, Wang Y, Jones T, et al. COVID-19 mortality and exposure to airborne PM_{2.5}: A lag time correlation. *Science of the Total Environment*. 2022;806:151286. doi:10.1016/j.scitotenv.2021.151286
46. Tung NH, Cheng PW, Chi KH, et al. Particulate matter and SARS-CoV-2: A possible model of COVID-19 transmission. *Science of the Total Environment*. 2021;750:141532. doi:10.1016/j.scitotenv.2020.141532

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.