


# Oncogene-induced senescence is a crucial antitumor defense mechanism of human endometrial stromal cells

A.L. Toropov<sup>1</sup>, P.I. Deryabin<sup>1</sup>, A.N. Shatrova<sup>2</sup>, A.V. Borodkina<sup>1</sup>, \*


<sup>1</sup>Mechanisms of cellular senescence group, Institute of Cytology of the Russian Academy of Sciences, Tikhoretsky Ave. 4, 194064, Saint-Petersburg, Russia

<sup>2</sup>Laboratory of intracellular membranes dynamic, Institute of Cytology of the Russian Academy of Sciences, Tikhoretsky Ave. 4, 194064, Saint-Petersburg, Russia

\*Correspondence to Aleksandra V. Borodkina, Tikhoretsky ave. 4, 194064, St-Petersburg, Russia; [borodkina618@gmail.com](mailto:borodkina618@gmail.com)



**Supplementary Figure 1.** Full-length blots presented in the article. (A) Blots for Figure 4A, (B) Blots for Figure 4D, (C) Blots for Figure 4E, (D) Blots for Figure 6D, (E) Blots for Figure 6B.



**Supplementary Figure 2.** Densitometry quantitation for Western blot analysis performed in the study. Estimated optical densities of bands were normalized by sum, each protein / protein modification were analyzed in 3 replicates. (A, B, C) Quantitation results for blots presented at Figure 4. (D, E) Quantitation results for blots presented at Figure 6. Data are presented as mean  $\pm$  SD. ns  $p \geq 0.05$ , \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.005$  using Welch's t-test.