
The predicate of the current mathematical knowledge increases the constructive mathematics what is impossible for the empirical sciences

[Apoloniusz Tyszk](#)^{*}

Posted Date: 31 July 2023

doi: 10.20944/preprints202307.2063.v1

Keywords: composite numbers of the form $2^{2^n}+1$, constructive algorithms, current mathematical knowledge, decidable sets $X \subseteq \mathbb{N}$, epistemic notions, informal notions, known algorithms, known elements of \mathbb{N} , primes of the form n^2+1 , primes of the form $n!+1$, primes of the form $2^{2^n}+1$



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

The Predicate of the Current Mathematical Knowledge Substantially Increases the Constructive Mathematics What Is Impossible for the Empirical Sciences

Apoloniusz Tyszk

Hugo Kołłątaj University, Balicka 116B, 30-149 Kraków, Poland; rttyszka@cyf-kr.edu.pl

Abstract: This is an expanded and revised version of the article: A. Tyszk, *Statements and open problems on decidable sets* $\mathcal{X} \subseteq \mathbb{N}$, Pi Mu Epsilon J. 15 (2023), no. 8, 493-504. The main results were presented at the 25th Conference Applications of Logic in Philosophy and the Foundations of Mathematics, see <http://applications-of-logic.uni.wroc.pl/XXV-Konferencja-Zastosowania-Logiki-w-Filozofii-i-Podstawach-Matematyki>. We assume that the current mathematical knowledge is a finite set of statements which is time-dependent. Nicolas D. Goodman observed that epistemic notions increase the understanding of mathematics without changing its content. We explain the distinction between algorithms whose existence is provable in ZFC and constructively defined algorithms which are currently known. By using this distinction, we obtain non-trivial statements on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that belong to constructive mathematics and refer to the current mathematical knowledge on \mathcal{X} . This and the next sentence justify the article title. For any empirical science, we can identify the current knowledge with that science because truths from the empirical sciences are not necessary truths but working models of truth from a particular context. Edmund Landau's conjecture states that the set $\mathcal{P}(n^2 + 1)$ of primes of the form $n^2 + 1$ is infinite. Landau's conjecture implies the following unproven statement Φ : $\text{card}(\mathcal{P}(n^2 + 1)) < \omega \Rightarrow \mathcal{P}(n^2 + 1) \subseteq [2, (((24!)!)!)]$. We heuristically justify the statement Φ . This justification does not yield the finiteness/infiniteness of $\mathcal{P}(n^2 + 1)$. We present a new heuristic argument for the infiniteness of $\mathcal{P}(n^2 + 1)$, which is not based on the statement Φ .

Keywords and phrases: composite numbers of the form $2^{2^n} + 1$; constructive algorithms; current mathematical knowledge; decidable sets $\mathcal{X} \subseteq \mathbb{N}$; epistemic notions; informal notions; known algorithms; known elements of \mathbb{N} ; primes of the form $n^2 + 1$; primes of the form $n! + 1$; primes of the form $2^{2^n} + 1$

2020 Mathematics Subject Classification: 03F55; 11A41

1. Introduction

This is an expanded and revised version of the article [18]. The main results of this article were presented at the 25th Conference Applications of Logic in Philosophy and the Foundations of Mathematics, see <http://applications-of-logic.uni.wroc.pl/XXV-Konferencja-Zastosowania-Logiki-w-Filozofii-i-Podstawach-Matematyki>. We assume that the current mathematical knowledge is a finite set of statements which is time-dependent. Nicolas D. Goodman in [5] observed that epistemic notions increase the understanding of mathematics without changing its content. We present non-trivial statements on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that belong to constructive mathematics and refer to the current mathematical knowledge on \mathcal{X} . This and the next sentence justify the article title. For any empirical science, we can identify the current knowledge with that science because truths from the empirical sciences are not necessary truths but working models of truth from a particular context.

2. Basic definitions

Algorithms always terminate. Semi-algorithms may not terminate. There is the distinction between *existing algorithms* (i.e. algorithms whose existence is provable in ZFC) and *known algorithms* (i.e. algorithms whose definition is constructive and currently known), see [2,12], [14, p. 9]. A definition of an integer n is called *constructive*, if it provides a known algorithm with no input that returns n . Definition 1 applies to sets $\mathcal{X} \subseteq \mathbb{N}$ whose infiniteness is false or unproven.

Definition 1. We say that a non-negative integer k is a known element of \mathcal{X} , if $k \in \mathcal{X}$ and we know an algebraic expression that defines k and consists of the following signs: 1 (one), + (addition), − (subtraction), · (multiplication), ^ (exponentiation with exponent in \mathbb{N}), ! (factorial of a non-negative integer), ((left parenthesis),) (right parenthesis).

The set of known elements of \mathcal{X} is finite and time-dependent, so cannot be defined in the formal language of classical mathematics. Let t denote the largest twin prime that is smaller than (((((((9!)!)!)!)!)!)!)!. The number t is an unknown element of the set of twin primes.

Definition 2. Conditions (1)–(5) concern sets $\mathcal{X} \subseteq \mathbb{N}$.

- (1) A known algorithm with no input returns an integer n satisfying $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$.
- (2) A known algorithm for every $k \in \mathbb{N}$ decides whether or not $k \in \mathcal{X}$.
- (3) No known algorithm with no input returns the logical value of the statement $\text{card}(\mathcal{X}) = \omega$.
- (4) There are many elements of \mathcal{X} and it is conjectured, though so far unproven, that \mathcal{X} is infinite.
- (5) \mathcal{X} is naturally defined. The infiniteness of \mathcal{X} is false or unproven. \mathcal{X} has the simplest definition among known sets $\mathcal{Y} \subseteq \mathbb{N}$ with the same set of known elements.

Condition (3) implies that no known proof shows the finiteness/infiniteness of \mathcal{X} . No known set $\mathcal{X} \subseteq \mathbb{N}$ satisfies Conditions (1)–(4) and is widely known in number theory or naturally defined, where this term has only informal meaning.

Example 1. The set $\mathcal{X} = \mathcal{P}_{n^2+1}$ satisfies Condition (3).

Let $[\cdot]$ denote the integer part function.

Example 2. The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } \left[\frac{((((((((9!)!)!)!)!)!)!)!}{\pi} \right] \text{ is odd} \\ \emptyset, & \text{otherwise} \end{cases}$$

does not satisfy Condition (3) because we know an algorithm with no input that computes $\left[\frac{((((((((9!)!)!)!)!)!)!)!}{\pi} \right]$. The set of known elements of \mathcal{X} is empty. Hence, Condition (5) fails for \mathcal{X} .

Example 3. ([2,12], [14, p. 9]). The function

$$\mathbb{N} \ni n \xrightarrow{h} \begin{cases} 1, & \text{if the decimal expansion of } \pi \text{ contains } n \text{ consecutive zeros} \\ 0, & \text{otherwise} \end{cases}$$

is computable because $h = \mathbb{N} \times \{1\}$ or there exists $k \in \mathbb{N}$ such that

$$h = (\{0, \dots, k\} \times \{1\}) \cup (\{k+1, k+2, k+3, \dots\} \times \{0\})$$

No known algorithm computes the function h .

Example 4. The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if the continuum hypothesis holds} \\ \emptyset, & \text{otherwise} \end{cases}$$

is decidable. This \mathcal{X} satisfies Conditions (1) and (3) and does not satisfy Conditions (2), (4), and (5). These facts will hold forever.

Statement 1. Condition (1) remains unproven for $\mathcal{X} = \mathcal{P}_{n^2+1}$.

Proof. For every set $\mathcal{X} \subseteq \mathbb{N}$, there exists an algorithm $\text{Alg}(\mathcal{X})$ with no input that returns

$$n = \begin{cases} 0, & \text{if } \text{card}(\mathcal{X}) \in \{0, \omega\} \\ \max(\mathcal{X}), & \text{otherwise} \end{cases}$$

This n satisfies the implication in Condition (1), but the algorithm $\text{Alg}(\mathcal{P}_{n^2+1})$ is unknown because its definition is ineffective. \square

3. Main results

Edmund Landau's conjecture states that the set \mathcal{P}_{n^2+1} of primes of the form $n^2 + 1$ is infinite, see [15,16,20].

Statement 2. The statement

$$\exists n \in \mathbb{N} (\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, n+3])$$

remains unproven in ZFC and classical logic without the law of excluded middle.

Let $f(1) = 10^6$, and let $f(n+1) = f(n)^{f(n)}$ for every positive integer n .

Statement 3. The set

$$\mathcal{X} = \{k \in \mathbb{N} : (10^6 < k) \Rightarrow (f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$$

satisfies Conditions (1)-(4). Condition (5) fails for \mathcal{X} .

Proof. Condition (4) holds as $\mathcal{X} \supseteq \{0, \dots, 10^6\}$ and the set \mathcal{P}_{n^2+1} is conjecturally infinite. Due to known physics we are not able to confirm by a direct computation that some element of \mathcal{P}_{n^2+1} is greater than $f(10^6)$, see [9]. Thus Condition (3) holds. Condition (2) holds trivially. Since the set

$$\{k \in \mathbb{N} : (10^6 < k) \wedge (f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$$

is empty or infinite, Condition (1) holds with $n = 10^6$. Condition (5) fails as the set of known elements of \mathcal{X} equals $\{0, \dots, 10^6\}$. \square

Statements 4 and 6 provide stronger examples.

Conjecture 1. ([1, p. 443], [6]). There are infinitely many primes of the form $k! + 1$.

For a non-negative integer n , let $\rho(n)$ denote $29.5 + \frac{11!}{3n+1} \cdot \sin(n)$.

Statement 4. The set

$$\mathcal{X} = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains more than } \rho(n) \text{ primes of the form } k! + 1\}$$

satisfies Conditions (1)–(5) except the requirement that \mathcal{X} is naturally defined. $501893 \in \mathcal{X}$. Condition (1) holds with $n = 501893$. $\text{card}(\mathcal{X} \cap [0, 501893]) = 159827$. $\mathcal{X} \cap [501894, \infty) = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains at least 30 primes of the form } k! + 1\}$.

Proof. For every integer $n \geq 11!$, 30 is the smallest integer greater than $\rho(n)$. By this, if $n \in \mathcal{X} \cap [11!, \infty)$, then $n + 1, n + 2, n + 3, \dots \in \mathcal{X}$. Hence, Condition (1) holds with $n = 11! - 1$. We explicitly know 24 positive integers k such that $k! + 1$ is prime, see [4]. The inequality $\text{card}(\{k \in \mathbb{N} \setminus \{0\} : k! + 1 \text{ is prime}\}) > 24$ remains unproven. Since $24 < 30$, Condition (3) holds. The interval $[-1, 11! - 1]$ contains exactly three primes of the form $k! + 1$: $1! + 1, 2! + 1, 3! + 1$. For every integer $n > 503000$, the inequality $\rho(n) > 3$ holds. Therefore, the execution of the following MuPAD code

```
m:=0:
for n from 0.0 to 503000.0 do
if n<1!+1 then r:=0 end_if:
if n>=1!+1 and n<2!+1 then r:=1 end_if:
if n>=2!+1 and n<3!+1 then r:=2 end_if:
if n>=3!+1 then r:=3 end_if:
if r>29.5+(11!/(3*n+1))*sin(n) then
m:=m+1:
print([n,m]):
end_if:
end_for:
```

displays the all known elements of \mathcal{X} . The output ends with the line $[501893.0, 159827]$, which proves Condition (1) with $n = 501893$ and Condition (4) with $\text{card}(\mathcal{X}) \geq 159827$. \square

Definition 3. Conditions (1a)–(5a) concern sets $\mathcal{X} \subseteq \mathbb{N}$.

(1a) A known algorithm with no input returns an integer n satisfying $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$.

(2a) A known algorithm for every $k \in \mathbb{N}$ decides whether or not $k \in \mathcal{X}$.

(3a) No known algorithm with no input returns the logical value of the statement $\text{card}(\mathcal{X}) < \omega$.

(4a) There are many elements of \mathcal{X} and it is conjectured, though so far unproven, that \mathcal{X} is finite.

(5a) \mathcal{X} is naturally defined. The finiteness of \mathcal{X} is false or unproven. \mathcal{X} has the simplest definition among known sets $\mathcal{Y} \subseteq \mathbb{N}$ with the same set of known elements.

Statement 5. The set

$$\mathcal{X} = \left\{ n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains more than } 6.5 + \frac{10^6}{3n+1} \cdot \sin(n) \text{ squares of the form } k! + 1 \right\}$$

satisfies Conditions (1a)–(5a) except the requirement that \mathcal{X} is naturally defined. $95151 \in \mathcal{X}$. Condition (1a) holds with $n = 95151$. $\text{card}(\mathcal{X} \cap [0, 95151]) = 30311$. $\mathcal{X} \cap [95152, \infty) = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains at least 7 squares of the form } k! + 1\}$.

Proof. For every integer $n > 10^6$, 7 is the smallest integer greater than $6.5 + \frac{10^6}{3n+1} \cdot \sin(n)$. By this, if $n \in \mathcal{X} \cap (10^6, \infty)$, then $n + 1, n + 2, n + 3, \dots \in \mathcal{X}$. Hence, Condition (1a) holds with $n = 10^6$. It is conjectured that $k! + 1$ is a square only for $k \in \{4, 5, 7\}$, see [19, p. 297]. Hence, the inequality $\text{card}(\{k \in \mathbb{N} \setminus \{0\} : k! + 1 \text{ is a square}\}) > 3$ remains unproven. Since $3 < 7$, Condition (3a) holds. The interval $[-1, 10^6]$ contains exactly three squares of the form $k! + 1$: $4! + 1, 5! + 1, 7! + 1$. Therefore, the execution of the following MuPAD code

```

m:=0:
for n from 0.0 to 1000000.0 do
if n<25 then r:=0 end_if:
if n>=25 and n<121 then r:=1 end_if:
if n>=121 and n<5041 then r:=2 end_if:
if n>=5041 then r:=3 end_if:
if r>6.5+(1000000/(3*n+1))*sin(n) then
m:=m+1:
print([n,m]):
end_if:
end_for:

```

displays the all known elements of \mathcal{X} . The output ends with the line [95151.0, 30311], which proves Condition (1a) with $n = 95151$ and Condition (4a) with $\text{card}(\mathcal{X}) \geq 30311$. \square

To formulate Statement 6 and its proof, we need some lemmas. For a non-negative integer n , let $\theta(n)$ denote the largest integer divisor of $10^{10^{10}}$ smaller than n . For a non-negative integer n , let $\theta_1(n)$ denote the largest integer divisor of 10^{10} smaller than n .

Lemma 1. For every integer $j > 10^{10^{10}}$, $\theta(j) = 10^{10^{10}}$. For every integer $j > 10^{10}$, $\theta_1(j) = 10^{10}$.

Lemma 2. For every integer $j \in (6553600, 7812500]$, $\theta(j) = 6553600$.

Proof. 6553600 equals $2^{18} \cdot 5^2$ and divides $10^{10^{10}}$. $7812500 < 2^{24}$. $7812500 < 5^{10}$. We need to prove that every integer $j \in (6553600, 7812500)$ does not divide $10^{10^{10}}$. It holds as the set

$$\{2^u \cdot 5^v : (u \in \{0, \dots, 23\}) \wedge (v \in \{0, \dots, 9\})\}$$

contains 6553600 and 7812500 as consecutive elements. \square

Lemma 3. The number $6553600^2 + 1$ is prime.

Proof. The following PARI/GP ([10]) command

```
isprime(6553600^2+1,{flag=2})
```

returns 1. This command performs the APRCL primality test, the best deterministic primality test algorithm ([21, p. 226]). It rigorously shows that the number $6553600^2 + 1$ is prime. \square

In the next lemmas, the execution of the command `isprime(n , {flag=2})` proves the primality of n . Let κ denote the function

$$\mathbb{N} \ni n \xrightarrow{\kappa} \underbrace{\text{the_exponent_of_2_in_the_prime_factorization_of_}n+1}_{\in \mathbb{N}} \in \mathbb{N}$$

Lemma 4. The set $\mathcal{X}_1 = \{n \in \mathbb{N} : (\theta_1(n) + \kappa(n))^2 + 1 \text{ is prime}\}$ is infinite.

Proof. Let $i = 142101504$. By the inequality $2^i \geq 2 + 10^{10}$ and Lemma 1, for every non-negative integer m , the number

$$\left(\theta_1\left(2^i \cdot (2m+1) - 1\right) + \kappa\left(2^i \cdot (2m+1) - 1\right)\right)^2 + 1 = \left(10^{10} + i\right)^2 + 1$$

is prime. \square

Before Open Problem 1, \mathcal{X} denotes the set $\{n \in \mathbb{N} : (\theta(n) + \kappa(n))^2 + 1 \text{ is prime}\}$.

Lemma 5. For every $n \in \mathcal{X} \cap (10^{10^{10}}, \infty)$ and for every non-negative integer j , $3^j \cdot (n+1) - 1 \in \mathcal{X} \cap (10^{10^{10}}, \infty)$.

Proof. By the inequality $3^j \cdot (n+1) - 1 \geq n$ and Lemma 1,

$$\theta(3^j \cdot (n+1) - 1) + \kappa(3^j \cdot (n+1) - 1) = 10^{10^{10}} + \kappa(n) = \theta(n) + \kappa(n)$$

□

Lemma 6. $\text{card}(\mathcal{X}) \geq 629450$.

Proof. By Lemmas 2 and 3, for every even integer $j \in (6553600, 7812500]$, the number $(\theta(j) + \kappa(j))^2 + 1 = (6553600 + 0)^2 + 1$ is prime. Hence,

$$\{2k : k \in \mathbb{N}\} \cap (6553600, 7812500] \subseteq \mathcal{X}$$

Consequently,

$$\text{card}(\mathcal{X}) \geq \text{card}(\{2k : k \in \mathbb{N}\} \cap (6553600, 7812500]) = \frac{7812500 - 6553600}{2} = 629450$$

□

Lemma 7. $10242 \in \mathcal{X}$ and $10242 \notin \mathcal{X}_1$.

Proof. The number $10240 = 2^{11} \cdot 5$ divides $10^{10^{10}}$. Hence, $\theta(10242) = 10240$. The number $(\theta(10242) + \kappa(10242))^2 + 1 = (10240 + 0)^2 + 1$ is prime. The set

$$\{2^u \cdot 5^v : (u \in \{0, \dots, 10\}) \wedge (v \in \{0, \dots, 10\})\}$$

contains 10000 and 12500 as consecutive elements. Hence, $\theta_1(10242) = 10000$. The number $(\theta_1(10242) + \kappa(10242))^2 + 1 = (10000 + 0)^2 + 1 = 17 \cdot 5882353$ is composite. □

Statement 6. The set \mathcal{X} satisfies Conditions (1)–(5) except the requirement that \mathcal{X} is naturally defined.

Proof. Condition (2) holds trivially. Let δ denote $10^{10^{10}}$. By Lemma 5, Condition (1) holds for $n = \delta$. Lemma 5 and the unproven statement $\mathcal{P}_{n^2+1} \cap [\delta^2 + 1, \infty) \neq \emptyset$ show Condition (3). The same argument and Lemma 6 yield Condition (4). By Lemma 4, the set \mathcal{X}_1 is infinite. Since Definition 1 applies to sets $\mathcal{X} \subseteq \mathbb{N}$ whose infiniteness is false or unproven, Condition (5) holds except the requirement that \mathcal{X} is naturally defined. □

The set \mathcal{X} satisfies Condition (5) except the requirement that \mathcal{X} is naturally defined. It is true because \mathcal{X}_1 is infinite by Lemma 4 and Definition 1 applies only to sets $\mathcal{X} \subseteq \mathbb{N}$ whose infiniteness is false or unproven. Ignoring this restriction, \mathcal{X} still satisfies the same identical condition due to Lemma 7.

Proposition 1. No set $\mathcal{X} \subseteq \mathbb{N}$ will satisfy Conditions (1)–(4) forever, if for every algorithm with no input, at some future day, a computer will be able to execute this algorithm in 1 second or less.

Proof. The proof goes by contradiction. We fix an integer n that satisfies Condition (1). Since Conditions (1)–(3) will hold forever, the semi-algorithm in Figure 1 never terminates and sequentially prints the following sentences:

$$n+1 \notin \mathcal{X}, n+2 \notin \mathcal{X}, n+3 \notin \mathcal{X}, \dots \quad (\text{T})$$




Figure 1. Semi-algorithm that terminates if and only if \mathcal{X} is infinite.

The sentences from the sequence (T) and our assumption imply that for every integer $m > n$ computed by a known algorithm, at some future day, a computer will be able to confirm in 1 second or less that $(n, m] \cap \mathcal{X} = \emptyset$. Thus, at some future day, numerical evidence will support the conjecture that the set \mathcal{X} is finite, contrary to the conjecture in Condition (4). \square

The physical limits of computation ([9]) disprove the assumption of Proposition 1.

Open Problem 1. Is there a set $\mathcal{X} \subseteq \mathbb{N}$ which satisfies Conditions (1)–(5)?

Open Problem 1 asks about the existence of a year $t \geq 2023$ in which the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})$$

will hold for some $\mathcal{X} \subseteq \mathbb{N}$. For every year $t \geq 2023$ and for every $i \in \{1, 2, 3\}$, a positive solution to Open Problem i in the year t may change in the future. Currently, the answers to Open Problems 1–5 are negative.

4. Satisfiable conjunctions which consist of Conditions (1)–(5) and their negations

The set $\mathcal{X} = \mathcal{P}_{n^2+1}$ satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})$$

The set $\mathcal{X} = \{0, \dots, 10^6\} \cup \mathcal{P}_{n^2+1}$ satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

The numbers $2^{2^k} + 1$ are prime for $k \in \{0, 1, 2, 3, 4\}$. It is open whether or not there are infinitely many primes of the form $2^{2^k} + 1$, see [8, p. 158] and [13, p. 74]. It is open whether or not there are infinitely many composite numbers of the form $2^{2^k} + 1$, see [8, p. 159] and [13, p. 74]. Most mathematicians believe that $2^{2^k} + 1$ is composite for every integer $k \geq 5$, see [7, p. 23].

The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\}, & \text{otherwise} \end{cases}$$

satisfies the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

Open Problem 2. Is there a set $\mathcal{X} \subseteq \mathbb{N}$ that satisfies the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})?$$

The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\} \cup \\ \{n \in \mathbb{N} : n \text{ is the sixth prime number of the form } 2^{2^k} + 1\}, & \text{otherwise} \end{cases}$$

satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

Open Problem 3. Is there a set $\mathcal{X} \subseteq \mathbb{N}$ that satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})?$$

It is possible, although very doubtful, that at some future day, the set $\mathcal{X} = \mathcal{P}_{n^2+1}$ will solve Open Problem 2. The same is true for Open Problem 3. It is possible, although very doubtful, that at some future day, the set $\mathcal{X} = \{k \in \mathbb{N} : 2^{2^k} + 1 \text{ is composite}\}$ will solve Open Problem 1. The same is true for Open Problems 2 and 3.

Table 1 shows satisfiable conjunctions of the form

$$\#(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \#(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \#(\text{Condition 5})$$

where $\#$ denotes the negation \neg or the absence of any symbol. Table 1 differs from Table 1 in [18] for three sets \mathcal{X} . These sets \mathcal{X} have the index *new*.

Table 1. Five satisfiable conjunctions.

	$(\text{Cond. 2}) \wedge (\text{Cond. 3}) \wedge (\text{Cond. 4})$	$(\text{Cond. 2}) \wedge \neg(\text{Cond. 3}) \wedge (\text{Cond. 4})$
$(\text{Cond. 1}) \wedge (\text{Cond. 5})$	Open Problem 1	Open Problem 2
$(\text{Cond. 1}) \wedge \neg(\text{Cond. 5})$	$\mathcal{X}_{\text{new}} = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains more than } 29.5 + \frac{11!}{3n+1} \cdot \sin(n) \text{ primes of the form } k! + 1\}$	$\mathcal{X}_{\text{new}} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\}, & \text{otherwise} \end{cases}$
$\neg(\text{Cond. 1}) \wedge (\text{Cond. 5})$	$\mathcal{X} = \mathcal{P}_{n^2+1}$	Open Problem 3
$\neg(\text{Cond. 1}) \wedge \neg(\text{Cond. 5})$	$\mathcal{X} = \{0, \dots, 10^6\} \cup \mathcal{P}_{n^2+1}$	$\mathcal{X}_{\text{new}} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\} \cup \{n \in \mathbb{N} : n \text{ is the sixth prime number of the form } 2^{2^k} + 1\}, & \text{otherwise} \end{cases}$

Definition 4. We say that an integer n is a threshold number of a set $\mathcal{X} \subseteq \mathbb{N}$, if $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$.

If a set $\mathcal{X} \subseteq \mathbb{N}$ is empty or infinite, then any integer n is a threshold number of \mathcal{X} . If a set $\mathcal{X} \subseteq \mathbb{N}$ is non-empty and finite, then the all threshold numbers of \mathcal{X} form the set $[\max(\mathcal{X}), \infty) \cap \mathbb{N}$.

Open Problem 4. Is there a known threshold number of \mathcal{P}_{n^2+1} ?

Open Problem 4 asks about the existence of a year $t \geq 2023$ in which the implication $\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq (-\infty, n]$ will hold for some known integer n .

Let \mathcal{T} denote the set of twin primes.

Open Problem 5. Is there a known threshold number of \mathcal{T} ?

Open Problem 5 asks about the existence of a year $t \geq 2023$ in which the implication $\text{card}(\mathcal{T}) < \omega \Rightarrow \mathcal{T} \subseteq (-\infty, n]$ will hold for some known integer n .

5. Number-theoretic statements Ψ_n

Let $f(1) = 2$, $f(2) = 4$, and let $f(n+1) = f(n)!$ for every integer $n \geq 2$. Let \mathcal{U}_1 denote the system of equations $\{x_1! = x_1$. For an integer $n \geq 2$, let \mathcal{U}_n denote the following system of equations:

$$\begin{cases} x_1! &= x_1 \\ x_1 \cdot x_1 &= x_2 \\ \forall i \in \{2, \dots, n-1\} \ x_i! &= x_{i+1} \end{cases}$$

Lemma 8. For every positive integer n , the system \mathcal{U}_n has exactly two solutions in positive integers x_1, \dots, x_n , namely $(1, \dots, 1)$ and $(f(1), \dots, f(n))$.

Let B_n denote the following system of equations:

$$\{x_j! = x_k : j, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$$

For every positive integer n , no known system $\mathcal{S} \subseteq B_n$ with a finite number of solutions in positive integers x_1, \dots, x_n has a solution $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$ satisfying $\max(x_1, \dots, x_n) > f(n)$. For every positive integer n and for every known system $\mathcal{S} \subseteq B_n$, if the finiteness/infiniteness of the set

$$\{(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n : (x_1, \dots, x_n) \text{ solves } \mathcal{S}\}$$

is unknown, then the statement

$$\exists x_1, \dots, x_n \in \mathbb{N} \setminus \{0\} ((x_1, \dots, x_n) \text{ solves } \mathcal{S}) \wedge (\max(x_1, \dots, x_n) > f(n))$$

remains unproven.

For a positive integer n , let Ψ_n denote the following statement: *if a system $\mathcal{S} \subseteq B_n$ has at most finitely many solutions in positive integers x_1, \dots, x_n , then each such solution (x_1, \dots, x_n) satisfies $x_1, \dots, x_n \leq f(n)$* . The statement Ψ_n says that for subsystems of B_n with a finite number of solutions, the largest known solution is indeed the largest possible. The statement $\forall n \in \mathbb{N} \setminus \{0\} \Psi_n$ is dubious, see [17, p. 70].

Theorem 1. *For every statement Ψ_n , the bound $f(n)$ cannot be decreased.*

Proof. It follows from Lemma 8 because $\mathcal{U}_n \subseteq B_n$. \square

Theorem 2. *For every integer $n \geq 2$, the statement Ψ_{n+1} implies the statement Ψ_n .*

Proof. If a system $\mathcal{S} \subseteq B_n$ has at most finitely many solutions in positive integers x_1, \dots, x_n , then for every integer $i \in \{1, \dots, n\}$ the system $\mathcal{S} \cup \{x_i! = x_{n+1}\}$ has at most finitely many solutions in positive integers x_1, \dots, x_{n+1} . The statement Ψ_{n+1} implies that $x_i! = x_{n+1} \leq f(n+1) = f(n)!$. Hence, $x_i \leq f(n)$. \square

Theorem 3. *Every statement Ψ_n is true with an unknown integer bound that depends on n .*

Proof. For every positive integer n , the system B_n has a finite number of subsystems. \square

6. A special case of the statement Ψ_9 applies to the conjecture that $\text{card}(\mathcal{P}_{n^2+1}) = \omega$

Let \mathcal{A} denote the following system of equations:

$$\left\{ \begin{array}{lcl} x_2! & = & x_3 \\ x_3! & = & x_4 \\ x_5! & = & x_6 \\ x_8! & = & x_9 \\ x_1 \cdot x_1 & = & x_2 \\ x_3 \cdot x_5 & = & x_6 \\ x_4 \cdot x_8 & = & x_9 \\ x_5 \cdot x_7 & = & x_8 \end{array} \right.$$

Lemma 9. *For every positive integers x and y , $x! \cdot y = y!$ if and only if*

$$(x+1=y) \vee (x=y=1)$$

Lemma 9 and the diagram in Figure 2 explain the construction of the system \mathcal{A} .




Figure 2. Construction of the system \mathcal{A} .

Lemma 10. (Wilson's theorem, [3, p. 89]). For every integer $x \geq 2$, x is prime if and only if x divides $(x-1)! + 1$.

Lemma 11. For every integer $x_1 \geq 2$, the system \mathcal{A} is solvable in positive integers x_2, \dots, x_9 if and only if $x_1^2 + 1$ is prime. In this case, the integers x_2, \dots, x_9 are uniquely determined by the following equalities:

$$\begin{aligned} x_2 &= x_1^2 \\ x_3 &= (x_1^2)! \\ x_4 &= ((x_1^2)!)! \\ x_5 &= x_1^2 + 1 \\ x_6 &= (x_1^2 + 1)! \\ x_7 &= \frac{(x_1^2)! + 1}{x_1^2 + 1} \\ x_8 &= (x_1^2)! + 1 \\ x_9 &= ((x_1^2)! + 1)! \end{aligned}$$

Proof. By Lemma 9, for every integer $x_1 \geq 2$, the system \mathcal{A} is solvable in positive integers x_2, \dots, x_9 if and only if $x_1^2 + 1$ divides $(x_1^2)! + 1$. Hence, the claim of Lemma 11 follows from Lemma 10. \square

Lemma 12. There are only finitely many tuples $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$, which solve the system \mathcal{A} and satisfy $x_1 = 1$. It is true as every such tuple (x_1, \dots, x_9) satisfies $x_1, \dots, x_9 \in \{1, 2\}$.

Proof. The equality $x_1 = 1$ implies that $x_2 = x_1 \cdot x_1 = 1$. Hence, $x_3 = x_2! = 1$. Therefore, $x_4 = x_3! = 1$. The equalities $x_5! = x_6$ and $x_5 = 1 \cdot x_5 = x_3 \cdot x_5 = x_6$ imply that $x_5, x_6 \in \{1, 2\}$. The equalities $x_8! = x_9$ and $x_8 = 1 \cdot x_8 = x_4 \cdot x_8 = x_9$ imply that $x_8, x_9 \in \{1, 2\}$. The equality $x_5 \cdot x_7 = x_8$ implies that $x_7 = \frac{x_8}{x_5} \in \left\{\frac{1}{1}, \frac{1}{2}, \frac{2}{1}, \frac{2}{2}\right\} \cap (\mathbb{N} \setminus \{0\}) = \{1, 2\}$. \square

Conjecture 2. The statement Ψ_9 is true when is restricted to the system \mathcal{A} .

Theorem 4. Conjecture 2 proves the following implication: if there exists an integer $x_1 \geq 2$ such that $x_1^2 + 1$ is prime and greater than $f(7)$, then the set \mathcal{P}_{n^2+1} is infinite.

Proof. Suppose that the antecedent holds. By Lemma 11, there exists a unique tuple $(x_2, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^8$ such that the tuple (x_1, x_2, \dots, x_9) solves the system \mathcal{A} . Since $x_1^2 + 1 > f(7)$, we obtain that $x_1^2 \geq f(7)$. Hence, $(x_1^2)! \geq f(7)! = f(8)$. Consequently,

$$x_9 = ((x_1^2)! + 1)! \geq (f(8) + 1)! > f(8)! = f(9)$$

Conjecture 2 and the inequality $x_9 > f(9)$ imply that the system \mathcal{A} has infinitely many solutions $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$. According to Lemmas 11 and 12, the set \mathcal{P}_{n^2+1} is infinite. \square

Landau's conjecture implies the following unproven statement Φ :

$$\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, (((24!)!)!)!]$$

Theorem 5 heuristically justifies the statement Φ . This justification does not yield the finiteness/infiniteness of \mathcal{P}_{n^2+1} .

Theorem 5. Conjecture 2 implies the statement Φ .

Proof. It follows from Theorem 4 and the equality $f(7) = (((24!)!)!)!$. \square

Theorem 6. The statement Φ implies Conjecture 2.

Proof. By Lemmas 11 and 12, if positive integers x_1, \dots, x_9 solve the system \mathcal{A} , then

$$(x_1 \geq 2) \wedge (x_5 = x_1^2 + 1) \wedge (x_5 \text{ is prime})$$

or $x_1, \dots, x_9 \in \{1, 2\}$. In the first case, Lemma 11 and the statement Φ imply that the inequality $x_5 \leq (((24!)!)!)! = f(7)$ holds when the system \mathcal{A} has at most finitely many solutions in positive integers x_1, \dots, x_9 . Hence, $x_2 = x_5 - 1 < f(7)$ and $x_3 = x_2! < f(7)! = f(8)$. Continuing this reasoning in the same manner, we can show that every x_i does not exceed $f(9)$. \square

Lemma 13. $\log_2(\log_2(\log_2(\log_2(\log_2(\log_2(\log_2(((24!)!)!)!))))) \approx 1.42298$.

Proof. We ask Wolfram Alpha at <http://wolframalpha.com>. \square

Statement 7. Conditions (2)–(5) hold for $\mathcal{X} = \mathcal{P}_{n^2+1}$. The statement Φ implies Condition (1) for $\mathcal{X} = \mathcal{P}_{n^2+1}$ and does not falsify Conditions (2)–(5).

Proof. Conditions (2), (3), and (5) hold trivially. The set \mathcal{P}_{n^2+1} is conjecturally infinite. There are 2199894223892 primes of the form $n^2 + 1$ in the interval $[2, 10^{28})$, see [16]. These two facts imply Condition (4). The statement Φ implies that Condition (1) holds for $\mathcal{X} = \mathcal{P}_{n^2+1}$ with $n = (((24!)!)!)!$. By Lemma 13, due to known physics we are not able to confirm by a direct computation that some element of \mathcal{P}_{n^2+1} is greater than $f(7) = (((24!)!)!)!$, see [9]. Hence, the statement Φ does not falsify Conditions (2)–(5). \square

Proving Landau's conjecture will disprove Statement 7. We do not conjecture that

$$(\text{Conditions (1)–(5) hold for } \mathcal{X} = \mathcal{P}_{n^2+1}) \wedge \Phi$$

7. A new heuristic argument for the infiniteness of \mathcal{P}_{n^2+1}

The system \mathcal{A} contains four factorials and four multiplications. Let \mathcal{F} denote the family of all systems $\mathcal{S} \subseteq B_9$ which contain at most four factorials and at most four multiplications.

Among known systems $\mathcal{S} \in \mathcal{F}$, the following system \mathcal{C}

$$\left\{ \begin{array}{lcl} x_1! & = & x_2 \\ x_2 \cdot x_9 & = & x_1 \\ x_2 \cdot x_2 & = & x_3 \\ x_3 \cdot x_3 & = & x_4 \\ x_4 \cdot x_4 & = & x_5 \\ x_5! & = & x_6 \\ x_6! & = & x_7 \\ x_7! & = & x_8 \end{array} \right.$$

attains the greatest solution in positive integers x_1, \dots, x_9 and has at most finitely many solutions in $(\mathbb{N} \setminus \{0\})^9$. Only the tuples $(1, \dots, 1)$ and $(2, 2, 4, 16, 256, 256!, (256!)!, ((256!)!)!, 1)$ solve \mathcal{C} and belong to $(\mathbb{N} \setminus \{0\})^9$.

For every known system $\mathcal{S} \in \mathcal{F}$, if the finiteness of the set

$$\{(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9 : (x_1, \dots, x_9) \text{ solves } \mathcal{S}\}$$

is unproven and conjectured, then the statement

$$\exists x_1, \dots, x_9 \in \mathbb{N} \setminus \{0\} ((x_1, \dots, x_9) \text{ solves } \mathcal{S}) \wedge (\max(x_1, \dots, x_9) > ((256!)!)!)$$

remains unproven.

Let Γ denote the statement: *if the system \mathcal{A} has at most finitely many solutions in positive integers x_1, \dots, x_9 , then each such solution (x_1, \dots, x_9) satisfies $x_1, \dots, x_9 \leq ((256!)!)!$* . The number $46^{512} + 1$ is prime ([11]) and greater than $256!$, see also [13, p. 239] for the primality of $150^{2048} + 1$. Hence, the statement Γ is equivalent to the infiniteness of \mathcal{P}_{n^2+1} . It heuristically justifies the infiniteness of \mathcal{P}_{n^2+1} in a sophisticated way.

References

1. C. K. Caldwell and Y. Gallot, *On the primality of $n! \pm 1$ and $2 \times 3 \times 5 \times \dots \times p \pm 1$* , Math. Comp. 71 (2002), no. 237, 441–448, <http://doi.org/10.1090/S0025-5718-01-01315-1>.
2. J. Case and M. Ralston, *Beyond Rogers' non-constructively computable function*, in: *The nature of computation*, Lecture Notes in Comput. Sci., 7921, 45–54, Springer, Heidelberg, 2013, http://link.springer.com/chapter/10.1007/978-3-642-39053-1_6.
3. M. Erickson, A. Vazzana, D. Garth, *Introduction to number theory*, 2nd ed., CRC Press, Boca Raton, FL, 2016.
4. *Factorial prime*, http://en.wikipedia.org/wiki/Factorial_prime.
5. N. D. Goodman, *The knowing mathematician*, Synthese 60 (1984), no. 1, 21–38, <http://link.springer.com/article/10.1007/BF00485616>; reprinted in: H. Leblanc, E. Mendelson, A. Orenstein, *Foundations: Logic, Language, and Mathematics*, Springer, 1984, 21–38, <http://link.springer.com/book/10.1007/978-94-017-1592-8>.
6. *Is $n! + 1$ often a prime?*, <http://math.stackexchange.com/questions/853085/is-n-1-often-a-prime>.
7. J.-M. De Koninck and F. Luca, *Analytic number theory: Exploring the anatomy of integers*, American Mathematical Society, Providence, RI, 2012.
8. M. Křížek, F. Luca, L. Somer, *17 lectures on Fermat numbers: from number theory to geometry*, Springer, New York, 2001.

9. S. Lloyd, *Ultimate physical limits to computation*, Nature 406 (2000), 1047–1054, <http://doi.org/10.1038/35023282>.
10. PARI/GP online documentation, http://pari.math.u-bordeaux.fr/dochtml/html/Arithmetic_functions.html.
11. X. M. Pi, *Searching for generalized Fermat primes (Chinese)*, J. Math. (Wuhan) 18 (1998), no. 3, 276–280.
12. R. Reitzig, *How can it be decidable whether π has some sequence of digits?*, <http://cs.stackexchange.com/questions/367/how-can-it-be-decidable-whether-pi-has-some-sequence-of-digits>.
13. P. Ribenboim, *The little book of bigger primes*, 2nd ed., Springer, New York, 2004.
14. H. Rogers, Jr., *Theory of recursive functions and effective computability*, 2nd ed., MIT Press, Cambridge, MA, 1987.
15. N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, A002496, *Primes of the form $n^2 + 1$* , <http://oeis.org/A002496>.
16. N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, A083844, *Number of primes of the form $x^2 + 1 < 10^n$* , <http://oeis.org/A083844>.
17. A. Tyszka, *A common approach to three open problems in number theory*, Discrete Math. Lett. 12 (2023), 66–72.
18. A. Tyszka, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$* , Pi Mu Epsilon J. 15 (2023), no. 8, 493–504.
19. E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics*, 2nd ed., Chapman & Hall/CRC, Boca Raton, FL, 2002.
20. Wolfram MathWorld, *Landau's Problems*, <http://mathworld.wolfram.com/LandausProblems.html>.
21. S. Y. Yan, *Number theory for computing*, 2nd ed., Springer, Berlin, 2002.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.