

The predicate of the current mathematical knowledge substantially increases the constructive and informal mathematics and why it cannot be adapted to any empirical science

[Apoloniusz Tyszk](#)^{*}

Posted Date: 15 August 2023

doi: 10.20944/preprints202307.2063.v3

Keywords: composite numbers of the form $22n + 1$; constructive algorithms; current mathematical knowledge; decidable sets $X \subseteq \mathbb{N}$; epistemic notions, informal notions, known algorithms, known elements of $\mathbb{N}$; primes of the form $n^2 + 1$; primes of the form $n! + 1$; primes of the form $22n + 1$



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

The Predicate of the Current Mathematical Knowledge Substantially Increases the Constructive and Informal Mathematics and Why It Cannot Be Adapted to Any Empirical Science

Apoloniusz Tyszk

Hugo Kołłątaj University, Balicka 116B, 30-149 Kraków, Poland; rttyszka@cyf-kr.edu.pl

Abstract: We assume that the current mathematical knowledge is a finite set of statements which is time-dependent. In every branch of mathematics, the set of all knowable truths is the set of all theorems. This set exists independently of our current scientific knowledge. Nicolas D. Goodman observed in *Synthese* that epistemic notions increase the understanding of mathematics without changing its content as a formal science. This content remains also unchanged when the current mathematical knowledge leads to new conjectures. We explain the distinction between algorithms whose existence is provable in *ZFC* and constructively defined algorithms which are currently known. By using this distinction, we obtain non-trivial statements on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that belong to constructive and informal mathematics and refer to the current mathematical knowledge on $\mathcal{X}$. This and the next sentence justify the article title. For any empirical science, we can identify the current knowledge with that science because truths from the empirical sciences are not necessary truths but working models of truth from a particular context.

Keywords: constructive algorithms; current mathematical knowledge; epistemic notions; informal notions; known algorithms; known elements of $\mathbb{N}$

MSC: 03A05; 03F55

1. Why such title of the article

This is a shortened and revised version of [15]. This section is new compared to [15]. The main mathematical results of this article were presented at the 25th Conference Applications of Logic in Philosophy and the Foundations of Mathematics, see <http://applications-of-logic.uni.wroc.pl/XXV-Konferencja-Zastosowania-Logiki-w-Filozofii-i-Podstawach-Matematyki>. We assume that the current mathematical knowledge is a finite set of statements which is time-dependent. In every branch of mathematics, the set of all knowable truths is the set of all theorems. This set exists independently of our current scientific knowledge. Nicolas D. Goodman in [4] observed that epistemic notions increase the understanding of mathematics without changing its content as a formal science. This content remains also unchanged when the current mathematical knowledge leads to new conjectures. We explain the distinction between algorithms whose existence is provable in *ZFC* and constructively defined algorithms which are currently known. By using this distinction, we obtain non-trivial statements on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that belong to constructive and informal mathematics and refer to the current mathematical knowledge on $\mathcal{X}$. This and the next sentence justify the article title. For any empirical science, we can identify the current knowledge with that science because truths from the empirical sciences are not necessary truths but working models of truth from a particular context, see [16] (p. 610).

2. Basic definitions and examples

Algorithms always terminate. Semi-algorithms may not terminate. There is the distinction between *existing algorithms* (i.e. algorithms whose existence is provable in ZFC) and *known algorithms* (i.e. algorithms whose definition is constructive and currently known), see [2,10] and [12] (p. 9). A definition of an integer n is called *constructive*, if it provides a known algorithm with no input that returns n . Definition 1 applies to sets $\mathcal{X} \subseteq \mathbb{N}$ whose infiniteness is false or unproven.

Definition 1. We say that a non-negative integer k is a known element of $\mathcal{X}$, if $k \in \mathcal{X}$ and we know an algebraic expression that defines k and consists of the following signs: 1 (one), + (addition), − (subtraction), · (multiplication), ^ (exponentiation with exponent in $\mathbb{N}$), ! (factorial of a non-negative integer), ((left parenthesis),) (right parenthesis).

The set of known elements of $\mathcal{X}$ is finite and time-dependent, so cannot be defined in the formal language of classical mathematics. Let t denote the largest twin prime that is smaller than (((((((9!)!)!)!)!)!)!)!. The number t is an unknown element of the set of twin primes.

Definition 2. Conditions (1)–(5) concern sets $\mathcal{X} \subseteq \mathbb{N}$.

- (1) A known algorithm with no input returns an integer n satisfying $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$.
- (2) A known algorithm for every $k \in \mathbb{N}$ decides whether or not $k \in \mathcal{X}$.
- (3) No known algorithm with no input returns the logical value of the statement $\text{card}(\mathcal{X}) = \omega$.
- (4) There are many elements of $\mathcal{X}$ and it is conjectured, though so far unproven, that $\mathcal{X}$ is infinite.
- (5) $\mathcal{X}$ is naturally defined. The infiniteness of $\mathcal{X}$ is false or unproven. $\mathcal{X}$ has the simplest definition among known sets $\mathcal{Y} \subseteq \mathbb{N}$ with the same set of known elements.

Condition (3) implies that no known proof shows the finiteness/infiniteness of $\mathcal{X}$. No known set $\mathcal{X} \subseteq \mathbb{N}$ satisfies Conditions (1)–(4) and is widely known in number theory or naturally defined, where this term has only informal meaning.

Example 1. The set $\mathcal{X} = \mathcal{P}_{n^2+1}$ satisfies Condition (3).

Let $[\cdot]$ denote the integer part function.

Example 2. The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } \left[\frac{(((((9!)!)!)!)!)!}{\pi} \right] \text{ is odd} \\ \emptyset, & \text{otherwise} \end{cases}$$

does not satisfy Condition (3) because we know an algorithm with no input that computes $\left[\frac{(((((9!)!)!)!)!)!}{\pi} \right]$. The set of known elements of $\mathcal{X}$ is empty. Hence, Condition (5) fails for $\mathcal{X}$.

Example 3. ([2,10] and [12] p. 9). The function

$$\mathbb{N} \ni n \xrightarrow{h} \begin{cases} 1, & \text{if the decimal expansion of } \pi \text{ contains } n \text{ consecutive zeros} \\ 0, & \text{otherwise} \end{cases}$$

is computable because $h = \mathbb{N} \times \{1\}$ or there exists $k \in \mathbb{N}$ such that

$$h = (\{0, \dots, k\} \times \{1\}) \cup (\{k+1, k+2, k+3, \dots\} \times \{0\})$$

No known algorithm computes the function h .

Example 4. The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if the continuum hypothesis holds} \\ \emptyset, & \text{otherwise} \end{cases}$$

is decidable. This $\mathcal{X}$ satisfies Conditions (1) and (3) and does not satisfy Conditions (2), (4), and (5). These facts will hold forever.

3. Main results

Edmund Landau's conjecture states that the set $\mathcal{P}_{n^2+1}$ of primes of the form $n^2 + 1$ is infinite, see [13,14,18].

Statement 1. Condition (1) remains unproven for $\mathcal{X} = \mathcal{P}_{n^2+1}$.

Proof. For every set $\mathcal{X} \subseteq \mathbb{N}$, there exists an algorithm $\text{Alg}(\mathcal{X})$ with no input that returns

$$n = \begin{cases} 0, & \text{if } \text{card}(\mathcal{X}) \in \{0, \omega\} \\ \max(\mathcal{X}), & \text{otherwise} \end{cases}$$

This n satisfies the implication in Condition (1), but the algorithm $\text{Alg}(\mathcal{P}_{n^2+1})$ is unknown because its definition is ineffective. $\square$

Statement 2. The statement

$$\exists n \in \mathbb{N} (\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, n+3])$$

remains unproven in ZFC and classical logic without the law of excluded middle.

Let $f(1) = 10^6$, and let $f(n+1) = f(n)^{f(n)}$ for every positive integer n .

Statement 3. The set

$$\mathcal{X} = \{k \in \mathbb{N} : (10^6 < k) \Rightarrow (f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$$

satisfies Conditions (1)-(4). Condition (5) fails for $\mathcal{X}$.

Proof. Condition (4) holds as $\mathcal{X} \supseteq \{0, \dots, 10^6\}$ and the set $\mathcal{P}_{n^2+1}$ is conjecturally infinite. Due to known physics we are not able to confirm by a direct computation that some element of $\mathcal{P}_{n^2+1}$ is greater than $f(10^6)$, see [8]. Thus Condition (3) holds. Condition (2) holds trivially. Since the set

$$\{k \in \mathbb{N} : (10^6 < k) \wedge (f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$$

is empty or infinite, Condition (1) holds with $n = 10^6$. Condition (5) fails as the set of known elements of $\mathcal{X}$ equals $\{0, \dots, 10^6\}$. $\square$

Statements 4 and 7 provide stronger examples.

Conjecture 1. ([1] (p. 443) and [5]). There are infinitely many primes of the form $k! + 1$.

For a non-negative integer n , let $\rho(n)$ denote $29.5 + \frac{11!}{3n+1} \cdot \sin(n)$.

Statement 4. The set

$$\mathcal{X} = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains more than } \rho(n) \text{ primes of the form } k! + 1\}$$

satisfies Conditions (1)–(5) except the requirement that $\mathcal{X}$ is naturally defined. $501893 \in \mathcal{X}$. Condition (1) holds with $n = 501893$. $\text{card}(\mathcal{X} \cap [0, 501893]) = 159827$. $\mathcal{X} \cap [501894, \infty) = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains at least 30 primes of the form } k! + 1\}$.

Proof. For every integer $n \geq 11!$, 30 is the smallest integer greater than $\rho(n)$. By this, if $n \in \mathcal{X} \cap [11!, \infty)$, then $n + 1, n + 2, n + 3, \dots \in \mathcal{X}$. Hence, Condition (1) holds with $n = 11! - 1$. We explicitly know 24 positive integers k such that $k! + 1$ is prime, see [3]. The inequality $\text{card}(\{k \in \mathbb{N} \setminus \{0\} : k! + 1 \text{ is prime}\}) > 24$ remains unproven. Since $24 < 30$, Condition (3) holds. The interval $[-1, 11! - 1]$ contains exactly three primes of the form $k! + 1$: $1! + 1, 2! + 1, 3! + 1$. For every integer $n > 503000$, the inequality $\rho(n) > 3$ holds. Therefore, the execution of the following MuPAD code

```
m:=0:
for n from 0.0 to 503000.0 do
if n<1!+1 then r:=0 end_if:
if n>=1!+1 and n<2!+1 then r:=1 end_if:
if n>=2!+1 and n<3!+1 then r:=2 end_if:
if n>=3!+1 then r:=3 end_if:
if r>29.5+(11!/(3*n+1))*sin(n) then
m:=m+1:
print([n,m]):
end_if:
end_for:
```

displays the all known elements of $\mathcal{X}$. The output ends with the line $[501893.0, 159827]$, which proves Condition (1) with $n = 501893$ and Condition (4) with $\text{card}(\mathcal{X}) \geq 159827$. $\square$

Definition 3. Conditions (1a)–(5a) concern sets $\mathcal{X} \subseteq \mathbb{N}$.

(1a) A known algorithm with no input returns an integer n satisfying $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$.

(2a) A known algorithm for every $k \in \mathbb{N}$ decides whether or not $k \in \mathcal{X}$.

(3a) No known algorithm with no input returns the logical value of the statement $\text{card}(\mathcal{X}) < \omega$.

(4a) There are many elements of $\mathcal{X}$ and it is conjectured, though so far unproven, that $\mathcal{X}$ is finite.

(5a) $\mathcal{X}$ is naturally defined. The finiteness of $\mathcal{X}$ is false or unproven. $\mathcal{X}$ has the simplest definition among known sets $\mathcal{Y} \subseteq \mathbb{N}$ with the same set of known elements.

Statement 5. The set

$$\mathcal{X} = \left\{ n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains more than } 6.5 + \frac{10^6}{3n+1} \cdot \sin(n) \text{ squares of the form } k! + 1 \right\}$$

satisfies Conditions (1a)–(5a) except the requirement that $\mathcal{X}$ is naturally defined. $95151 \in \mathcal{X}$. Condition (1a) holds with $n = 95151$. $\text{card}(\mathcal{X} \cap [0, 95151]) = 30311$. $\mathcal{X} \cap [95152, \infty) = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains at least 7 squares of the form } k! + 1\}$.

Proof. For every integer $n > 10^6$, 7 is the smallest integer greater than $6.5 + \frac{10^6}{3n+1} \cdot \sin(n)$. By this, if $n \in \mathcal{X} \cap (10^6, \infty)$, then $n + 1, n + 2, n + 3, \dots \in \mathcal{X}$. Hence, Condition (1a) holds with $n = 10^6$. It is conjectured that $k! + 1$ is a square only for $k \in \{4, 5, 7\}$, see [17] (p. 297). Hence, the inequality $\text{card}(\{k \in \mathbb{N} \setminus \{0\} : k! + 1 \text{ is a square}\}) > 3$ remains unproven. Since $3 < 7$, Condition (3a) holds. The interval $[-1, 10^6]$ contains exactly three squares of the form $k! + 1$: $4! + 1, 5! + 1, 7! + 1$. Therefore, the execution of the following MuPAD code

```

m:=0:
for n from 0.0 to 1000000.0 do
if n<25 then r:=0 end_if:
if n>=25 and n<121 then r:=1 end_if:
if n>=121 and n<5041 then r:=2 end_if:
if n>=5041 then r:=3 end_if:
if r>6.5+(1000000/(3*n+1))*sin(n) then
m:=m+1:
print([n,m]):
end_if:
end_for:

```

displays the all known elements of $\mathcal{X}$. The output ends with the line [95151.0, 30311], which proves Condition (1a) with $n = 95151$ and Condition (4a) with $\text{card}(\mathcal{X}) \geq 30311$. $\square$

Statement 6. *The set*

$$\mathcal{X} = \{k \in \mathbb{N} : \text{card}([-1, k] \cap \mathcal{P}_{n^2+1}) < 10^{10000}\}$$

satisfies the conjunction

$$\neg(\text{Condition 1a}) \wedge (\text{Condition 2a}) \wedge (\text{Condition 3a}) \wedge (\text{Condition 4a}) \wedge (\text{Condition 5a})$$

To formulate Statement 7 and its proof, we need some lemmas. For a non-negative integer n , let $\theta(n)$ denote the largest integer divisor of $10^{10^{10}}$ smaller than n . For a non-negative integer n , let $\theta_1(n)$ denote the largest integer divisor of 10^{10} smaller than n .

Lemma 1. *For every integer $j > 10^{10^{10}}$, $\theta(j) = 10^{10^{10}}$. For every integer $j > 10^{10}$, $\theta_1(j) = 10^{10}$.*

Lemma 2. *For every integer $j \in (6553600, 7812500]$, $\theta(j) = 6553600$.*

Proof. 6553600 equals $2^{18} \cdot 5^2$ and divides $10^{10^{10}}$. $7812500 < 2^{24}$. $7812500 < 5^{10}$. We need to prove that every integer $j \in (6553600, 7812500)$ does not divide $10^{10^{10}}$. It holds as the set

$$\{2^u \cdot 5^v : (u \in \{0, \dots, 23\}) \wedge (v \in \{0, \dots, 9\})\}$$

contains 6553600 and 7812500 as consecutive elements. $\square$

Lemma 3. *The number $6553600^2 + 1$ is prime.*

Proof. The following PARI/GP ([9]) command

```
isprime(6553600^2+1,{flag=2})
```

returns 1. This command performs the APRCL primality test, the best deterministic primality test algorithm ([19], p. 226). It rigorously shows that the number $6553600^2 + 1$ is prime. $\square$

In the next lemmas, the execution of the command `isprime( $n$ , {flag=2})` proves the primality of n . Let κ denote the function

$$\mathbb{N} \ni n \xrightarrow{\kappa} \underbrace{\text{the_exponent_of_2_in_the_prime_factorization_of_}n+1}_{\in \mathbb{N}} \in \mathbb{N}$$

Lemma 4. The set $\mathcal{X}_1 = \{n \in \mathbb{N} : (\theta_1(n) + \kappa(n))^2 + 1 \text{ is prime}\}$ is infinite.

Proof. Let $i = 142101504$. By the inequality $2^i \geq 2 + 10^{10}$ and Lemma 1, for every non-negative integer m , the number

$$\left(\theta_1\left(2^i \cdot (2m+1) - 1\right) + \kappa\left(2^i \cdot (2m+1) - 1\right)\right)^2 + 1 = \left(10^{10} + i\right)^2 + 1$$

is prime. $\square$

Before Open Problem 1, $\mathcal{X}$ denotes the set $\{n \in \mathbb{N} : (\theta(n) + \kappa(n))^2 + 1 \text{ is prime}\}$.

Lemma 5. For every $n \in \mathcal{X} \cap \left(10^{10^{10}}, \infty\right)$ and for every non-negative integer j , $3^j \cdot (n+1) - 1 \in \mathcal{X} \cap \left(10^{10^{10}}, \infty\right)$.

Proof. By the inequality $3^j \cdot (n+1) - 1 \geq n$ and Lemma 1,

$$\theta\left(3^j \cdot (n+1) - 1\right) + \kappa\left(3^j \cdot (n+1) - 1\right) = 10^{10^{10}} + \kappa(n) = \theta(n) + \kappa(n)$$

$\square$

Lemma 6. $\text{card}(\mathcal{X}) \geq 629450$.

Proof. By Lemmas 2 and 3, for every even integer $j \in (6553600, 7812500]$, the number $(\theta(j) + \kappa(j))^2 + 1 = (6553600 + 0)^2 + 1$ is prime. Hence,

$$\{2k : k \in \mathbb{N}\} \cap (6553600, 7812500] \subseteq \mathcal{X}$$

Consequently,

$$\text{card}(\mathcal{X}) \geq \text{card}(\{2k : k \in \mathbb{N}\} \cap (6553600, 7812500]) = \frac{7812500 - 6553600}{2} = 629450$$

$\square$

Lemma 7. $10242 \in \mathcal{X}$ and $10242 \notin \mathcal{X}_1$.

Proof. The number $10240 = 2^{11} \cdot 5$ divides $10^{10^{10}}$. Hence, $\theta(10242) = 10240$. The number $(\theta(10242) + \kappa(10242))^2 + 1 = (10240 + 0)^2 + 1$ is prime. The set

$$\{2^u \cdot 5^v : (u \in \{0, \dots, 10\}) \wedge (v \in \{0, \dots, 10\})\}$$

contains 10000 and 12500 as consecutive elements. Hence, $\theta_1(10242) = 10000$. The number $(\theta_1(10242) + \kappa(10242))^2 + 1 = (10000 + 0)^2 + 1 = 17 \cdot 5882353$ is composite. $\square$

Statement 7. The set $\mathcal{X}$ satisfies Conditions (1)–(5) except the requirement that $\mathcal{X}$ is naturally defined.

Proof. Condition (2) holds trivially. Let δ denote $10^{10^{10}}$. By Lemma 5, Condition (1) holds for $n = \delta$. Lemma 5 and the unproven statement $\mathcal{P}_{n^2+1} \cap [\delta^2 + 1, \infty) \neq \emptyset$ show Condition (3). The same argument and Lemma 6 yield Condition (4). By Lemma 4, the set $\mathcal{X}_1$ is infinite. Since Definition 1 applies to sets $\mathcal{X} \subseteq \mathbb{N}$ whose infiniteness is false or unproven, Condition (5) holds except the requirement that $\mathcal{X}$ is naturally defined. $\square$

The set $\mathcal{X}$ satisfies Condition (5) except the requirement that $\mathcal{X}$ is naturally defined. It is true because $\mathcal{X}_1$ is infinite by Lemma 4 and Definition 1 applies only to sets $\mathcal{X} \subseteq \mathbb{N}$ whose infiniteness is false or unproven. Ignoring this restriction, $\mathcal{X}$ still satisfies the same identical condition due to Lemma 7.

Proposition 1. No set $\mathcal{X} \subseteq \mathbb{N}$ will satisfy Conditions (1)–(4) forever, if for every algorithm with no input, at some future day, a computer will be able to execute this algorithm in 1 second or less.

Proof. The proof goes by contradiction. We fix an integer n that satisfies Condition (1). Since Conditions (1)–(3) will hold forever, the semi-algorithm in Figure 1 never terminates and sequentially prints the following sentences:

$$n+1 \notin \mathcal{X}, n+2 \notin \mathcal{X}, n+3 \notin \mathcal{X}, \dots \quad (\text{T})$$




Figure 1. Semi-algorithm that terminates if and only if $\mathcal{X}$ is infinite.

The sentences from the sequence (T) and our assumption imply that for every integer $m > n$ computed by a known algorithm, at some future day, a computer will be able to confirm in 1 second or less that $(n, m] \cap \mathcal{X} = \emptyset$. Thus, at some future day, numerical evidence will support the conjecture that the set $\mathcal{X}$ is finite, contrary to the conjecture in Condition (4). $\square$

The physical limits of computation ([8]) disprove the assumption of Proposition 1.

Open Problem 1. Is there a set $\mathcal{X} \subseteq \mathbb{N}$ which satisfies Conditions (1)–(5)?

Open Problem 1 asks about the existence of a year $t \geq 2023$ in which the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})$$

will hold for some $\mathcal{X} \subseteq \mathbb{N}$. For every year $t \geq 2023$ and for every $i \in \{1, 2, 3\}$, a positive solution to Open Problem i in the year t may change in the future. Currently, the answers to Open Problems 1–5 are negative.

4. Satisfiable conjunctions which consist of Conditions (1)–(5) and their negations

The set $\mathcal{X} = \mathcal{P}_{n^2+1}$ satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})$$

The set $\mathcal{X} = \{0, \dots, 10^6\} \cup \mathcal{P}_{n^2+1}$ satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

The numbers $2^{2^k} + 1$ are prime for $k \in \{0, 1, 2, 3, 4\}$. It is open whether or not there are infinitely many primes of the form $2^{2^k} + 1$, see [7, p. 158] and [11, p. 74]. It is open whether or not there are infinitely many composite numbers of the form $2^{2^k} + 1$, see [7, p. 159] and [11, p. 74]. Most mathematicians believe that $2^{2^k} + 1$ is composite for every integer $k \geq 5$, see [6, p. 23].

The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\}, & \text{otherwise} \end{cases}$$

satisfies the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

Open Problem 2. Is there a set $\mathcal{X} \subseteq \mathbb{N}$ that satisfies the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})?$$

The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\} \cup \\ \{n \in \mathbb{N} : n \text{ is the sixth prime number of the form } 2^{2^k} + 1\}, & \text{otherwise} \end{cases}$$

satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

Open Problem 3. Is there a set $\mathcal{X} \subseteq \mathbb{N}$ that satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})?$$

It is possible, although very doubtful, that at some future day, the set $\mathcal{X} = \mathcal{P}_{n^2+1}$ will solve Open Problem 2. The same is true for Open Problem 3. It is possible, although very doubtful, that at some future day, the set $\mathcal{X} = \{k \in \mathbb{N} : 2^{2^k} + 1 \text{ is composite}\}$ will solve Open Problem 1. The same is true for Open Problems 2 and 3.

Table 1 shows satisfiable conjunctions of the form

$$\#(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \#(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \#(\text{Condition 5})$$

where # denotes the negation $\neg$ or the absence of any symbol.

Table 1. Five satisfiable conjunctions.

	$(\text{Cond. 2}) \wedge (\text{Cond. 3}) \wedge (\text{Cond. 4})$	$(\text{Cond. 2}) \wedge \neg(\text{Cond. 3}) \wedge (\text{Cond. 4})$
$(\text{Cond. 1}) \wedge (\text{Cond. 5})$	Open Problem 1	Open Problem 2
$(\text{Cond. 1}) \wedge \neg(\text{Cond. 5})$	$\mathcal{X} = \{n \in \mathbb{N} : \text{the interval } [-1, n] \text{ contains more than } 29.5 + \frac{11!}{3n+1} \cdot \sin(n) \text{ primes of the form } k! + 1\}$	$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\}, & \text{otherwise} \end{cases}$
$\neg(\text{Cond. 1}) \wedge (\text{Cond. 5})$	$\mathcal{X} = \mathcal{P}_{n^2+1}$	Open Problem 3
$\neg(\text{Cond. 1}) \wedge \neg(\text{Cond. 5})$	$\mathcal{X} = \{0, \dots, 10^6\} \cup \mathcal{P}_{n^2+1}$	$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } 2^{2^{f(9^9)}} + 1 \text{ is composite} \\ \{0, \dots, 10^6\} \cup \{n \in \mathbb{N} : n \text{ is the sixth prime number of the form } 2^{2^k} + 1\}, & \text{otherwise} \end{cases}$

Definition 4. We say that an integer n is a threshold number of a set $\mathcal{X} \subseteq \mathbb{N}$, if $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$.

If a set $\mathcal{X} \subseteq \mathbb{N}$ is empty or infinite, then any integer n is a threshold number of $\mathcal{X}$. If a set $\mathcal{X} \subseteq \mathbb{N}$ is non-empty and finite, then the all threshold numbers of $\mathcal{X}$ form the set $[\max(\mathcal{X}), \infty) \cap \mathbb{N}$.

Open Problem 4. Is there a known threshold number of $\mathcal{P}_{n^2+1}$?

Open Problem 4 asks about the existence of a year $t \geq 2023$ in which the implication $\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq (-\infty, n]$ will hold for some known integer n .

Let $\mathcal{T}$ denote the set of twin primes.

Open Problem 5. Is there a known threshold number of $\mathcal{T}$?

Open Problem 5 asks about the existence of a year $t \geq 2023$ in which the implication $\text{card}(\mathcal{T}) < \omega \Rightarrow \mathcal{T} \subseteq (-\infty, n]$ will hold for some known integer n .

References

1. C. K. Caldwell and Y. Gallot, *On the primality of $n! \pm 1$ and $2 \times 3 \times 5 \times \dots \times p \pm 1$* , Math. Comp. 71 (2002), no. 237, 441–448, <http://doi.org/10.1090/S0025-5718-01-01315-1>.
2. J. Case and M. Ralston, *Beyond Rogers’ non-constructively computable function*, in: *The nature of computation*, Lecture Notes in Comput. Sci., 7921, 45–54, Springer, Heidelberg, 2013, http://link.springer.com/chapter/10.1007/978-3-642-39053-1_6.
3. *Factorial prime*, http://en.wikipedia.org/wiki/Factorial_prime.
4. N. D. Goodman, *The knowing mathematician*, Synthese 60 (1984), no. 1, 21–38, <http://link.springer.com/article/10.1007/BF00485616>; reprinted in: H. Leblanc, E. Mendelson, A. Orenstein, *Foundations: Logic, Language, and Mathematics*, Springer, 1984, 21–38, <http://link.springer.com/book/10.1007/978-94-017-1592-8>.

5. *Is $n! + 1$ often a prime?*, <http://math.stackexchange.com/questions/853085/is-n-1-often-a-prime>.
6. J.-M. De Koninck and F. Luca, *Analytic number theory: Exploring the anatomy of integers*, American Mathematical Society, Providence, RI, 2012.
7. M. Křížek, F. Luca, L. Somer, *17 lectures on Fermat numbers: from number theory to geometry*, Springer, New York, 2001.
8. S. Lloyd, *Ultimate physical limits to computation*, *Nature* 406 (2000), 1047–1054, <http://doi.org/10.1038/35023282>.
9. PARI/GP online documentation, http://pari.math.u-bordeaux.fr/dochtml/html/Arithmetic_functions.html.
10. R. Reitzig, *How can it be decidable whether π has some sequence of digits?*, <http://cs.stackexchange.com/questions/367/how-can-it-be-decidable-whether-pi-has-some-sequence-of-digits>.
11. P. Ribenboim, *The little book of bigger primes*, 2nd ed., Springer, New York, 2004.
12. H. Rogers, Jr., *Theory of recursive functions and effective computability*, 2nd ed., MIT Press, Cambridge, MA, 1987.
13. N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, A002496, *Primes of the form $n^2 + 1$* , <http://oeis.org/A002496>.
14. N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, A083844, *Number of primes of the form $x^2 + 1 < 10^n$* , <http://oeis.org/A083844>.
15. A. Tyszk, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$* , *Pi Mu Epsilon J.* 15 (2023), no. 8, 493–504.
16. B. L. J. Webb, *Science, Truth, and Meaning: From Wonder to Understanding*, World Scientific, Singapore, 2021.
17. E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics*, 2nd ed., Chapman & Hall/CRC, Boca Raton, FL, 2002.
18. Wolfram MathWorld, *Landau's Problems*, <http://mathworld.wolfram.com/LandausProblems.html>.
19. S. Y. Yan, *Number theory for computing*, 2nd ed., Springer, Berlin, 2002.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.