

SUPPLEMENTARY MATERIAL

“Variability in bioactivities among wild populations of *Baccharis crispa* Spreng. (Asteraceae) from Córdoba, Argentina: screening of cytotoxic and antiviral activity”

Giuliana Lingua^{1,2}, Ana Guadalupe Chaves³, Juan Javier Aguilar¹, Florencia Martinez^{1,2}, Lorena Torres³, Enrique Lucini⁴, Marta Contigiani¹, Susana Nuñez Montoya^{5,6}, Brenda S. Konigheim^{1,2,*}

¹ *Universidad Nacional de Córdoba, Facultad de Ciencias Médicas, Instituto de Virología “Dr. J. M. Vanella”- Argentina. Enfermera Gordillo S/N, Ciudad Universitaria, X5000HUA Córdoba, Argentina.*

² *Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.*

³ *Universidad Nacional de Córdoba, Facultad de Ciencias Agropecuaria, Cátedra de Genética- Argentina. Av. Valparaíso S/N Ciudad Universitaria, X5000HUA Córdoba, Argentina.*

⁴ *Universidad Nacional de Córdoba, Facultad de Ciencias Agropecuaria, Cátedra de Microbiología Agrícola- Argentina. Av. Valparaíso S/N Ciudad Universitaria, X5000HUA Córdoba, Argentina.*

⁵ *Universidad Nacional de Córdoba, Facultad de Ciencias Químicas, Dpto. de Ciencias Farmacéuticas, Farmacognosia- Edificio de Ciencias 2, Ciudad Universitaria. Córdoba, Argentina.*


⁶ *CONICET, Instituto Multidisciplinario de Biología Vegetal.*

Legends of Supplemental File (Figure)




Figure S1: Curves of cellular viability percentage vs. concentration of the ethanolic (EtOH), chloroform (CHCl₃), and aqueous (Aqu) extracts for the four populations of *B. crista* on Vero cells, evaluated by NR uptake assay. Each result is expressed as $\bar{x} \pm SE$ of three independent experiments performed in triplicate. A-Villa del Parque Population (VP); B-Tala Cañada Population (TC); C-San Gerónimo Population (SG); D-Puesto Pedernera Population (PP).

Figure S2: Curves of percentage of inhibition of each viral model vs. the concentration of the ethanolic (EtOH), chloroform (CHCl₃), and aqueous (Aqu) extracts of the four populations of *B. crista*. Each result is expressed as $\bar{x} \pm SE$ of three independent experiments performed in triplicate. A-Villa del Parque Population (VP); B-Tala Cañada Population (TC); C-San Gerónimo Population (SG); D-Puesto Pedernera Population (PP).

Figure S1



C- San Gerónimo Population



D- Puesto Pedernera Population












Figure S2


A-Villa del Parque Population



B-Tala Cañada Population



C-San Gerónimo Population



D-Puesto Pedernera Population

