

Article

Not peer-reviewed version

Cross-Sectional Anatomy and Computed Tomography of the Coelomic Cavity in the Atlantic Puffin (Aves, Alcidae, Fratercula arctica).

José Raduan Jaber ^{*} , Marcos Fumero-Hernández ^{*} , Juan Alberto Corbera , Inmaculada Morales , Manuel Amador , Gregorio Ramírez Zarzosa , Mario Encinoso

Posted Date: 13 August 2023

doi: 10.20944/preprints202308.0964.v1

Keywords: Coelomic cavity; computed tomography; anatomical sections; seabirds; puffin.



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Cross-Sectional Anatomy and Computed Tomography of the Coelomic Cavity in the Atlantic Puffin (Aves, *Alcidae, Fratercula arctica*)

José Raduan Jaber ^{1,*}, Marcos Fumero-Hernández ^{2,*}, Juan Alberto Corbera ², Inmaculada Morales ², Manuel Amador ³, Gregorio Ramírez Zarzosa ⁴ and Mario Encinosa ²

¹ Department of Morphology Faculty of Veterinary Medicine, University of Las Palmas de Gran Canaria, Trasmontaña, Arucas, 35413 Las Palmas, Spain; joseraduan.jaber@ulpgc.es (J.R.J.).

² Veterinary Clinical Hospital, Faculty of Veterinary Medicine, University of Las Palmas de Gran Canaria, Trasmontaña, Arucas, 35413 Las Palmas, Spain; marcos.fumero101@alu.ulpgc.es (M.F.H.); juan.corbera@ulpgc.es (J.A.C.S.); inmaculada.morales@ulpgc.es (I.M.); mcinoso@gmail.com (M.E.).

³ Dirección Insular de Medio Ambiente, Consejería de Área de Medio Ambiente, Clima, Energía y Conocimiento of the Cabildo de Gran Canaria, Gran Canaria, Spain; mcamador@grancanaria.com (M.A.).

⁴ Department of Animal Medicine and Surgery, Veterinary Faculty, University of Murcia, Campus de Espinardo, Murcia, 30100; grzar@um.es (G.R.).

* Correspondence: joseraduan.jaber@ulpgc.es (J.R.J.); marcos.fumero101@alu.ulpgc.es (M.F.H.)

Simple Summary: Birds constitute an important group within the new companion animals. Since their popularity has been increasing, it is more common to find them in homes, zoos and wildlife centres around the world, evidencing the need for clinicians, biologists and researchers to have a deep knowledge of their anatomy, pathology and physiology to provide adequate medical care.

Abstract: In birds, unlike mammals, there is no complete separation between the thoracic and abdominal cavities. Instead, they have the coelomic cavity where most main organs are found. Therefore, an adequate knowledge of the anatomy of the coelomic cavity is of great importance for veterinarians, biologists and the scientific community. This study aimed to evaluate the coelomic cavity anatomy in the Atlantic puffins (*Fratercula Arctica*) using anatomical sections and computed tomography images.

Keywords: coelomic cavity; computed tomography; anatomical sections; seabirds; puffin

1. Introduction

Birds constitute an enormously diverse taxonomic group, including thousands of species worldwide [1]. These animals have historically caused the fascination of humans, who have adopted them as companion animals. The possession of birds as pets is a growing trend, and their presence in zoos, wildlife and veterinary centres is also increasingly common [2]. For all these reasons, knowledge of the anatomy and pathophysiology of these animals has become one of the objectives of current veterinary medicine.

Imaging diagnosis is an essential tool for bird clinicians since many psittacine species exhibit a special ability to mask disease signs [3]. Conventional radiology is the method most widely used as an auxiliary diagnostic technique in avian medicine due to its low cost, non-invasive nature and the fact that it is found in most veterinary centres [4-6]. However, modern imaging techniques such as magnetic resonance (MR) and computed tomography (CT) make it possible to obtain three-dimensional images with little distortion and even simulations of dynamic processes of different organs and vascular structures in a minimally invasive way [7-9]. Consequently, its use in anatomical studies has spread, and even smaller versions, such as micro-CT, have been created for animal research [10-21]. These anatomical studies are essential not only from the clinical point of view but

also for endangered species conservation, enabling adequate knowledge of their anatomical characteristics and behaviour patterns [21-23].

Diverse literature can be found regarding the use of diagnostic imaging in the study of the coelomic cavity of birds, however, as far as we know, there are no studies that deal with the anatomy of the avian coelom using the Atlantic puffin as a model [24-27]. The Atlantic puffin (*Fratercula arctica*) is a medium-sized marine bird of the Alcidae family with characteristic black and white colouration. It lives mainly in the North Atlantic Ocean, where it faces a multitude of threats, many of them linked to human activity. All this has led it to be classified as a vulnerable species on the list of threatened species of the International Union for Conservation of Nature (IUCN) [28-29].

The aim of this paper was to study the coelomic cavity of the Atlantic puffin using anatomical sections and CT images, which could serve as an anatomical model for pathologic studies of other phylogenetically related birds.

2. Materials and Methods

2.1. Animals

In the present study, a total of 20 Atlantic puffin (*Fratercula arctica*) carcasses were utilized. The avian specimens displayed an average mass of 0.251 kg (with a range of 0.185-0.310 kg) and an average length, measured from the beak tip to the base of the tail, of 19.85 cm (with a range of 16-24 cm). The study group comprised young animals that were provided by the Consejería de Área de Medio Ambiente, Clima, Energía y Conocimiento of the Cabildo de Gran Canaria following a massive stranding event on the island's coastline. Subsequently, all enrolled puffins for the study were sectioned and scanned at the Veterinary Hospital of Las Palmas de Gran Canaria University (Canary Islands, Spain).

2.2. CT Technique

For the CT examinations, the carcasses were thawed at room temperature for 12 hours. Sequential transverse CT slices were acquired using a 16-slice helical CT scanner (Toshiba Astelion, Canon Medical System®, Tokyo, Japan). The animals were symmetrically positioned in dorsal recumbency on the stretcher, with craniocaudal entry. A standard clinical protocol was followed, employing the following parameters: 120 kVp, 80 mA, 512 × 512 acquisition matrix, 1809 × 858 field of view, pitch of 0.94 and a gantry rotation of 1.5. The obtained images had a slice thickness of 0.6 mm. To enhance the assessment of the coelomic structures on CT, three different CT Windows settings were applied by adjusting the window widths (WW) and window levels (WL): A bone window setting (WW = 1500; WL = 300), a soft tissue window setting (WW = 248; WL = 123), and a lung windows setting (WW = 1400; WL = -500). No significant variations in CT density or anatomy were detected within the coelomic cavity of the birds used in this investigation. Finally, all these images were uploaded to an image viewer (OsiriX MD, Apple, Cupertino, CA, USA) to facilitate data manipulation and analysis.

2.3. Anatomical sections

Following the acquisition of CT images, the imaged cadavers were frozen at -80°C for 72 hours. Subsequently, eight specimens were sectioned in the transverse plane. To accomplish this, parallel sections, each one cm thick, were generated using an electric band saw. Finally, after carefully moistening the sections obtained with water, removing the artefacts (feathers and sand) with Adson forceps and identifying them, they were photographed on both sides.

For further examination and identification of internal structures in anatomical cross-sections and CT images, three puffins were dissected to expose the coelomic cavity and visualize the location of the organs. The dissection facilitated the accurate identification and correlation of anatomical features observed in anatomical sections and CT images.

2.4. Anatomic identification

For the purpose of identifying and labelling the cross section along with the corresponding CT images, we used reference materials, such as textbooks and articles from scientific journals focused on avian anatomy [30-32]. Furthermore, to ensure precise anatomical interpretation of the coelomic structures, we also used different anatomical preparations provided by the Department of Anatomy of the Faculty of Veterinary Medicine, University of Las Palmas de Gran Canaria. These additional resources proved beneficial in enhancing our understanding and accuracy in interpreting the anatomical features within the coelomic cavity.

3. Results

Different figures revealing the anatomical structures of the Atlantic puffin coelomic cavity are presented (Figures 1–12). Figure 1 comprises a compilation of various dissections displaying the main structures within the coelomic cavity. In Figure 2, a sagittal multiplanar reconstruction (MPR) volume rendering is presented, wherein each line and corresponding number (I–IX) represents approximately the level of the following anatomical and CT transverse planes. Figures 3–11 consisted of four individual images for each case: A) an anatomical cross-section, B) a pulmonary CT window, C) a soft tissue CT window, and D) a bone CT window. These images are presented in a rostrocaudal progression, starting from the lungs (Figure 3) to the cloaca levels (Figure 11). Finally, Figure 12 is composed of three images: A) an anatomical dissection, B,C) dorsal MPR volume rendering images in pulmonary CT window at different levels.




Figure 1. Gross dissection image of the coelomic cavity of the *Fratercula arctica* (A), images of the cardiovascular (B), and digestive (C) structures. StM: Sternotracheal Muscle. Mlc: Longissimus colli muscle. Tr: Trachea. Sy: Syrinx. CEAS': Right Cervical Air Sac. CEAS: Left Cervical Air Sac. LBT: Left Brachiocephalic Trunk. RBT: Right Brachiocephalic Trunk. LJV: Left Jugular vein. CCA: Common Carotid artery. LCrVc: Left cranial vena cava. H: Heart. RA: Right Atrium. LA: Left Atrium. Pr: Pericardium. CS: Coronary sinus. LHL: Left Hepatic lobe. RHL: Right Hepatic Lobe. HPL: Hepatopericardial Ligament. DD: Descending Duodenum. AD: Ascending Duodenum. P: Pancreas. C: Caecum. RAS: Right abdominal air sac. Vn: Ventriculus. SI: small intestine. IPC: Intestinal Peritoneal Cavity.




Figure 2. Sagittal MPR volumen rendering image of the body of an Atlantic puffin. The lines and numbers (I–IX) represent the approximate level of the following transverse cross-sections and CT images.




Figure 3. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the trachea corresponding to line I in Figure 1. SC: Spinal cord. V: Vertebra (Vertebral body). ShM: Scapulohumeralis muscle. SCM: Scapulohumeral Caudal muscle. RL: Right lung. LL: Left lung. Mlc: Longissimus colli muscle. Os: Oesophagus. Tr: Trachea. CTAS: Cranial thoracic air sac (left). CTAS': Cranial thoracic air sac (right). CAS: Clavicular Air Sac. LBT: Left Brachiocephalic Trunk. RBT: Right Brachiocephalic Trunk. CrVc: Cranial cava vein. St: Sternum. PM: Pectoral Muscle (thoracobrachialis muscle). ScM: Supracoracoid Muscle. IcM: Intercostal muscle. Hu: Humerus. Rb: Ribs.




Figure 4. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the heart corresponding to line II in Figure 1. SC: Spinal cord. V: Vertebra (Vertebral body). ShM: Scapulohumeralis muscle. SCM: Scapulohumeral Caudal muscle. RL: Right lung. LL: Left lung. Pv: Pulmonary vessels. Os: Oesophagus. Tr: Trachea. LPB: Left Primary Bronchus. RPB: Right Primary Bronchus. CTAS: Cranial thoracic air sac (left). CAS: Clavicular Air Sac. LBT: Left Brachiocephalic Trunk. RBT: Right Brachiocephalic Trunk. H: Heart. RA: Right Atrium. LA: Left Atrium. RV: Right Ventricle. LV: Left Ventricle. St: Sternum. PM: Pectoral Muscle (thoracobrachialis muscle). ScM: Supracoracoid Muscle. IcM: Intercostal muscle. Hu: Humerus. Rb: Ribs.




Figure 5. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the cranial thoracic air sacs corresponding to line III in figure 1. SC: Spinal cord. V: Vertebra (Vertebral body). ShM: Scapulohumeralis muscle. SCM: Scapulohumeral Caudal muscle. RL: Right lung. LL: Left lung. Pv: Pulmonary vessels. Os: Oesophagus. LPB: Left Primary Bronchus. RPB: Right Primary Bronchus. CTAS': Cranial thoracic air sac (right). CTAS: Cranial thoracic air sac (left). H: Heart. St: Sternum. C: Coracoid bone. PM: Pectoral Muscle. ScM: Supracoracoid Muscle. IcM: Intercostal muscle. Hu: Humerus. Rb: Ribs.




Figure 6. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the liver (hepatic lobes) corresponding to line IV in figure 1. SC: Spinal cord. V: Vertebra (Vertebral body). LDM: Longissimus Dorsi Muscle. RL: Right lung. LL: Left lung. RPA: Right pulmonary artery. LPA: Left pulmonary artery. G: Gonad. CTAS: Cranial thoracic air sac (left). CTAS': Cranial thoracic air sac (right). Pr: Proventriculus. RHL: Right hepatic lobe. LHL: Left hepatic lobe. H: Heart (ventricle). SCM: Scapulohumeral caudal muscle. St: Sternum. ScM: Supracoracoid muscle. PM: Pectoral Muscle. IcM: Intercostal muscle. Hu: Humerus. Rb: Ribs.




Figure 7. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the caudal thoracic air sac corresponding to line V in figure 1. SC: Spinal cord. V: Vertebra (Vertebral body and spinal process). LDM: Longissimus Dorsi Muscle. ShM: Scapulohumeral muscle. RL: Right lung. LL: Left lung. CdTAS: Caudal thoracic air sac (right). CdTAS': Caudal thoracic air sac (left). Pr: Proventriculus. RHL: Right hepatic lobe. LHL: Left hepatic lobe. RV: Right ventricle. St: Sternum. ScM: Supracoracoid muscle. PM: Pectoral Muscle. IcM: Intercostal muscle. Hu: Humerus. Rb: Ribs.




Figure 8. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the kidneys corresponding to line VI in figure 1. SC: Spinal cord. V: Vertebra (spinal process). RK: Right kidney. LK: Left kidney. DM: Dorsal Mesenterium. LAS: Left abdominal Sac. RAS: Right abdominal Sac. ASW: Air Sac Wall. I: Intestine. P: Pancreas. Vn: Ventriculus. CCdM: Crassus caudodorsalis muscle. CCM: Crassus cranioventralis muscle. IPC: Intestinal Peritoneal Cavity. LVC: Left Ventral Hepatic Peritoneal Cavity. Lv: Liver. St: Sternum. PM: Pectoral Muscle. IcM: Intercostal muscle. Hu: Humerus. RaU: Radius and Ulna. Rb: Ribs. LDM: Longissimus Dorsi Muscle.




Figure 9. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the ventriculus corresponding to line VII in figure 1. SC: Spinal cord. V: Vertebra (vertebral body and spinal process). RK: Right kidney. LK: Left kidney. RU: Right ureter. LU: Left ureter. Lv: Liver. RAS: Right abdominal air sac. LAS: Left abdominal air sac. I: Intestine. Vn: Ventriculus. In: Ingesta. St: Sternum. PM: Pectoral Muscle (thoracic part). Rb: Ribs.




Figure 10. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the caudal end of the right and left kidneys corresponding to line VIII in figure 1. SC: Spinal cord. V: Vertebra (Vertebral body). Syn: Synsacrum. F: Femur. RK: Right kidney. LK: Left kidney. RU: Right ureter. RAS: Right abdominal air sac. LAS: Left abdominal air sac. P: Pancreas. I: Intestine. SI: Small intestine (duodenum). St: Sternum. PM: Pectoral Muscle (thoracic part). IcM: Intercostal muscle. Rb: Ribs.




Figure 11. Transverse cross-section image (A), pulmonary (B), soft tissue (C), and bone (D) CT images of the coelomic cavity of the *Fratercula arctica* at the level of the cloaca corresponding to line IX in figure 1. V: Vertebra. Clm: Cloacal muscles (Musculus levator cloacae+Musculus transversus cloacae). Cl: Cloaca. IPC: Intestinal Peritoneal Cavity. Pu: Pubis. PM: Pectoral muscle (abdominal portion). Mic: Musculus levator caudae.




Figure 12. Gross dissection image of the coelomic cavity of the *Fratercula arctica* (A), dorsal MPR volume rendering images in pulmonary CT window of the coelomic cavity at level of hepatic lobes (B), and kidneys (C). Mlc: Longissimus colli muscle. CEAS': Right Cervical Air Sac. CEAS: Left Cervical Air Sac. Tr: Trachea. Sy: Syrinx. Os: Oesophagus. LL: Left lung. RL: Right lung. Pv: Pulmonary vessels. H: Heart. LHL: Left Hepatic lobe. RHL: Right Hepatic Lobe. Gb: Gallbladder. DD: Descending Duodenum. AD: Ascending Duodenum. Vn: Ventriculus. P: Pancreas. C: Caecum. RC: Right Caecum. LC: Left Caecum. Hu: Humerus. F: Femur. Syn: Synsacrum. Rb: Rib.

3.1. Anatomical Dissections and Cross-Sections

Anatomical dissections (Figures 1A–C and 12A), as well as transverse (Figures 3A–11A) cross-sections of the coelomic cavity are presented. All these figures were essential in facilitating the identification of the main organs of the respiratory, circulatory, digestive, and urinary systems within this cavity. Consequently, we identified the puffin's heart, which exhibits an oval-shaped with a sharply pointed apex located along the central axis of the coelomic cavity and cranial to the liver (Figure 1A–C). Moreover, the heart chambers, including the right and left atrium and the right and left ventricles, were observed. Additionally, these images were crucial in visualizing significant blood vessels, such as the left and right brachiocephalic trunks, the common carotid artery, the left cranial vena cava, and the left jugular vein (Figure 1A,B). The right and left lungs are visually discernible in a craniodorsal location, ventral to the thoracic vertebra and lateral to the ribs. The anatomical dissections and transverse cross-sections also aided in distinguishing the trachea (Figure 1A,B). This structure coursed in a median position into the coelomic cavity until bifurcating into the right and left main bronchi (Figures 3A–5A). In contrast, we only could identify the syrinx in specific dissected images (Figures 1B and 12A). Dorsal to the trachea, we visualized other formations, such as the oesophagus, the longissimus colli and the sternotracheal muscles (Figures 1A,B and 12A).

The dissected and transverse images played a crucial role in observing the walls of the air sac. As a result, the cross sections provided an opportunity to observe the topographic distribution of the different air sacs, including the clavicular, the cervical, the thoracic and the abdominal air sacs (Figures 1A,B, 3A–10A and 12A). These anatomical preparations also facilitated the observation of the puffin's liver, which consists of the right and the left hepatic lobes, whose cranioventral segments surrounded the heart (Figures 1A,C and 12A). Both lobes exhibited similar sizes and were larger than other organs within the coelomic cavity. At the level of the visceral surface of the right lobe of the liver, a membranous structure corresponding to the gallbladder was distinguishable (Figure 12A). Moreover, the medial border of the hepatic lobes showed the hepatopericardial ligament, which connects the liver with the heart apex (Figure 1C). Caudally and closely related to the visceral surface of the left hepatic lobe, we distinguished the muscular portion of the puffin's stomach, known as the

ventriculus, was identified (Figures 1A,C and 12A). Its identification was facilitated by the presence of a thick muscular layer, as well as by the visualization of two muscles, the *crassus caudodorsalis* and the *crassus cranoventralis*, which delineate the dorsal and ventral borders of the *ventriculus* (Figures 8A and 9A). Besides, these sections also allowed us to observe its glandular part, the *proventriculus*, which shared a similar location at the left side of the coelomic cavity after the junction with the oesophagus and the *ventriculus* (Figures 6A and 7A). Other components of the digestive system, such as the small and large intestine, were also visible in these anatomic sections. The small intestine (*duodenum*) was better identified in the dissected image caudal to the liver (Figures 1A,C and 12A). It showed a U-shaped *ansa duodeni* comprising a descending segment, the *pars descendens*, and an ascending portion, the *pars ascendens* (Figures 1C and 12A). More caudally, we could distinguish the terminal intestine (*caecum*), which was well developed in the puffin, exhibiting a distinctive colouration (Figures 1C and 12A). Between these two segments and close to the duodenal loops, we observed the pancreas that lies within the mesoduodenum, displaying a pale yellow colouration (Figures 1C, 8A, 10A and 12A). Moreover, the utilization of transverse sections was crucial in discerning the presence of paired kidneys situated laterally to the spine and embedded dorsally in excavations of the synsacrum (Figures 8A, 9A and 10A). Interestingly, these transverse sections allowed us to visualize the ventral surface of the right and left kidneys in close contact with the paired abdominal air sacs (Figures 8A and 9A). Furthermore, the caudal pole of the kidney housed the ureters, as depicted in Figures 9A and 10A. These structures were distinguished emerging from the ventral aspect of the kidneys. Concerning the genital organs, we could distinguish a round single yellow structure that closely resembled the ovary (Figure 6A).

In the caudal part of the coelomic cavity, we could identify an excretory passage for the digestive and urogenital systems known as the cloaca, it was surrounded by cloacae muscles (*musculus levator cloacae, and musculus transversus cloacae*), which allow the cloaca to become larger for different functions, including egg laying, copulation and defecation. (Figure 11A). In addition, various bony structures related to the coelomic cavity were observed, including the vertebrae, ribs, sternum, thoracic and pelvic limbs, as well as the lateral parts of the pubis. These bone formations were associated to specific muscles, such as the different parts of the pectoral muscle (*sternobrachialis*, thoracic and abdominal portions), the supracoracoid muscle, the *scapulohumeralis* muscle, the *scapulohumeral caudal* muscle, and the *longissimus dorsi* muscle (Figures 3A–12A).

3.2. Computed Tomography Images

The CT images that closely corresponded to their anatomical sections were selected (Figures 3B–D–11B–D and 12 B,C). These CT images provided additional morphological and tomographic details regarding the coelomic structures when compared with the corresponding cross-sections. The CT images acquired using the pulmonary window setting (Figures 3B–11B and 12B,C) provided excellent visualization of various bones and their associated muscles, which were also seen with the soft tissue and bone window settings. Consequently, we observed different bony structures, such as the thoracic vertebrae, ribs, sternum, humerus, femur and pubis, as well as various muscles associated with these bony structures, including the *scapulohumeralis*, the *scapulohumeral caudal*, the *longissimus colli*, the *longissimus dorsi*, the *pectoral*, the *supracoracoid*, and the *intercostal* muscles. In addition, the essential components of the respiratory tract, such as the trachea, the syrinx, the tracheal bifurcation, and the course of the right and left main bronchi, the pulmonary vessels, as well as the pulmonary parenchyma could also be distinguished due to their higher attenuation (Figures 3B–7B and 12B,C). Interestingly, the lung parenchyma was well-defined, showing a honeycomb-like pattern in all planes, exhibiting an irregular aspect in all puffins (Figures 3B–7B). Moreover, the right and left primary bronchi were visualized entering into the ventromedial aspect of the right and left lungs (Figures 4B, 5B and 12C). Furthermore, other intrathoracic structures, including the heart and large vessels, such as the two brachiocephalic trunks, were well visualized (Figures 4B–5B). In addition, the walls of the clavicular, thoracic and abdominal air sacs could be distinguished (Figures 3B–10B). Concerning the digestive structures, this specific CT window effectively delineated the oesophagus, the right and left hepatic lobe, the glandular and the muscular parts of the puffin stomach, different intestinal segments

and the cloaca (Figures 5B–11B). In addition, the dorsal CT images were pretty helpful in observing the bifurcation of the caecum in the right and left caecum (Figure 12C).

The CT images obtained using the soft tissue windows (Figures 3C–11C) also provided satisfactory identification of the bony and muscle structures that comprise the coelomic cavity. Besides, this CT window allowed the identification of the oesophagus located ventrally to the vertebral body (Figures 3C–5C). Additionally, adequate visualization of the heart was distinguished with this CT window. Therefore, we could observe the right ventricle surrounded by the right and left hepatic lobe (Figures 5C–7C), as well as the main arteries, such as the right and left brachiocephalic trunks, which appeared with intermediate attenuation (Figure 4C). Moreover, the lumen of the proventriculus and the ventricular folds of the puffin stomach were seen with excellent detail in the left portion of the coelomic cavity (Figures 6C–9C).

The CT images acquired with the use of the bone window setting (Figures 3D–11D) demonstrated excellent differentiation between the bones and the soft tissues within the coelomic cavity. In contrast, the heart chambers and the main arteries exhibited an intermediate CT attenuation (Figures 3D–7D). Additionally, various digestive structures, including the oesophagus, the hepatic lobes, different intestinal sections and the cloaca, as well as the glandular and the muscular part of the puffin stomach were also observed with intermediate attenuation (Figures 3D–11D). Interestingly, this CT window enabled the observation of the left abdominal air sac covering the ventricular surface, which adjoins the left ventral hepatic peritoneal cavity on the left side of the ventriculus (Figure 8D). Besides, the crassus cranioventralis was discernible by its anatomical location ventral to the ventricle, as well as by its round shape and intermediate attenuation (Figure 8D).

4. Discussion

Birds possess a complex anatomy, very different from mammals, as they lack a distinct separation between the thorax and the abdomen, resulting in the coelomic cavity. Consequently, this cavity has a complex organization that encloses organs of the circulatory system, as well as the respiratory, digestive and urogenital apparatus [24–25, 30–34].

The conventional approach to studying animal anatomy, which involves dissections for visualizing organs and internal structures, has undergone a profound transformation with the irruption of modern imaging techniques such as MRI and CT, which are the most advanced diagnostic imaging modalities available for identifying normal anatomy and abnormalities in the internal organs of live animals [9]. In contrast to conventional imaging methods, such as radiography or ultrasonography, these modern modalities enable better morphologic characterization of the coelomic cavity by providing views of body sections from multiple tomographic planes, thereby delivering images with exceptional anatomic resolution without tissue superimposition, high contrast between different structures, and excellent tissue differentiation that permits the assessment of spatial relationships between different organs. Disadvantages of these procedures include their high cost and the need to sedate and possibly restrain the bird [4–6, 21, 25–27].

In the present study, coelomic CT images were assessed using pulmonary, soft tissue, and bone window settings. The employment of the pulmonary window setting yielded notable benefits in the differentiation of the respiratory tract and diverse vascular structures. As a result, the lung parenchyma, the course of the two pulmonary arteries supplying each lung, and the paired brachiocephalic trunks were well distinguished with this window setting. Comparable findings have been reported in previous CT investigations performed on other species, such as neonatal foals [35], reptiles [36] and different avian species [25–27]. Meanwhile, the application of the soft tissue window provided additional insights into puffin cardiac topography and its associated blood vessels. However, the combination of the different CT window settings was not useful for the visualization of the four cardiac chambers, which was due to the absence of contrast media administration. Although there is limited literature on the use of contrast media in birds [25–27], its application has proven to enhance and facilitate the visualization of cardiovascular structures and air sacs [25–26]. Despite the constraint posed by the use of cadaveric specimens, precluding the administration of intravenous contrast medium, the pulmonary CT window facilitated visualization of the wall of the

thoracic and abdominal sacs. Similar to previous studies [25], the combination of the aforementioned pulmonary CT window with the bone window setting revealed distinct visualization of the ventriculus of the stomach, whose content exhibited an amalgamation of air and dense materials. Besides, this particular CT window was quite helpful in distinguishing the muscular layer of puffin ventriculus, and the relation between the bird heart and the right and left hepatic lobes. In contrast to other avian species, such as parrots [25], whose right lobe is generally bigger than the left, the puffin liver presented lobes of similar size. In addition, they showed gallbladder, which was only distinguishable in the dissected images. This fact could be due to the lack of contrast administration that also could impede the visualization of other organs, including the spleen, the ureters or the pancreas.

In avian species, the sexes are separated, consisting of either male or female genital organs. Nevertheless, several avian species do not show phenotypic sexual dimorphism and consequently, making it necessary to employ endoscopy or DNA testing for definitive gender determination [32]. In this study, we identified a single round structure adjacent to the caudal margin of the left lung that closely resembled to the left ovary. However, the small size of the puffins suggested that they were juvenile specimens and, therefore, sexually immature animals, which could explain the challenge encountered in distinguishing the gonads in these animals. As stated in other reports described in birds [25], the gonads are clearly visible only in sexually mature individuals of large species.

This CT-based anatomical study has confirmed the suitability of cadavers for evaluating diverse anatomical patterns. However, it is essential to consider the absence of blood flow in dead animals when comparing the results with those obtained from live specimens using contrast media [6, 17, 18, 25]. Results from the present study demonstrated that utilization of frozen anatomical sections and dissections was beneficial in identifying various coelomic structures observed on transverse CT images and ensured accurate correlations.

5. Conclusions

The current investigation elucidates the spatial arrangement of the different organs within the coelomic cavity of puffins, achieved through the integration of diverse CT window settings in conjunction with anatomical cross-sections. The outcomes underscore the efficacy of CT imaging for the comprehensive study of avian anatomical structure, obtaining high-quality images that could serve as a reference to evaluate birds with pathologies at the coelom level. Notably, it is imperative to emphasize that the use of CT as a diagnostic modality in live birds requires the administration of anaesthesia, a procedure that carries potential complications and mortality risks. Consequently, the utilization of this technique in live avians demands judicious consideration and rigorous justification by the attending clinician.

Author Contributions: Conceptualization, J.R.J., M.F. and M.E.; methodology, M.E., J.R.J. and M.F.; investigation, J.R.J., M.F., J.A., G.R., I.M.F. and M.E.; resources, M.A. writing—original draft preparation, J.R.J. and M.F.; writing—review and editing, J.R.J., M.F., M.A. and M. E. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: In this study, we did not need ethical approval since the puffins were coming to our hospital to rule out by CT any cause of stranding. From these studies, we obtained the information presented in our paper. As happens in domestic mammals, just informed consent from the owner allowed us to carry out this study. Therefore, El Cabildo Insular de Gran Canaria was informed that all animal identity information obtained from this study was treated as confidential to the extent permitted by law and only used for research or teaching purposes.

Informed Consent Statement: Written informed consent has been obtained from El Cabildo Insular de Gran Canaria to publish this paper.

Data Availability Statement: The information is available at "<https://accedacris.ulpgc.es>", accessed on 25 June 2023.

Acknowledgments: In loving memory of Alvaro Domingo Rodriguez Garcia. We also thank Concepción Mingot, Carmen Mingot, Emilia Mingot, Nicolas Aquino, Ayesh Mohamad, Marisa Mohamad and Jamal Jaber for their support and constructive comments. Moreover, we also thank to the Consejería de Área de Medio Ambiente, Clima, Energía y Conocimiento of the Cabildo Insular de Gran Canaria to provide the animals of this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Barrowclough, G.F.; Cracraft, J.; Klicka, J.; Zink, R.M. How Many Kinds of Birds Are There and Why Does It Matter?. *PLOS ONE* **2016**, 11 (11).
2. American Veterinary Medical Association. Available online: <https://www.avma.org/resources-tools/reports-statistics/us-pet-ownership-statistics> (accessed on 1 June 2023).
3. Raftery, A. *BSAVA Manual of Psittacine Birds*. 2nd ed.; British small animal veterinary association: Gloucester, U.K, 2005; pp. 35–49.
4. Krautwald-Junghanns, M.E.; Pees, M.; Reese, S.; Tully, T. *Diagnostic imaging of exotic pets*. 1st ed.: Schlütersche: Hannover, Germany, 2011.
5. McMillan, M. C. *Imaging techniques*. Wingers Publishing: Lake Worth, Florida, U.S.A. 1994; pp. 246–326.
6. Zeeland, Y. R. A.; Schoemaker, N. J.; Hsu, E. W. *Current therapy in avian medicine and surgery*; Elsevier: St. Louis, U.S.A. 2016; pp. 531–549.
7. Haacke, E.M.; Brown, R.W.; Thompson, M.R.; Vankatesan, R. *Magnetic Resonance Imaging*. John Wiley & Sons: New York, U.S.A., 1999; pp. 331–380 and 669–702.
8. Beuf, O.; Jaillon, F.; Saint-Jalmes, H. Small-animal MRI: signal-to-noise ratio comparison at 7 and 15 T with multiple-animal acquisition strategies. *Mag Reson Mater Phy* **2006**, 19, 202–208.
9. Lauridsen, H.; Hansen, K.; Wang, T.; Agger, P.; Andersen, J.L.; Knudsen, P.S.; Rasmussen, A.S.; Uhrenholt, L.; Pedersen, M. Inside Out: Modern Imaging Techniques to Reveal Animal Anatomy. *PLOS ONE* **2011**, 6(3).
10. Dinley, J.; Hawkins, L.; Paterson, G.; Ball, A.D.; Sinclair, I.; Sinnott-Jones, P.; Lanham, S. Micro-computed X-ray tomography: a new non-destructive method of assessing sectional, fly-through and 3D imaging of a soft-bodied marine worm. *J Microsc* **2010**, 238, 123–133.
11. Ziegler, A.; Faber, A.; Mueller, S.; Bartolomaeus, T. Systematic comparison and reconstruction of sea urchin (*Echinoidea*) internal anatomy: a novel approach using magnetic resonance imaging. *BMC Biol* **2008**, 6, 1–15.
12. Rieppel, O.; Gauthier, J.; Maisano, J. Comparative morphology of the dermal palate in squamate reptiles, with comments on phylogenetic implications. *Zool J Linn Soc* **2008**, 152, 131–152.
13. Digital Fish Library. A collaborative project at the University of California, San Diego funded by the National Science Foundation's (NSF) Biological Infrastructure program. Available online: www.digitalfishlibrary.org (accessed on 8 June 2023).
14. Webb, E.; Yuan, M.; Lemoine, N.; Wang, Y. Imaging in animal models. *Integr Cancer Sci Ther* **2016**, 3, 428–431.
15. Schambach, S.J.; Bag, S.; Schilling, L.; Groden, C.; Brockmann, M.A. Application of micro-CT in small animal imaging. *Methods* **2010**, 50, 2–13.
16. Farrag, M.; Pukale, D.D.; Leipzig, N.D. Micro-computed tomography utility for estimation of intraparenchymal spinal cord cystic lesions in small animals. *Neural Regen Res* **2021**, 16(11), 2293–2298.
17. Delk, K.W.; Mejia-Fava, J.; Jiménez, D.A.; Kent, M.; Myrna, K.; Mayer, J.; Divers, S. Diagnostic imaging of peripheral vestibular disease in a Chinese goose (*Anser cygnoides*). *J Avian Med Surg* **2014**, 28, 31–37.
18. Fleming, G.J.; Lester, N.V.; Stevenson, R.; Silver, X.S. High field strength (4.7 T) magnetic resonance imaging of hydrocephalus in an African Grey parrot (*Psittacus erithacus*). *Vet Radiol Ultrasound* **2003**, 44, 542–545.
19. Jirak, D.; Janacek, J.; Kear, B. P. A combined MR and CT study for precise quantitative analysis of the avian brain. *Sci. Rep* **2015**, 5(1), 1–7.
20. Romagnano, A.; Shiroma, J.T.; Heard, D.J.; Johnson, R.D.; Schiering, M.R.; Mladinich, M.S. Magnetic resonance imaging of the brain and coelomic cavity of the domestic pigeon (*Columba livia domestica*). *Vet Radiol Ultrasound* **1996**, 37, 431–440.

21. Fumero-Hernández, M.; Encinosa, M.; Ramírez, A.S.; Morales, I.; Suárez Pérez, A.; Jaber, J.R. A Cadaveric Study Using Computed Tomography for Measuring the Ocular Bulb and Scleral Skeleton of the Atlantic Puffin (Aves, Alcidae, *Fratercula arctica*). *Animals* **2023**, *13* (15): 2418.
22. Hall, M.I. The anatomical relationships between the avian eye, orbit and sclerotic ring: Implications for inferring activity patterns in extinct birds. *J. Anat* **2008**, *212*, 781–794.
23. Andrade, S.B.D.; Araujo, N.L.L.C.D.; Raposo, A.C.S.; Muramoto, C.; Oriá, A.P. Morphometric descriptive report of scleral ossicle rings, by ultrasound and computed tomography, in three Testudines specimens. *Ciência Rural* **2023**, *53*, 1–10.
24. Orosz, S. E.; Toal, R.L. Tomographic anatomy of the golden eagle (*Aquila chrysaetos*). *J. Zoo Wildl. Med* **1992**, *23*, 39–46.
25. Veladiano, I.A.; Banzato, T.; Bellini, L.; Montani, A.; Catania, S.; Zotti, A. Normal computed tomographic features and reference values for the coelomic cavity in pet parrots. *BMC Vet.Res* **2016**, *12* (1), 1–9.
26. da Silva, J.P.; Rahal, S.C.; Castiglioni, M.C.R.; Baldissera Gonçalves, R.A.; Doiche, D.P.; Moresco, A.; Mamprim, M.J.; Vulcano, L.C. Radiography and computed tomography of the heart and lower respiratory tract in toco toucans (*Ramphastos toco*). *Anat Histol Embryol* **2020**, *49*(4), 541–549.
27. Pepperberg, I.M.; Howell, K.S.; Banta, P.A.; Patterson, D.K.; Meisser, M. Measurement of grey parrot (*Psittacus erithacus*) trachea via magnetic resonance imaging, dissection, and electron beam computed tomography. *J Morphol* **1998**, *238*(1), 81–91.
28. IUCN. The IUCN Red List of Threatened Species. Version 2022-2. Available online: <https://www.iucnredlist.org> (accessed on 2 June 2023).
29. Painter, J. *Fratercula arctica*. Available online: https://animaldiversity.org/accounts/Fratercula_arctica/ (accessed on 30 May 2023).
30. Sisson, S.; Getty, R.; Grossman, J.D. *Anatomía de los animales domésticos*, 5th ed.; Elsevier Masson: Issy les Moulineaux , France, 1982.
31. MacLellan, J. *Atlas en color de anatomía de las aves*. McGraw-Hill Interamericana: Madrid, Spain, 1992.
32. König, E.H.; Liebich, H.G.; Bragulla, H. *Anatomie und Propädeutik des Geflügels*. Schattauer: Stuttgart, German, 2001.
33. Isaac, I.; Richardson, J.; Liuti, T.; Longo, M. Safety of intravenous iodinated contrast medium injection in rabbits undergoing conscious computed tomography. *Veterinary Record Open* **2022**, *9*(1), e31.
34. Seamon, A.B.; Hofmeister, E.H.; Divers, S.J. Outcome following inhalation anesthesia in birds at a veterinary referral hospital: 352 cases (2004–2014). *J Am Vet Med Assoc* **2017**, *251*(7), 814–817.
35. Arencibia, A.; Corbera JA.; Ramírez, G.; Díaz-Bertrana, M.L.; Pitti, L.; Morales, M.; Jaber, JR. Anatomical Assessment of the Thorax in the Neonatal Foal Using Computed Tomography Angiography, Sectional Anatomy, and Gross Dissections. *Animals* **2020**, *10*(6):1045.
36. Banzato, T.; Selleri, P.; Veladiano, IA.; Zotti, A. Comparative evaluation of the cadaveric and computed tomographic features of the coelomic cavity in the green iguana (*Iguana iguana*), black and white tegu (*Tupinambis merianae*) and bearded dragon (*Pogona vitticeps*). *Anat Histol Embryol* **2013**, *42*(6):453–60.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.