

Review

Not peer-reviewed version

---

# Various Biomimetics as Antifungals

---

[Elena Efremenko](#)<sup>\*</sup>, [Aysel Aslanli](#), Nikolay Stepanov, [Olga Senko](#), [Olga Maslova](#)

Posted Date: 28 August 2023

doi: 10.20944/preprints202308.1824.v1

Keywords: antifungal peptides; enzyme-like activity; growth inhibition; antifungal activity; fungal metabolites; quorum response molecules; combined antifungals



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

# Various Biomimetics as Antifungals

Elena Efremenko \*, Aysel Aslanli, Nikolay Stepanov, Olga Senko and Olga Maslova

Faculty of Chemistry, Lomonosov Moscow State University, Lenin Hills 1/3, Moscow 119991, Russia; elena\_efremenko@list.ru (E.E.); ayselaskanli@mail.ru (A.A.); na.stepanov@gmail.com (N.S.); senkoov@gmail.com (O.S.); olga.maslova.rabota@gmail.com (O.M.)

\* Correspondence: elena\_efremenko@list.ru; Tel.: +7-(495)-939-3170; Fax: +7-(495)-939-5417

**Abstract:** Biomimetics, which are similar to natural compounds that play an important role in the metabolism, manifestation of functional activity and reproduction of various fungi, have a pronounced attraction in the current search for new effective antifungals. Actual trends in the development of this area of research indicate that unnatural amino acids can be used as such biomimetics, including those containing halogen atoms; compounds similar to nitrogenous bases embedded in the nucleic acids synthesized by fungi; peptides imitating fungal analogues; molecules similar to natural substrates of numerous fungal enzymes and Quorum Sensing signaling molecules of fungi and yeast; etc. Most part of the review is devoted to the analysis of semi-synthetic and synthetic antifungal peptides and their targets of action. This review is aimed at combining and systematizing the current scientific information accumulating in this area of research, developing various antifungals with an assessment of the effectiveness of the created biomimetics and the possibility of combining them with other antimicrobial substances to reduce cell resistance and improve antifungal effects.

**Keywords:** antifungal peptides; enzyme-like activity; growth inhibition; antifungal activity; fungal metabolites; quorum response molecules; combined antifungals

## 1. Introduction

Currently, there is an urgent problem of efficient prevention of plants, animal and human diseases triggered by pathogenic fungi [1–3]. Infections that cause plant diseases annually affect food crops and destroy the harvested crop during storage, which entails the formation of serious social and economic problems. In addition, among plant protection products there are many fungicides which are toxic to humans, as well as those that have a negative impact on agrobiocenoses, causing environmental pollution and the accumulation of remaining amounts of the pesticides in agricultural products [4]. Filamentous fungi (*Aspergillus*, *Cladosporium*, *Penicillium*, *Stachybotrys*, *Fusarium*) synthesizing various mycotoxins are particularly dangerous, both at the stage of crop growth and at the stage of its storage [5].

With fungal (yeast) contaminations of humans and animals, special difficulties arise in the treatment of invasive infections characterized by high mortality [6]. Currently, various antifungals are used to combat such fungal pathogens as polyene antibiotics (nistatin, amphotericin B), azoles (fluconazole), modified lipopeptides (echinocandins), allylamines (terbinafine), etc. [7,8]. Most of them are synthetic compounds, since then there are concerns about their high toxicity due to limited water solubility and the need to use their relatively high concentrations, as well as about the decrease in the effectiveness of their action due to the emerging resistance in fungi. [9].


The interest exists to the natural compounds possessing strong antifungal activity with low enough toxicity to other eukaryotic cells, however most of such compounds appeared to be biodegradable by fungal strains. The fungal pathogens can use almost any natural substrates, due to their ability to synthesize various hydrolases and oxidoreductases with a wide substrate specificity of action [10,11]. Having a unique ability to adapt to changing environmental conditions, fungi can overcome the effects of various antifungal agents. In addition, the frequent and irregular use of fungicides has led to the emergence of resistant strains [12]. At the same time, such resistance of fungi

at the present stage of research of these microorganisms is also associated with the presence of a "Quorum Sensing" (QS) mechanism similar to bacterial populations [13].

In this regard, the development of new antifungal agents is an urgent task and there is a huge interest in biomimetics, that is, substances imitating their natural analogues. Thus, natural antifungal peptides (AFPs) have demonstrated unprecedented advantages as selective biomaterials for obtaining effective biologics [14]. Therefore, AFPs obtained from natural sources (plants, fungi, bacteria), which can be chemically modified or artificially synthesized to improve their antifungal properties and reduce side effects are of great interest. When obtaining AFPs as biomimetics, the possibility of increasing their stability and bioavailability turns out to be very attractive.

In addition to AFPs, the synthesis of a number of other compounds and materials forming various surfaces imitating natural analogues that can be attributed to biomimetics is also interesting from scientific and practical point of view. The antifungal effect of these compounds and materials is not limited to the destruction of the cell wall and membranes of fungi, but affects their various metabolically important pathways, growth and spore formation [15–17]. Of great interest is the possible synergistic effect among such biomimetics with antifungal drugs, which allows to reduce the doses of individual substances used [18,19].


Thus, the purpose of this review was to analyze and summarize the main current trends in research on the production of antifungal biomimetics using various approaches (Figure 1) based on damage to the integrity of cells and individual organelles, disruption of synthetic processes in fungal cells, inhibition of the activity of various enzymatic systems, etc. A separate emphasis in the review was made on comparing the antifungal effect that various variants of biomimetics have on different fungal cells, and their studied combinations in order to identify the most effective and promising ones for further improvement and application.



**Figure 1.** Current trends in the development of biomimetic antifungals analyzed in this Review.

## 2. Antifungal peptides (AFPs)

Antimicrobial peptides (AMPs) produced by living organisms attract particular interest as new antimicrobial agents with a wide range of biotargets (bacteria, fungi, parasites) capable of replacing blockbuster antibiotics [20]. To date, there are a number of databases for documenting AMPs [21]. In this review, we considered information from four of them: the Antimicrobial Peptides Database (APD3), the Collection of Antimicrobial Peptides (CAMP<sub>R3</sub>), the Database of Antimicrobial Activity and Structure of Peptides (DBAASP), and the Database of Antimicrobial Peptides (dbAMPs) [22–25] (Figure 2).



**Figure 2.** Composition of AMP databases according to the source of peptides with antifungal activity. Total number of AFPs in each database was taken as 100%: 1,311, 1,144, 5,968 and 5,540 for APD3, CAMP<sub>R3</sub>, DBAASP and dbAMPs, respectively.

APD3 is a system dedicated to discovery timeline, glossary, nomenclature, classification, structure, information search, prediction, design, statistics and tools of AMPs and beyond. As of August 2023, the APD3 contains totally 3,569 AMPs.

The dbAMP database created by Dr. Lee's team contains information about 28,709 different types of AMPs in 3044 organisms, including experimentally verified AMPs. The information from such protein databases as UniProt, NCBI, Protein Data Bank was used for the construction of dbAMP.

DBAASP database contains 20,876 AMPs and information about their chemical structure, amino acid sequences, target species and object, hemolytic and cytotoxic activities of peptides. In addition, DBAASP provides a tool for the *in silico* prediction/design of new AMPs.

CAMP<sub>R3</sub> contains information on the AMP sequences, protein definition, accession numbers, activity, source organism, target organisms, protein family descriptions, N and C terminal modifications. Currently, the database contains 8164 peptide sequences and also provides tools for sequence alignment, pattern creation, pattern and HMM-based search.

The peptides in these databases can be classified into four categories: (i) natural AMPs; (ii) predicted peptides, which are predicted by machine learning or other technologies and tested to be active; (iii) synthetic peptides derived from natural AMPs and (iv) patented AMPs. Thus, the described AMP databases are an effective tool for the analysis, prediction and design of new peptides with desired properties, particularly AFPs.

As of August 2023, the proportion of AFPs among all AMPs reported in the APD3, CAMP<sub>R3</sub>, DBAASP and dbAMPs databases was 36.7 %, 14 %, 28.6 % and 19.3 %, respectively (Figure 2).

Interestingly, the ratio of synthetic AFPs (SAFPs) to natural AFPs varied depending on the database under consideration and was the maximum (68.1 %) in the DBAASP database.

According to the data of Figure 2 and Table 1 [26–41]), a number of AMPs with antifungal activity have already been identified, which can be obtained from various natural sources. Due to the possibility of obtaining of SAFP capable of imitating various natural peptides, such biomimetics can find effective application in practice and are notable objects for current developments and investigations (Table 2 [32,34,37,42–58]).

It should be noted that AFPs can be classified by total charge, secondary structure, mechanism of action, etc. In this review, when discussing AFPs, the main attention was paid to their origin (natural, semi-synthetic or synthetic), the object of their influence, the mechanism of their influence on fungal cells and the level of antifungal effect. Such information was summarized in Tables 1 and 2.

**Table 1.** Natural antimicrobial peptides with antifungal activity.

Antifungal peptide	Origin	Molecular weight [Reference]	Antifungal effect
<b>Mechanism of action: inhibition of chitin biosynthesis</b>			
Nikkomycin Z	fungi	495.4 Da [26]	0.5-64 mg/L (yeasts, fungi) <sup>a</sup>
<b>Mechanism of action: destabilization of plasma membrane, pore formation, cell wall damage</b>			
Magainin 2	frog	2466.9 Da [27,28]	6.25 µM ( <i>Saccharomyces cerevisiae</i> , <i>Trichosporon beigelii</i> , <i>Candida albicans</i> ) <sup>a</sup> 60-100 mg/L ( <i>Penicillium digitatum</i> ; <i>Alternaria solani</i> ; <i>Phytophthora infestans</i> ) <sup>a</sup>
Halictine Hal-2	sweat bee	1452.85 Da [29]	1.6-25 µM ( <i>Candida</i> spp.) <sup>a</sup>
Halocidin	ascidian	3445.1 Da [30]	Effect was not estimated
Polymyxin B	bacteria	1301.6 Da [31]	16-256 mg/L (multi-drug-resistant fungal strains) <sup>a</sup>
Colistin	bacteria	1155.4 Da [31]	
Defensin DefMT3	ticks	1613.1 Da [32]	4 µM ( <i>Fusarium culmorum</i> ; <i>F. graminearum</i> ) <sup>b</sup>
Indolicidin	bovine	1906.3 Da [33]	12.5-50 mg/L ( <i>C. albicans</i> ) <sup>a</sup>
Lactoferrampin B	bovine	2389.8 Da [34]	0.7-39 µM ( <i>C. albicans</i> ) <sup>c</sup>
Lactoferricin B	bovine	3125.8 Da [34]	0.31-400 mg/L (yeasts) <sup>a</sup> 4-32 µM (fungi) <sup>a</sup>
Lactoferricin H	human	5513 Da [34]	10 mg/L ( <i>C. albicans</i> ) <sup>a</sup>
Leg2	chickpea legumin hydrolysates	2157.6 Da [35]	125-250 µM ( <i>S. cerevisiae</i> , <i>Zygosaccharomyces bailii</i> ) <sup>a</sup>
LL-37	human	4493.3 Da [36]	4-64 µM ( <i>Candida</i> spp.) <sup>a</sup>
<b>Mechanism of action: cell/spore lysis, cell wall perturbations</b>			
Cecropin B	silkworm	3835.7 Da [38]	0.9 mg/L ( <i>C. albicans</i> ) <sup>a</sup> 160-320 mg/L ( <i>F. solani</i> ) <sup>a</sup>
Osmotin	plant	24285.3 Da [38]	4-25 mg/L (fungi) <sup>a</sup>
Stomoxyn	stable fly	4474.2 Da [39]	0.8-50 µM (yeasts); 0.4-7 µM (fungi) <sup>a</sup> 50-100 µM ( <i>A. fumigatus</i> ) <sup>a</sup>
Temporin B	frog	1391.8 Da [40]	1.4-4 µM ( <i>Candida</i> spp.) <sup>a</sup>
Temporin G		1457.8 Da [41]	8-128 µM (yeasts/fungi) <sup>a</sup>

<sup>a</sup>MIC, minimal inhibiting concentration; <sup>b</sup>IC<sub>50</sub>, half maximal inhibitory concentration; <sup>c</sup>LC<sub>50</sub>, 50% lethal concentration.



**Table 2.** Synthetic / (semi)synthetic antimicrobial peptides with antifungal activity.

Antifungal peptide	Molecular weight; [Reference]	Antifungal effect
<b>Mechanism of action: inhibition of 1,3-<math>\beta</math>-D-glucan synthase</b>		
Anidulafungin	1140.2 Da [42–44]	0.06–0.25 mg/L ( <i>Candida</i> spp.) <sup>a</sup> 0.015–32 mg/L (fungi) <sup>a</sup>
Caspofungin	1093.3 Da [42–44]	0.25–4 mg/L ( <i>Candida</i> spp.) <sup>a</sup>
Micafungin	1270.3 Da [42–44]	0.015–4 mg/L ( <i>Candida</i> spp.) <sup>a</sup>
<b>Mechanism of action: destabilization of plasma membrane, pore formation, cell wall damage</b>		
Lf(1–11) H	1317.5 Da [37]	> 12.5 mg/L ( <i>Candida</i> sp.); 80–160 mg/L ( <i>F. solani</i> ) <sup>a</sup> 4.3 $\mu$ M ( <i>A. fumigatus</i> ) <sup>a</sup>
Lfchimera (bLfcin/Lfampin)	4422 Da [34]	6.25 mg/L ( <i>C. parapsilosis</i> ) <sup>a</sup>
$\gamma$ -core DefMT3	1611.8 Da [32]	1–2 $\mu$ M ( <i>F. culmorum</i> ; <i>F. graminearum</i> ) <sup>b</sup>
Brilacidin <sup>d</sup>	936.9 Da [45]	2.5–>80 $\mu$ M ( <i>C. neoformans</i> , <i>C. albicans</i> , <i>C. auris</i> , <i>A. fumigatus</i> ) <sup>a</sup>
RcAlb-PepII	637.77 Da [46,47]	17–250 $\mu$ M ( <i>Candida</i> spp.) <sup>a</sup> 0.04 mg/L ( <i>Cryptococcus neoformans</i> ) <sup>a</sup>
Halictine Hal-2 derivatives	1471 Da [48]	0.5–1 $\mu$ M ( <i>Candida</i> spp., <i>S. cerevisiae</i> ) <sup>a</sup>
di-K19Hc	4115.1 Da [49]	< 4 mg/L ( <i>C. albicans</i> ); <16 mg/L ( <i>Aspergillus</i> sp.) <sup>a</sup>
Pexiganan/MSI-78	2478.2 Da [37]	10–80 mg/L ( <i>F. solani</i> ) <sup>a</sup>
PepGAT	1044.18 Da [46,50,51]	40–80 mg/L ( <i>Candida</i> spp., <i>P. digitatum</i> ) <sup>a</sup>
PepKAA	1238.44 Da [46,50,51]	
Peptide 77-3	994.2 Da [52,53]	3.5–5 mg/L ( <i>A. flavus</i> , <i>A. parasiticus</i> ) <sup>a</sup>
KK14	144.2 Da [54]	6.25–100 mg/L (fungi) <sup>a</sup>
D4E1	2079.4 Da [52,53]	7.75 $\mu$ M ( <i>A. flavus</i> ); 0.60 $\mu$ M ( <i>V. dahliae</i> ) <sup>b</sup> 13.02 $\mu$ M ( <i>C. destructivum</i> ) <sup>b</sup>
PAF26	991.2 Da [55]	4–6 $\mu$ M ( <i>P. digitatum</i> ) <sup>a</sup>
<b>Mechanism of action: production of reactive oxygen species, cell wall degradation</b>		
Octominin	2652.2 Da [56]	50 mg/L ( <i>C. albicans</i> ) <sup>a</sup>
Mo-CBP <sub>3</sub> -PepI	893.12 Da [57]	2.2 $\mu$ M ( <i>C. albicans</i> ) <sup>c</sup>
<b>Mechanism of action: Cell/spore lysis, cell wall perturbations</b>		
Osm-pepA	3050.5 Da [58]	40 $\mu$ M ( <i>S. cerevisiae</i> ) <sup>a</sup> 20 $\mu$ M ( <i>Pichia pastoris</i> ) <sup>a</sup>

<sup>a</sup>MIC, minimal inhibiting concentration; <sup>b</sup>IC<sub>50</sub>, half maximal inhibitory concentration; <sup>c</sup>IC<sub>90</sub>, concentration that inhibited yeast growth by 90%; <sup>d</sup>Non-peptide mimetic of host defense peptides.

Despite the fact that the mechanism of action of AMPs in relation to fungal cells is not as well studied as in relation to bacteria, the basic principle of their antifungal action is similar to antibacterial one and most often consists in a violation of the functions and integrity of the cell wall and plasma membrane. Due to this principle of action of AMPs, the development of resistance to them in microorganisms is considered to become almost impossible. AFPs, in addition to mentioned effect on fungal cells, are able to specifically inhibit membrane proteins,  $\beta$ -1,3-glucan and chitin synthases, thereby contributing to the formation of defects in the cell wall, or inhibit H<sup>+</sup>-ATPase, causing an apoptosis-like process. Some AFPs can also induce intracellular generation of reactive oxygen species (ROS) destroying various biomolecules (lipids, proteins, nucleic acids, etc.) by their active oxidation [59,60] (Table 1).

Semi-synthetic AFPs are chemically modified natural peptides, while preserving the active centers of the origin molecule, in order to achieve optimal properties [61]. Echinocandins (anidulafungin, caspofungin, micafungin), having a lipopeptide nature, are members of the "youngest" clinically used group of semi-synthetic AFPs. Echinocandins consist of a cyclic hexapeptide nucleus with a lipid side chain. The peptide lipid tail anchors the lipopeptide in the cell membrane next to the target enzyme. These peptides act as non-competitive inhibitors of a key enzyme in the synthesis of  $\beta$ -1,3-glucan. Unlike semi-synthetic AFPs, SAFP are obtained entirely by chemical synthesis. Most often, a solid-phase method is used for the synthesis of SAFP, based on the addition of one amino acid at one step of synthesis, which allows investigators to study the role of each amino acid in the synthesized sequence.

When analyzing examples of currently created semi-synthetic and synthetic AFPs, the result of studying their properties and mechanisms of action, as well as comparing them with natural analogues, is always interesting and useful. It appeared that targets of action of AFPs can be different and that depends on their chemical structure.

The mechanism of action and activity of a series of synthetic analogues of the halictine HAL-2 peptide (from the venom of the wild bee *Halictus sexcinctus*) was investigated in relation to cells *Candida* spp. It has been found, that halictines can rapidly permeabilize cell membranes and cause the leakage of cytosolic components, and that their mode of action is likely to depend on the plasma-membrane sterols. The pre-treatment with the inhibitors of sterol synthesis (terbinafine and fluconazole) resulted in a significant reduction of peptides efficacy, while their killing efficacy increased when combined with amphotericin B [48].

It was shown, that peptides naturally derived from milk protein lactoferrin, such as lactoferricin (Lfcin), lactoferrampin (Lfampin) and Lf(1–11) and some modified or synthesized peptides can also exhibit strong antifungal activity. In particular, antifungal activity of bovine lactoferrin-derived Lfcin was investigated against a wide range of fungal species and it was shown, that bLfcin demonstrates strongly enhanced antimicrobial activity compared to lactoferrin [34,62–64].

The cacao osmotin-like protein (TcOsm1) -derived peptides, named Osm-pepA and Osm-pepB inhibited growth of yeasts (*Saccharomyces cerevisiae* S288C and *Pichia pastoris* X-33) and spore germination of the phytopathogenic fungi *Fusarium* f. sp. *glycines* and *Colletotrichum gossypii*. Osm-pepA was more efficient than Osm-pepB [58].

A synthetic analogue of antimicrobial peptide halocidin, di-K19Hc, has been exhibited improved antifungal activity against a panel of fungi including several strains of *Aspergillus* and *Candida* [49].

The small antimicrobial peptide PAF26 (Ac-RKKWFW-NH<sub>2</sub>) demonstrated multiple detrimental effects on the filamentous fungi *Penicillium digitatum*, which ultimately resulted in permeation and killing the growing cells [55].

The full-length *Neosartorya* (*Aspergillus*) *fischeri* AMPs and novel rationally designed  $\gamma$ -core peptide derivatives  $\gamma^{\text{NFAP-opt}}$  and  $\gamma^{\text{NFAP-optGZ}}$  exhibited high efficacy by inhibiting the growth of the agriculturally relevant filamentous ascomycetes in vitro [65].

Small peptide, RcAlb-PepII, designed based on the primary structure of Rc-2S-Alb, a 2S albumin from the seed cake of *Ricinus communis*, strongly inhibited the growth of *Klebsiella pneumoniae* and *Candida parapsilosis*, and induced morphological alterations in their cell surface. The peptide also degraded and reduced the biofilm formation in *C. parapsilosis* and in *K. pneumonia* cells [47].

SAFPs PepGAT, PepKAA, and Mo-CBP3-PepI demonstrated strong inhibition of *P. digitatum* growth. All peptides targeted the fungal membrane leading to pore formation, loss of internal content and death. The induction of high levels of ROS was also a mechanism employed by some peptides [51,66].

The antifungal activity of two peptides, called Mo-CBP3-PepI (CPIAQRCC) and Mo-CBP3-PepIII (AIQRCC), designed based on the structure of Mo-CBP3, a chitin binding protein purified from *Moringa oleifera* seeds, was evaluated against *C. albicans* and *C. parapsilosis* biofilms [67]. Eight SAMPs were tested regarding their antifungal potential against *C. neoformans* and five SAMPs showed an inhibitory effect on *C. neoformans* growth at low concentrations. Peptides induced many

morphological alterations such as cell membrane, wall damage, and loss of internal content on *C. neoformans* cells [46].

Investigation of the effect of four antimicrobial peptides - PPD1 (FRLHF), 66-10 (FRLKFH), 77-3 (FRLKFHF) and D4E1 (FKLRKIKVRLRAKIKL) on the aflatoxin production by *A. flavus* and *A. parasiticus* suggested that AMPs at near minimum inhibitory concentrations (MIC) were effectively inhibiting aflatoxins, without hindering the growth of the fungi. At higher concentrations, these peptides exerted fungicidal action on *A. flavus* [52,53].

Antimicrobial peptides polymyxin B and colistin (polymyxin E) exhibited an antifungal activity against 11 MDR yeast and filamentous fungal strains including strains belonging to the *Candida*, *Cryptococcus* and *Rhodotorula* yeast genera, along with others belonging to the *Aspergillus*, *Fusarium*, *Scedosporium*, *Lichtheimia* and *Rhizopus* with MICs ranging from 16 to 128 µg/ml, except for the *Aspergillus* species [31].

In addition to the methods of chemical modification and synthesis of AMPs molecules, the use of methods of computer molecular design played a significant role in obtaining synthetic AMPs with the desired characteristics. This made it possible to combine information about the chemical properties and biological activity acquired by new peptides and the amino acid sequences present in them. This, in turn, made it possible to develop methods for predicting and evaluating the antifungal potential of synthesized sequences in silico [68].

Using an easy step-by-step way to choose, characterize and test potential sequences to be assayed as synthetic AMPs, two peptides (PepGAT and PepKAA) with antimicrobial potential against *Candida* spp., including activity against biofilms and without any hemolytic effects, were identified and characterized [50].

Using a simple and reliable method a set of an anti-*Candida* peptide CGA-N12 analogues were rationally designed and seven CGA-N12 analogues with significantly improved antifungal activity against *C. tropicalis* were screened [69].

The rationally designed and synthesized new structural class of dipeptides Trp-His(1-Bn)-OMe/NHBn and tripeptides His(1-Bn)-Trp-His(1-Bn)-OMe/NHBn possessing modified amphiphilic histidine along with hydrophobic tryptophan residues demonstrated promising antifungal activity with membrane lytic action against *C. neoformans* [70].

A designed and synthesized synthetic peptide of 23 amino acids Octominin (1GWLIRGAIHAGKAIHGLIHRRRH23) from a defense protein 3 cDNA sequence of Octopus minor showed an inhibitory effect against *Candida albicans* by causing ultrastructural cell wall deformities [56].

Ultrashort peptide H-Orn-Orn-Trp-Trp-NH<sub>2</sub> (O3TR) inhibited the growth of the filamentous fungi *Fusarium culmorum*, *Penicillium expansum* and *A. niger* and yeasts *Saccharomyces cerevisiae*, *Zygosaccharomyces bailii*, *Z. rouxii*, *Debaryomyces hansenii* and *Kluyveromyces lactis*. The addition of a C12 fatty-acid chain tail at the N-terminus of these peptide improved its antifungal activity by 2 - 8-fold in relation to different fungi [71].

A C14 - residue peptide named KK14, with the sequence KKFFRAWWAPRFLK - NH<sub>2</sub>, was designed and inhibited conidial germination and fungal growth of food contaminants. The substitution of a Pro residue with Arg increased the helical content of the peptide not only its antifungal activity but also its cytotoxicity. The insertion of an unnatural bulky residue β - diphenylalanine or a full denantiomerization overall increased the antifungal potency [54].

Inspired by antimicrobial peptides different side chain methionine, leucine and tyrosine based polymethacrylates (pH responsive cationic biocompatible polymers) have been designed and their fungicidal activities have been investigated against *A. niger*, demonstrating the effective inhibition of hyphal growth and distortion of conidiophores [72].


The antifungal activity of a series of peptides with a varying number of lysine and tryptophan residue repeats (KW<sub>n</sub>-NH<sub>2</sub>) was confirmed against *C. albicans* [73].

It has been shown that a number of AFPs can exhibit not only antifungal activity, but also inhibit production of mycotoxins synthesized by fungi as molecules for their self-defense [52,59]. It turned out that in some cases, individual AFPs may lose their ability to inhibit the growth of fungi, but retain



their effectiveness in inhibiting the biosynthesis of mycotoxins by fungi. For instance, the treatment of *F. graminearum* cells with the reduced form of the  $\gamma$ -core of the tick defensin DefMT3 (TickCore3, TC3), decreases growth of the fungal cells and abrogates mycotoxin (trichothecene B) production. The oxidation of TC3 leads to loss of its growth inhibitory activity, while the anti-mycotoxin activity is retained [74].

Summarizing the information presented, it can be concluded that various methods of chemical modification of natural peptides, including phosphorylation, cyclization, halogenation, etc. [60], can be used to obtain semi-synthetic AFPs (Figure 3). However, in addition to the same methods, computer molecular analysis and the design of the peptides being created play an important role in the design of synthetic AFPs, which makes it possible to proceed not from the natural basis that can only be modified, but to obtain the synthesis of new compounds with the desired properties and taking into account the use of key amino acid motifs (Figure 3). At the same time, of course, all new AFPs must be checked not only for the effectiveness of the antifungal effect obtained from them, but also to evaluate them for toxicity, including prior properties [75].



**Figure 3.** Approaches used in the development of biomimetic AFPs.

Thus, the use of various AFPs for their modifications described in this section is an effective approach to solving the problem of pathogenicity and resistance of fungi. The fact that these peptides, as natural molecules, make it possible to create a number of different biomimetics/peptidomimetics based on them, which can be used as antifungals, allows us to conclude that there is a need for significant development of rational design methods today. The use of organic synthesis methods makes it possible to construct SAFP with the most effective amino acid sequences and desired characteristics as well as changed mechanisms (targets) of action.

### 3. Biomimetics with antifungal activity of non-peptide nature

In addition to AFPs, various molecules can act as biomimetics with antifungal action, which play a significant role in maintaining the viability and metabolic activity of fungal cells. Such molecules may include non-canonical amino acids potentially involved in the construction of structural proteins and enzymes, analogues of nitrogenous bases involved in the formation of nucleic acids (RNA and DNA), analogues of hormones involved in the formation of membrane structures; mimetics of QS molecules of fungi etc.

The development of various materials and tools that acquire antifungal properties due to the inclusion of various biomimetics in their content is now coming to the forefront of scientific and practical research. At the same time, the action of biomimetics inside of such developed materials can be divided into those that (Figure 1, Tables 3–5):

- prevent adhesion of spores and vegetative cells of fungi, as well as their subsequent invasive development;
  - affect the membranes and cell walls of fungi, showing fungicidal activity, which leads to cell death;
  - change the morphology and metabolic activity of fungal cells, inhibiting the synthesis of mycotoxins as a means of protecting fungi, as well as the most important enzymes of cells and QS molecules involved in the development of resistance of fungi to various influences;
  - inhibit cell growth and spore formation, reducing the biosynthesis of adenosine triphosphate (ATP) and nucleic acids.
- Further, a number of these main effects of different biomimetics on fungi will be considered separately.

3.1. Biomimetics with a destructive effect on fungal cells

Polyene antibiotics are the most well-known effective and widely used compounds with antifungal action. Nystatin (polyene macrolide synthesized by bacteria of the genus *Streptomyces*) is one of the classic antifungal antibiotics. However, there is a traditional interest in biomimetics, which are chemically modified nystatin molecules, the proposed mechanism of action of which is binding to ergosterol in the structure of lipid bilayers of fungal membranes. In this case, extra-membrane aggregates are formed that inhibit the transport of amino acids and glucose through the plasma membrane of cells [76]. Finally, polyene biomimetics contribute to the formation of macropores in the cell membrane and leads to their destruction.

The reason for the creation of new biomimetics is that nystatin itself has poor water solubility, instability to light, acidic air and heating, which limits its use. A significant amount of research is aimed at finding ways to avoid these disadvantages of natural nystatin. Chitosan, gelatin, alginate and other polymers are being studied for its modification by conjugation and improvement of mentioned characteristics [77,78]. Selenium, bismuth, and copper are often used to modify biomimetics based on polyene macrolides, which make it possible to enhance their antifungal effect [78–80].

A special place in the ongoing research takes the study of the possibility of increasing the specificity of the toxic effects of the obtained biomimetics on fungal cells. Thus, the obtained nystatin amides demonstrated increased water solubility, reduced toxicity in relation to human kidney cells (HEK293) while maintaining inhibition of ergosterol biosynthesis by fungal cells [76].

In the case of another polyene macrocyclic antibiotic (amphotericin B) with antifungal action, isolated from extract of *Aloe vera* leaf, the effectiveness of such modification was shown, which leads to the formation of its nanoparticles with slow release of the main substance, which causes the death of yeast cells and mycelial fungi [81,82] (Table 3 [16,72,82–88]).

**Table 3.** Biomimetics as antifungals with effect on fungal membranes and cell walls.

Biomimetics as antifungals	Target of action	*Antifungal effect
Amphotericin B nano-aggregates [82]	Inhibition of ergosterol biosynthesis for membrane formation and provoking lysis of cells	MIC (mg/L): <i>Candida spp.</i> (0.125–0.5); <i>A. fumigatus</i> (1.0)
Biomimetic nanopillar Si-containing surfaces [16]	Rupture of coat and inner membrane of spores leading to cell death	4-fold reduction of amounts of attached spores and approximately 9-fold reduction of viable conidia of <i>Aspergillus brasiliensis</i> on biomimetic surface

Methacryloyloxyethyl ester monomers with tyrosine, methionine and leucine [72]	Destruction of the cell membrane	MIC (mg/L): <i>A. niger</i> (0.16)
Synthetic peptides from predicted cysteine-rich peptides of tomato (mimicking $\gamma$ -core regions of cysteine-rich peptides of <i>Solanum lycopersicum</i> ) [83]	Charge of the derived peptides is positive, favoring interactions with the membranes of the pathogens. Inducing permeabilization and disruption of the fungal membranes	IC <sub>50</sub> ( $\mu$ M): <i>Cryptococcus neoformans</i> (5.1-11.5); <i>Fusarium culmorum</i> (42.1-126.7); <i>F. oxysporum</i> (43.8-165.8); <i>F. solani</i> (47.5-138.8); <i>F. verticillioides</i> (99.8-152.0)
Itraconazole and difluorinated-curcumin containing chitosan nanoparticles in hydrogel [84]	Synergistic antifungal activity composed of increased permeation through fungal cell wall and membrane and lethal action of difluorinated curcumin	EC <sub>50</sub> (mg/L): <i>Trichophyton mentagrophytes</i> (150)
Chitosan/polyethylene oxide (CPO) [85]	Antifungal effect similar to voriconazole (production of intracellular ROS and increasing of permeability of plasma membrane)	<i>C. albicans</i> cells: diameter of inhibition area by CPO - 25-27 mm (agar disc diffusion method); Control: inhibition by voriconazole – 27 mm
N-alkylated glycine oligomers (peptoids) [86]	Suppressed formation of hyphae resulted in changes in cell and organelle morphology, most dramatically in the nucleus and nucleolus, and in the number, size, and location of lipidic bodies	MIC ( $\mu$ M): <i>C. albicans</i> (3.0-13.0)
Metal-organic framework (Ce-MOF) with enzyme-like activity of catalase, superoxide dismutase, and peroxidase [87]	Production of ROS and inhibition of fungal growth	40 $\mu$ g/mL Ce-MOF gives 93.3–99.3% growth inhibition of <i>Aspergillus flavus</i> , <i>A. niger</i> , <i>A. terreus</i> , <i>C. albicans</i> , <i>Rhodotorula glutinis</i>
Iodine-doped carbon dots (IDCDs) with peroxidase-like activity [88]	Production of ROS	The 90% decrease in number of <i>C. albicans</i> CFU is observed in presence of 2.72 g/L IDCDs under visible light irradiation over 2 h in presence of 0.5 mM H <sub>2</sub> O <sub>2</sub>

\*MIC - minimum inhibitory concentration; EC<sub>50</sub> - half maximal effective concentration.

Often the use of various biomimetics in the form of nanoparticles (often by encapsulating the main active substances) makes it possible to increase their antifungal activity [84].

Peptoids are considered as perspective peptidomimetics consisting of N-alkylated glycine oligomers  $(-RN-CH_2-CO-)_n$ . The compounds possess capability of forming a wide variety of secondary structures, including:  $\alpha$ -helix, polyproline type I helix, and type I and type III  $\beta$ -turns. Peptoids have high resistance to proteolysis and demonstrate antifungal activity. It was revealed that peptoids cause disruption of the membrane and DNA molecules of *C. albicans* yeast cells [86]

Synthesis of biomimetics based on  $\gamma$ -core regions of cysteine-rich peptides of *Solanum lycopersicum* allowed obtaining antifungal compounds that permeabilize fungal membranes of yeast-like fungus *Cryptococcus neoformans*, fungi of the genera *Fusarium*, *Bipolaris* and *Botrytis* [83].

Fungicidal activity was established in a number of biomimetic pH-responsive amphiphilic polymers based on amino acids found in antimicrobial peptides (tyrosine, methionine, leucine) synthesized via reversible addition fragmentation chain transfer polymerization technique. Such polymers violated the structural integrity of hyphae. Treatment with cationic polymers induced distribution of sterol components within plasma membrane lipid raft, followed by inhibition of the growth of hyphae. The interactions of fungal cells with polymers subsequently have prompted the altered cellular signaling resulting in failure of polarized cell growth. [72]. As a result, there was a complete loss of cellular integrity due to the emerging membrane depolarization and subsequent cell lysis. Among all amino acids, polymers with methionine showed higher activity than with leucine and tyrosine.

Interesting, from the point of view of antifungal effects, were the development of various materials, contact with which leads to a decrease in the number of living fungal cells. It was shown that by enhancing hydrophobicity or hydrophilicity, as well as changing the topography of the surface of materials into which, for example, nanostructured Si is introduced, it is possible to simulate the surface of dragonfly wings [16]. Such biomimicking materials possess antifungal activity due to prevention of attaching of *A. brasiliensis* fungal spores and conidia and the contact of fungal cells with such materials results in rupture of coat and inner membrane of spores leading to cell death.

New biomimetic nanofibers based on chitosan with polyethylene oxide acting against *C. albicans* have been developed. The antifungal effect was similar to known voriconazole [85]. The integration of peptides into synthetic macromolecular systems is attractive for obtaining materials with antifungal properties. Thus, molecular assembly of a cationic polymer based on peptides with anionic polyoxometallates leads to the formation of a well-defined structure.

A composite based on Phe-Ala dipeptide polymer and Keggin phosphotungstate was developed. Two amino acids, phenylalanine and alanine, were chosen to produce the dipeptide because of their amphiphilic properties and their ability to form higher-order self-assembling structures in water [89]. The resulting composite showed fungicidal activity against *A. niger* cells by increasing the local charge in the polymer. Significant deviation in cellular integrity and complete alteration in cell morphology was observed. The branches mycelia became flattened and stalked together, while no presence of conidiophores was noticed.

The cerium-based organometallic framework (Ce-MOF) turned out to be very interesting because of its high (93.3-99.3%) effectiveness in inhibiting the growth of vegetative cells of *A. flavus*, *A. niger*, *A. terreus*, *C. albicans* and *Rhodotorula glutinis* due to the generation of a high concentration of ROS [87]. Ce-MOF also caused extensive (up to 77.4%) deformation of the conidiophores of fungi with spores. The antifungal activity of Ce-MOF was due to its ability to act as a biomimetic of several oxidative enzymes at once (catalase, superoxide dismutase, and peroxidase). The observed morphological changes consisted in violation of the integrity of the cell walls of fungi and the release of cell contents as a result of the effect of ROS generated by Ce-MOF on the cells.

Iodine-doped carbon dots (IDCDs) were synthesized as peroxidase mimics for the antifungals against *Candida albicans*. The antifungal activity of the IDCDs was 90% in the presence of exogenous  $H_2O_2$  (0.5 mM) and under visible light irradiation for 2 h. The IDCDs promotes the formation of ROS, including hydroxyl radicals ( $\cdot OH$ ), which can successfully inhibit the growth of *C. albicans* [88]. This material imitated the action of peroxidase used for yeast disinfection. In relation to the enzyme the peroxidase-like action of this biomimetic material was more effective due to the presence of iodine in its content.

Despite the great interest in biomimetics that cause the death of fungal cells, there remains a general problem of their practical application, which is associated with the need for their toxicity to other eukaryotic cells [90]. The search for the most highly specific active substances with a minimum level of toxicity for non-fungal cells remains relevant.

### 3.2. Biomimetics with inhibition effect on fungal proteins/enzymes

To inhibit the enzymatic activity of fungi, synthetic substrates, playing role of mimetics, or specific inhibitors can be used to reduce the metabolic activity of fungi or to carry out their catalytic or physiological functions, reducing the biosynthesis of a number of metabolites and preventing their growth (Table 4) [15,91–98]. Biomimetic substrates should, on the one hand, inhibit the widest possible range of fungal enzymes, and on the other hand, not be inhibitors for enzymes of plant and animal origins.

**Table 4.** Biomimetics as antifungals with effect on fungal proteins/enzymes.

<b>Biomimetics as antifungals</b>	<b>Target of action</b>	<b>*Antifungal effect</b>
Iridoid alkaloids biomimetic of camptothecin [91]	Strong inhibitory effects against mycelial growth and spore germination. Disturbing the replication and transcription of DNA by binding to topoisomerase I, inhibiting of ergosterol biosynthesis	LC <sub>50</sub> (mg/L): <i>F. graminearum</i> (34.5); <i>Rhizoctonia solani</i> (18); <i>Botrytis cinerea</i> (26)
Sulfonyl hydrazide derivatives containing the 1,2,3,4-tetrahydroquinoline [92]	Inhibition of laccase activity	EC <sub>50</sub> (mg/L): <i>Sclerotinia sclerotiorum</i> (3.32), <i>Valsa mali</i> (2.78)
3-aryl-isoquinoline derivatives [93]	Inhibition of succinate dehydrogenase activity	EC <sub>50</sub> (mg/L): <i>Physalospora piricola</i> (3.7)
Hymexazol glycosides [15]	Inhibition of chitinase, produced by fungi	EC <sub>50</sub> (mg/L): <i>Alternaria alternata</i> (1.58)
N-1-(β-d-ribofuranosyl) benzimidazole derivatives [94]	Inhibition of fungal cytochrome P <sub>450</sub> 3A-dependent C14-α-demethylase which is responsible for the conversion of lanosterol to ergosterol and ergosterol biosynthesis	MIC (mg/L): <i>A. flavus</i> (0.8); <i>A. niger</i> (1.6); <i>F. oxysporum</i> (3.1); <i>C. albicans</i> (0.8)
pCF2Ser peptide as substrate mimetic [95]	Inhibition of Cdc14 phosphatases; formation of defective conidiation and ascospores, reducing cell virulence	Inhibitory constant (Ki) against fungal phosphatase Cdc14 homologs - 3-19 μM
Synthetic oxime-derivatives of resorcyate aminopyrazole [96]	Selective inhibition of chaperone Hsp90 functions	EC <sub>50</sub> (μM): <i>Candida neoformans</i> (0.040); <i>C. albicans</i> (0.011)
Dextran-coated Gd-based nanoparticles (NPs) as phosphatase-like nanozyme [97]	Selective hydrolysis of the terminal high-energy phosphate bonds in ATP	464 mg/L NPs increases ethanol yield per 17%. The characteristics of NPs: Km, Vmax, and Ea were 29.4 μM, 7.17 × 10 <sup>-7</sup> M/s, and 29.34 kJ/mol, respectively
Nanopillars of poly(methyl methacrylate) like cicada wing surface topography [98]	Superhydrophobic surfaces with reduced adsorption capability of proteins needed for adhesion of fungal spores	100% removal of 10 <sup>5</sup> spores of <i>Fusarium oxysporum</i> on the surface with antifungal activity

\*MIC - minimum inhibitory concentration; EC<sub>50</sub> - half maximal effective concentration.



New fungicides based on 4-chlorocinnamaldehyde theosemicarbazide (PMDD-5Y) and 1,2,3,4-tetrahydroquinoline inhibiting fungal laccases were obtained. Antifungal activity of the new compounds was confirmed against *Sclerotinia sclerotiorum* and *Valsa mali* cells [92].

Biomimetic synthesis of iridoid alkaloids resulted in obtaining of compounds with antifungal activity similar to camptothecin, which is a topoisomerase inhibitor, a natural quinoline alkaloid. These iridoid alkaloids disrupted DNA replication and transcription by binding to topoisomerase I, and then interrupted mycelial cell division and inhibited ergosterol biosynthesis [91].

Hymexazol (HIM) can inhibit spore germination of fungi by combining aluminum and iron ions or can interfere with the RNA or DNA biosynthesis of pathogenic fungi. Modification of HYM by amino sugars significantly stimulates plant growth, but causes morphological changes in the mycelium *Alternaria alternata*: plasma membranes are separated from cell walls, cytoplasm becomes vacuolized, and mitochondria, lysosomes and other organelles decrease in size in cells, the mechanical strength of cell walls is significantly reduced. Glycosylated HB suppresses the activity of chitin synthase G involved in chitin biosynthesis, which leads to these changes in cells and limits their growth [15].

Imidazole is a structural element of many important biological compounds: the essential amino acid histidine, purine bases in the structures of nucleic acids (DNA and RNA), and ATP. Biomimetics obtained as derivatives of imidazole-containing compounds have an inhibitory effect on many chemical reactions in living cells. Many of these compounds are widely distributed in cells and predominantly in a protein-bound state [99].

Such interactions lead to the inhibition of bound proteins and enzymes. Namely, inhibition of the ergosterol synthesis by direct binding to sterol 14 $\alpha$ -demethylase (CYP51) is the main function of the imidazole derivatives as antifungals. The membrane-binding enzyme CYP51 is localized in outer membrane of endoplasmic reticulum and catalyzes the removal of methyl group at the 14-th carbon atom [100]. Since recently there has been a decrease in the effectiveness of the inhibitory effect of antifungals based on azoles in relation to CYP51 due to mutational changes in the enzyme [101], then the synthesis of new derivatives continues.

Among the biomimetics obtained as imidazole derivatives, many compounds are known today that contain several halogen atoms (F or Cl) to enhance their antifungal effect [102]. For example, fluconazole and albaconazole contain two F-atoms, whereas itraconazole contains two Cl atoms.

Synthetic aminopyrazole substituted resorcylatamides have been studied as inhibitors of the functioning of the chaperone protein Hsp90 [96]. Hsp90s function supports protein quality control mechanisms, productive folding, and the stability of conformationally labile proteins, many involved in key signaling cascades. The chaperoning by Hsp90 of its so-called client proteins is ATP-dependent and coordinated by a suite of cochaperones and accessory factors that impart client selectivity and help regulate progression through the chaperoning cycle. Hsp90 isoform, based on semisynthetic oxime-derivatization of the resorcylate macrocycle natural products radicicol (1) and monocillin I (a substance synthesized by the fungi *Monocillium nordinii*, and other microorganisms). The modification made it possible to obtain selective inhibitors of fungal chaperones that have minimal negative effects on human chaperones.

To inhibit fungal phosphatase Cdc14, which is a key regulator of mitosis in yeast and fungal pathogens, a peptide substrate mimetic containing a non-hydrolyzable  $\alpha,\alpha$ -difluoromethylene phosphonoserine (pCF2Ser) residue instead of pSer was developed and investigated [95]. The inhibition constant (K<sub>i</sub>) for the new pCF2Ser peptide was determined for a number of Cdc14 homologues present in *Saccharomyces cerevisiae*, *Cercospora zeina*, *Magnaporthe oryzae*, *Rhizoctonia solani*, and *Puccinia striiformis* cells, which depended on the cell type and was in the range of 3-19  $\mu$ M. It was found that *Fusarium graminearum* and *Aspergillus flavus* fungus cells devoid of active Cdc14 did not form conidia and ascospores and could not infect plants [95].

Lanosterol 14 $\alpha$ -demethylase is a key enzyme in the biosynthesis of ergosterol in fungi. This enzyme is a target for action of new antifungals. New N-1-( $\beta$ -d-ribofuranosyl) benzimidazole derivatives were obtained and their fungicidal activity against several pathogenic fungal strains was evaluated (*Aspergillus flavus*, *A. niger*, *F. oxysporum* and *C. albicans*) [94]. The maximum inhibitory

efficacy was found against *A. flavus* and *C. albicans*, which turned out to be better than that known for the fungicide fluconazole against the same cells. It is interesting to note, that hydrophilicity of the resulting compound was the determining factor for enhanced antifungal activity, while hydrophobic compounds showed increased antibacterial activity.

Succinate dehydrogenase provides electrons for aerobic respiratory circuits in the mitochondria of eukaryotic fungal cells. New 3-aryl-isoquinoline derivatives based on sanguinarine (heleritrine and berberine) inhibiting this enzyme have been synthesized [93]. The fungicidal activity of the compounds obtained at a concentration of 50 mg/L was higher in comparison with the antimicrobial action of sanguinarine, which inhibits mycelial growth of *Alternaria solani*, *A. alternata* and *Phylospora piricola*. Molecular docking showed that the obtained compounds completely overlap the active center of succinate dehydrogenase, interacting with negatively charged amino acid residues of the chain A of this enzyme (Thr58, Ala61, Glu398, Leu415, Leu418, etc.), and thus inhibit the enzyme, significantly affecting the metabolism of fungal cells.

Nanoparticles of phosphatase-like nanozyme based on gadolinium covered by dextran was synthesized as mimetic of enzymatic catalysts to intensify the dephosphorylation of ATP, an important biomolecule reflecting the energy status of living cells [103].

The use of dextran made it possible to increase the biocompatibility and endocytosis of nanoparticles to yeast cells. The resulting samples selectively catalyzed the hydrolysis of terminal high-energy phosphate bonds in the ATP molecule, led to a decrease in the concentration of this important substance in the yeast cells of *S. cerevisiae* and increased the yield of ethanol during glucose fermentation [97].

In fact, there was a stimulation of the transition of cells from active growth under aerobic conditions to glycolysis under anaerobic conditions with a rapid decrease in growth rate. A decrease in the concentration of intracellular ATP and the accumulation of an increased concentration of ethanol in the medium with cells resulted in inhibition of cell metabolism.

It is known that *Aspergillus fumigatus* and *F. oxysporum* fungi form spores that are transported with air masses and adhere to various biological surfaces through protein-protein and protein-peptide interactions involving membrane-bound fungal proteins. Studies of fungicidal properties of biomimetic nanostructured surfaces made of poly(methyl methacrylate) with characteristics close to "cicada wing" in relation to these fungi have shown that such materials prevent the adhesion of spores and their germination with the subsequent development of fungal infections [98]. The same materials proved to be effective against the adhesion of ascospores of *S. cerevisiae* yeast cells.

Thus, such materials counteract the adhesion of fungal protein and prevent them from getting on the bio-surfaces.

### 3.3. Biomimetics of metabolites with effect on growth and metabolic activity of fungi

Various fungal metabolites become important targets of antifungal exposure [104]. The use of the created biomimetics is aimed at their competition with the substituted molecules, which affects the metabolic activity and growth of fungal cells [105]. The well-known metabolomics of fungi enable developing of approaches to the decrease in fungal resistance and adaptation to various stress conditions (ionizing radiation, pH change, hypoxic stress, solar ultraviolet radiation, presence of toxic compounds, nutrient stress, oxidative stress, cold-heat stress) (Table 5) [17,89,106–113].

**Table 5.** Biomimetics as antifungals with effect on the growth and metabolic activity of fungi.

<b>Biomimetics as antifungals</b>	<b>Target of action</b>	<b>*Antifungal effect</b>
Formyl phloroglucinol meroterpenoids [106]	Reducing of hyphae elongation and filamentation due to blocking of potential outflow of fungal substrates	MIC <sub>50</sub> (mg/L): <i>C. albicans</i> (8.7), <i>C. glabrata</i> (13.5)
4-fluorophenylalanine (FPA) [107]	Incorporation in proteins and inhibition of cell growth	Twofold decrease in growth rate of <i>Sacharomyces cerevisiae</i> by 500 mg/L FPA
Phe-Ala dipeptide polymer/polyoxometalate composite [89]	Deformation of conidial heads and the appearance of indistinguishable sterigmata; smooth cell walls of hyphae become completely depressed and destroyed; spores become wrinkled	MIC (mg/L): <i>A. niger</i> (230)
2-(2-hydroxypropyl) phenol [108]	Inhibition of respiration, causing decrease in ATP concentration and metabolic activity	EC <sub>50</sub> (μg/mL): <i>Rhizoctonia cerealis</i> (1.0); <i>Pythium aphanidermatum</i> (20.3); <i>V. mali</i> (14.9); <i>Botrytis cinerea</i> (23.5)
2- deoxyglucose [109; 110]	Violation of glycolysis and ATP biosynthesis	> 2 times increase in the doubling time of <i>S. cerevisiae</i> cells
L-pyroglutamic acid 4-chiral hydroxyl sulfonyl ester derivatives [17]	Inhibition of biosynthesis of trichothecenes	61.6% inhibition of <i>Fusarium graminearum</i> growth by 100 mg/L
Fraxinellone [111]	Changes in lipopolysaccharide-induced DNA binding activity and reduced translation	EC <sub>50</sub> (mg/L): <i>Alternaria longipes</i> (64.2); <i>Curvularia lunata</i> (123.3)
1-amino-1-(4-imidazole)methylphosphonic acid [112]	Inhibition of various enzymes, especially proteases	MIC (mg/L): <i>Rhodotorula mucilaginosa</i> (1024), <i>A. niger</i> (5000)
{[(2-hydroxy-4-nitrophenyl) amino] (thiophen-3-yl) methyl} phosphonic acid (5N3TPA) and {[(2-hydroxy-4-methylphenyl) amino] (thiophen-3-yl) methyl} phosphonic acid (5M3TPA) [113]	Inhibition of fungal enzymes	78.42% and 50% inhibition of <i>Fusarium oxysporum</i> and <i>Alternaria alternata</i> growth by 100 mg/L of 5N3TPA and 5M3TPA, respectively

\*MIC - minimum inhibitory concentration; EC<sub>50</sub> - half maximal effective concentration.

In recent decades, considerable attention has been paid to the development of antimicrobial and antitumor agents in the form of natural and non-canonical amino acids derivatives [114,115]. It has been established that such unnatural amino acids can be incorporated by yeast cells into their proteins, which leads to a noticeable inhibition of cell growth (Table 5) [107]. It was shown, that up to 35-40% of phenylalanine can be replaced by its fluorine-containing analogue in cell biomass of yeast *Saccharomyces cerevisiae*. The introduction of amino acids with halogen atoms to the media with

fungi results in biosynthesis of peptides and proteins with their electron-withdrawing features and non-natural steric-effects, because the fluorination influences the folding properties, changing the secondary structure propensity of peptides and protein segments containing aliphatic halogenated amino acids. In addition, halogen atoms have an impact on the proteolytic stability of the peptides [116]

One of the other approaches in the molecular design of antifungal active substances is associated with the introduction of nitrogen-containing heterocyclic fragments into the structure of compounds. Tetrazole cycle is one of the most promising in this context. NH-disubstituted tetrazoles are sufficiently strong acids, comparable in acidity to carboxylic acids and rather weak Brensted bases. High metabolic stability and the ability to overcome biological membranes ensure the successful use of the tetrazole cycle as a bioisoster of some functional groups in biologically active compounds. Tetrazolyl analogues of L-glutamic acid, L-tyrosine and L-ornithine as potential amino acid biomimetics have been synthesized and characterized.

Under environmental conditions, halogenated amino acids are synthesized by some fungi and required enzymes to introduce the halogen to protein or peptide structures within a post-translational modification. Non-proteinogenic amino acids may have diverse physiological functions and regulate functioning of metabolic cycles. The properties of halogenated AMPs, both native and synthetic, and their possible contribution to the growing problem of antibiotic resistance of microorganisms are currently investigated [116].

Formyl phloroglucinol meroterpenoids are characteristic metabolites of the plant genus *Eucalyptus* and exhibit fungicidal activity against various *Candida* species, including strains resistant to fluconazole. Terpenoids were found to be potential efflux pump substrates of fungi. Their fungicidal activity enhances by conjugating them with triphenyl phosphonium cations aimed at mitochondria. In this study, terpenoids were modified with phloroglucinol groups and it was shown that the compounds obtained are most active against *Candida albicans* and *C. glabrata* – 13.5) and are comparable in efficacy with fluconazole. In addition, the compounds obtained at concentrations of 25 and 100 mg/L suppressed the growth of yeast biofilms by 66% and 90%, respectively [106].

2-Allylphenol is a synthetic fungicide that mimics ginkgo from *Ginkgo biloba*, effectively used against various fungal pathogens (*V. mali*, *Botrytis cinerea*, *Drechslera turcica*, *Rhizoctonia ceramicis*, *Sphaerotheca aphansis*). The antifungal mechanism of action includes a decrease in ATP biosynthesis and inhibition of cell respiration. As a result of such an impact, protoplasm outflow occurs from hyphae, which after a few hours begin to collapse [108].

Natural limonoid derivatives with antifungal activity can be isolated from plant sources and used against pathogenic fungus *Cladosporium cucumerinum* to reduce the rate of fungal growth [117]. Fraxinellone is the most effective antifungal compound from this group [118], however, this compound and many other lemonoid derivatives can be obtained in a semy-synthetic or synthetic process [111,119]. The synthesis of potential antifungals representing biomimetics of natural limonoids bases on natural enzyme-catalyzed conversion of squalene to lanosterol with participation of oxidosqualene cyclase and squalene epoxidase (monooxidase). Lanosterol is a tetracyclic triterpenoid and is the compound from which all fungal steroids are derived.

Secondary metabolites of plants in the form of terpenoids, alkaloids and various organic acids can exhibit fungicidal properties. Thus, L-pyroglutamic acid isolated from the perennial herbaceous plant *Disporopsis aspersa* demonstrates antifungal effect against *Phytophthora infestans* and *Psilocybe cubensis*. L-pyroglutamic acid is an analog and potential precursor of glutamate, which is necessary for the biosynthesis of glutathione. L-pyroglutamic acid reduces the accumulation of trichothecenes in fungi of the genus *Fusarium*. Studies of the expression of genes that induce the biosynthesis of trichothecenes have shown that the production of mycotoxins is inhibited by L-pyroglutamic acid at the transcription level. A number of derivatives of L-pyroglutamic acid from L-hydroxyproline were synthesized. It turned out that the reaction of L-pyroglutamic acid with 4-chlorophenol to form an ester enhances its antifungal activity. Among the 31 synthesized compounds at a concentration of 100 mg/L, the maximum inhibition level (61.6%) of the fungal growth of *F. graminearum*, comparable to the commercial fungicide chlorothalonil, were derivatives obtained by the introduction of

naphthalene or alkane by hydroxyl group. The electron-withdrawing groups on the aromatic ring facilitated the inhibitory effect [17].

In studies of the fungal metabolism, structural analogs of substrates are used, which make it possible to "close" one of the possible metabolic pathways for the study of alternative pathways. Thus, as a result of studies of glycolysis and related processes in yeast cells, it was found that deoxyglucose (2-DG), imitating glucose, as well as mannose, changes not only the rate of glycolysis, but also many other processes (glycosylation of proteins, ATP production, regulation of signaling pathways), causing energy depletion of cells and generation of stress in them [109]. As a result, the rate of cell doubling is reduced by more than 2 times. Despite the fact that in the course of such studies it was found that yeast cells consuming 2-DG are able to mutate and develop resistance to this substrate [110], the use of 2-DG as an inhibitor of yeast metabolism can be further investigated under certain conditions, in particular, in combinations with other antifungals.

Phosphonates are compounds with diverse biological activity, including antimicrobial one [120,121]. The aminophosphonates, being structural analogues of natural amino acids, inhibit the activity of various enzymes, especially fungal proteases [112,121]. Due to this property, aminophosphonic acids containing heterocyclic functional groups are used as pharmaceutical preparations or pesticides. However, the presence of a heterocycle reduces the toxicity of these compounds compared to aromatic analogues [112]. Recently, it was revealed that a number of enzymes of different microorganisms are capable of degradation and transformation of similar compounds [120,122], since then further possible use of these compounds can also be considered in the presence of other antifungals in combinations with them to improve results.

Currently, various aminophosphonic acids containing a thiophene ring are actively synthesized, which exhibit antifungal activity against fungi and yeast cells [112,113]. The presence of active functional groups in the aromatic ring, P=O and P-OH groups of  $\alpha$ -aminophosphonates and the thiophene ring enhances the antifungal activity of these compounds. Molecular docking of newly synthesized aminophosphonic acids confirmed the possibility of notable inhibition of proteases.

Like other ascomycete fungi, *F. oxysporum* secretes an  $\alpha$ -pheromone, a small peptide that functions as a chemoattractant and as a QS molecule. Three of the 10 amino acid residues of the  $\alpha$ -pheromone are tryptophan residues, composed a side chain with a high affinity for lipid bilayers. The use of lipid mimetics based on phospholipids (analogues of the cell membrane bilayer) allows the binding of this QS molecule and the formation of an intramolecular disulfide bond between two cysteine residues during the interaction. It should be noted that the oxidized version of the  $\alpha$ -pheromone had no biological activity as a chemoattractant and a QS molecule [123].

This effect leads to a weakening of the QS, which plays an important role in the functioning of fungal and yeast populations [124] and ensuring a more effective action of the antimicrobial agents used simultaneously.

Thus, the processes that make up the metabolism of filamentous fungi and yeast, and the study of fungal metabolomics can serve as a source of new ideas for creating effective antifungals. Substances regulating the metabolic activity of microbial cells, including end products of their metabolism, can significantly affect the survival and energetic status of fungal cells. The largest number of antifungals being developed in this area relate to inhibitors of one or another stage of cell metabolism.

#### 4. Combination of antifungal peptides with each other and/or with antifungal drugs

Despite expectations of the absence of resistance formation in fungi to the effects of new antifungals, including AFPs [125,126], azoles [127] and 2-DG [109], researchers note such cases. One of the effective solutions to this problem today is the combined use of various antifungals. Such possible combinations can include various traditional antifungal agents and AFPs (Table 6) [18,37,45,57,67,128–140]. Such combinations make it possible to joint different mechanisms (targets) of the effects of antifungals and significantly decrease their toxicity by improving the effectiveness of the action and reducing the doses applied in comparison with individual substances.



Table 6. Combination of AFPs with different antifungal agents.

Components of combined antifungals		Target [Reference]	Antifungal effect
AFP as the 1 <sup>st</sup> component	2 <sup>nd</sup> component		
Mo-CBP <sub>3</sub> -PepI	Nystatin	<i>C. albicans</i> [57]	0.13 <sup>a</sup>
Mo-CBP <sub>3</sub> -PepII			
Mo-CBP <sub>3</sub> -PepIII	Itraconazole	<i>C. parapsilosis</i> [67]	82 % <sup>b</sup>
RcAlb-PepIII		<i>C. neoformans</i> [128]	96 % <sup>b</sup>
MSI-78	Voriconazole	<i>Fusarium solani</i> [37]	84.1 % <sup>b</sup>
hLf(1-11)			0.34 <sup>a</sup>
Cecropin B			0.21 <sup>a</sup>
MSI-78			0.17 <sup>a</sup>
hLf(1-11)	Amphotericin B		0.37 <sup>a</sup>
Cecropin B			0.31 <sup>a</sup>
Brilacidin (non-peptide mimetic of host defense peptides)	Caspofungin	<i>Aspergillus fumigatus</i> [45]	0.28 <sup>a</sup>
	Voriconazole		1.0 <sup>a</sup>
	Geldanamycin		0.64 <sup>a</sup>
Neosartorya fischeri AFPs (NFAP)	NFAP2	<i>Botrytis cinerea</i> ,	1.25 <sup>a</sup>
NFAP	$\gamma^{\text{NFAP-opt}}$	<i>Cladosporium herbarum</i>	0.28 - 1.50 <sup>a</sup>
NFAP2	$\gamma^{\text{NFAP-opt}}$	[129]	0.31-1.5 <sup>a</sup>
P256 and P256	Amphotericin B	<i>C. albicans</i> [130]	0.28 <sup>a</sup>
Fengycin	Surfactin	<i>Rhizopus solonifer</i> [131]	5 <sup>a</sup>
ToAP2	NDBP-5.7	<i>C. albicans</i> [132]	0.75 <sup>a</sup>
ToAP2			0.18 <sup>a</sup>
NDBP-5.7	Amphotericin B	<i>C. albicans</i> ; [133]	0.18 <sup>a</sup>
gH625M			0.5-0.8 <sup>a</sup>
L-His(2-adamantyl)-L-Trp-L-His(2-phenyl)-OMe		<i>C. neoformans</i> [19]	0.28 <sup>a</sup>
L-Trp-L-His(1-biphenyl)-NHBzl			0.28 <sup>a</sup>
	Fluconazole	<i>C. neoformans</i> [134]	1.04 <sup>a</sup>
L-His[1-(4-n-butylphenyl)]-L-Trp-L-His[1-(4-n-butylphenyl)]-NHBzl	Amphotericin B		0.31 <sup>a</sup>
			0.75 <sup>a</sup>
DP-23 peptoid	Fluconazole	<i>A. flavus</i> , <i>A. niger</i> [18]	0.16-0.38 <sup>a</sup>
SPO peptoid		<i>C. albicans</i> [135]	
$\gamma$ -AA peptide MW5			$\leq 0.5$ <sup>a</sup>
ToAP2		<i>C. albicans</i> [132]	0.5 <sup>a</sup>
NDBP-5.7			0.56 <sup>a</sup>
KW-23		<i>C. albicans</i> [136]	0.37-0.60 <sup>a</sup>
HLopt2 (mimic of human lactoferrin)		<i>Candida</i> spp. [137]	2-125 mg/L <sup>c</sup>
gH625M	Flucytosine	<i>C. albicans</i> [133]	0.30 <sup>a</sup>
			0.20 <sup>a</sup>
14-helical $\beta$ -peptide	Isoamyl alcohol	<i>C. albicans</i> [138]	4 mg/L <sup>d</sup>
Lactofungin	Amphotericin B	<i>C. albicans</i> , <i>C. glabrata</i> , <i>C. neoformans</i> , <i>C. deuterogattii</i> [139]	0.16-0.28 <sup>a</sup>
TP10-NH <sub>2</sub> (analog of transporan 10)	Ciprofloxacin or Levofloxacin	<i>Candida</i> spp. [140]	6.3-100 $\mu\text{M}$ <sup>c</sup>

<sup>a</sup>FICI: Fractional inhibitory concentration index. FICI values suggest antagonistic (>4.0), indifferent (>1 to <4), additive (>0.5–1) and synergistic ( $\leq 0.5$ ) effect; <sup>b</sup>Percent of biofilm and growth inhibition; <sup>c</sup>MIC/MIC<sub>50</sub> — minimum inhibitory concentration (complete inhibition/at least 50% inhibition); <sup>d</sup>Minimal biofilm prevention concentrations (MBPC).

The synergistic effect of combined antifungals makes it possible to effectively combat yeast biofilms of the genus *Candida* both at the stage of their formation and already for the degradation of mature biofilms. Combination of AFPs Mo-CBP<sub>3</sub>-PepI and Mo-CBP<sub>3</sub>-PepIII with nystatin and itraconazole against *Candida* species biofilms resulted in a 2 to 4-fold improvement of antibiofilm activity of antifungals [57,67]. The combination of Mo-CBP<sub>3</sub>-PepIII and RcAlb-PepIII synthetic peptides with itraconazole enhanced the activity of the latter by 10-fold against *C. neoformans* [128]. Likewise, synthetic histidine-containing amphipathic peptides enhanced the activity of amphotericin B against *Cryptococcus neoformans* by 4 to 16-fold [134]. When combined with lactoferrin-derived synthetic peptide lactofungin, the minimum inhibitory concentration of amphotericin B for *Candida* spp. and *C. neoformans* decreased 4 times [139].

The antifungal activity of two *N. fischeri* peptides NFAP and NFAP2 and their  $\gamma$ -core peptide derivatives ( $\gamma$ NFAP-opt,  $\gamma$ NFAP2-opt), was tested in vitro against *Botrytis*, *Cladosporium*, and *Fusarium* spp. Synergistic mechanism of action was observed when NFAP or NFAP2 was applied in combination with  $\gamma$ NFAP-opt. The investigated proteins and peptides did not show any toxicity to tomato plant leaves, except for  $\gamma$ NFAP2-opt [129].

*Fusarium* infections have been associated with high mortality rates due to the lack of effective treatment strategy. The in vitro activity of AFPs MSI-78, h-Lf1-11 and cecropin B combined with amphotericin B or voriconazole were tested against ten *Fusarium solani* strains. All AFPs demonstrated synergistic mechanism of action when combined with conventional antifungals [37].

The combination of brilacidin (synthetic, nonpeptidic, small molecule mimetic of defensin) with caspofungin has a synergism that is able to affect *A. fumigatus* viability through multiple mechanisms of action, encompassing functional changes/depolarization of the microorganism cell membrane, interference to calcineurin signaling, and misexpression of the cell wall integrity pathway, and prevents  $\beta$ -1,3-glucan biosynthesis [45].

The use of mimetic peptides DP-23 (short lipopeptide) and SPO (N-substituted polyglycine) in combination with fluconazole provided a synergistic effect on *A. niger* and *A. flavus* cells [18].

Peptides P255 and P256 obtained from hexapeptide PDF 26 were combined with amphotericin B and investigated against *C. albicans*. Both peptides showed a synergistic effect with a polyene antibiotic, while P256 showed stronger antifungal effect than P255. Peptides violated the integrity of the cell wall, increased membrane permeability, disrupted cell morphology and caused intracellular changes: they affected the expression of genes for replication and repair of fungal DNA, biosynthesis of cell wall components and ergosterol. They also increased the production of ROS in cells and bound to the genomic DNA of fungi [130].

$\gamma$ -AA peptides are a class of peptidomimetics with a definite folded structure and resistance to proteolytic hydrolysis. It has been shown that lipo- $\gamma$ -AA peptide MW5 significantly increases the efficacy of fluconazole action against azole-resistant *C. albicans* CARG5 cells. The peptide destroys the cell membrane and provokes the production of ROS [135].

Synthetic peptidomimetics of antimicrobial  $\alpha$ -peptides exhibit fungicidal activity as a result of hydrophobic and electrostatic interactions with cell membranes, which lead to permeabilization and subsequent cell death. These peptidomimetics were more stable than their natural counterparts and are not degraded by proteases.  $\beta$ -peptides were obtained structurally constructed on the basis of natural  $\alpha$ -helical AMPs. These mimetics possessed a fungicidal effect against *C. albicans* cells and inhibited the formation of yeast biofilms. The hydrophobicity of the  $\beta$ -peptides directly correlates with their antifungal properties and a narrow range of concentrations at which  $\beta$ -peptides effectively kill *C. albicans* cells without lysis of erythrocytes. The combination of these peptides with isoamyl alcohol reduced their MIC for inhibiting biofilms by 4 times [138].

Conjugates of fluoroquinolone antibiotics (ciprofloxacin or levofloxacin) with the TP10-NH<sub>2</sub> peptide penetrating into yeast cells showed antifungal effect against different strains of genus *Candida*. At the same time, the synergistic effect of these conjugates was not revealed in experiments *C. glabrata* cells [140].

Antifungal effect against *Candida* spp. cells was investigated for conjugates obtained on the basis of a modified fragment of lactoferrin HLOpt2 and ciprofloxacin, levofloxacin and fluconazole. Three

different nutrient media were used for these purposes. There was no activity against four different *Candida* species for the substances studied on the RPMI (mimics physiological conditions) medium. The use of BP (1% peptone) and BHI/100 (0.034% brain-heart infusion) media resulted in inhibition of cell growth, especially *C. tropicalis*, whereas *C. glabrata* cells were the most resistant among all tested strains [137].

The antifungal activity of cationic antimicrobial peptides ToAP2, from a cDNA library of the scorpion *Tityus obscurus* venom gland and NDBP-5.7, from a cDNA library of the scorpion *Opisthacanthus cayaporum* venom gland was demonstrated against *C. albicans* cells. Both peptides affected membrane permeability and caused such changes in the morphology of fungal cells as cell wall deformations and disruption of ultrastructural cell organization. Both peptides showed synergism with amphotericin B, and demonstrated synergistic and additive effect in combination with fluconazole Top 2 and NBSP-5.7, respectively [132].

Unique ultra-short peptide KW23 had a positive effect on standard and resistant *Candida* species, showing powerful synergistic antimicrobial activity in combination with fluconazole. Interesting, that effect of this combination was additive with respect to the resistant strain. In addition, the peptide showed a low enough toxicity to human erythrocytes [136].

Persistent cells that can tolerate lethal concentrations of antimicrobials and grow again after their removal, being found in different microbial populations. It is assumed that the appearance of persistent yeast cells involves different regulations of genes correlating with the pathways of ergosterol biosynthesis (ERG1 and ERG25) and  $\beta$ -1,6-glucan (SKN1 and KRE1). Special studies have shown that membranes of cells in biofilms contain a lower concentration of ergosterol, mainly in the deepest layers of the biofilm, in comparison with planktonic yeast cells. It is likely that cells from mature biofilms require less ergosterol to maintain membrane fluidity, and this is also confirmed by the limited effectiveness of biomimetics aimed at inhibiting ergosterol biosynthesis. In this regard, the possibility of destroying resistant persistent-derived biofilms of *C. albicans* with gH625-M peptide, which is an analogue of viral membranotropic peptide gH625, was investigated [133]. The combination of gH625-M with various antifungals (fluconazole, 5-flucytosine, amphotericin B) demonstrated a synergistic effect when acting on persistent cells in biofilms at relatively low doses of the antifungals (Table 6).

Among antifungals that can be used in effective combination with AFPs, metal-containing compounds and enzymes are discussed [141,142]. The main mechanism of the antifungal action of metals is the triggering of generation and accumulation of ROS. Enzymes are of special interest due to their wide range of substrate specificity of action and various mechanisms of antifungal actions.

Combination of metalloenzyme such as hexahistidine-containing organophosphorus hydrolase (His<sub>6</sub>-OPH), capable of hydrolyzing signal molecules of yeast QS, with AFPs (polymyxins, bacitracin, Lfcin) resulted in a notable improvement of antimicrobial efficiency of action of the AFPs (up to 8.5 times) against different yeast species, such as *Saccharomyces cerevisiae*, *Candida* sp., *Trichosporon beigeei*, etc. It is interesting, that His<sub>6</sub>-OPH had an increased catalytic efficiency of action in the hydrolysis of its substrates (QS molecules) when it was introduced in combination with AFPs [143,144]. Thus, combination of the AFPs with enzymes possessing antifungal activity is a perspective trend in the development of efficient antifungals.

To determine the effect of combining two antifungals and interpreting the results obtained, a fractional inhibitory concentration index (FICI) range (from 0.5 to 4) is usually applied. The synergistic effect corresponds to FICI values < 0.5 and is achieved through a combination of different mechanisms of action on antifungals. From the studies analyzed in this section, it follows that only in the case of a combination of fengycin with surfactin, an antagonistic effect was revealed (FICI = 4), which is an indicator of the incompatibility of the mechanisms of action of these compounds with each other [131].

In the case of a combination of brilacidin with conventional fungicides, AFPs of *Neosartorya fischeri*, among themselves and histidine-containing amphipathic peptides with fluconazole, an additive effect (FICI > 1) was marked, which means that these antifungals do not enhance the effectiveness of each other action [45,129,134]. In the case of all other analyzed combinations, a

synergistic effect was revealed. The most effective variants were obtained in the case of combinations of synthetic peptides Mo-CBP3-PepI and Mo-CBP3-PepII with nystatin (FICI = 0.13) [57,67]. It is interesting to note that when combining two synthetic peptides, MSI-78 and hLf(1-11), with a natural peptide, Cecropin B, as well as with voriconazole or with amphotericin B, the strongest synergistic effect was achieved with a natural peptide, but not with considered successful antifungals like azole and polyene antibiotic [37].

When developing biomimetics and some combinations with them, special attention is paid to minimizing their toxicity (or its complete absence) in relation to human/animal/plant cells, as well as to the type of antifungal effect (mechanism of action) on cells in order to avoid the development of resistance [76,145]. At the initial stage of such investigations, toxicity is assessed using modern computer modeling methods [146]. Erythrocytes are used as experimental models in vitro and in vivo [147].

Taking into account the fact that a minimal number of cases of fungal resistance have been detected with respect to SAFP, as well as the minimum level of toxicity shown in these peptides for other cells, they seem promising candidates in the development of modern antifungals.

## 5. Conclusions

Analysis of the data in Tables 1 and 2 showed that the molecular weight of SAFP is often comparable to the molecular weight of natural AFP, it may be more/less than it, however not it, but the structure of the substances affects their antifungal properties. Most natural phases turn out to be effective at slightly higher concentrations compared to SAFP.

Discussing the concentration ranges recommended for the use of biomimetics, it should be noted that it is quite difficult to compare this indicator for different groups of drugs, since in some studies different indicators of their effectiveness are used (MIC, LC<sub>50</sub>, EC<sub>50</sub>, and up (Tables 1-6), presented in different units of measurement. Nevertheless, trying to assess the concentrations of different biomimetics necessary for their antifungal effect, we consider that it is possible to conditionally arrange the groups of preparations in the following row with increasing compared parameter: Synthetic / (semi)synthetic AMPs with antifungal activity < Biomimetics as antifungals with effect on fungal membranes and cell walls < Natural AMPs with antifungal activity < Biomimetics as antifungals with effect on fungal proteins/enzymes < Biomimetics as antifungals with effect on the growth and metabolic activity of fungi. To estimate the effectiveness of the newly developed antifungals, widespread test cultures are mainly used, depending on the intended application, for example, yeast of the genus *Candida*, mycelial fungi of the genus *Aspergillus* or *Fusarium* (Tables 1-6).

In the perspective, new SAFP are expected to appear with a wider range of antifungal effects, as well as peptides that can have a targeted and selective effect on the cells of potentially dangerous fungal cultures for humans and animals, as well as on fungal phytopathogens such, for example, as representatives of the genera *Acremonium*, *Absidia*, *Paecilomyces*, *Scopulariopsis*. A great potential of specifically acting antifungal agents can be disclosed through proteomic studies of fungal pathogens and the search for key proteins and enzymes, the inhibition of which can give a maximum and rapid effect without the formation of fungal resistance [148]. The creation of combinations of peptides and new antifungal agents may soon yield significant results in the impact not only on resistant strains, but also on heterogeneous biofilms [133,136]. Moreover, combinations can reduce the toxicity of individual components in relation to human cells [136,137].

Already developed combinations of SAFP with other antifungal agents, as it turned out, can effectively affect resistant strains of fungi even those which are in the state of QS (biofilms). Such combined antifungals seem to be a worthy solution to one of the important health and food safety problems associated with the resistance of fungi to antimicrobials used.

**Author Contributions:** Conceptualization, E.E.; investigation, E.E., A.A., N.S., O.S. and O.M.; data curation E.E. and A.A.; writing—original draft preparation, A.A., N.S., O.S., O.M. and E.E.; writing—review and editing, A.A. and E.E.; supervision, E.E. All authors have read and agreed to the published version of the manuscript.

**Funding:** The research was funded by Russian Science Foundation (23-14-00092).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Kainz, K.; Bauer, M.A.; Madeo, F.; Carmona-Gutierrez, D. Fungal infections in humans: the silent crisis. *Microb. Cell* **2020**, *7*, 143. doi: 10.15698/mic2020.06.718
2. Seyedmousavi, S.; Bosco, S.D.M.; De Hoog, S.; Ebel, F.; Elad, D.; Gomes, R.R.; Jacobsen, I.D.; Jensn, H.E.; Martel, A.; Mignon, B.; Pasmans, F.; Pieckova, E.; Rodrigues, A.S.; Singh, K.; Vicente, V.A.; Wibbelt, G.; Wiederhold, N.P.; Guillot, J. Fungal infections in animals: a patchwork of different situations. *Med. Mycol.* **2018**, *56*, S165-S187. doi:10.1093/mmy/myx104
3. Doehlemann, G.; Ökmen, B.; Zhu, W.; Sharon, A. Plant pathogenic fungi. *Microbiol. Spectrum* **2017**, *5*, 1-23. doi:microbiolspec.funk-0023-20
4. Ons, L.; Bylemans, D.; Thevissen, K.; Cammue, B.P. Combining biocontrol agents with chemical fungicides for integrated plant fungal disease control. *Microorganisms*, **2020**, *8*, 1930. doi:10.3390/microorganisms8121930
5. Iqbal, S.Z. Mycotoxins in food, recent development in food analysis and future challenges; a review. *Curr. Opin. Food Sci.* **2021**, *42*, 237-247. doi:10.1016/j.cofs.2021.07.003
6. Jenks, J.D.; Cornely, O.A.; Chen, S.C.A.; Thompson III, G.R.; Hoenigl, M. Breakthrough invasive fungal infections: Who is at risk? *Mycoses* **2020**, *63*, 1021-1032. doi:10.1111/myc.13148
7. Gintjee, T.J.; Donnelly, M.A.; Thompson, G.R. Aspiring antifungals: review of current antifungal pipeline developments. *J. Fungi* **2020**, *6*, 28. doi:10.3390/jof6010028
8. Campoy, S.; Adrio, J.L. Antifungals. *Biochem. Pharmacol.* **2017**, *133*, 86-96. doi:10.1016/j.bcp.2016.11.019
9. Houšť, J.; Spižek, J.; Havlíček, V. Antifungal drugs. *Metabolites* **2020**, *10*, 106. doi:10.3390/metabo10030106
10. El-Gendi, H.; Saleh, A.K.; Badierah, R.; Redwan, E. M.; El-Maradny, Y.A.; El-Fakharany, E.M. A comprehensive insight into fungal enzymes: Structure, classification, and their role in mankind's challenges. *J. Fungi* **2021**, *8*, 23. doi: 10.3390/jof8010023
11. Duarte, A.W.F.; Barato, M.B.; Nobre, F.S.; Polezel, D.A.; de Oliveira, T.S.; dos Santos, J.A.; Rodrigues, A.; Sette, L.D. Production of cold-adapted enzymes by filamentous fungi from King George Island, Antarctica. *Polar Biol.* **2018**, *41*, 2511-2521. doi:10.1007/s00300-018-2387-1
12. Geddes-McAlister, J.; Shapiro, R.S. New pathogens, new tricks: emerging, drug-resistant fungal pathogens and future prospects for antifungal therapeutics. *Ann. N. Y. Acad. Sci.* **2019**, *1435*, 57-78. doi:10.1111/nyas.13739
13. Tian, X.; Ding, H.; Ke, W.; Wang, L. Quorum sensing in fungal species. *Annu. Rev. Microbiol.* **2021**, *75*, 449-469. doi:10.1146/annurev-micro-060321-045510
14. Fernández de Ullivarri, M.; Arbulu, S.; Garcia-Gutierrez, E.; Cotter, P.D. Antifungal peptides as therapeutic agents. *Front. Cell. Infect. Microbiol.* **2020**, *10*, 105. doi:10.3389/fcimb.2020.00105
15. Gao, K.; Qin, Y.; Wang, L.; Li, X.; Liu, S.; Xing, R.; Yu, H.; Chen, X.; Li, P. Design, synthesis, and antifungal activities of hymexazol glycosides based on a biomimetic strategy. *J. Agric. Food. Chem.* **2022**, *70*, 9520-9535. doi:10.1021/acs.jafc.2c02507
16. Linklater, D.P.; Le, P.H.; Aburto-Medina, A.; Crawford, R.J.; Maclaughlin, S.; Juodkazis, S.; Ivanova, E.P. Biomimetic nanopillar silicon surfaces rupture fungal spores. *Int. J. Mol. Sci.* **2023**, *24*, 1298. doi:10.3390/ijms24021298
17. Ai, L.; Fu, S.; Li, Y.; Zuo, M.; Huang, W.; Huang, J.; Jin, Z.; Chen, Y. Natural products-based: synthesis and antifungal activity evaluation of novel L-pyrroglutamic acid analogues. *Front. Plant Sci.* **2022**, *13*, 1102411. doi:10.3389/fpls.2022.1102411
18. Sharma, L.; Bisht, G.S. Synergistic effects of short peptides and antibiotics against bacterial and fungal strains. *J. Pept. Sci.* **2023**, *29*, e3446. doi:10.1002/psc.3446
19. Aaghaz, S.; Sharma, K.; Maurya, I. K.; Rudramurthy, S.M.; Singh, S.; Kumar, V.; Tikoo, K.; Jain, R. Anticryptococcal activity and mechanistic studies of short amphipathic peptides. *Arch. Pharm.* **2023**, *356*, 2200576. doi:10.1002/ardp.202200576
20. Konakbayeva, D.; Karlsson, A.J. Strategies and opportunities for engineering antifungal peptides for therapeutic applications. *Curr Opin Biotechnol.* **2023**, *81*, 102926. doi:10.1016/j.copbio.2023.102926



21. Ramazi, S.; Mohammadi, N.; Allahverdi, A.; Khalili, E.; Abdolmaleki, P. A review on antimicrobial peptides databases and the computational tools. *Database* **2022**, 2022, baac011. doi:10.1093/database/baac011
22. Wang, G.; Li, X.; Wang, Z. APD3: The antimicrobial peptide database as a tool for research and education. *Nucleic Acids Res.* **2016**, *44*, D1087–D1093. doi:10.1093/nar/gkv1278
23. Waghu, F.H.; Barai, R.S.; Gurung, P.; Idicula-Thomas, S. CAMPR3: a database on sequences, structures and signatures of antimicrobial peptides. *Nucleic Acids Res.* **2016**, *44*, D1094–D1097. doi:10.1093/nar/gkv1051
24. Pirtskhalava, M.; Armstrong, A.A.; Grigolava, M.; Chubinidze, M.; Alimbarashvili, E.; Vishnepolsky, B.; Gabrielian, A.; Rosenthal, A.; Hurt, D.E.; Tartakovsky, M. DBAASP v3: Database of antimicrobial/cytotoxic activity and structure of peptides as a resource for development of new therapeutics. *Nucleic Acids Res.* **2021**, *49*, D288–D297. doi:10.1093/nar/gkaa991
25. Jhong, J.H.; Yao, L.; Pang, Y.; Li, Z.; Chung, C.R.; Wang, R.; Li, S.; Li, W.; Luo, M.; Ma, R.; et al. dbAMP 2.0: Updated resource for antimicrobial peptides with an enhanced scanning method for genomic and proteomic data. *Nucleic Acids Res.* **2022**, *50*, D460–D470. doi:10.1093/nar/gkab1080
26. Bentz, M.L.; Nunnally, N.; Lockhart, S.R.; Sexton, D.J.; Berkow, E.L. Antifungal activity of nikkomycin Z against *Candida auris*. *J. Antimicrob. Chemother.* **2021**, *76*, 1495–1497. doi:10.1093/jac/dkab052
27. Park, Y.; Lee, D.G.; Hahm, K.S. HP (2–9)-magainin 2 (1–12), a synthetic hybrid peptide, exerts its antifungal effect on *Candida albicans* by damaging the plasma membrane. *J. Pept. Sci.* **2004**, *10*, 204–209. doi:10.1002/psc.489
28. Alan, A.R. Sensitivity of bacterial and fungal plant pathogens to the lytic peptides, MSI-99, magainin II, and cecropin B. *Mol. Plant-Microbe Interact.* **2002**, *15*, 701–708. doi:10.1094/MPMI.2002.15.7.701
29. Kočendová, J.; Vaňková, E.; Volejníková, A.; Nešuta, O.; Buděšínský, M.; Socha, O.; Hájek, M.; Hadravová, R.; Čeřovský, V. Antifungal activity of analogues of antimicrobial peptides isolated from bee venoms against vulvovaginal *Candida* spp. *FEMS Yeast Res.* **2019**, *19*, foz013. doi:10.1093/femsyr/foz013
30. Jang, W.S.; Kim, K.N.; Lee, Y.S.; Nam, M.H.; Lee, I.H. Halocidin: a new antimicrobial peptide from hemocytes of the solitary tunicate, *Halocynthia aurantium*. *FEBS Lett.* **2002**, *521*, 81–86. doi: 10.1016/s0014-5793(02)02827-2
31. Yousfi, H.; Ranque, S.; Rolain, J.M.; Bittar, F. In vitro polymyxin activity against clinical multidrug-resistant fungi. *Antimicrob. Resist. Infect. Control* **2019**, *8*, 1–10. doi:10.1186/s13756-019-0521-7
32. Tonk, M.; Cabezas-Cruz, A.; Valdés, J.J.; Rego, R.O.; Grubhoffer, L.; Estrada-Peña, A.; Vilcinskis, A.; Kotsyfakis, M.; Rahnamaeian, M. Ixodes ricinus defensins attack distantly-related pathogens. *Dev. Comp. Immunol.* **2015**, *53*, 358–365. doi:10.1016/j.dci.2015.08.001
33. Lee, D.G.; Kim, H.K.; Am Kim, S.; Park, Y.; Park, S.C.; Jang, S.H.; Hahm, K.S. Fungicidal effect of indolicidin and its interaction with phospholipid membranes. *Biochem. Biophys. Res. Commun.* **2003**, *305*, 305–310. doi:10.1016/S0006-291X(03)00755-1
34. Fernandes, K.E.; Carter, D.A. The antifungal activity of lactoferrin and its derived peptides: mechanisms of action and synergy with drugs against fungal pathogens. *Front. Microbiol.* **2017**, *8*, 2. doi:10.3389/fmicb.2017.00002
35. Heymich, M.-L.; Nißl, L.; Hahn, D.; Noll, M.; Pischetsrieder, M. Antioxidative, antifungal and additive activity of the antimicrobial peptides Leg1 and Leg2 from Chickpea. *Foods* **2021**, *10*, 585. doi:10.3390/foods10030585
36. Rather, I.A.; Sabir, J.S.M.; Asseri, A.H.; Ali, S. Antifungal activity of human cathelicidin LL-37, a membrane disrupting peptide, by triggering oxidative stress and cell cycle arrest in *Candida auris*. *J. Fungi* **2022**, *8*, 204. doi:10.3390/jof8020204
37. Denardi, L.B.; Weiblen, C.; Ianiski, L.B.; Stibbe, P.C.; Santurio, J.M. Activity of MSI-78, h-Lf1-11 and cecropin B antimicrobial peptides alone and in combination with voriconazole and amphotericin B against clinical isolates of *Fusarium solani*. *J. Med. Mycol.* **2021**, *31*, 101119. doi:10.1016/j.mycmed.2021.101119
38. de Freitas, C.D.T.; de Souza Lopes, J.L.; Beltramini, L.M.; de Oliveira, R.S.B.; Oliveira, J.T.A.; Ramos, M.V. Osmotin from *Calotropis procera latex*: new insights into structure and antifungal properties. *Biochim. Biophys. Acta Biomembr.* **2011**, *1808*, 2501–2507. doi:10.1016/j.bbamem.2011.07.014
39. Landon, C.; Meudal, H.; Boulanger, N.; Bulet, P.; Vovelle, F. Solution structures of stomoxyn and spinigerin, two insect antimicrobial peptides with an  $\alpha$ -helical conformation. *Biopolymers* **2006**, *81*, 92–103. doi:10.1002/bip.20370
40. Kakar, A.; Holzknecht, J.; Dubrac, S.; Gelmi, M.L.; Romanelli, A.; Marx, F. New perspectives in the antimicrobial activity of the amphibian temporin B: peptide analogs are effective inhibitors of *Candida albicans* growth. *J. Fungi* **2021**, *7*, 457. doi:10.3390/jof7060457
41. D'Auria, F.D.; Casciaro, B.; De Angelis, M.; Marcocci, M.E.; Palamara, A.T.; Nencioni, L.; Mangoni, M.L. Antifungal activity of the frog skin peptide temporin G and its effect on *Candida albicans* virulence factors. *Int. J. Mol. Sci.* **2022**, *23*, 6345. doi:10.3390/ijms23116345
42. Mroczńska, M.; Brillowska-Dąbrowska, A. Review on current status of echinocandins use. *Antibiotics* **2020**, *9*, 227. doi:10.3390/antibiotics9050227

43. Antachopoulos, C.; Meletiadiis, J.; Sein, T.; Roilides, E.; Walsh, T.J. Comparative in vitro pharmacodynamics of caspofungin, micafungin, and anidulafungin against germinated and nongerminated *Aspergillus* conidia. *Antimicrob. Agents Chemother.* **2008**, *52*, 321-328. doi:10.1128/AAC.00699-07
44. Kovács, R.; Tóth, Z.; Locke, J.B.; Forgács, L.; Kardos, G.; Nagy, F.; Borman, A.M.; Majoros, L. Comparison of in vitro killing activity of rezafungin, anidulafungin, caspofungin, and micafungin against four *Candida auris* clades in RPMI-1640 in the absence and presence of human serum. *Microorganisms* **2021**, *9*, 863. doi:10.3390/microorganisms9040863
45. Dos Reis, T.F.; de Castro, P.A.; Bastos, R.W.; Pinzan, C.F.; Souza, P.F.; Ackloo, S.; Hossain, M.A.; Drewry, D.H.; Alkhazraji, S.; Ibrahim, A.S.; Jo, H.; Lightfoot, J.D.; Adams, E.M.; Fuller, K.K.; degrade, W.F.; Goldman, G. H. (2023). A host defense peptide mimetic, brilacidin, potentiates caspofungin antifungal activity against human pathogenic fungi. *Nat. Commun.* **2023**, *14*, 2052. doi:10.1038/s41467-023-37573-y
46. Aguiar, T.K.B.; Neto, N.A.S.; Freitas, C.D.T.; Silva, A.F.B.; Bezerra, L.P.; Malveira, E.A.; Branco, L.A.C.; Mesquita, F.P.; Goldman, G.H.; Alencar, L.M.R.; Oliveira, J.T.A.; Santos-Oliveira, R.; Souza, P.F.N. Antifungal potential of synthetic peptides against *Cryptococcus neoformans*: mechanism of action studies reveal synthetic peptides induce membrane-pore formation, DNA degradation, and apoptosis. *Pharmaceutics* **2022**, *14*, 1678. doi:10.3390/pharmaceutics14081678
47. Dias, L.P.; Souza, P.F.; Oliveira, J.T.; Vasconcelos, I.M.; Araújo, N.M.; Tilburg, M.F.; Guedes, M.I.F.; Carneiro, R.F.; Lopes, J.L.S.; Sousa, D.O. RcAlb-PepII, a synthetic small peptide bioinspired in the 2S albumin from the seed cake of *Ricinus communis*, is a potent antimicrobial agent against *Klebsiella pneumoniae* and *Candida parapsilosis*. *Biochim. Biophys. Acta Biomembr.* **2020**, *1862*, 183092. doi:10.1016/j.bbamem.2019.183092
48. Kodedová, M.; Sychrová, H. Synthetic antimicrobial peptides of the halictines family disturb the membrane integrity of *Candida* cells. *Biochim. Biophys. Acta Biomembr.* **2017**, *1859*, 1851-1858. doi:10.1016/j.bbamem.2017.06.005
49. Jang, W.S.; Kim, H.K.; Lee, K.Y.; Am Kim, S.; Han, Y.S.; Lee, I.H. Antifungal activity of synthetic peptide derived from halocidin, antimicrobial peptide from the tunicate, *Halocynthia aurantium*. *FEBS Letters* **2006**, *580*, 1490-1496. doi:10.1016/j.febslet.2006.01.041
50. Souza, P.F.; Marques, L.S.; Oliveira, J.T.; Lima, P.G.; Dias, L.P.; Neto, N.A.; Lopes, F.E.S.; Sousa, J.S.; Silva, A.F.B.; Carneiro, R.F.; Lopes, J.L.S.; Ramos, M.V.; Freitas, C. D. Synthetic antimicrobial peptides: from choice of the best sequences to action mechanisms. *Biochimie* **2020**, *175*, 132-145. doi:10.1016/j.biochi.2020.05.016
51. Lima, P.G.; Freitas, C.D.; Oliveira, J.T.; Neto, N.A.; Amaral, J.L.; Silva, A.F.; Sousa, J.S.; Franco, O.L.; Souza, P.F.N. Synthetic antimicrobial peptides control *Penicillium digitatum* infection in orange fruits. *Food Res. Int.* **2021**, *147*, 110582. doi:10.1016/j.foodres.2021.110582
52. Devi, M.S.; Sashidhar, R.B. Antiaflatoxigenic effects of selected antifungal peptides. *Peptides* **2019**, *115*, 15-26. doi:10.1016/j.peptides.2019.02.006
53. Devi, S.M.; Raj, N.; Sashidhar, R.B. Efficacy of short-synthetic antifungal peptides on pathogenic *Aspergillus flavus*. *Pestic. Biochem. Physiol.* **2021**, *174*, 104810. doi:10.1016/j.pestbp.2021.104810
54. Thery, T.; Shwaiki, L.N.; O'Callaghan, Y.C.; O'Brien, N.M.; Arendt, E.K. (2019). Antifungal activity of a de novo synthetic peptide and derivatives against fungal food contaminants. *J. Pept. Sci.* **2019**, *25*, e3137. doi:10.1002/psc.3137
55. Munoz, A.; López-García, B.; Marcos, J.F. Studies on the mode of action of the antifungal hexapeptide PAF26. *Antimicrob. Agents Chemother.* **2006**, *50*, 3847-3855. doi:10.1128/aac.00650-06
56. Nikapitiya, C.; Dananjaya, S.H.S.; Chandrarathna, H.P.S.U.; De Zoysa, M.; Whang, I. Octominin: A novel synthetic anticandidal peptide derived from defense protein of *Octopus minor*. *Mar. Drugs* **2020**, *18*, 56. doi:10.3390/md18010056
57. Lima, P.G.; Souza, P.F.N.; Freitas, C.D.T.; Oliveira, J.T.A.; Dias, L.P.; Neto, J.X.S.; Vasconcelos, I.M.; Lopes, J.L.S.; Sousa, D.O.B. Anticandidal activity of synthetic peptides: mechanism of action revealed by scanning electron and fluorescence microscopies and synergism effect with nystatin. *J. Pept. Sci.* **2020**, *26*, e3249. doi:10.1002/psc.3249
58. Falcao, L.L.; Silva-Werneck, J.O.; Ramos, A.D.R.; Martins, N.F.; Bresso, E.; Rodrigues, M.A.; Bemquerer, M.P.; Marcellino, L.H. Antimicrobial properties of two novel peptides derived from *Theobroma cacao* osmotin. *Peptides* **2016**, *79*, 75-82. doi:10.1016/j.peptides.2016.03.006
59. Martínez-Culebras, P.V.; Gandía, M.; Garrigues, S.; Marcos, J.F.; Manzanares, P. Antifungal peptides and proteins to control toxigenic fungi and mycotoxin biosynthesis. *Int. J. Mol. Sci.* **2021**, *22*, 13261. doi:10.3390/ijms222413261
60. Struyfs, C.; Cammue, B.P.A.; Thevissen, K. Membrane-interacting antifungal peptides. *Front Cell Dev Biol.* **2021**, *9*, 649875. doi: 10.3389/fcell.2021.649875
61. Huan, Y.; Kong, Q.; Mou, H.; Yi, H. Antimicrobial Peptides: Classification, design, application and research progress in multiple fields. *Front. Microbiol.* **2020**, *11*, 582779. doi: 10.3389/fmicb.2020.582779
62. Wang, B.; Timilsena, Y.P.; Blanch, E.; Adhikari, B. Lactoferrin: Structure, function, denaturation and digestion. *Crit. Rev. Food Sci. Nutr.* **2019**, *59*, 580-596. doi:10.1080/10408398.2017.1381583

63. Andrés, M.T.; Acosta-Zaldívar, M.; Fierro, J.F. Antifungal mechanism of action of lactoferrin: identification of H<sup>+</sup>-ATPase (P3A-type) as a new apoptotic-cell membrane receptor. *Antimicrob. Agents Chemother.* **2016**, *60*, 4206-4216. doi:10.1128/aac.03130-15
64. Gruden, Š.; Poklar Ulrih, N. Diverse mechanisms of antimicrobial activities of lactoferrins, lactoferricins, and other lactoferrin-derived peptides. *Int. J. Mol. Sci.*, **2021**, *22*, 11264. doi:10.3390/ijms222011264
65. Tóth, L.; Váradi, G.; Boros, É.; Borics, A.; Ficze, H.; Nagy, I.; Tóth, G.K.; Rákhely, G.; Marx, F.; Galgóczy, L. Biofungicidal potential of *Neosartorya (Aspergillus) fischeri* antifungal protein NFAP and novel synthetic -core peptides. *Front. Microbiol.* **2020**, *11*, 820. doi: 10.3389/fmicb.2020.00820
66. Oliveira, J.T.; Souza, P.F.; Vasconcelos, I.M.; Dias, L.P.; Martins, T.F.; Van Tilburg, M.F.; Guedes, M.I.F.; Sousa, D.O. Mo-CBP3-PepI, Mo-CBP3-PepII, and Mo-CBP3-PepIII are synthetic antimicrobial peptides active against human pathogens by stimulating ROS generation and increasing plasma membrane permeability. *Biochimie* **2019**, *157*, 10-21. doi:10.1016/j.biochi.2018.10.016
67. Bezerra, L.P.; Freitas, C.D.T.; Silva, A.F.B.; Amaral, J.L.; Neto, N.A.S.; Silva, R.G.G.; Parra, A.L.C.; Goldman, G.H.; Oliveira, J.T.A.; Mesquita, F.P.; et al. Synergistic antifungal activity of synthetic peptides and antifungal drugs against *Candida albicans* and *C. parapsilosis* biofilms. *Antibiotics* **2022**, *11*, 553. doi: 10.3390/antibiotics11050553
68. Cardoso, M.H.; Orozco, R.Q.; Rezende, S.B.; Rodrigues, G.; Oshiro, K.G.N.; Cândido, E.S.; Franco, O.L. (2020) Computer-aided design of antimicrobial peptides: Are we generating effective drug candidates? *Front. Microbiol.* **2020**, *10*, 3097. doi: 10.3389/fmicb.2019.03097
69. Li, R.; Wu, J.; He, F.; Xu, Q.; Yin, K.; Li, S.; Li, W.; Wei, A.; Zhang, L.; Zhang, X.-H.; Zhang, B. Rational design, synthesis, antifungal evaluation and docking studies of antifungal peptide CGA-N12 analogues based on the target CtkRE9. *Bioorg. Chem.* **2023**, *132*, 106355. doi:10.1016/j.bioorg.2023.106355
70. Sharma, K.; Aaghaz, S.; Maurya, I.K.; Rudramurthy, S.M.; Singh, S.; Kumar, V.; Tikoo, K.; Jain, R. Antifungal evaluation and mechanistic investigations of membrane active short synthetic peptides-based amphiphiles. *Bioorg. Chem.* **2022**, *127*, 106002. doi:10.1016/j.bioorg.2022.106002
71. Thery, T.; O'Callaghan, Y.; O'Brien, N.; Arendt, E.K. Optimisation of the antifungal potency of the amidated peptide H-Orn-Orn-Trp-Trp-NH<sub>2</sub> against food contaminants. *Int. J. Food Microbiol.* **2018**, *265*, 40-48doi:10.1016/j.ijfoodmicro.2017.10.024
72. Datta, L.P.; Mukherjee, T.; Das, T.K. Biomimetic pH responsive amphiphilic polymers: Solution property dependent antifungal mechanism. *React. Funct. Polym.* **2021**, *165*, 104937. doi:10.1016/j.reactfunctpolym.2021.104937
73. Ramamourthy, G.; Park, J.; Seo, C.; J. Vogel, H.; Park, Y. Antifungal and antibiofilm activities and the mechanism of action of repeating lysine-tryptophan peptides against *Candida albicans*. *Microorganisms* **2020**, *8*, 758. doi:10.3390/microorganisms8050758
74. Leanne-Rialland, V.; Cabezas-Cruz, A.; Atanasova, V.; Chereau, S.; Ponts, N.; Tonk, M.; Vivinskis, A.; Ferrer, N.; Valdes, J.J.; Richard-Forget, F. Tick defensin  $\gamma$ -core reduces *Fusarium graminearum* growth and abrogates mycotoxins production with high efficiency. *Sci. Rep.* **2021**, *11*, 7962. doi:10.1038/s41598-021-86904-w
75. Efremenko, E.; Aslanli, A.; Lyagin, I. Advanced situation with recombinant toxins: Diversity, production and application purposes. *Int. J. Mol. Sci.* **2023**, *24*, 4630. doi:10.3390/ijms24054630
76. Tevyashova, A.; Efimova, S.; Alexandrov, A.; Omelchuk, O.; Ghazy, E.; Bychkova, E.; Zatonsky, G.; Grammatikova, N.; Dezhenskova, L.; Solovieva, S.; Ostroumova, O. Semisynthetic amides of amphotericin B and nystatin A1: a comparative study of *in vitro* activity/toxicity ratio in relation to selectivity to ergosterol membranes. *Antibiotics* **2023**, *12*, 151. doi:10.3390/antibiotics12010151
77. Enache, A.C.; Cojocar, C.; Samoilă, P.; Bele, A.; Bostanaru, A.C.; Mares, M.; Harabagiu, V. Evaluation of physically and/or chemically modified chitosan hydrogels for proficient release of insoluble nystatin in simulated fluids. *Gels* **2022**, *8*, 495. doi:10.3390/gels8080495
78. Abid, S.; Uzair, B.; Niazi, M.B.K.; Fasim, F.; Bano, S.A.; Jamil, N.; Batool, R.; Sajjad, S. Bursting the virulence traits of MDR strain of *Candida albicans* using sodium alginate-based microspheres containing nystatin-loaded MgO/CuO nanocomposites. *Int. J. Nanomedicine* **2021**, *16*, 1157-1174. doi:10.2147/IJN.S282305
79. Nile, S.H.; Thombre, D.; Shelar, A.; Gosavi, K.; Sangshetti, J.; Zhang, W.; Sieniawska, E.; Patil, R.; Kai, G. Antifungal properties of biogenic selenium nanoparticles functionalized with nystatin for the inhibition of *Candida albicans* biofilm formation. *Molecules* **2023**, *28*, 1836. doi:10.3390/molecules28041836
80. El-Batal, A.I.; Nada, H.G.; El-Behery, R.R.; Gobara, M.; El-Sayyad, G.S. Nystatin-mediated bismuth oxide nano-drug synthesis using gamma rays for increasing the antimicrobial and antibiofilm activities against some pathogenic bacteria and *Candida species*. *RSC Adv.* **2020**, *10*, 9274-9289. doi:10.1039/C9RA10765G
81. Lin, Y.; Yin, Q.; Zhuge, D.; Hu, Y.; Yang, X.; Tian, D.; Li, L.; Wang, H.; Liu, S.; Weng, C.; Zhang, X.; Wen, B.; Wang, F.; Yan, L.; Chen, M.; Wang, L.; Chen, Y. Enhanced targeting, retention, and penetration of amphotericin B Through a biomimetic strategy to treat against vulvovaginal candidiasis. *Adv. Ther.* **2023**, *6*, 2200086. doi:10.1002/adtp.202200086

82. Zia, Q.; Mohammad, O.; Rauf, M.A.; Khan, W.; Zubair, S. Biomimetically engineered Amphotericin B nano-aggregates circumvent toxicity constraints and treat systemic fungal infection in experimental animals. *Sci. Rep.* **2017**, *7*, 11873. doi:10.1038/s41598-017-11847-0
83. Slezina, M.P.; Istomina, E.A.; Kulakovskaya, E.V.; Abashina, T.N.; Odintsova, T.I. Synthetic oligopeptides mimicking  $\gamma$ -core regions of cysteine-rich peptides of *Solanum lycopersicum* possess antimicrobial activity against human and plant pathogens. *Curr. Issues Mol. Biol.* **2021**, *43*, 1226-1242. doi:10.3390/cimb43030087
84. Kesharwani, P.; Fatima, M.; Singh, V.; Sheikh, A.; Almalki, W.H.; Gajbhiye, V.; Sahebkar, A. Itraconazole and difluorinated-curcumin containing chitosan nanoparticle loaded hydrogel for amelioration of onychomycosis. *Biomimetics* **2022**, *7*, 206. doi:10.3390/biomimetics7040206
85. Ionescu, O.M.; Iacob, A.T.; Mignon, A.; Van Vlierberghe, S.; Baican, M.; Danu, M.; Ibănescu, C.; Simionescu, N.; Profire, L. Design, preparation and in vitro characterization of biomimetic and bioactive chitosan/polyethylene oxide based nanofibers as wound dressings. *Int. J. Biol. Macromol.* **2021**, *193*, 996-1008. doi:10.1016/j.ijbiomac.2021.10.166
86. Uchida, M.; McDermott, G.; Wetzler, M.; Le Gros, M.A.; Myllys, M.; Knoechel, C.; Barron, A.E.; Larabell, C.A. Soft X-ray tomography of phenotypic switching and the cellular response to antifungal peptoids in *Candida albicans*. *Proc. Natl. Acad. Sci. U.S.A.*, **2009**, *106*, 19375-19380. doi:10.1073/pnas.0906145106
87. Abdelhamid, H.N.; Mahmoud, G.A.E.; Sharmouk, W. A cerium-based MOFzyme with multi-enzyme-like activity for the disruption and inhibition of fungal recolonization. *J. Mater. Chem. B* **2020**, *8*, 7548-7556. doi:10.1039/D0TB00894J
88. Li, X.; Wu, X.; Yuan, T.; Zhu, J.; Yang, Y. Influence of the iodine content of nitrogen-and iodine-doped carbon dots as a peroxidase mimetic nanozyme exhibiting antifungal activity against *C. albicans*. *Biochem. Eng. J.* **2021**, *175*, 108139. doi:10.1016/j.bej.2021.108139
89. Biswas, S.; Priya Datta, L.; Kumar Das, T. Peptide core containing polymer-polyoxometalate hybrid as novel antifungal agent. *J. Mol. Eng. Mater.*, **2022**, *10*, 2240004. doi:10.1142/S2251237322400044
90. Ivanov, M.; Ćirić, A.; Stojković, D. Emerging antifungal targets and strategies. *Int. J. Mol. Sci.* **2022**, *23*, 2756. doi:10.3390/ijms23052756
91. Xia, Q.; Tian, H.; Li, Y.; Yu, X.; Zhang, W.; Wang, Q. Biomimetic synthesis of iridoid alkaloids as novel leads for fungicidal and insecticidal agents. *J. Agric. Food Chem.* **2020**, *68*, 12577-12584. doi:10.1021/acs.jafc.0c04885
92. Zhang, X.; Xu, H.; Su, H.; Yang, X.; Sun, T.; Lu, X.; Shi, F.; Duan, H.; Liu, X.; Ling, Y. Design, synthesis, and biological activity of novel fungicides containing a 1, 2, 3, 4-tetrahydroquinoline scaffold and acting as laccase inhibitors. *J. Agric. Food Chem.* **2022**, *70*, 1776-1787. doi:10.1021/acs.jafc.1c06595
93. Chen, W.; Zhang, R.; Chen, Y.; Yu, P.; Lan, Y.; Xu, H.; Lei, S. Design, synthesis and mechanism study of novel natural-derived isoquinoline derivatives as antifungal agents. *Mol. Divers.*, **2023**, *27*, 1011-1022. doi:10.1007/s11030-022-10463-z
94. Chaurasia, H.; Singh, V.K.; Mishra, R.; Yadav, A.K.; Ram, N.K.; Singh, P.; Singh, R.K. Molecular modelling, synthesis and antimicrobial evaluation of benzimidazole nucleoside mimetics. *Bioorg. Chem.* **2021**, *115*, 105227. doi:10.1016/j.bioorg.2021.105227
95. DeMarco, A.G.; Milholland, K.L.; Pendleton, A.L.; Whitney, J.J.; Zhu, P.; Wesenberg, D.T.; Nambiar, M.; Pepe, A.; Paula, S.; Chmielewski, J.; Wisecaver, J.H. Conservation of Cdc14 phosphatase specificity in plant fungal pathogens: implications for antifungal development. *Sci. Rep.* **2020**, *10*, 12073. doi:10.1038/s41598-020-68921-3
96. Huang, D.S.; LeBlanc, E.V.; Shekhar-Guturja, T.; Robbins, N.; Krysan, D.J.; Pizarro, J.; Whitesell, L.; Cowen, L.E.; Brown, L.E. Design and synthesis of fungal-selective resorcyate aminopyrazole Hsp90 inhibitors. *J. Med. Chem.* **2019**, *63*, 2139-2180. doi:10.1021/acs.jmedchem.9b00826
97. Xiong, Y.; Su, L.; Peng, Y.; Zhao, S.; Ye, F. Dextran-coated Gd-based ultrasmall nanoparticles as phosphatase-like nanozyme to increase ethanol yield via reduction of yeast intracellular ATP level. *J. Colloid Interface Sci.*, **2022**, *627*, 405-414. doi:10.1016/j.jcis.2022.07.036
98. Rosenzweig, R.; Marshall, M.; Parivar, A.; Ly, V.K.; Pearlman, E.; Yee, A.F. Biomimetic nanopillared surfaces inhibit drug resistant filamentous fungal growth. *ACS Appl. Bio Mater.*, **2019**, *2*, 3159-3163. doi:10.1021/acsabm.9b00290
99. Dalefield, R. Human Pharmaceuticals. In *Veterinary toxicology for Australia and New Zealand*, Dalefield, R., Eds.; Elsevier: Amsterdam, Netherlands, 2017; Chapter 12, pp. 155-201. doi:10.1016/B978-0-12-420227-6.00036-0
100. Millikan, L.E. Current concepts in systemic and topical therapy for superficial mycoses. *Clin. Dermatol.* **2010**, *28*, 212-216. doi:10.1016/j.clindermatol.2009.12.016
101. Parker, J.E.; Warrilow, A.G.S.; Price, C.L.; Mullins, J.G.L.; Kelly, D.E.; Kelly, S.L. Resistance to antifungals that target CYP51. *J. Chem. Biol.* **2014**, *7*, 143. doi:10.1007/s12154-014-0121-1
102. Leadbeater, A.J. Plant health management: fungicides and antibiotics. In *Encyclopedia of Agriculture and Food Systems*, Van Alfen, N.K., Eds; Academic Press: N-Y., USA, 2014, pp. 408-424. doi:10.1016/B978-0-444-52512-3.00179-0



103. Efremenko, E.; Senko, O.; Stepanov, N.; Maslova, O.; Lomakina, G.; Ugarova, N. Luminescent analysis of ATP: Modern objects and processes for sensing. *Chemosensors* **2022**, *10*, 493. doi:10.3390/chemosensors10110493
104. Batchuluun, B.; Pinkosky, S.L.; Steinberg, G.R. Lipogenesis inhibitors: therapeutic opportunities and challenges. *Nat. Rev. Drug Discov.*, **2022**, *21*, 283-305. doi:10.1038/s41573-021-00367-2
105. Li, G.; Jian, T.; Liu, X.; Lv, Q.; Zhang, G.; Ling, J. Application of metabolomics in fungal research. *Molecules* **2022**, *27*, 7365. doi:10.3390/molecules27217365
106. Zhong, L.F.; Shang, Z.C.; Sun, F.J.; Zhu, P.H.; Yin, Y.; Kong, L.Y.; Yang, M.H. Anticandidal formyl phloroglucinol meroterpenoids: Biomimetic synthesis and in vitro evaluation. *Bioorg. Chem.*, **2020**, *104*, 104248. doi:10.1016/j.bioorg.2020.104248.
107. Pojtkov, A.E.; Efremenko, E.N.; Varfolomeev, S.D. Unnatural amino acids in enzymes and proteins. *J. Molecul. Catal. B: Enzym.*, **2000**, *10*, 47-55. doi:10.1016/s1381-1177(00)00107-7
108. Qu, T.; Gao, S.; Li, J.; Hao, J.J.; Ji, P. Synthesis and antifungal activity of 2-allylphenol derivatives against fungal plant pathogens. *Pestic. Biochem. Physiol.*, **2017**, *135*, 47-51. doi:10.1016/j.pestbp.2016.06.006
109. Laussel, C.; Léon, S. Cellular toxicity of the metabolic inhibitor 2-deoxyglucose and associated resistance mechanisms. *Biochem. Pharmacol.* **2020**, *182*, 114213. doi:10.1016/j.bcp.2020.114213
110. Hellemann, E.; Walker, J.L.; Lesko, M.A.; Chandrashekarappa, D.G.; Schmidt, M.C.; O'Donnell, A.F.; Durrant, J.D. Novel mutation in hexokinase 2 confers resistance to 2-deoxyglucose by altering protein dynamics. *PLoS Comput. Biol.* **2022**, *18*, e1009929. doi:10.1371/journal.pcbi.1009929
111. Trudeau, S.; Morken, J. Short and efficient total synthesis of fraxinellone limonoids using the stereoselective oshima-utimoto reaction. *Org. Lett.* **2005**, *7*, 5465-5468. doi:10.1021/ol0522687
112. Serafin-Lewañczuk, M.; Brzezińska-Rodak, M.; Lubiak-Kozłowska, K.; Majewska, P.; Klimek-Ochab, M.; Olszewski, T.K.; Żymańczyk-Duda, E. Phosphonates enantiomers receiving with fungal enzymatic systems. *Microb. Cell Factories* **2021**, *20*, 81. doi:10.1186/s12934-021-01573-8
113. Tlidjane, H.; Chafai, N.; Chafaa, S.; Bensouici, C.; Benbouguerra, K. New thiophene-derived  $\alpha$ -aminophosphonic acids: Synthesis under microwave irradiations, antioxidant and antifungal activities, DFT investigations and SARS-CoV-2 main protease inhibition. *J. Mol. Struct.* **2022**, *1250*, 131853. doi:10.1016/j.molstruc.2021.131853
114. Varfolomeev, S.D.; Aliev, T.K.; Efremenko, E.N. Postgenomic chemistry: New problems and challenges. *Pure Appl. Chem.*, **2004**, *76*, 1781-1798. doi:10.1351/pac200476091781.
115. Maslova, O.V.; Senko, O.V.; Efremenko, E.N. Aspartic and glutamic acids polymers: preparation and applications in medicinal chemistry and pharmaceuticals. *Russ. Chem. Bull.* **2018**, *67*, 614-623. doi:10.1007/s11172-018-2115-6
116. Mardirossian, M.; Rubini, M.; Adamo, M.F.A.; Scocchi, M.; Saviano, M.; Tossi, A.; Gennaro, R.; Caporale, A. Natural and synthetic halogenated amino acids—structural and bioactive features in antimicrobial peptides and peptidomimetics. *Molecules* **2021**, *26*, 7401. doi:10.3390/molecules26237401
117. Zhao, W.; Wolfender, J.L.; Hostettmann, K.; Xu, R.; Qin, G. Antifungal alkaloids and limonoid derivatives from *Dictamnus dasycarpus*. *Phytochemistry* **1998**, *47*, 7-11. doi:10.1016/S0031-9422(97)00541-4
118. Wang, M.; Zhang, J.; Qian, Y.; Li, J.; Ji, Z. The study on fungicidal activity of *Dictamnus dasycarpus*. *Agrochemicals* **2006**, *45*, 739-741. doi:10.16820/j.cnki.1006-0413.2006.11.007
119. Yang, X.-J.; Dong, Q.-M.; Wang, M.-R.; Tang, J.-J. Semi-synthesis of C-ring cyclopropyl analogues of fraxinellone and their insecticidal activity against *Mythimna separate walker*. *Molecules* **2020**, *25*, 1109. doi:10.3390/molecules25051109
120. Varfolomeev, S.D.; Efremenko, E.N. *Organophosphorus Neurotoxins*; RIOR: Moscow, Russia, 2020; p. 380. ISBN 978-5-369-02026-5. doi:10.29039/02026-5.
121. Kirouani, I.; Hellal, A.; Haddadi, I.; Layaida, H.; Madani, A.; Madani, S.; Haroun, M.F.; Rachida, D.; Touafri, L.; Bensouici, C. Effect of the phosphonomethylene moiety on the structural, vibrational, energetic, thermodynamic and optical proprieties of ((Phenylcarbamoylmethyl-phosphonomethyl-amino)-methyl)-phosphonic acid: DFT investigation. *J. Mol. Struct.* **2020**, *1215*, 128193. doi:10.1016/j.molstruc.2020.128193
122. Lyagin, I.V.; Andrianova, M.S.; Efremenko, E.N. Extensive hydrolysis of phosphonates as unexpected behaviour of the known His<sub>6</sub>-organophosphorus hydrolase. *Appl. Microbiol. Biotechnol.* **2016**, *100*, 5829-5838. doi:10.1007/s00253-016-7407-x
123. Partida-Hanon, A.; Maestro-López, M.; Vitale, S.; Turrà, D.; Di Pietro, A.; Martínez-del-Pozo, Á.; Bruix, M. Structure of fungal  $\alpha$  mating pheromone in membrane mimetics suggests a possible role for regulation at the water-membrane interface. *Front. Microbiol.*, **2020**, *11*, 1090. doi:10.3389/fmicb.2020.01090.
124. Doley, K.; Thomas, S.; Borde, M. External signal-mediated overall role of hormones/pheromones in fungi. In *Fungal reproduction and growth*, Sultan, S., Singh, G.K.S., Eds.; In techOpen, London, United Kingdom, 2022, pp. 73-83. doi:10.5772/intechopen.101154
125. Jung, S.I.; Finkel, J.S.; Solis, N.V.; Chaili, S.; Mitchell, A.P.; Yeaman, M.R.; Filler, S.G. Bcr1 functions downstream of Ssd1 to mediate antimicrobial peptide resistance in *Candida albicans*. *Eukaryot. Cell* **2013**, *12*, 411-419. doi:10.1128/EC.00285-12



126. Pappas, P.G.; Lionakis, M.S.; Arendrup, M.C.; Ostrosky-Zeichner, L.; Kullberg, B.J. Invasive candidiasis. *Nat. Rev. Dis. Primers*. **2018**, *4*, 18026. doi:10.1038/nrdp.2018.26
127. Pimienta, D.A.; Cruz Mosquera, F.E.; Palacios Velasco, I.; Giraldo Rodas, M.; Oñate-Garzón, J.; Liscano, Y. Specific focus on antifungal peptides against azole resistant *Aspergillus fumigatus*: Current status, challenges, and future perspectives. *J. Fungi* **2023**, *9*, 42. doi:10.3390/jof9010042
128. Aguiar, T.K.B.; Feitosa, R.M.; Neto, N.A.S.; Malveira, E.A.; Gomes, F.I.R.; Costa, A.C.M.; Freitas, C.D.T.; Mesquita, F.P.; Souza, P.F.N. Giving a hand: synthetic peptides boost the antifungal activity of itraconazole against *Cryptococcus neoformans*. *Antibiotics* **2023**, *12*, 256. doi:10.3390/antibiotics12020256
129. Tóth, L.; Poór, P.; Ördög, A.; Váradi, G.; Farkas, A.; Papp, C.; Bende, G.; Tóth, G.K.; Rákhely, G.; Marx, F.; Galgóczy, L. The combination of *Neosartorya (Aspergillus) fischeri* antifungal proteins with rationally designed  $\gamma$ -core peptide derivatives is effective for plant and crop protection. *BioControl* **2022**, *67*, 249–262. doi:10.1007/s10526-022-10132-y
130. Ding, K.; Shen, P.; Xie, Z.; Wang, L.; Dang, X. In vitro and in vivo antifungal activity of two peptides with the same composition and different distribution. *Comp. Biochem. Physiol. C: Toxicol. Pharmacol.* **2022**, *252*, 109243. doi:10.1016/j.cbpc.2021.109243
131. Tao, Y.; Bie, X.; Lv, F.; Zhao, H.; Lu, Z. Antifungal activity and mechanism of fengycin in the presence and absence of commercial surfactin against *Rhizopus stolonifer*. *J. Microbiol.* **2011**, *49*, 146–150. doi:10.1007/s12275-011-0171-9
132. do Nascimento Dias, J.; de Souza Silva, C.; de Araújo, A.R.; Souza, J.M.T.; de Holanda Veloso Junior, P.H.; Cabral, W.F.; da Glória da Silva, M.; Eaton, P.; de Souza de Almeida Leite, J.R.; Nicola, A.M.; Albuquerque, P. Mechanisms of action of antimicrobial peptides ToAP2 and NDBP-5.7 against *Candida albicans* planktonic and biofilm cells. *Sci. Rep.* **2020**, *10*, 10327. doi:10.1038/s41598-020-67041-2
133. Galdiero, E.; de Alteriis, E.; De Natale, A.; D'Alterio, A.; Siciliano, A.; Guida, M.; Lombardi, L.; Falanga, A.; Galdiero, S., 2020. Eradication of *Candida albicans* persister cell biofilm by the membranotropic peptide gH625. *Sci. Rep.* **2020**, *10*, 5780. doi:10.1038/s41598-020-62746-w
134. Sharma, K.; Aaghaz, S.; Maurya, I.K.; Sharma, K.K.; Singh, S.; Rudramurthy, S.M.; Kumar, V.; Tikoo, K.; Jain, R. Synthetic amino acids-derived peptides target *Cryptococcus neoformans* by inducing cell membrane disruption. *Bioorg. Chem.* **2023**, *130*, 106252. doi:10.1016/j.bioorg.2022.106252
135. Zhang, X.; Wang, M.; Zhu, X.; Peng, Y.; Fu, T.; Hu, C.H.; Cai, J.; Liao, G. Development of Lipo- $\gamma$ -AA Peptides as potent antifungal agents. *J. Med. Chem.* **2022**, *65*, 8029–8039. doi:10.1021/acs.jmedchem.2c00595
136. Darwish, R.M.; Salama, A.H. A pilot study on ultrashort peptide with fluconazole: A promising novel anticandidal combination. *Vet. World* **2023**, *16*, 1284–1288. doi:10.14202/vetworld.2023.1284-1288
137. Ptaszyńska, N.; Gucwa, K.; Olkiewicz, K.; Łęgowska, A.; Okońska, J.; Ruczyński, J.; Gitlin-Domagalska, A.; Dębowski, D.; Milewski, S.; Rolka, K. Antibiotic-based conjugates containing antimicrobial HLOpt2 peptide: design, synthesis, antimicrobial and cytotoxic activities. *ACS Chem. Biol.* **2019**, *14*, 2233–2242. doi:10.1021/acschembio.9b00538
138. Rodríguez-López, A.D.L.; Lee, M.R.; Wang, N.B.; Dunn, K.K.; Sanchez, H.; Raman, N.; Andes, D.R.; Lynn, D.M.; Palecek, S.P. Small-molecule morphogenesis modulators enhance the ability of 14-helical  $\beta$ -peptides to prevent *Candida albicans* biofilm formation. *Antimicrob. Agents Chemother.* **2019**, *63*, 10–1128. doi:10.1128/aac.02653-18
139. Fernandes, K.E.; Payne, R.J.; Carter, D.A. Lactoferrin-derived peptide lactofungin is potently synergistic with amphotericin B. *Antimicrob. Agents Chemother.* **2020**, *64*, 10–1128. doi:10.1128/aac.00842-20
140. Ptaszyńska, N.; Gucwa, K.; Olkiewicz, K.; Heldt, M.; Serocki, M.; Stupak, A.; Martynow, D.; Dębowski, D.; Gitlin-Domagalska, A.; Lica, J.; Łęgowska, A.; Sławomir Milewski, S.; Rolka, K. Conjugates of ciprofloxacin and levofloxacin with cell-penetrating peptide exhibit antifungal activity and mammalian cytotoxicity. *Int. J. Mol. Sci.* **2020**, *21*, 4696. doi:10.3390/ijms21134696
141. Lyagin, I.; Aslanli, A.; Domnin, M.; Stepanov, N.; Senko, O.; Maslova, O.; Efremenko, E. Metal nanomaterials and hydrolytic enzyme-based formulations for improved antifungal activity. *Int. J. Mol. Sci.* **2023**, *24*, 11359. doi:10.3390/ijms241411359
142. Efremenko, E.; Stepanov, N.; Aslanli, A.; Lyagin, I.; Senko, O.; Maslova, O. Combination of enzymes with materials to give them antimicrobial features: Modern trends and perspectives. *J. Funct. Biomater.* **2023**, *14*, 64. doi:10.3390/jfb14020064
143. Aslanli, A.; Domnin, M.; Stepanov, N.; Efremenko, E. “Universal” Antimicrobial combination of bacitracin and His-OPH with lactonase activity, acting against various bacterial and yeast cells. *Int. J. Mol. Sci.* **2022**, *23*, 9400. doi:10.3390/ijms23169400
144. Aslanli, A.; Domnin, M.; Stepanov, N.; Efremenko, E. Synergistic antimicrobial action of lactoferrin-derived peptides and quorum quenching enzymes. *Int. J. Mol. Sci.* **2023**, *24*, 3566. doi:10.3390/ijms24043566
145. Bezerra, L.P.; Silva, A.F.; Santos-Oliveira, R.; Alencar, L.M.; Amaral, J.L.; Neto, N.A.; Silva, R.G.; Belém, M.O.; de Andrade, C.R.; Oliveira, J.T.; Freitas, C.D. Combined antibiofilm activity of synthetic peptides and antifungal drugs against *Candida spp.* *Future Microbiol.* **2022**, *17*, 1133–1146. doi:10.2217/fmb-2022-0053

146. Khabbaz, H.; Karimi-Jafari, M.H.; Saboury, A.A.; BabaAli, B. Prediction of antimicrobial peptides toxicity based on their physico-chemical properties using machine learning techniques. *BMC Bioinform.* **2021**, *22*, 1-11. doi:10.1186/s12859-021-04468-y
147. Greco, I.; Molchanova, N.; Holmedal, E.; Jenssen, H.; Hummel, B.D.; Watts, J.L.; Joakim Hakansson, J.; Hansen, P.R.; Svenson, J. Correlation between hemolytic activity, cytotoxicity and systemic in vivo toxicity of synthetic antimicrobial peptides. *Sci. Rep.* **2020**, *10*, 13206. doi:10.1038/s41598-020-69995-9
148. Hong, K.; Wang, L.; Johnpaul, A.; Song, Y.; Guo, L.; Xie, X.; Lv, C.; Ma, C. Response of *Saccharomyces cerevisiae* var. *diastaticus* to nerol: Evaluation of antifungal potential by inhibitory effect and proteome analyses, *Food Chem.* **2023**, *403*, 134323. doi:10.1016/j.foodchem.2022.134323

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.