

Review

Not peer-reviewed version

PPAR Gamma Receptor: A Novel Target to Improve Morbidity in Preterm Babies

[Suresh Victor](#) *, [Ben Forbes](#) , [Anne Greenough](#) , A David Edwards

Posted Date: 1 September 2023

doi: 10.20944/preprints202309.0060.v1

Keywords: pioglitazone; prematurity; Bronchopulmonary dysplasia; lung, brain; adiponectin; PPAR gamma



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

PPAR Gamma Receptor: A Novel Target to Improve Morbidity in Preterm Babies

Suresh Victor ^{1,*}, Ben Forbes ², Anne Greenough ³ and A David Edwards ¹

¹ Centre for the Developing Brain, Department of Perinatal Imaging and Health, School of Biomedical Engineering and Imaging Sciences, King's College London, St Thomas' Hospital, London, United Kingdom SE1 7EH; suresh.victor@kcl.ac.uk

² Institute of Pharmaceutical Science, King's College London, Franklin-Wilkins Building, Stamford Street, London, United Kingdom SE1 9NH; ben.forbes@kcl.ac.uk

³ Department of Women and Children's Health, School of Life Course and Population Sciences, King's College London, Neonatal Intensive Care Centre, Floor 4, Golden Jubilee Wing, King's College Hospital, Denmark Hill, Brixton, London, United Kingdom SE5 9RS; anne.greenough@kcl.ac.uk

* Correspondence: suresh.victor@kcl.ac.uk

Abstract: Three-quarters of a million babies are born extremely preterm (< 28 weeks' gestation) world-wide with devastating outcomes: 20% die in the newborn period, a further 35% develop bronchopulmonary dysplasia (BPD) and 10% suffer from cerebral palsy. Pioglitazone, a Peroxisome Proliferator Activated Receptor Gamma (PPAR γ) agonist, may reduce the incidence of BPD and improve neurodevelopment in extreme preterm babies. Pioglitazone exerts an anti-inflammatory action mediated through Nuclear Factor-kappa B repression. PPAR γ signalling is underactive in preterm babies as adiponectin remains low during the neonatal period. In newborn animal models, pioglitazone is shown to be protective against BPD, necrotising enterocolitis, and lipopolysaccharide-induced brain injury. Single Nucleotide Polymorphisms of PPAR γ are associated with inhibited preterm brain development and impaired neurodevelopment. Pioglitazone was well tolerated by the foetus in reproductive toxicology experiments. Bladder cancer, bone fractures, and macular oedema seen rarely in adults may be avoided with a short treatment course. Other effects of pioglitazone including improved glycaemic control and lipid metabolism may provide added benefit in prematurity. Currently, there is no formulation of pioglitazone suited for administration to preterm babies. A liquid formulation of pioglitazone needs to be developed before clinical trials. The potential benefits are likely to outweigh any anticipated safety concerns.

Keywords: pioglitazone; prematurity; Bronchopulmonary dysplasia; lung; brain; adiponectin; PPAR gamma

1. Introduction

Prematurity accounts for 11% of all births and is the leading global cause of death and disability under five years of age [1]. Bronchopulmonary dysplasia (BPD) also known as chronic lung disease (defined as supplemental oxygen dependency at 36 weeks postmenstrual age) is seen in around 55% of extremely preterm babies. Around 10,000-15,000 new cases are diagnosed annually in the USA and a further 1250 in the UK. Recurrent respiratory symptoms requiring treatment are common, particularly in those who had BPD, even at school age and in adults, as well as lung function abnormalities and exercise intolerance [2]. Recent evidence suggests that affected individuals are at risk of the premature onset of Chronic Obstructive Pulmonary Disease [3]. Nearly 10% of preterm infants suffer from cerebral palsy with the risk doubling with BPD (OR: 2.10; 95% CI 1.57-2.82) [4].

Over 30% of survivors following preterm birth experience neurocognitive and socio-emotional problems from early life lasting into adulthood [5]. Psychiatric disorders are present in around 25%

of preterm adolescents [6,7]. Prematurity leads to seventy-five million disability adjusted life years per annum worldwide.

Improvements in mortality and morbidity in preterm babies have so far been achieved through improvements in intensive care management. There is currently no pharmacological treatment available for preterm infants during the newborn period that aims to prevent BPD and improve neurodevelopmental outcomes. Caffeine, when administered to preterm babies reduced BPD and improved neurodevelopment demonstrating that pharmacological treatment for BPD and neuroprotection is viable postnatally [8]. The CAP trial found a higher patient survival rate without neurodevelopmental disability (cognitive delay, cerebral palsy, severe hearing loss or bilateral blindness) at a corrected age of 18 to 21 months in infants within the caffeine group compared with those in the placebo group (59.8% versus 53.8%; adjusted OR: 0.77, 95%CI: 0.64-0.93, P = 0.008) [8].

There is an urgent need to develop treatment strategies that reduce devastating outcomes in premature babies. We propose that pioglitazone, a Peroxisome Proliferator Activated Receptor Gamma (PPAR γ) agonist, may reduce the incidence of death or BPD and improve neurodevelopment in extreme preterm babies. In this review, we summarise the supporting evidence in non-clinical and clinical studies, the safety considerations, and the pharmaceutical challenges in the translation of this novel treatment strategy.

2. Preclinical studies using PPAR γ agonists in preterm disease models

PPAR γ 1, PPAR γ 2, and PPAR γ 3 are the three isoforms of PPAR γ . While PPAR γ 2 is expressed predominantly in adipose tissue, PPAR γ 1 and PPAR γ 3 are expressed in adipose tissue, heart, kidney, pancreas, spleen, intestine, colon epithelial cells and skeletal muscle. In the nervous system, PPAR γ is expressed in neurons and glia, including microglia and astrocytes, in multiple brain regions [9,10]. PPAR γ is also expressed in lung epithelium, submucosa and airway smooth muscle [11]. The 15-deoxy prostaglandin J₂ (15d-PGJ₂) is a natural ligand for PPAR γ that binds and activates PPAR γ . Thiazolidinediones or glitazones (e.g. pioglitazone and rosiglitazone) are a class of synthetic compounds that function as high affinity agonists for PPAR γ . Studies have been conducted in various animal models of preterm diseases showing benefit (Figure 1).




Figure 1. Activation of PPAR γ may have potential beneficial effects (green arrows) as well as adverse effects (red arrows) in preterm babies. Adverse effects may be avoided by short duration of treatment. BPD: Bronchopulmonary Dysplasia; NEC: Necrotising Enterocolitis; ROP: Retinopathy of Prematurity.

2.1. PPAR γ agonists and BPD

PPAR γ expression has been reported in several pulmonary cell types, including inflammatory, mesenchymal, and epithelial cells [12]. To define the physiological role of PPAR γ within the lung, Simon et al. (2006) generated lung epithelium-specific PPAR γ -deficient mice by breeding the CCre transgenic line with mice harbouring loxP sites flanking exon 2 of the PPAR γ gene [13]. Airway epithelial cell PPAR γ -depleted mice displayed enlarged airspaces resulting from insufficient postnatal lung maturation. The increase in airspace size was accompanied by alterations in lung physiology, including increased lung volumes and decreased tissue resistance similar to BPD.

Wang et al. (2009) administered either diluent or rosiglitazone (0.1 to 8 mg/kg intraperitoneal once daily for either one day or 7 days) to newborn Sprague-Dawley rat pups [14]. The pups were then sacrificed, and the lungs were examined for specific markers of alveolar epithelial, mesenchymal, and vascular maturation and lung morphometry. Systemically administered rosiglitazone significantly enhanced lung maturation without affecting serum electrolytes, blood glucose, blood gases, plasma cholesterol, triglycerides and serum cardiac troponin levels. There was no significant effect on blood glucose at either day one or day 7, for any of the doses of rosiglitazone examined except for the highest dose used, i.e., 8 mg/kg for 7 days, which resulted in statistically significantly lower glucose values compared with controls (84 ± 15 mg/dL versus 127 ± 3 mg/dL (mean \pm SD), ($P < 0.05$, rosiglitazone versus control). Similar effects were noted when 15d-PGJ₂ was administered instead of rosiglitazone.

Rehan et al. (2006) exposed newborn Sprague Dawley rat pups to control (21% O₂), hyperoxia alone (95% O₂ for 24 hours), or hyperoxia with rosiglitazone (95% O₂ for 24 hours + rosiglitazone, 3 mg/kg, administered intraperitoneal) [15]. In lungs from hyperoxic animals, septal thickness was reduced significantly (40% versus 15%) compared with control lungs ($P < 0.01$, 95% versus 21% O₂). Quantitative analysis of alveolar number demonstrated a significant reduction (50 versus 15) in average radial alveolar counts in hyperoxic animals compared with controls. A significant decrease in the expression of lipogenic markers, and a significant increase in the expression of myogenic markers in the hyperoxia-alone group were observed. Exposure to hyperoxia alone markedly decreased PPAR γ , and markedly increased alpha smooth muscle actin expression. These hyperoxia-induced morphologic, molecular, and immunohistochemical changes were completely prevented by rosiglitazone.

Lee et al. (2020) administered dexamethasone or pioglitazone (2-doses, 0.3-3 mg/kg bolus intraperitoneal injection) to pregnant Sprague-Dawley rat dams on embryonic day 18 and 19 and delivered by caesarean section on day twenty [16]. Antenatal pioglitazone increased alveolar epithelial and mesenchymal maturation markers equally in males and females; in contrast, antenatal dexamethasone had sex-specific effects. Dexamethasone (0.25 mg/kg intraperitoneal injection on embryonic days 18 and 19) significantly increased surfactant phospholipid synthesis, and alveolar epithelial and mesenchymal maturation markers in both males and females. However, this increase was more prominent in females. Additionally, unlike dexamethasone, antenatal pioglitazone did not increase hippocampal apoptosis.

Richter et al. (2016) injected pregnant rabbits with saline or rosiglitazone (3 mg/kg) 48 and 24 hours prior to preterm delivery at 28 days of gestation (term = 31 days) [17]. The pups were held in normoxia (21% O₂) or hyperoxia (>95% O₂), and assessment was done at three different time points (1 hour, 24 hours and 7 days). Rosiglitazone significantly increased expression of Vascular Endothelial Growth Factor (VEGF), and surfactant protein B immediately after delivery. On day seven, there was a more mature lung parenchymal architecture in rosiglitazone-exposed pups.

Morales et al. (2014) administered one-day old Sprague-Dawley rat pups PPAR γ agonists rosiglitazone (3 mg/kg), pioglitazone (3 mg/kg), or the diluent, via nebulization every 24 hours; animals were exposed to 21% or 95% O₂ for up to 72 hours [18]. Rosiglitazone and pioglitazone enhanced lung maturation in both males and females, as evidenced by the increased expression of markers of alveolar epithelial and mesenchymal maturation at 72 hours after initial nebulisation. This approach also protected against hyperoxia-induced lung injury.

In one-day-old rats exposed to 95% O₂, aerosolized/ intraperitoneal pioglitazone (1-3 mg/kg; plasma concentration: 0.52-0.99 ± 0.15 µg/mL) once daily for three days alone or in combination with surfactant significantly increased markers of lung maturation, significantly decreased markers of lung inflammation, and prevented lung injury [19].

One-day-old rats induced through intra-amniotic delivery of lipopolysaccharide (LPS) and postnatal hyperoxia (80% for 7 days) and treated with intraperitoneal rosiglitazone (3 mg/kg for 14 days) had significantly smaller and more complex airspaces, larger alveolar surface area, greater pulmonary vascular density, restored levels of VEGF and its receptor and decreased right ventricular hypertrophy [20].

2.2. PPAR γ agonists and preterm brain injury

Pioglitazone represses two signalling pathways associated with preterm brain injury, (a) the Nuclear Factor-kappa B (NF- κ B) signalling pathway which promotes inflammation mediated injury [21], and (b) the Wnt canonical signalling pathway which inhibits developmental myelination [22]. The NF- κ B activation in microglia plays a significant role in the pathogenesis of hypoxic ischemic injury of the immature brain, and its prophylactic inhibition offers significant neuroprotection [23].

Pioglitazone (20 mg/kg single dose intraperitoneal) significantly improved LPS induced neuro-behavioural and physiological disturbances including loss of weight, hypothermia, righting reflex, wire hanging manoeuvre, negative geotaxis, and hind limb suspension in post-natal day five rats²⁴. Pioglitazone attenuated LPS induced loss of oligodendrocytes, reduction of mitochondrial complex one activity, increase in thiobarbituric acid-reactive substance content, and increase in microglial activation and inflammatory responses. Pioglitazone attenuated the LPS-induced increase of IL (Interleukin)-1 β in both serum and brain (P< 0.05) and increase of IL-6 and TNF (Tumour Necrosis Factor)- α levels in the serum of rats (P< 0.05) [24].

Retinal ischemia/reperfusion injury is considered to be associated with retinopathy of prematurity (ROP). In a mouse model of retinal ischemia/reperfusion, pioglitazone promoted the survival of retinal cells in ganglion cell layer following retinal ischemia/reperfusion injury (P< 0.05) by suppressing ischemia/reperfusion-induced activation of Toll-like receptor 4 and NLR family pyrin domain containing 3 (NLRP3) inflammasomes via inhibiting NF- κ B and p38 phosphorylation [25]. Pioglitazone was one of the novel candidate compounds identified for the treatment of ROP using computational drug-gene association analysis [26].

2.3. PPAR agonists and necrotising enterocolitis

Swiss-Webster mice were randomized to receive sham (control) or ischemia-reperfusion injury to the gut induced by transient occlusion of superior mesenteric artery for 45 min with variable periods of reperfusion. Ischemia-reperfusion injury resulted in early induction of PPAR γ expression and activation of NF- κ B in small intestine. Pre-treatment with PPAR γ - agonist, 15d-PGJ2 (2 mg/kg), 45 min prior to injury attenuated intestinal NF- κ B response and ischemia-reperfusion induced gut injury. Activation of PPAR γ demonstrated a protective effect on small bowel during ischemia-reperfusion induced gut injury [27].

Corsini et al. (2016) studied preterm rats in which necrotising enterocolitis was induced using the hypoxia-hypothermia model [28]. The treatment group (n = 30) received enteral pioglitazone (10 mg/kg/day) for 72 h and the control group (n = 30) did not. Animals were sacrificed 96 h after birth. Necrotising enterocolitis was diagnosed evaluating histological ileum changes, and messenger RNA (mRNA) levels of IL-4, IL-12, IL-6, IL-10, Interferon- γ , and TNF- α cytokines were measured. Necrotising enterocolitis occurrence was higher in the control group (18/30; 60%) than in the treatment group (5/30; 16.7%) and was more severe. Proinflammatory IL-12 and Interferon- γ mRNA levels were significantly lower in the treatment group than in the control group; conversely, the anti-inflammatory IL-4 mRNA level was significantly higher in the treatment group than in the control group.

3. Clinical studies on PPAR γ activity in premature babies

3.1. Adiponectin concentrations in premature babies

Adiponectin is a robust biomarker of PPAR γ activity. Existing literature on plasma adiponectin concentrations in preterm newborn babies suggest that PPAR γ signalling may be underactive in preterm newborn babies. Kajantie et al. (2004) measured plasma adiponectin concentrations by ELISA in the cord vein of 197 infants [29]. Of them, 122 were born preterm (22 to 32 weeks' gestation), and 75 at term (49 from a healthy and 26 from a diabetic pregnancy, with similar findings, and thus all data from term infants were pooled). At birth, preterm babies have low plasma adiponectin concentrations ($n = 122$; $19.9 \pm 10.6 \mu\text{g/mL}$) when compared to term babies ($n = 75$; $33.7 \pm 13.6 \mu\text{g/mL}$) [29]. Mean adiponectin concentrations increased from less than 1 g/ml at 24 weeks' gestation to approximately 20 g/ml at term. Preterm females had 57% higher adiponectin concentrations (0 to 146%; $P = 0.05$) than preterm males. Adiponectin levels were lower in preterm infants with recent (<12 hours) exposure to maternal betamethasone but were unrelated to mode of delivery, preeclampsia, or impaired umbilical artery flow [29].

Hansen-Pupp et al. (2015) analysed adiponectin concentrations in cord blood at birth, and peripheral blood at 72 h, day 7, and then weekly until postmenstrual age of 40 weeks in 52 preterm babies born at 26 ± 1.9 weeks gestational age [30]. The mean adiponectin concentration increased from $6.8 \pm 4.4 \mu\text{g/mL}$ at 72 h to $37.4 \pm 22.2 \mu\text{g/mL}$ at three weeks. Mean adiponectin concentration during days 3 to 21 ($21.4 \pm 12 \mu\text{g/mL}$) correlated with gestational age at birth ($r = 0.46$, $P = 0.001$), birth weight ($r = 0.71$, $P = 0.001$) and birth weight Standard Deviation Score (SDS) ($r = 0.42$, $P = 0.003$). Furthermore, mean adiponectin concentration during days 3 to 21 correlated with weight SDS, length SDS, and head circumference SDS at 35 weeks corrected gestational age ($r = 0.62$, 0.65, and 0.62, respectively; all $P = 0.001$). Peak concentrations at 3 weeks of age ($n = 52$; $37.4 \pm 22.2 \mu\text{g/mL}$) did not correlate with gestational age but positively correlated with catch-up growth ($\beta = 0.021$; CI: 0.001–0.041; $P = 0.04$) [30].

3.2. PPAR γ signalling in brain imaging genetics studies

In a candidate gene study of thirteen genes from seventy-two preterm infants, Single Nucleotide Polymorphisms (SNPs) in the genes FADS2 and ARVCF were significantly associated with fractional anisotropy (FA) in white matter extracted by Tract Based Spatial Statistics (TBSS). FADS2 is involved with lipid metabolism, including PPAR signalling [31]. A pathway-based genome-wide imaging genomics analysis was carried out on the same cohort, using the Pathways sparse Reduced Rank Regression (sRRR) machine learning approach with genome-wide SNP (Single Nucleotide Polymorphisms) genotyping and a reduced version of the same TBSS phenotype. This showed that the PPAR signalling pathway was the top ranked pathway in the model including adjustment for both gestational age and post menstrual age [32].

A SNP-based genome-wide imaging genomics analysis was carried on a large independent cohort of 271 preterm infants, using sRRR method with genome-wide SNP genotyping and a probabilistic tractography phenotype incorporating FA. This detected an association between SNPs in the PPAR γ gene and the imaging phenotype fully adjusted for gestational age, post menstrual age and ancestry [33].

Meirhaeghe et al. (2007) genotyped two independent cross-sectional studies from Northern Ireland ($n = 382$ and 620) for Pro12Ala polymorphism of the PPAR γ 2 [34]. In combined populations, the PPAR γ 2 Ala12 allele was associated ($P = 0.03$) with lower birth weight, primarily caused by shorter gestational duration ($P = 0.04$). The frequency of Ala12 allele carriers was higher ($P = 0.027$) in the group of individuals born before term (35%, $n = 60$) than in the group of individuals born at term (22%, $n = 942$). The odds ratios (95% CI) of preterm birth for Ala12 allele carriers were 1.9 (1.1 - 3.4), $P = 0.022$, and 4.2 (1.9 - 9.7), $P = 0.0006$ (adjusted for sex, maternal age, and study), when considering 37 or 35 weeks of pregnancy as a threshold for preterm birth, respectively.

The association between PPAR γ 2 Pro12Ala polymorphism and neurodevelopment at 18–24 months of age was assessed in two groups of European infants, 155 born before 33 weeks of gestation

and 180 born later than 36 weeks of gestation [35]. The Ala allele of the Pro12Ala polymorphism was noted in 25% of the preterm infants and 20% of the term infants. The Ala allele of PPAR γ 2 was significantly associated with adverse cognitive ($P = 0.019$), language ($P = 0.03$), and motor development ($P = 0.036$) at 18 to 24 months of age after taking into consideration the duration of ventilation, gender, and index of multiple deprivation scores, but without correction for potential shared ancestry. There was no association between the PPAR γ 2 Pro12Ala polymorphism and neurodevelopment in term infants.

3.3. Clinical trials of PPAR γ agonists in preterm babies

No clinical trials have been conducted using PPAR γ agonists in preterm babies. Pioglitazone has been trialled in children and adults for its neuroprotective action and found to be safe. Improvements in behaviour were demonstrated on 4 to 12-year-old children ($n = 44$; 30mg OD) with Autistic Spectral Disorders after 10 weeks of treatment with pioglitazone [36].

4. Safety considerations for using pioglitazone in premature babies

The effect of pioglitazone on preterm lung disease have been demonstrated in animal models at plasma concentrations of $0.52 - 0.99 \pm 0.15 \mu\text{g/mL}$ [18]. This target plasma concentrations can be achieved at dose ranges currently licensed in adults for diabetic treatment. Efficacy in animal experiments has also been demonstrated with short course of treatment lasting up to 14 days [20]. Studies using samples of airway secretions have demonstrated that lung inflammation peaks between 7 and 10 days after birth in premature babies and is exacerbated by both antenatal and nosocomial infections [37]. Duration of treatment for two to three weeks from soon after birth may therefore be sufficient for the prevention of bronchopulmonary dysplasia in preterm babies. A much shorter duration of treatment in comparison to current licensed treatment for adult diabetes needs to be taken into consideration when evaluating the safety of pioglitazone in premature babies.

4.1. Increased insulin sensitivity

Pioglitazone hydrochloride is an antidiabetic agent that depends on the presence of insulin for its mechanism of action. Pioglitazone decreases insulin resistance in the peripheral muscle and liver, resulting in increased insulin-dependent glucose disposal and decreased hepatic glucose output, respectively. It is not an insulin secretagogue [38].

Hyperglycaemia is often seen in premature newborn babies and is likely to be due to decreased insulin production. Since pioglitazone enhances the effects of circulating insulin (by decreasing insulin resistance), it does not lower blood glucose in animal models that lack endogenous insulin. Hypoglycaemia is therefore not expected in preterm babies when treated with pioglitazone alone. However, there is a risk of hypoglycaemia in premature babies when combined with insulin treatment. Since the threshold for starting insulin treatment in premature babies is set around twelve mmol/L and hypoglycaemia is defined as less than 2.5 mmol/L, there is sufficient safety margin to titrate the insulin treatment to maintain normoglycemia.

4.2. Effect on lipid metabolism

In pharmacodynamic studies of both monotherapy and combination therapy, treatment with pioglitazone was associated with decreased blood free fatty acids and triglyceride concentrations and increased blood HDL-C concentrations [38]. Preterm infants on total parenteral nutrition often have increased level of blood triglyceride concentrations resulting in temporary cessation of lipid supplementation until triglycerides levels decrease [39]. A reduction in blood triglyceride concentrations with pioglitazone treatment may be beneficial to preterm babies, with improved lipid metabolism resulting in improved weight gain.

4.3. Weight gain

Pioglitazone is associated with weight gain after 24 weeks of treatment, possibly due to oedema [38]. In addition, pioglitazone significantly decreases visceral (abdominal) fat stores while increasing extra-abdominal fat. The reduction in visceral fat correlates with improved hepatic and peripheral tissue insulin sensitivity.

In preterm babies, peak adiponectin concentrations on day 21 after birth was associated with increased discharge weight, suggesting that weight gain may be due to reasons other than oedema in preterm babies [30]. Concerns related to excess weight gain or oedema may not be an issue with a short duration of treatment.

4.4. Rare side effects

Bladder cancer, fractures in women, and macular oedema have been reported rarely in adult diabetic patients [38]. These have been associated with prolonged treatment (> 1 year). Rare transient elevations in liver enzymes and very rare clinically apparent liver injury have been noted.

4.5. Reproduction toxicity in animals

Several animal studies have been conducted by Takeda Pharmaceuticals for the licensing of pioglitazone [38]. Pioglitazone is non-teratogenic in both rats and rabbits. High doses were associated with maternal toxicity (e.g., placental pathology, weight loss), implantation losses, reduced foetal size, live birth weight and survival ratios, and delayed pup development. It is suggested that the adverse effects on the offspring result from disturbances in maternal physiology rather than a direct toxic effect.

Six pregnant rats per group, received twice daily oral doses of 0, 50, 100, 200 or 400 mg/kg of pioglitazone on gestational days six through to 20 days. Based on the results of this study, the no observable adverse effect level (NOAEL) was determined to be less than 50 mg/kg pioglitazone.

Thirty-three to thirty-five rat pups received enteral doses of 0, 20, 40, 80 mg/kg/day of pioglitazone on gestational days six through to 17 days. The NOAEL for developmental toxicity was determined to be 20 mg/kg based on the reductions in embryo viability, litter size and pup survival in the 40 mg/kg group. The NOAEL for F₁ generation rats was 40 mg/kg based on reduced body weights for males and reduced body weight gains (gestational period) for females in the 80 mg/kg group.

Five pregnant rabbits received oral doses of 0, 5, 40, 100 or 400 mg/kg/day pioglitazone on gestational days six through to eighteen. The top dose was reduced to 160 mg/kg due to high mortality. Based on the results of this study, the NOAEL for maternal and developmental toxicity was determined to be closer to 40 mg/kg.

Fifteen to eighteen rabbits were given enteral doses of 0, 40, 80 or 160 mg/kg/day pioglitazone on gestational days six through to eighteen. Mild signs of maternal toxicity were noted at 80 and 160 mg/kg/day. The NOAEL for maternal toxicity was 40 mg/kg/day. Slight embryotoxicity was noted at 160 mg/kg/day.

A standard series of reproduction toxicity studies was conducted with pioglitazone in rats and rabbits. In fertility studies there was no effect on copulation, impregnation, or fertility index. Pioglitazone is non-teratogenic in both rats and rabbits. High doses were associated with maternal toxicity (e.g., placental pathology, weight loss), implantation losses, reduced foetal size, live birth weight and survival ratios, and delayed pup development. It is suggested that the adverse effects on the offspring result from disturbances in maternal physiology rather than a direct toxic effect. This is supported by differences in maternal and foetal NOAEL.

5. Pharmaceutical challenges in creating a formulation suited for premature babies

Pioglitazone is currently available as a licensed medicine in tablet form. A formulation suitable for administration to premature babies is not commercially available. The major challenge for developing a liquid formulation is that pioglitazone freebase and its hydrochloride salt, which is used

in pharmaceutical products, have extremely low aqueous solubilities (< 1 mg/mL). In animal experiments, pioglitazone has been dissolved in surfactant. This limits the duration of treatment as usually not more than two doses of surfactant are administered in preterm babies. Several solubility studies on pioglitazone have been reported. Most of the excipients used in these studies are not suitable for use in preterm babies [40-43]. The choice of excipients is restricted by the limited experience in the safety profiles of excipients in newborn babies and children. Furthermore, the concentration of these excipients cannot exceed the daily consumption limits set by regulatory bodies.

Creating a solution formulation of pioglitazone at a concentration that enables delivery of the dose within the fluid restrictions for pre-term babies is a pharmaceutical challenge, especially with the limited portfolio of co-solvents and solubilisers that have regulatory acceptance for use in children. The pioglitazone concentration must be such that dosing per kilogram body weight in extremely low birth weight babies (< 1000 grams birth weight) is not impractical due to too small measurable volumes and not too large a volume that bigger babies (> 1500 grams birth weight) cannot tolerate. This applies to both intravenous and oral preparations.

Different routes of administration, such as intravenous (injection), endotracheal (tube), inhalation (nebuliser) can be considered in the newborn period when infants are not tolerating enteral feeds. Oral formulations, e.g. solutions or suspensions, can be used for subsequent maintenance treatment when enteral feeds are tolerated. When considering oral formulations such as suspensions, the viscosity of the formulation and particle size needs to be sufficiently low to allow for administration through size five French gastric tubes.

Considerations for liquid pharmaceutical formulations include the chemical and microbiological stability. Chemical stability of the pioglitazone needs to be verified to remain within pharmacopeial limits for the developed formulation when stored in its final container under specified conditions, e.g. room or fridge temperature. Preservatives are normally required to assure the microbiological quality of multiple dose liquid preparations. However, for many preservatives there is still limited data regarding the levels of safe exposure in children of different ages. The need to preserve a paediatric preparation and the choice of the preservative system at the lowest concentration feasible should be justified in terms of benefit-risk balance.

6. Future Perspectives and Scope

There is an urgent need to develop treatment strategies that reduce devastating outcomes in premature babies. We propose that a PPAR γ agonist may reduce the incidence of death or BPD and improve neurodevelopment in extreme preterm babies. The availability of a safe, well-tolerated agonist for PPAR γ signalling, pioglitazone, allows the exploration of a repurposing strategy to test this approach in the clinic. We are currently developing formulation options for pioglitazone that will be suitable for administration to preterm babies. This will allow us to conduct early phase clinical trials to determine the safety and tolerance of pioglitazone in preterm newborn babies. In an online UK survey conducted by us in Jan 2023 of 207 parents who had a premature baby born before 28 weeks of gestational age, 94% agreed that preventing lung and brain disease is an important treatment outcome of preterm babies. Ninety-six percent stated they would have agreed to participate in a randomised, placebo controlled, early phase clinical trial of a prophylactic treatment even if their child may not have developed the disease. Furthermore, 74% said that the potential side effects of the treatment (e.g. oedema, abnormal liver function and increased risk of hypoglycaemia) would not have worried them.

7. Conclusions

Prematurity has a devastating impact on individuals, families and society. There is an urgent need to develop treatment strategies that improve the lung and brain outcomes in premature newborn babies. Pioglitazone, a PPAR γ agonist, may reduce the incidence of death or BPD and improve neurodevelopment in extreme preterm babies. A stable liquid formulation of pioglitazone suited for preterm babies needs to be developed before this treatment can progress to the clinic. The potential benefits are likely to outweigh any anticipated safety concerns.

Author Contributions: Conceptualization: SV; Writing—original draft preparation, SV; Writing—review and editing, AG, BF, ADE; funding acquisition, SV. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Medical Research Council Impact Acceleration Award, grant number MR/X502923/1

Institutional Review Board Statement: Not applicable

Informed Consent Statement: Not applicable

Data Availability Statement: Not applicable

Acknowledgments: Nil

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Blencowe, H.; Cousens, S.; Oestergaard, M. Z.; Chou, D.; Moller, A. B.; Narwal, R.; Adler, A.; Vera Garcia, C.; Rohde, S.; Say, L.; et al. National, Regional, and Worldwide Estimates of Preterm Birth Rates in the Year 2010 with Time Trends since 1990 for Selected Countries: A Systematic Analysis and Implications. *The Lancet* **2012**, *379*, 2162–2172. [https://doi.org/10.1016/S0140-6736\(12\)60820-4](https://doi.org/10.1016/S0140-6736(12)60820-4).
2. Dassios, T.; Greenough, A. Long-Term Sequelae of BPD. In *Respiratory Diseases of the Newborn Infant (ERS monograph)*; Sinha IP, Bhatt JM, Cleator A, Eds.; European Respiratory Society: Sheffield, 2021; pp 68–78.
3. Harris, C.; Bisquera, A.; Zivanovic, S.; Lunt, A.; Calvert, S.; Marlow, N.; Peacock, J. L.; Greenough, A. Postnatal Dexamethasone Exposure and Lung Function in Adolescents Born Very Prematurely. *PLoS One* **2020**, *15*, e0237080. <https://doi.org/10.1371/journal.pone.0237080>.
4. Gou, X.; Yang, L.; Pan, L.; Xiao, D. Association between Bronchopulmonary Dysplasia and Cerebral Palsy in Children: A Meta-Analysis. *BMJ Open* **2018**, *8*, e020735. <https://doi.org/10.1136/bmjopen-2017-020735>.
5. Hack, M. Adult Outcomes of Preterm Children. *J Dev Behav Pediatr* **2009**, *30*, 460–470. <https://doi.org/10.1097/DBP.0b013e3181ba0fba>.
6. Johnson, S.; Wolke, D. Behavioural Outcomes and Psychopathology during Adolescence. *Early Human Development* **2013**, *89*, 199–207. <https://doi.org/10.1016/j.earlhumdev.2013.01.014>.
7. Nosarti, C.; Reichenberg, A.; Murray, R. M.; Cnattingius, S.; Lambe, M. P.; Yin, L.; MacCabe, J.; Rifkin, L.; Hultman, C. M. Preterm Birth and Psychiatric Disorders in Young Adult Life. *Arch Gen Psychiatry* **2012**, *69*, E1–8. <https://doi.org/10.1001/archgenpsychiatry.2011.1374>.
8. Schmidt, B.; Roberts, R. S.; Davis, P.; Doyle, L. W.; Barrington, K. J.; Ohlsson, A.; Solimano, A.; Tin, W. Long-Term Effects of Caffeine Therapy for Apnea of Prematurity. *N Engl J Med* **2007**, *357*, 1893–902. <https://doi.org/10.1056/NEJMoa073679>.
9. Villapol, S. Roles of Peroxisome Proliferator-Activated Receptor Gamma on Brain and Peripheral Inflammation. *Cellular and Molecular Neurobiology* **2018**, *38*, 121–132. <https://doi.org/10.1007/s10571-017-0554-5>.
10. Warden, A.; Truitt, J.; Merriman, M.; Ponomareva, O.; Jameson, K.; Ferguson, L. B.; Mayfield, R. D.; Harris, R. A. Localization of PPAR Isotypes in the Adult Mouse and Human Brain. *Sci Rep* **2016**, *6*, 27618. <https://doi.org/10.1038/srep27618>.
11. Hwang, S. J.; Kim, J. H.; Shim, J. W.; Kim, D. S.; Jung, H. L.; Park, M. S.; Lee, W. Y.; Kim, S. Y.; Shim, J. Y. Peroxisome Proliferator-Activated Receptor-Gamma Expression in the Lung Tissue of Obese Rats. *Yonsei Med J* **2011**, *52*, 495–501. <https://doi.org/10.3349/ymj.2011.52.3.495>.
12. Wei, J.; Bhattacharyya, S.; Varga, J. Peroxisome Proliferator-Activated Receptor γ : Innate Protection from Excessive Fibrogenesis and Potential Therapeutic Target in Systemic Sclerosis. *Curr Opin Rheumatol* **2010**, *22*, 671–676. <https://doi.org/10.1097/BOR.0b013e32833de1a7>.
13. Simon, D. M.; Arikian, M. C.; Srisuma, S.; Bhattacharya, S.; Tsai, L. W.; Ingenito, E. P.; Gonzalez, F.; Shapiro, S. D.; Mariani, T. J. Epithelial Cell PPAR γ Contributes to Normal Lung Maturation. *FASEB J* **2006**, *20*, 1507–9. <https://doi.org/10.1096/fj.05-5410fje>.

14. Wang, Y.; Santos, J.; Sakurai, R.; Shin, E.; Cerny, L.; Torday, J. S.; Rehan, V. K. Peroxisome Proliferator-Activated Receptor γ Agonists Enhance Lung Maturation in a Neonatal Rat Model. *Pediatr Res* **2009**, *65*, 150–155. <https://doi.org/10.1203/PDR.0b013e3181938c40>.
15. Rehan, V. K.; Wang, Y.; Patel, S.; Santos, J.; Torday, J. S. Rosiglitazone, a Peroxisome Proliferator-Activated Receptor- γ Agonist, Prevents Hyperoxia-Induced Neonatal Rat Lung Injury in Vivo. *Pediatr Pulmonol* **2006**, *41*, 558–69. <https://doi.org/10.1002/ppul.20407>.
16. Lee, C.; Sakurai, R.; Shin, E.; Wang, Y.; Liu, J.; Rehan, V. K. Antenatal PPAR- γ Agonist Pioglitazone Stimulates Fetal Lung Maturation Equally in Males and Females. *Am J Physiol Lung Cell Mol Physiol* **2020**, *319*, L435–L443. <https://doi.org/10.1152/ajplung.00376.2018>.
17. Richter, J.; Toelen, J.; Nagatomo, T.; Jimenez, J.; Vanoirbeek, J.; Deprest, J. Transplacental Administration of Rosiglitazone Attenuates Hyperoxic Lung Injury in a Preterm Rabbit Model. *Fetal Diagn Ther* **2016**, *39*, 297–305. <https://doi.org/10.1159/000439199>.
18. Morales, E.; Sakurai, R.; Husain, S.; Paek, D.; Gong, M.; Ibe, B.; Li, Y.; Husain, M.; Torday, J. S.; Rehan, V. K. Nebulized PPAR γ Agonists: A Novel Approach to Augment Neonatal Lung Maturation and Injury Repair in Rats. *Pediatr Res* **2014**, *75*, 631–640. <https://doi.org/10.1038/pr.2014.8>.
19. Sakurai, R.; Lee, C.; Shen, H.; Waring, A. J.; Walther, F. J.; Rehan, V. K. A Combination of the Aerosolized PPAR- γ Agonist Pioglitazone and a Synthetic Surfactant Protein B Peptide Mimic Prevents Hyperoxia-Induced Neonatal Lung Injury in Rats. *Neonatology* **2018**, *113*, 296–304. <https://doi.org/10.1159/000486188>.
20. Lee, H. J.; Lee, Y. J.; Choi, C. W.; Lee, J. A.; Kim, E. K.; Kim, H. S.; Kim, B. Il; Choi, J. H. Rosiglitazone, a Peroxisome Proliferator-Activated Receptor- γ Agonist, Restores Alveolar and Pulmonary Vascular Development in a Rat Model of Bronchopulmonary Dysplasia. *Yonsei Med J* **2014**, *55*, 99–106. <https://doi.org/10.3349/ymj.2014.55.1.99>.
21. Kaplan, J.; Nowell, M.; Chima, R.; Zingarelli, B. Pioglitazone Reduces Inflammation through Inhibition of NF-KB in Polymicrobial Sepsis. *Innate Immun* **2014**, *20*, 519–528. <https://doi.org/10.1177/1753425913501565>.
22. Vallée, A.; Lecarpentier, Y.; Guillevin, R.; Vallée, J. N. Demyelination in Multiple Sclerosis: Reprogramming Energy Metabolism and Potential PPAR γ Agonist Treatment Approaches. *Int J Mol Sci* **2018**, *19*, 1212. <https://doi.org/10.3390/ijms19041212>.
23. Zaghloul, N.; Kurepa, D.; Bader, M. Y.; Nagy, N.; Ahmed, M. N. Prophylactic Inhibition of NF-KB Expression in Microglia Leads to Attenuation of Hypoxic Ischemic Injury of the Immature Brain. *J Neuroinflammation* **2020**, *17*, 365. <https://doi.org/10.1186/s12974-020-02031-9>.
24. Yeh, J. H.; Wang, K. C.; Kaizaki, A.; Lee, J. W.; Wei, H. C.; Tucci, M. A.; Ojeda, N. B.; Fan, L. W.; Tien, L. T. Pioglitazone Ameliorates Lipopolysaccharide-Induced Behavioral Impairment, Brain Inflammation, White Matter Injury and Mitochondrial Dysfunction in Neonatal Rats. *Int J Mol Sci* **2021**, *22*, 6306. <https://doi.org/10.3390/ijms22126306>.
25. Zhang, Y. L.; Wang, R. B.; Li, W. Y.; Xia, F. Z.; Liu, L. Pioglitazone Ameliorates Retinal Ischemia/Reperfusion Injury via Suppressing NLRP3 Inflammasome Activities. *Int J Ophthalmol* **2017**, *10*, 1812–1818. <https://doi.org/10.18240/ijo.2017.12.04>.
26. Xie, E. F.; Hilkert Rodriguez, S.; Xie, B.; D’Souza, M.; Reem, G.; Sulakhe, D.; Skondra, D. Identifying Novel Candidate Compounds for Therapeutic Strategies in Retinopathy of Prematurity via Computational Drug-Gene Association Analysis. *Front Pediatr* **2023**, *11*, 1151239. <https://doi.org/10.3389/fped.2023.1151239>.
27. Baregamian, N.; Mourot, J. M.; Ballard, A. R.; Evers, B. M.; Chung, D. H. PPAR- γ Agonist Protects against Intestinal Injury during Necrotizing Enterocolitis. *Biochem Biophys Res Commun* **2009**, *379*, 423–427. <https://doi.org/10.1016/j.bbrc.2008.11.155>.
28. Corsini, I.; Polvani, S.; Tarocchi, M.; Tempesti, S.; Marroncini, G.; Generoso, M.; Bresci, C.; Gozzini, E.; Bianconi, T.; Galli, A.; et al. Peroxisome Proliferator-Activated Receptor- γ Agonist Pioglitazone Reduces the Development of Necrotizing Enterocolitis in a Neonatal Preterm Rat Model. *Pediatr Res* **2017**, *81*, 364–368. <https://doi.org/10.1038/pr.2016.214>.
29. Kajantie, E.; Hytinantti, T.; Hovi, P.; Andersson, S. Cord Plasma Adiponectin: A 20-Fold Rise between 24 Weeks Gestation and Term. *J Clin Endocrinol Metab* **2004**, *89*, 4031–4036. <https://doi.org/10.1210/jc.2004-0018>.
30. Hansen-Pupp, I.; Hellgren, G.; Hård, A. L.; Smith, L.; Hellström, A.; Löfqvist, C. Early Surge in Circulatory Adiponectin Is Associated with Improved Growth at near Term in Very Preterm Infants. *J Clin Endocrinol Metab* **2015**, *100*, 2380–2387. <https://doi.org/10.1210/jc.2015-1081>.

31. Boardman, J. P.; Walley, A.; Ball, G.; Takousis, P.; Krishnan, M. L.; Hughes-Carre, L.; Aljabar, P.; Serag, A.; King, C.; Merchant, N.; et al. Common Genetic Variants and Risk of Brain Injury After Preterm Birth. *Pediatrics* **2014**, *133*, e1655–1663. <https://doi.org/10.1542/peds.2013-3011>.
32. Krishnan, M. L.; Wang, Z.; Silver, M.; Boardman, J. P.; Ball, G.; Counsell, S. J.; Walley, A. J.; Montana, G.; Edwards, A. D. Possible Relationship between Common Genetic Variation and White Matter Development in a Pilot Study of Preterm Infants. *Brain Behav* **2016**, *6*, e00434. <https://doi.org/10.1002/brb3.434>.
33. Krishnan, M. L.; Wang, Z.; Aljabar, P.; Ball, G.; Mirza, G.; Saxena, A.; Counsell, S. J.; Hajnal, J. V.; Montana, G.; Edwards, A. D. Machine Learning Shows Association between Genetic Variability in PPARG and Cerebral Connectivity in Preterm Infants. *Proc Natl Acad Sci U S A* **2017**, *114*, 13744–13749. <https://doi.org/10.1073/pnas.1704907114>.
34. Meirhaeghe, A.; Boreham, C. A. G.; Murray, L. J.; Richard, F.; Smith, G. D.; Young, I. S.; Amouyel, P. A Possible Role for the PPARG Pro12Ala Polymorphism in Preterm Birth. *Diabetes* **2007**, *56*, 494–498. <https://doi.org/10.2337/db06-0915>.
35. Victor, S.; Chew, A.; Falconer, S. Pro12Ala Polymorphism of Peroxisome Proliferator Activated Receptor Gamma 2 May Be Associated with Adverse Neurodevelopment in European Preterm Babies. *Brain Behav* **2021**, *11*, e2256. <https://doi.org/10.1002/brb3.2256>.
36. Ghaleiha, A.; Rasa, S. M.; Nikoo, M.; Farokhnia, M.; Mohammadi, M. R.; Akhondzadeh, S. A Pilot Double-Blind Placebo-Controlled Trial of Pioglitazone as Adjunctive Treatment to Risperidone: Effects on Aberrant Behavior in Children with Autism. *Psychiatry Res* **2015**, *229*, 181–187. <https://doi.org/10.1016/j.psychres.2015.07.043>.
37. Davies, P. L.; Spiller, O. B.; Beeton, M. L.; Maxwell, N. C.; Remold-O'Donnell, E.; Kotecha, S. Relationship of Proteinases and Proteinase Inhibitors with Microbial Presence in Chronic Lung Disease of Prematurity. *Thorax* **2010**, *65*, 256–51. <https://doi.org/10.1136/thx.2009.116061>.
38. US Food and Drug Administration. Drug Approval Package Actos (Pioglitazone Hydrochloride) Tablets Home Page. Available online: https://www.accessdata.fda.gov/drugsatfda_docs/nda/99/021073A_Actos.cfm (accessed on 29 August 2023).
39. Sinclair, R.; Schindler, T.; Lui, K.; Bolisetty, S. Hypertriglyceridaemia in Extremely Preterm Infants Receiving Parenteral Lipid Emulsions. *BMC Pediatr* **2018**, *18*, 348. <https://doi.org/10.1186/s12887-018-1325-2>.
40. Jouyban, A.; Soltanpour, S. Solubility of Pioglitazone Hydrochloride in Binary and Ternary Mixtures of Water, Propylene Glycol, and Polyethylene Glycols 200, 400, and 600 at 298.2 K. *AAPS PharmSciTech* **2010**, *11*, 1713–1717. <https://doi.org/10.1208/s12249-010-9551-4>.
41. Seedher, N.; Kanojia, M. Micellar Solubilization of Some Poorly Soluble Antidiabetic Drugs: A Technical Note. *AAPS PharmSciTech* **2008**, *9*, 431–436. <https://doi.org/10.1208/s12249-008-9057-5>.
42. Seedher, N.; Kanojia, M. Co-Solvent Solubilization of Some Poorly-Soluble Antidiabetic Drugs Solubilization Antidiabetic Drugs. *Pharm Dev Technol* **2009**, *14*, 185–192. <https://doi.org/10.1080/10837450802498894>.
43. Pandit, V.; Gorantla, R.; Devi, K.; Pai, R. S.; Sarasija, S. Preparation and Characterization of Pioglitazone Cyclodextrin Inclusion Complexes. *J Young Pharm* **2011**, *3*, 267–274. <https://doi.org/10.4103/0975-1483.90234>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.