

Review

Not peer-reviewed version

Neurologic Injury and Post-Traumatic Stress Disorder

[Wiley Braxton Gillam V](#) , [Nikhil Godbole](#) , Shourya Sangam , Alyssa DeTommaso , [Marco Foreman](#) , [Brandon Lucke-Wold](#) *

Posted Date: 4 September 2023

doi: 10.20944/preprints202309.0100.v1

Keywords: Post-Traumatic Stress Disorder, neurologic injury, traumatic brain injury, subarachnoid hemorrhage, stroke



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

Neurologic Injury and Post-Traumatic Stress Disorder

Wiley Gillam ¹, Nikhil Godbole ², Shourya Sangam ³, Alyssa DeTommaso ⁴, Marco Foreman ¹ and Brandon Lucke-Wold ^{5,*}

¹ College of Medicine, University of Florida, Gainesville, FL 32610, USA

² School of Medicine, Tulane University, New Orleans, LA 70112, USA

³ College of Liberal Arts and Sciences, University of Florida, Gainesville, FL 32603, USA

⁴ College of Medicine, University of Central Florida, Orlando, FL 32827, USA

⁵ Department of Neurosurgery, University of Florida, Gainesville, FL 32610, USA

* Correspondence: correspondence: brandon.lucke-wold@neurosurgery.ufl.edu

Abstract: The present review aimed to identify through what means neurologic injury can predispose individuals to Post-Traumatic Stress Disorder (PTSD). In recent years, comprehensive studies have helped to clarify which structures in the central nervous system (CNS) that, when damaged, lead to distinct PTSD symptoms—namely dissociative reactions or flashbacks. Our review narrowed its focus to three common neurologic injuries including traumatic brain injury (TBI), subarachnoid hemorrhage (SAH), and stroke. Beyond discussing the potential mechanisms by which neurotrauma may lead to PTSD, we summarized our current understanding of the pathophysiology of the disorder and discussed predicted associations between the limbic system and PTSD. In particular, the role of noradrenergic neuromodulatory signaling on the HPA axis as it pertains to fear memory recall needs to be further explored to better understand its effects on limbic structures in PTSD patients. This paper also described how damage to the CNS can alter the function of the limbic system and result in characteristic symptoms of PTSD like intrusive memories and acute psychological distress. Further, our review intended to draw attention to a lack of PTSD prognosis in TBI, SAH, and stroke patients who could benefit from early treatment. PTSD symptoms often compound with pre-existing issues further deteriorating health outcomes for these patients. It is ultimately our goal to clarify the relationship between neurotrauma and PTSD such that earlier diagnoses and appropriate treatment are observed in clinic.

Keywords: Post-Traumatic Stress Disorder; neurologic injury; traumatic brain injury; subarachnoid hemorrhage; stroke

1. Introduction: Neurotrauma Preceding Post-Traumatic Stress Disorder

Post-Traumatic Stress Disorder (PTSD) is characterized by a wide-ranging set of symptoms including but not limited to intrusive and distressing memories, dissociative reactions, and serious psychological distress in response to stimuli that resemble a previously experienced traumatic event [1]. Common to all individuals diagnosed with PTSD is exposure to trauma. However, this trauma may be physical or emotional, suggesting that the disorder does not necessitate neurologic injury. Though it may not be inherent to all cases, many studies indicate a correlation between traumatic brain injury (TBI) and PTSD [2,3,4,5,6]. One meta-analysis found a 13.5% prevalence rate of PTSD in civilians who had been diagnosed with mild traumatic brain injury (mTBI) [2]. Another study found that greater than 17% of United States Army soldiers with mTBI returning from Operation Enduring Freedom (OEF) in Afghanistan and Operation Iraqi Freedom (OIF) in Iraq screened positive for PTSD [4].

In addition to TBI, stroke and subarachnoid hemorrhage may predispose to PTSD. Several studies document the development of PTSD symptoms in individuals following stroke [7,8,9]. These symptoms can be assessed using several measures. The present review focused primarily on studies

which utilized the gold-standard technique, Clinician Administered PTSD Scale (CAPS) as found in the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition* (DSM-V) [7,10]. A significant proportion of stroke and subarachnoid hemorrhage patients presented with PTSD symptoms a year following the event suggesting PTSD treatment may prove beneficial for such individuals [7,10]. It should be noted that some of the studies included in the present review used DSM-IV which arguably has a less restrictive concept of PTSD. One significant discrepancy can be found where DSM-V defines trauma thus precluding any subjective interpretations of the type of events – actual or threatened death, sexual violence, or serious injury – which must precede diagnosis [11].

Although it appears evident that neurotrauma can be involved in the development of PTSD in patients, it is unclear how the extent of an injury affects symptoms. One study claims that PTSD severity scales with the level of involvement in combat events, indicating the possibility of a dosage-dependent relationship between trauma exposure and the disorder [2]. An alternative explanation is that increased exposure to combat events increases the likelihood of a more serious neurologic injury thereby worsening PTSD symptoms. However, there is little evidence for this theory. In fact, some data suggests that mild TBI (mTBI) is more predictive of PTSD symptoms than moderate to severe TBI [12]. There are two important considerations to make when trying to interpret this result: 1) The sample sizes for most studies which have drawn this conclusion have been quite small and may not accurately represent the large population of patients suffering from both TBI and PTSD. 2) Moderate to severe TBI is associated with longer durations of post-traumatic amnesia which is inversely related to intrusive memories due to limited trauma memory encoding. Together, these factors further complicate our current understanding of how neurotrauma may be linked to PTSD.

2. The Pathophysiology of Post-Traumatic Stress Disorder

The major brain structures which have proved to play some role in PTSD symptoms include the prefrontal cortex (PFC), hippocampus, amygdala, and other downstream elements such as the locus coeruleus (LC) and hypothalamus [13,14,15]. A critical component of the prefrontal cortex is negatively regulating amygdala function to prevent hyperactivity. It has been found that PTSD patients exhibit reduced activity in the medial prefrontal cortex (mPFC) and anterior cingulate cortex (ACC) during presentation of trauma-related and non-related aversive stimuli [16,17]. Further, they have been observed with decreased volume in the ventromedial prefrontal cortex (vmPFC) and ACC [16,17,18,19,20], and even to utilize altered pathways from the ACC to the amygdala [21]. All these changes in the PFC suggest that amygdala dysregulation may be involved in the pathophysiology of PTSD by erasure of the conventionally strict regulation of fear responses [13].

Noradrenaline is a neuromodulatory neurotransmitter system capable of producing a rapid and coordinated response to acute stress. Studies have found increased levels of urinary noradrenaline (NA) and plasma NA in PTSD patients compared to controls indicating the likelihood of increased noradrenergic tone in these individuals [22,23,24,25]. The first convincing evidence, however, was found when concentrations of NA in hourly CSF samples (collected over a 6h period) were found to be higher in male combat veterans compared to healthy controls [26]. Based on these findings, Pietrzak and colleagues conducted a study to determine whether a reduction in the Norepinephrine Transporter (NET) could be responsible for the increased noradrenergic tone in PTSD patients [27]. The researchers found evidence for a significant reduction in the density of NET labeled in the LC in PTSD patients compared to controls [27]. It is important to note that this article has since been retracted. As such, further research is still needed to determine the legitimacy of these findings.

As previously mentioned, changes in the PFC found in PTSD patients have downstream impacts on the amygdala. Specifically, it has been found that the amygdala in those with PTSD is hyperactive during exposure to trauma-linked events [28]. A critical function of the amygdala is to stimulate the hippocampus during the formation of new memories related to fear-inducing events. A lack of proper control of the amygdala provides a possible explanation why PTSD patients experience intrusive memories from prior trauma. Although the results were similar for both trauma-exposed individuals and PTSD patients, meta-analyses observed statistically significant differences between these two groups and healthy controls [29]. Reduced cortisol levels and signaling in PTSD patients could be

responsible for stronger sympathetic nervous system activation and extreme lucidity of traumatic memories [30,31,32,33]. Notably, Robert Sapolsky claimed that under stress, the body releases glucocorticoids resulting in downregulation of glucocorticoid receptors thereafter leading to hippocampal atrophy in response to chronic stress [34]. Only a decade later, he described the relevance this theory has to PTSD whereby hyperactivation of this stress system in response to trauma could lead to cell death in the hippocampus and eventually cognitive dysfunction [35]. Particularly compelling evidence for hippocampal involvement in PTSD was found by several meta-analyses reporting a significant reduction in hippocampal volume in PTSD patients as compared to both healthy controls and trauma-exposed individuals without PTSD [36,37].

Norepinephrine and indirect signaling from the limbic system including the hippocampus, mPFC, and amygdala act on the paraventricular nucleus of the hypothalamus to trigger the release of corticotropin releasing hormone, or CRH [38]. The hypothalamic pituitary adrenal (HPA) axis which is initiated by CRH release is the single most critical hormonal facilitator of stress response in humans. Rodent models have demonstrated that PTSD-associated behaviors are observed in response to CRH injection [39] and that CRH receptor binding in the amygdala can produce fear responses [40]. Despite substantial evidence implicating CRH to high stress states and PTSD symptoms, results from other studies complicate the matter. Notably, CRH release in the bed nucleus of the stria terminalis has an inhibitory effect on neurons in the amygdala which results in a decreased fear response [41]. This clearly contradicts the idea that CRH release necessitates amygdala excitation. It is possible that CRH differentially affects the amygdala according to the pathway it uses but further evidence is needed to confirm this theory.

Several studies have also looked downstream of CRH to determine whether cortisol is involved in the development of PTSD. It has been found that lower concentrations of hair cortisol in military veterans prior to overseas deployment was positively correlated with PTSD symptoms post-deployment [42]. Likewise, police academy recruits with reduced increases in cortisol while watching videos of traumatic situations were more likely to demonstrate signs of long-term distress and decreased resilience after four years [43]. Together, these findings seem to indicate that lower cortisol concentrations predict PTSD symptoms following exposure to trauma. However, a meta-analysis demonstrated that heart rate was the only biological measure obtained post-trauma that was capable of predicting PTSD [44]. Additionally, even if low cortisol concentrations are associated with PTSD, no threshold cortisol concentration has been identified leaving little clinical applicability. It is evident that more research is needed to clarify the relationship between the HPA axis and PTSD. Figure 1 is a greatly simplified model that serves to clarify the complex pathophysiology of PTSD but it should be noted that the relationship between these neurological structures and PTSD is still widely contested.




Figure 1. Chronic or high acute stress experienced during and after trauma may directly and indirectly act on several limbic structures. (1) depicts altered regulation of the amygdala which may induce fearful behaviors observed in those with PTSD. Stress may also lead to hippocampal atrophy which could be responsible for many symptoms of PTSD including cognitive dysfunction, involuntary recurrent memories (flashbacks), and more (2). (3) describes how limbic structures such as the hippocampus or amygdala may act on the hypothalamus to activate the HPA axis by CRH release. These neuroendocrine disruptions in the body's primary stress response mechanism could also induce fearful behaviors.

3. TBI

Traumatic brain injury (TBI) is defined as the alteration of brain physiology and function secondary to an external force [45]. TBIs can range from mild, often referred to as "concussions", to moderate and severe forms, each bearing its distinguishing features and implications [46]. TBI has emerged as a significant area of investigation with respect to understanding its role in rendering individuals more susceptible to developing PTSD. Studies exploring the connection between TBI and PTSD in different populations have provided valuable insights into how TBI could affect the emergence and comorbidity of PTSD symptoms. A study published in 2008 by Hoge et al. found that mTBI experienced by soldiers deployed in Iraq was strongly associated with PTSD and physical health problems. This study suggests addressing PTSD is critical for improving the physical and psychological well-being of soldiers with TBI [47]. Another study shows that when black women experienced TBI as a result of intimate partner violence, there was a subsequent increase in the presence of both PTSD and depression. After accounting for other factors, the statistical analysis shows that the occurrence of TBI was associated with an increase in comorbid PTSD and depression symptoms [48]. In terms of children and adolescents, one study found that after experiencing a severe TBI, the percentage of those with PTSD increased to 13%, which is significantly higher than the general child population's lifetime prevalence of 7.8%. This difference in PTSD prevalence between children with severe TBI and those without suggests a strong correlation between TBI and the subsequent development of PTSD in children and adolescents [49].

Several potential mechanisms could explain why TBI might predispose different populations of individuals to PTSD. One theory is that TBI can lead to damage to the brain regions involved in emotion regulation, such as the amygdala and hippocampus, which are known to play a role in the development of PTSD [50]. TBI can damage the hippocampus, which may lead to impaired memory and emotion processing, both of which have been linked to PTSD [51]. Additionally, TBI has been

associated with altered activity in the amygdala, a brain region involved in the recognition and processing of fear, which is thought to play a key role in the development of PTSD [52]. Moreover, TBI can lead to dysregulation of neurotransmitters such as serotonin and norepinephrine, which have also been implicated in the development of PTSD [53]. Lastly, TBI can also lead to damage to the frontal lobe, which is involved in emotional regulation and impulse control, which could potentially contribute to the development of PTSD [52].




Figure 2. (A) Anatomical representation of the limbic system, which is composed of delicate neuronal structures including the anterior cingulate gyrus, thalamus, hippocampus, amygdala, and prefrontal cortex, among others. The limbic system plays vital roles related to memory, learning, and behavior through a complex interplay among the aforementioned structures. (B) Representation of these structures and their subsequent dysregulation following traumatic brain injury (TBI), which can be delivered in the form of external force and/or psychological insult. .

Expanding on the relationship between TBI and PTSD, another potential mechanism centers around how TBI can lead to feelings of vulnerability and powerlessness. The physical injury itself, resulting from a forceful impact on the head, can create a profound sense of helplessness in individuals who cannot care for themselves. This reliance on others for basic needs can exacerbate feelings of vulnerability and a loss of autonomy. TBI can also cause cognitive changes, which can disrupt a person's sense of capability and self-assurance. This emotional response, originating from the very nature of the trauma, could establish a foundation that strengthens the risk of developing PTSD [54]. In military personnel who undergo combat-related trauma, the added layer of exposure to life-threatening situations and witnessing the traumatic experiences of others can heighten these feelings of vulnerability and powerlessness, strengthening the link between TBI and the susceptibility to PTSD. For these individuals, the combination of TBI and the psychological complications of combat stressors becomes a convoluted interplay that shapes their psychological response. Furthermore, people with a history of emotional and physical abuse may also be more susceptible to developing PTSD following TBI due to their pre-existing issues with emotional regulation and stress management [55]. This means that TBI may exacerbate already present risk factors, making it more likely for individuals to develop PTSD. In essence, what occurs is a dynamic where TBI not only influences cognitive aspects but also intersects with pre-existing vulnerabilities. By heightening feelings of vulnerability and exacerbating pre-existing emotional regulation issues, TBI can act as an accelerant for the development of PTSD.

4. Subarachnoid Hemorrhage

Subarachnoid hemorrhage (SAH) is a severe neurological condition that is characterized by spontaneous bleeding into the subarachnoid space, which consists of the cerebrospinal fluid (CSF), major blood vessels, and cisterns [56]. In 80% of cases of SAH without preceding trauma, this begins with the rupture of an intracranial aneurysm, with other causes including vascular malformations [57,58]. The major presenting symptom of SAH is the onset of a sudden, severe headache that reaches

peak intensity within seconds [59]. The death rate of SAH is approximately 50% and this includes up to 18% of all aneurysmal SAH patients who die at home or in transit to a hospital [60]. About half of the surviving patients suffer from long-term effects and experience a subsequent decreased quality of life due to a significant risk of developing physical, memory, and neurocognitive impairments [61,62,63].




Figure 3. Subarachnoid hemorrhage (SAH) is caused in most cases by the rupture of an intracranial aneurysm, resulting in bleeding into the subarachnoid space (the space between the arachnoid mater and pia mater). 50% of SAH patients survive, of which half suffer from long term effects such as physical, memory, and neurocognitive impairment as well as potential PTSD symptoms.

Based on current clinical studies, the best predictor of a patient's neurological outcome is the level of impairment and extent of subarachnoid bleeding at the time of hospital admission [63,64]. A study that followed 415 eligible patients with SAH over a 90-day period after hospital discharge in Vietnam concluded that the Prognosis on Admission of Aneurysmal Subarachnoid Hemorrhage (PAASH) grading scale was most reliable in predicting poor outcomes [61]. However, predicting the overall outcomes of SAH remains imperfect because of the high variability of the clinical conditions, complications, and treatments administered [65,66,67]. For example, in the aforementioned study, a number of the patients experienced complications such as pneumonia and rebleeding, which greatly contributed to a high rate of poor outcomes [61]. Nonetheless, grading SAH patients upon admission is crucial to predicting their short and long-term health outcomes.

Emerging research findings suggest that survivors of SAH may be at higher risk of developing Post-Traumatic Stress Disorder (PTSD). The clinical symptoms and postictal events of aneurysmal SAH can be a significantly traumatic experience that leads to a PTSD diagnosis for some [68,69]. In a study that conducted PTSD assessments on 105 SAH patients at 3- and 13-months post-ictus, 37% of the patients met the requirements of a PTSD diagnosis during both assessment periods, which is four times the rate of PTSD in the general population [70]. The patients of this study also reported elevated

levels of fatigue and increased problems with their sleep, both of which are linked to PTSD. However, this study failed to demonstrate a clear pattern of PTSD symptoms over the course of data collection. Existing models have consistently shown differences in PTSD symptoms when the disorder arises from a life-threatening medical event versus a past traumatic external event, such as combat [71]. Furthermore, demographic factors like age, sex, educational level, and cognitive ability also affect individual PTSD symptoms [72,73,74,75,76].

In a study that examined the specific PTSD symptoms of aneurysmal SAH patients over the course of one year following the event, a variety of PTSD symptom patterns was observed. Researchers noted: about 50% of the patients did not show symptoms suggesting PTSD at any point, 15% showed recovery following PTSD symptoms at the first assessment, 9% exhibited symptoms suggesting PTSD during all assessments, and 13% experienced delayed symptoms [77]. The study's findings were, however, dependent on a self-report questionnaire used to evaluate PTSD symptoms (not as accurate as PTSD diagnosis by formal psychiatric review) and a multilevel analysis that could not completely explain differences in the course of PTSD [77]. Nevertheless, the finding that half of the patients did not experience any PTSD symptoms in the first year post SAH is consistent with similar studies of SAH patients [70,78,79].

These results reinforce that a majority of SAH patients do not initially show symptoms indicative of PTSD yet later receive a diagnosis. Thus, it is imperative to evaluate these individuals for PTSD symptoms throughout the first year following SAH diagnosis. Despite the increased availability of treatment options, up to 70% of individuals experiencing PTSD symptoms still do not pursue professional care for their condition [80]. Further research must be conducted to better understand which SAH patients are likely to experience chronic PTSD. Particularly, studies that can sufficiently account for factors such as coping style, psychiatric history, and levels of social support would help to better specify the differences in variation of PTSD symptoms over time.

5. Stroke

Ischemic and hemorrhagic strokes account for 11.8% of deaths worldwide [81]. Like PTSD, a stroke is unexpected, uncontrollable, and life-threatening. A meta-analysis of 1,138 participants suggested 25% of stroke survivors will develop posttraumatic stress symptoms and 1 in 9 will develop chronic PTSD [82]. Stroke-induced PTSD has been linked with worse long-term stroke outcomes, including recurrent stroke [82], greater disability, and nonadherence to medications [83]. This may result in poor compliance with medications intended for secondary prevention of recurrent stroke. Non-compliance with these medications may suggest that stroke-induced PTSD could be a novel risk factor for recurrent stroke. Stroke-associated PTSD is characterized by intrusive symptoms such as flashbacks of the stroke/TIA, persistent avoidance of stimuli associated with the stroke, altered arousal, increased irritability, and negative alterations in cognition and mood [84]. The literature suggests that the presence of these mental health disorders has a large impact on quality of life and functional outcomes after stroke. Stein et al. found that in a population of mild stroke survivors with no measurable differences in neurologic deficits, patients with PTSD had significantly reduced quality of life scores [85]. There is also evidence post-stroke psychiatric conditions may increase healthcare utilization and costs [86]. Mohd Zulkifly et al. found that in 147 stroke patients, the presence of post-traumatic stress symptoms was a significant predictor of functional disability [87].

Recent literature has made great strides in identifying pertinent risk factors. These include psychiatric comorbidities, neurotic coping strategies, low socioeconomic status, female sex, recurrent stroke, and past trauma and disability [84]. Multiple reports have correlated pre-stroke anxiety and depression diagnosis with PTSD occurrence [88]. Current research suggests the tendency to develop post-stroke PTSD may be linked to pre-stroke coping styles [89]. Individuals with repressor and low-anxious coping styles are less likely to develop PTSD compared to those with defensive and high-anxious coping styles [89]. Additionally, there may be increased prevalence in low socioeconomic populations, which have a propensity for recurrent strokes and poorer physical and mental health status [89,90]. Despite this trend, the literature lacks standardized consensus on the significance of

individual socioeconomic predictors. A cross-sectional study assessing traumatic response post-stroke found significant association in development of PTSD and was associated with female sex and comorbid anxiety disorder (80%) and depression (30-50%) [89]. This study did not find any associations between stroke severity [88]. The only significant association was the role of emotional experience and cognitive appraisal of the event itself [89]. To this end, the number of past traumatic events also conferred an exponential risk [91]. Number of strokes, negative affect, and cognitive appraisal was identified as a significant predictor of number and severity of PTSD symptoms [7]. Peritraumatic distress was also found to prospectively predict acute PTSD symptoms [7,92].




Figure 4. Several risk factors may be implicated in the development of post-stroke PTSD.

Currently, the pathophysiology of post-stroke PTSD remains poorly understood. Neuroimaging studies have implied selective brain atrophy may be implicated in PTSD pathophysiology. A prospective cohort study of first-ever TIA/post-stroke survivors found that probable PTSD development is linked with lower performance in cognitive and functional tests, as well as with pro-inflammatory states with elevated CRP. Stroke severity has also been associated with PTSD severity, specifically white matter lesion volume and hippocampal connectivity. Infarct volume and location were not. Lesions affecting connectivity are more likely to influence PTSD. Currently, there are no studies that have correlated stroke location with PTSD development. One study attempting to do this was inconclusive [93]. Although there is no established consensus in lesion localization and PTSD development, Rutovic et al. identified some suggestive relationships [94]. A higher incidence was found in patients with right-sided and brainstem strokes [94]. This was attributed to a tendency for delayed treatment and more pronounced disability [94]. However, again, this relationship does not distinguish between PTSD secondary to lesion location or that secondary to severe disability. Despite the absence of evidence correlating macroscopic findings, neuroendocrine dysregulation post-stroke has been associated with PTSD development. Pitman et al. found that a cortisol deficit may be implicated in PTSD pathogenesis [95]. Abnormal glucocorticoid levels have been found to interfere with memory, learning, stress adaptation, and resilience [96]. Furthermore, hydrocortisone injections post trauma were found to improve the trajectory of PTSD [97]. One caveat is these findings are not specific to pathogenesis of PTSD post-stroke. Further studies are needed in this area. Currently, metabolic associations between TIA/stroke and PTSD are unclear. Changes in N-acetylaspartate have been observed in both PTSD and in neural tissue fewer than three days after TIA [98]. Only correlative associations have been identified thus far.

Current targets for intervention include fostering social support (friend, spouse, partner) [89] and reducing social isolation. Other current management strategies include exposure therapy, which has demonstrated efficacy in treatment of combat veterans, and interventions to teach more effective coping skills [84]. In support of this, avoidance has been found to be the strongest predictor for post-traumatic stress symptoms [87]. This suggests that facing traumatic experiences by refining coping skills may have therapeutic benefits in this population. The literature on specifically treating PTSD in a post-stroke population is now showing some progress. Jiang et al. conducted a randomized controlled trial on 170 first-ever stroke patients and provided modest evidence that supportive psychological therapy can effectively treat post-stroke PTSD in its early stages [99]. A recent case report also showed clinically significant resolution of PTSD symptoms following 3 sessions of eye movement desensitization and reprocessing therapy [100]. This study is limited, however, due to assessment via self-report and its design as a case report. Current treatment efforts are focused on trialing existing PTSD therapies in a post-stroke population, but there is scant existing data for this. Therapies used in post-trauma PTSD should be evaluated in randomized controlled trials within the post-stroke population as well. Potentially effective medications include fluoxetine, paroxetine, sertraline, and venlafaxine [101]. Emerging treatment options include 3,4-methylenedioxymethamphetamine-assisted psychotherapy, virtual reality exposure therapy, ketamine, cannabinoids, prazosin, and repetitive transcranial magnetic stimulation [101]. Although not yet investigated in the post-stroke PTSD population, CBT has also been shown to be effective in the treatment of post-stroke mood disorders [85]. Additionally, traditional CBT may be challenging for stroke patients who have impaired mobility. To this end, the efficacy of telehealth therapy should be further researched.

The predominant limitation to the current data of the epidemiology is its reliance on questionnaires and scales which tend to overdiagnose PTSD as opposed to formal psychiatric interviews [84]. The development of PTSD post-stroke is also confounded by the disability conferred by severe stroke which may elicit a stress response of its own.

6. Conclusions

In sum, the pathophysiology of PTSD typically involves dysregulation of the limbic system, namely the amygdala and hippocampus [13,35]. Complicating the development of pharmacological treatment for PTSD is the many upstream elements which can result in PTSD symptoms if damaged or destroyed. A key point is that even when activity levels in the limbic system are normal, trauma to downstream elements like the hypothalamus can also contribute to PTSD by altering CRH and cortisol levels which are both major actors in the body's major stress responses [42,43]. Although external traumatic experiences like combat are often attributed as a prerequisite to PTSD, internal medical events such as TBI, subarachnoid hemorrhage, and stroke are also associated with the psychiatric disorder. Furthermore, despite being reported by several studies that the majority of TBI, SAH, and stroke patients do not develop PTSD symptoms, these instances of neurotrauma are each still moderately predictive of PTSD with an average rate of symptom development of 11-20% in TBI patients, 37% in SAH patients, and 25% in stroke patients [2,70,82]. Through various means, it is likely that traumatic injury to the central nervous system, particularly the limbic system, acts to exacerbate fear and intrusive memories. These findings support the idea that many neurotrauma patients may benefit from regular PTSD screening to prevent the worsening of symptoms if left untreated. Future studies should aim to clarify the impacts of altered noradrenergic signaling on the HPA axis. In addition to hyperactivity in the amygdala, it is not unlikely that hypersensitive CRH and cortisol responses to noradrenergic signaling may induce unusual stress behaviors like those seen in PTSD patients. Developing a concrete understanding of the physiological underpinnings of PTSD may allow for differential treatment of those who previously suffered neurotrauma from those who have not.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. American Psychiatric Association; Arlington, VA, USA: 2013. [[Google Scholar](#)]
2. Van Praag DLG, Cnossen MC, Polinder S, Wilson L, Maas AIR. Post-Traumatic Stress Disorder after Civilian Traumatic Brain Injury: A Systematic Review and Meta-Analysis of Prevalence Rates. *Journal of Neurotrauma*. 2019. Dec 1;36(23):3220–32. doi: 10.1089/neu.2018.5759 [[DOI](#)] [[PMC](#)] [[PubMed](#)]
3. Elder GA, Dorr NP, De Gasperi R, Gama Sosa MA, Shaughnessy MC, Maudlin-Jeronimo E, et al. Blast exposure induces post-traumatic stress disorder-related traits in a rat model of mild traumatic brain injury. *J Neurotrauma*. 2012;29(16):2564–2575. doi: 10.1089/neu.2012.2510. - [DOI](#) - [PMC](#) - [PubMed](#)
4. Kaplan GB, Leite-Morris KA, Wang L, Rumbika KK, Heinrichs SC, Zeng X, et al. Pathophysiological bases of comorbidity: traumatic brain injury and post-traumatic stress disorder. *J Neurotrauma*. (2018) 35(2):210–25. doi: 10.1089/neu.2016.4953 - [DOI](#) - [PubMed](#)
5. Wang, Z., Luo, C., Zhou, E. W., Sandhu, A. F., Yuan, X., Williams, G. E., Cheng, J., Sinha, B., Akbar, M., Bhattacharya, P., Zhou, S., Song, B. J., & Wang, X. (2023). Molecular Toxicology and Pathophysiology of Comorbid Alcohol Use Disorder and Post-Traumatic Stress Disorder Associated with Traumatic Brain Injury. *International journal of molecular sciences*, 24(10), 8805. doi: 10.3390/ijms24108805. - [DOI](#) - [PubMed](#)
6. O'Neil ME, et al. Associations among PTSD and postconcussive symptoms in the Long-Term Impact of Military-Relevant Brain Injury Consortium–Chronic Effects of Neurotrauma Consortium Prospective, longitudinal study cohort. *J Head Trauma Rehab*. 2021;36(6):E363–E372. doi: 10.1097/HTR.0000000000000665. - [DOI](#) - [PubMed](#)
7. Merriman C, Norman P, Barton J. Psychological correlates of PTSD symptoms following stroke. *Psychol Heal Med*. (2007) 12:592–602. doi: 10.1080/13548500601162747 - [DOI](#) - [PubMed](#)
8. Bruggemann L., Annoni J., Staub F., Von Steinbüchel N., Van der Linden M., Bogousslavsky J. (2006). Chronic posttraumatic stress symptoms after nonsevere stroke. *Neurology* 66, 513–516. doi: 10.1212/01.wnl.0000194210.98757.49 - [DOI](#) - [PubMed](#)
9. Sembi S, Tarrier N, O'Neill P, Burns A, Faragher B. Does post-traumatic stress disorder occur after stroke: a preliminary study. *Int J Geriatr Psychiatry*. 1998;13:315–322. doi: 10.1002/(sici)1099-1166(199805)13:5<315::aid-gps766>3.0.co;2-p - [DOI](#) - [PubMed](#)
10. Noble A.J., Baisch S., Covey J., Mukerji N., Nath F., Schenk T. Subarachnoid hemorrhage patients' fears of recurrence are related to the presence of posttraumatic stress disorder. *Neurosurgery*. 2011;69:323–332. doi: 10.1227/NEU.0b013e318216047e. - [DOI](#) - [PubMed](#)
11. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. American Psychiatric Association; Washington, DC, USA: 1994. - [Google Scholar](#)
12. Bryant, Richard. 2011. "State of the Art: Post-Traumatic Stress Disorder vs. Traumatic Brain Injury." *Dialogues in Clinical Neuroscience* 13 (3): 251–262. doi:10.31887/DCNS.2011.13.2/rbryant - [DOI](#) - [Google Scholar](#)
13. Souza RR, Noble LJ, McIntyre CK. Using the single prolonged stress model to examine the pathophysiology of PTSD. *Front Pharmacol*. (2017) 8:615. doi: 10.3389/fphar.2017.00615, PMID: - [DOI](#) - [PMC](#) - [PubMed](#)
14. Hendrickson R.C., Raskind M.A. Noradrenergic dysregulation in the pathophysiology of PTSD. *Exp. Neurol*. 2016;284:181–195. doi: 10.1016/j.expneurol.2016.05.014. - [DOI](#) - [PubMed](#)
15. Vieweg W.V., Julius D.A., Fernandez A., Beatty-Brooks M., Hettima J.M., Pandurangi A.K. Posttraumatic stress disorder: Clinical features, pathophysiology, and treatment. *Am. J. Med*. 2006;119:383–390. doi: 10.1016/j.amjmed.2005.09.027. - [DOI](#) - [PubMed](#)
16. Hamilton JP, Etkin A, Furman DJ, Lemus MG, Johnson RF, Gotlib IH. Functional Neuroimaging of Major Depressive Disorder: A Meta-Analysis and New Integration of Baseline Activation and Neural Response Data. *AJP*. 2012;169(7):693–703. doi: 10.1176/appi.ajp.2012.11071105. - [DOI](#) - [PubMed](#)
17. Gold A.L., Shin L.M., Orr S.P., Carson M.A., Rauch S.L., Macklin M.L., Lasko N.B., Metzger L.J., Dougherty D.D., Alpert N.M., et al. Decreased regional cerebral blood flow in medial prefrontal cortex during trauma-unrelated stressful imagery in Vietnam veterans with post-traumatic stress disorder. *Psychol. Med*. 2011;41:2563–2572. doi: 10.1017/S0033291711000730. - [DOI](#) - [PubMed](#)
18. Kasai K., Yamasue H., Gilbertson M., Shenton M., Rauch S., Pitman R. (2008). Evidence for acquired pregenual anterior cingulate gray matter loss from a twin study of combat-related posttraumatic stress disorder. *Biol. Psychiatry* 63 550–556. doi: 10.1016/j.biopsych.2007.06.022. - [DOI](#) - [PMC](#) - [PubMed](#)

19. Kitayama N., Quinn S., Bremner J.D. Smaller volume of anterior cingulate cortex in abuse-related posttraumatic stress disorder. *J. Affect. Disord.* 2006;90:171–174. doi: 10.1016/j.jad.2005.11.006. - [DOI](#) - [PMC](#) - [PubMed](#)
20. Shin L.M., Rauch S.L., Pitman R.K. Psychobiology of Posttraumatic Stress Disorder: A Decade of Progress. Volume 1071. Wiley-Blackwell; Hoboken, NJ, USA: 2006. Amygdala, medial prefrontal cortex, and hippocampal function in PTSD; pp. 67–79. doi: 10.1196/annals.1364.007 - [DOI](#) - [PubMed](#)
21. Kim SJ, Jeong DU, Sim ME, Bae SC, Chung A, Kim MJ, et al. Asymmetrically altered integrity of cingulum bundle in posttraumatic stress disorder. *Neuropsychobiology.* 2006;54:120–5. doi: 10.1159/000098262 - [PubMed](#) - [DOI](#)
22. Kosten T.R., Mason J.W., Giller E.L., Ostroff R.B., Harkness L. Sustained urinary norepinephrine and epinephrine elevation in post-traumatic stress disorder. *Psychoneuroendocrinology.* 1987;12:13–20. doi: 10.1016/0306-4530(87)90017-5. - [DOI](#) - [PubMed](#)
23. Liberzon I., Abelson, J. L., Flagel, S. B., Raz, J., & Young, E. (1999). Neuroendocrine and psychophysiologic responses in PTSD: A symptom provocation study. *Neuropsychopharmacology*, 21(1), 40–50. doi: 10.1016/S0893-133X(98)00128-6. - [DOI](#) - [PubMed](#)
24. Liberzon I, López JF, Flagel SB, Vázquez DM, Young EA. Differential regulation of hippocampal glucocorticoid receptors mRNA and fast feedback: relevance to post-traumatic stress disorder. *J. Neuroendocrinol.* 1999;11:11–17. doi: 10.1046/j.1365-2826.1999.00288.x. - [DOI](#) - [PubMed](#)
25. Cortisol and catecholamines in posttraumatic stress disorder: an epidemiologic community study. Young EA, Breslau N. *Arch Gen Psychiatry.* 2004;61:394–401. doi: 10.1001/archpsyc.61.4.394. [DOI](#) - [PubMed](#)
26. Geraciotti T.D., Baker D.G., Ekhtator N.N., West S.A., Hill K.K., Bruce A.B., Schmidt D., Rounds-Kugler B., Yehuda R., Keck P.E., et al. CSF norepinephrine concentrations in posttraumatic stress disorder. *Am. J. Psychiatry.* 2001;158:1227–1230. doi: 10.1176/appi.ajp.158.8.1227. - [DOI](#) - [PubMed](#)
27. Pietrzak RH, Gallezot JD, Ding YS et al (2013a) Association of posttraumatic stress disorder with reduced in vivo norepinephrine transporter availability in the locus coeruleus. *JAMA Psychiatry* 70:1199–1205. doi: 10.1001/jamapsychiatry.2013.399. - [DOI](#) - [PubMed](#) - [PMC](#)
28. Koenigs M, Grafman J. Posttraumatic stress disorder: the role of medial prefrontal cortex and amygdala. *Neuroscientist.* 2009;15:540–8. doi: 10.1177/1073858409333072. - [DOI](#) - [PubMed](#) - [PMC](#)
29. Klaassens ER, Giltay EJ, Cuijpers P, van Veen T, Zitman FG. Adulthood trauma and HPA-axis functioning in healthy subjects and PTSD patients: a meta-analysis. *Psychoneuroendocrinology.* 2012;37:317–31. doi: 10.1016/j.psyneuen.2011.07.003. - [DOI](#) [PubMed](#)
30. Finsterwald C, Alberini CM. 2014. Stress and glucocorticoid receptor-dependent mechanisms in long-term memory: from adaptive responses to psychopathologies. *Neurobiol. Learn. Mem.* 112, 17-29. doi: 10.1016/j.nlm.2013.09.017 - [DOI](#) - [PMC](#) - [PubMed](#)
31. Niermann HCM, Figner B, Tyborowska A, van Peer JM, Cillessen AHN, Roelofs K. Defensive freezing links Hypothalamic-Pituitary-Adrenal-axis activity and internalizing symptoms in humans. *Psychoneuroendocrinology.* 2017;82:83–90. doi: 10.1016/j.psyneuen.2017.05.001. - [DOI](#) - [PubMed](#)
32. van Stegeren A. H., Wolf O. T., Everaerd W., Rombouts S. A. (2008). Interaction of endogenous cortisol and noradrenaline in the human amygdala. *Prog. Brain Res.* 167 263–268. doi: 10.1016/S0079-6123(07)67020-4. - [DOI](#) - [PubMed](#)
33. Zuj D.V., et al., Endogenous cortisol reactivity moderates the relationship between fear inhibition to safety signals and posttraumatic stress disorder symptoms. *Psychoneuroendocrinology*, 2017. 78: p. 14–21. doi: 10.1016/j.psyneuen.2017.01.012 - [DOI](#) - [PubMed](#)
34. Sapolsky R.M., Krey L.C., McEwen B.S. The neuroendocrinology of stress and aging: the glucocorticoid cascade hypothesis. *Endocr. Rev.* 1986;7(3):284–301. doi: 10.1210/edrv-7-3-284. - [DOI](#) - [PubMed](#)
35. Sapolsky R.M. Why Stress Is Bad for Your Brain. *Science.* 1996;273:749–750. doi: 10.1126/science.273.5276.749. - [DOI](#) - [PubMed](#)
36. Nelson MD, Tumpap AM (2017) Posttraumatic stress disorder symptom severity is associated with left hippocampal volume reduction: a meta-analytic study. *CNS Spectr* 22(4):363–372. doi: 10.1017/s1092852916000833. - [DOI](#) - [PubMed](#)
37. O'Doherty D.C.M., Chitty K.M., Saddiqui S., Bennett M.R., Lagopoulos J. A systematic review and meta-analysis of magnetic resonance imaging measurement of structural volumes in posttraumatic stress disorder. *Psychiatry Res.* 2015;232(1):1–33. doi: 10.1016/j.psychresns.2015.01.002. - [DOI](#) - [PubMed](#)

38. Herman JP, et al. Regulation of the Hypothalamic-Pituitary-Adrenocortical Stress Response. *Compr Physiol.* 2016;6(2):603–21. doi: 10.1002/cphy.c150015. - [DOI](#) - [PubMed](#) - [PMC](#)
39. Laryea G., Arnett M.G., Muglia L.J. Behavioral Studies and Genetic Alterations in Corticotropin-Releasing Hormone (CRH) Neurocircuitry: Insights into Human Psychiatric Disorders. *Behav. Sci.* 2012;2:135–171. doi: 10.3390/bs2020135. - [DOI](#) - [PMC](#) - [PubMed](#)
40. Kolber B. J., Roberts M. S., Howell M. P., Wozniak D. F., Sands M. S., Muglia L. J. (2008). Central amygdala glucocorticoid receptor action promotes fear-associated CRH activation and conditioning. *Proc. Natl. Acad. Sci. U. S. A.* 105, 12004–12009. doi: 10.1073/pnas.0803216105. - [DOI](#) - [PMC](#) - [PubMed](#)
41. Haufier D., Nagy F. Z., Pare D. (2013). Neuronal correlates of fear conditioning in the bed nucleus of the stria terminalis. *Learn Mem.* 20 633–641. 10.1101/lm.031799.113. - [DOI](#) - [PMC](#) - [PubMed](#)
42. Pacella ML, Hruska B, Steudte-Schmiedgen S, et al. The utility of hair cortisol concentrations in the prediction of PTSD symptoms following traumatic physical injury. *Soc Sci Med* 2017; 175: 228–234. doi: 10.1016/j.socscimed.2016.12.046. [DOI](#) - [PubMed](#)
43. Galatzer-Levy IR, Steenkamp MM, Brown AD, Qian M, Inslicht S, Henn-Haase C, et al. Cortisol response to an experimental stress paradigm prospectively predicts long-term distress and resilience trajectories in response to active police service. *J Psychiatr Res.* 2014;56:36–42. doi: 10.1016/j.jpsychires.2014.04.020. [DOI](#) - [PMC](#) - [PubMed](#)
44. Morris, M. C., Hellman, N., Abelson, J. L., Rao, U., & Cortisol, H. R. (2016). Cortisol, heart rate, and blood pressure as early markers of PTSD risk: A systematic review and meta-analysis. *Clinical Psychology Review*, 49, 79–91. doi: 10.1016/j.cpr.2016.09.001. - [DOI](#) - [PMC](#) - [PubMed](#)
45. Maas A.I.R., Menon D.K., Adelson P.D., et al. Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol.* 2017;16(12):987–1048. doi: 10.1016/S1474-4422(17)30371-X. - [DOI](#) - [PubMed](#)
46. Traumatic Brain Injury (TBI). (2023, February 7). National Institute of Neurological Disorders and Stroke. <https://www.ninds.nih.gov/health-information/disorders/traumatic-brain-injury-tbi#toc-who-is-more-likely-to-get-a-traumatic-brain-injury-tbi>
47. Hoge CW, McGurk D, Thomas JL, Cox AL, Engel CC, Castro CA. Mild traumatic brain injury in U.S. Soldiers returning from Iraq. *N Engl J Med.* (2008) 358(5):453–63. doi: 10.1056/NEJMoa072972. - [DOI](#) - [PubMed](#)
48. Cimino AN Yi G, Patch M, Alter Y, Campbell JC, Gundersen KK, et al. . The effect of intimate partner violence and probable traumatic brain injury on mental health outcomes for black women. *J Aggr Maltreat Trauma.* (2019) 28:714–31. doi: 10.1080/10926771.2019.1587657. - [DOI](#) - [PMC](#) - [PubMed](#)
49. Schachar R.J., Park L.S., Dennis M. Mental health implications of traumatic brain injury (TBI) in children and youth. *J. Can. Acad. Child Adolesc. Psychiatry.* 2015;24(2):100–108. - [PMC](#) - [PubMed](#)
50. Bremner J. Traumatic stress: effects on the brain. *Dialogues Clin. Neurosci.* 2006;8(4):445–461. doi: 10.31887/dcns.2006.8.4/jbremner. - [DOI](#) - [PMC](#) - [PubMed](#)
51. Atkins CM. Decoding hippocampal signaling deficits after traumatic brain injury. *Transl. Stroke Res.* 2011;2:546–555. doi: 10.1007/s12975-011-0123-z. - [DOI](#) - [PMC](#) - [PubMed](#)
52. Wolf JA, Koch PF. Disruption of network synchrony and cognitive dysfunction after traumatic brain injury. *Front Syst Neurosci.* 2016;10:43. doi: 10.3389/fnsys.2016.00043. - [DOI](#) - [PMC](#) - [PubMed](#)
53. Abe K, Shimada R, Okada Y, Kibayashi K. Traumatic brain injury decreases serotonin transporter expression in the rat cerebrum. *Neurol Res.* (2016) 38:358–63. doi: 10.1080/01616412.2015.1110402. - [DOI](#) - [PubMed](#)
54. Bryson-Campbell M., Shaw L., O'Brien J., Holmes J., Magalhaes L. (2013). A scoping review on occupational and self identity after a brain injury. *Work* 44, 57–67. doi: 10.3233/WOR-2012-01561. - [DOI](#) - [PubMed](#)
55. Vasterling JJ, Jacob SN, Rasmusson A. Traumatic brain injury and posttraumatic stress disorder: conceptual, diagnostic, and therapeutic considerations in the context of co-occurrence. *J Neuropsychiatry Clin Neurosci.* (2018) 30:91–100. doi: 10.1176/appi.neuropsych.17090180. - [DOI](#) - [PubMed](#)
56. Shafique, S., & Rayi, A. (2022). Anatomy, Head and Neck, Subarachnoid Space. In *StatPearls*. StatPearls Publishing. - [PubMed](#)
57. Lawton M.T., Vates G.E. Subarachnoid hemorrhage. *N. Engl. J. Med.* 2017;377(3):257–266. doi: 10.1056/NEJMcp1605827. - [DOI](#) - [PubMed](#)

58. Ono, I., Kayahara, T., Kawashima, A., Okada, A., Miyamoto, S., Kataoka, H., Kurita, H., Ishii, A., & Aoki, T. (2023). Hypoxic microenvironment as a crucial factor triggering events leading to rupture of intracranial aneurysm. *Scientific reports*, 13(1), 5545. doi: 10.1038/s41598-023-32001-z. - [DOI](#) - [PubMed](#)
59. Deem, S., Diringer, M., Livesay, S., & Treggiari, M. M. (2023). Hemodynamic Management in the Prevention and Treatment of Delayed Cerebral Ischemia After Aneurysmal Subarachnoid Hemorrhage. *Neurocritical care*. doi: 10.1007/s12028-023-01738-w. - [DOI](#) - [PubMed](#)
60. Huang J, van Gelder JM. The probability of sudden death from rupture of intracranial aneurysms: A meta-analysis. *Neurosurgery*. 2002;51:1101–1105. doi: 10.1097/00006123-200211000-00001. - [DOI](#) - [PubMed](#)
61. Nguyen, T. A., Vu, L. D., Mai, T. D., Dao, C. X., Ngo, H. M., Hoang, H. B., Do, S. N., Nguyen, H. T., Pham, D. T., Nguyen, M. H., Nguyen, D. N., Vuong, H. T. T., Vu, H. D., Nguyen, D. D., Nguyen, L. Q., Dao, P. V., Vu, T. D., Nguyen, D. T., Tran, T. A., Pham, T. Q., ... Luong, C. Q. (2023). Predictive validity of the prognosis on admission aneurysmal subarachnoid haemorrhage scale for the outcome of patients with aneurysmal subarachnoid haemorrhage. *Scientific reports*, 13(1), 6721. doi: 10.1038/s41598-023-33798-5. - [DOI](#) - [PubMed](#)
62. Hop J.W., Rinkel G.J., Algra A., van Gijn J. Case-Fatality Rates and Functional Outcome After Subarachnoid Hemorrhage. *Stroke*. 1997;28:660–664. doi: 10.1161/01.STR.28.3.660. - [DOI](#) - [PubMed](#)
63. Connolly ES, Rabinstein AA, Carhuapoma JR et al (2012) Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American stroke association. *Stroke* 43:1711–1737. doi: 10.1161/STR.0b013e3182587839 - [DOI](#) - [PubMed](#)
64. Suarez JL. Diagnosis and management of subarachnoid hemorrhage. *Contin Lifelong Learn Neurol*. 2015;21(5):1263–87. doi: 10.1212/CON.0000000000000217. - [DOI](#) - [PubMed](#)
65. Luong CQ, et al. Clinical characteristics and factors relating to poor outcome in patients with aneurysmal subarachnoid hemorrhage in Vietnam: A multicenter prospective cohort study. *PLoS ONE*. 2021;16:e0256150. doi: 10.1371/journal.pone.0256150. - [DOI](#) - [PMC](#) - [PubMed](#)
66. Lord AS, Fernandez L, Schmidt JM, Mayer SA, Claassen J, Lee K, Connolly ES, Badjatia N (2012) Effect of rebleeding on the course and incidence of vasospasm after subarachnoid hemorrhage. *Neurology* 78:31-37. doi: 10.1212/WNL.0b013e31823ed0a4. - [DOI](#) - [PubMed](#) - [PMC](#)
67. Rosengart A.J., Schultheiss K.E., Tolentino J., Macdonald R.L. Prognostic factors for outcome in patients with aneurysmal subarachnoid hemorrhage. *Stroke*. 2007;38:2315–2321. doi: 10.1161/STROKEAHA.107.484360. - [DOI](#) - [PubMed](#)
68. Baisch SB; Schenk T; Noble AJ What is the cause of post-traumatic stress disorder following subarachnoid haemorrhage? Post-ictal events are key. *Acta Neurochir*. 2011, 153, 913–922. doi: 10.1007/s00701-010-0843-y. - [DOI](#) - [PubMed](#)
69. Hütter BO, Kreitschmann-Andermahr I. Subarachnoid hemorrhage as a psychological trauma: clinical article. *J Neurosurg*. (2014) 120:923–30. doi: 10.3171/2013.11.JNS121552 - [DOI](#) - [PubMed](#)
70. Noble A.J., Baisch S., Mendelow A.D., Allen L., Kane P., Schenk T. Posttraumatic stress disorder explains reduced quality of life in subarachnoid hemorrhage patients in both the short and long term. *Neurosurgery*. 2008;63:1095–1104. doi: 10.1227/01.NEU.0000327580.91345.78. - [DOI](#) - [PubMed](#)
71. Edmondson D. (2014). An enduring somatic threat model of posttraumatic stress disorder due to acute life-threatening medical events. *Soc. Personal. Psychol. Compass* 8, 118–134. doi: 10.1111/spc3.12089. - [DOI](#) - [PMC](#) - [PubMed](#)
72. Breslau N., Schultz L. Neuroticism and post-traumatic stress disorder: A prospective investigation. *Psychol. Med*. 2013;43:1697–1702. doi: 10.1017/S0033291712002632. - [DOI](#) - [PubMed](#)
73. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *J Consult Clin Psychol*. 2000;68:748–66. doi: 10.1037/0022-006X.68.5.748. - [DOI](#) - [PubMed](#)
74. Kessler RC, Sonnega A, Bromet E, et al. . Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995;52:1048–60. doi: 10.1001/archpsyc.1995.03950240066012. - [DOI](#) - [PubMed](#)
75. Parslow RA, Jorm AF. Pretrauma and posttrauma neurocognitive functioning and PTSD symptoms in a community sample of young adults. *Am J Psychiatry*. 2007;164:509-515. doi: 10.1176/ajp.2007.164.3.509. - [DOI](#) - [PubMed](#)
76. Reynolds K, Pietrazk RH, Mackenzie CS, Chou KL, Sareen J (2016) Post-traumatic stress disorder across the adult lifespan: Findings from a nationally representative survey, *Am J Geriatr Psychiatry* 24, 81–93. doi: 10.1016/j.jagp.2015.11.001. - [DOI](#) - [PubMed](#)

77. Wajer I.M.C.H., Smits A.R., Rinkel G.J.E., van Zandvoort M.J.E., Wijngaards-de Meij L., Visser-Meily J.M.A. Exploratory study of the course of posttraumatic stress disorder after aneurysmal subarachnoid hemorrhage. *Gen. Hosp. Psychiatry*. 2018;53:114–118. doi: 10.1016/j.genhosppsych.2018.03.004. - [DOI](#) - [PubMed](#)
78. Sheldrick R, Tarrier N, Berry E, Kincey J. Post-traumatic stress disorder and illness perceptions over time following myocardial infarction and subarachnoid haemorrhage. *Br J Health Psychol*. 2006;11(Pt 3):387–400. doi: 10.1348/135910705x71434. - [DOI](#) - [PubMed](#)
79. Depression and post-traumatic stress disorder after aneurysmal subarachnoid haemorrhage in relation to lifetime psychiatric morbidity. Hedlund M, Zetterling M, Ronne-Engström E, Carlsson M, Ekselius L. *Br J Neurosurg*. 2011;25:693–700. doi: 10.3109/02688697.2011.578769. - [DOI](#) - [PMC](#) - [PubMed](#)
80. Greenberg N., Brooks S., Dunn R. Latest developments in post-traumatic stress disorder: Diagnosis and treatment. *Br. Med. Bull*. 2015;114:147–155. doi: 10.1093/bmb/ldv014. - [DOI](#) - [PubMed](#)
81. Feigin, V. L., et al. (2014). Global and regional burden of stroke during 1990–2010: Findings from the global burden of Disease Study 2010. *Lancet*, 383(9913), 245–254. doi: 10.1016/s0140-6736(13)61953-4. - [DOI](#) - [PubMed](#) - [PMC](#)
82. Edmondson D, Richardson S, Fausett JK, Falzon L, Howard VJ, Kronish IM. Prevalence of PTSD in survivors of stroke and transient ischemic attack: a meta-analytic review. *PLoS One*. 2013 Jun 19;8(6):e66435. doi: 10.1371/journal.pone.0066435. - [DOI](#) - [PMC](#) - [PubMed](#)
83. Kronish IM, Edmondson D, Goldfinger JZ, Fei K, Horowitz CR. Posttraumatic stress disorder and adherence to medications in survivors of strokes and transient ischemic attacks. *Stroke*. 2012;43(8):2192–2197. doi: 10.1161/strokeaha.112.655209. - [DOI](#) - [PMC](#) - [PubMed](#)
84. Ferro J, Caeiro L, Figueira M. Neuropsychiatric sequelae of stroke. *Nat Rev Neurol*. (2016) 12:269–80. doi: 10.1038/nrneurol.2016.46. - [DOI](#) - [PubMed](#)
85. Stein L. A., Goldmann E., Zamzam A., Luciano J. M., Messé S. R., Cucchiara B. L., et al. . (2018). Association between anxiety, depression, and post-traumatic stress disorder and outcomes after ischemic stroke. *Front. Neurol*. 9:890. doi: 10.3389/fneur.2018.00890. - [DOI](#) - [PMC](#) - [PubMed](#)
86. Ghose S.S., Williams L.S., Swindle R.W. Depression and other mental health diagnoses after stroke increase inpatient and outpatient medical utilization three years poststroke. *Med. Care*. 2005;43:1259–1264. doi: 10.1097/01.mlr.0000185711.50480.13. - [DOI](#) - [PubMed](#)
87. Mohd Zulkifly, M. F. , Ghazali, S. E. , Che Din, N. , Desa, A. , & Raymond, A. A. (2015). The Ability of Recovery Locus of Control Scale (RLOC) and Post-traumatic Stress Symptoms (PTSS) to Predict the Physical Functioning of Stroke Patients. *Malaysian Journal of Medical Science*, 22(5), 31–41. - [PMC](#) - [PubMed](#)
88. Marques, P. T., Beltrami, L. P. B., Rosa, C. T., Massuda, R., Barbosa, F. J. L., Zetola, V. F., & Lange, M. C. (2020). Post-traumatic stress disorder in non-disabled ischemic stroke patients. *General Hospital Psychiatry*, 67, 150–151. doi: 10.1016/j.genhosppsych.2020.04.002. - [DOI](#) - [PubMed](#)
89. Ben Assayag E, Tene O, Korczyn AD, et al. Posttraumatic stress symptoms after stroke: the effects of anatomy and coping style. *Stroke*. 2022;53(6):1924–1933. doi: 10.1161/STROKEAHA.121.036635. - [DOI](#) - [PubMed](#)
90. Goldfinger J. Z., Edmondson D., Kronish I. M., Fei K., Balakrishnan R., Tuhim S., et al. . (2014). Correlates of post-traumatic stress disorder in stroke survivors. *J. Stroke Cerebrovasc. Dis*. 23, 1099–1105. doi: 10.1016/j.jstrokecerebrovasdis.2013.09.019. - [DOI](#) - [PMC](#) - [PubMed](#)
91. Favrole P., Jehel L., Levy P., Descombes S., Muresan I.-P., Manificier M.J., Alamowitch S. Frequency and predictors of post-traumatic stress disorder after stroke: A pilot study. *J. Neurol. Sci*. 2013;327:35–40. doi: 10.1016/j.jns.2013.02.001. - [DOI](#) - [PubMed](#)
92. Letamendia, C., Leblanc, N. J., Pariente, J., Simon, N. M., Thomas, C. L., Chabrol, H., ... Bui, E. (2012). Peritraumatic distress predicts acute posttraumatic stress disorder symptoms after a first stroke. *General Hospital Psychiatry*, 34(5), e11–3. doi: 10.1016/j.genhosppsych.2012.03.011. - [DOI](#) - [PubMed](#)
93. Gozzi SA, Wood AG, Chen J, Vaddadi K, Phan TG. Imaging predictors of poststroke depression: methodological factors in voxel-based analysis. *BMJ Open*. (2014) 4:e004948. doi: 10.1136/bmjopen-2014-004948. - [DOI](#) - [PMC](#) - [PubMed](#)
94. Rutovic S, Kadojic D, Dikanovic M, Solic K, Malojcic B. Prevalence and correlates of post-traumatic stress disorder after ischaemic stroke. *Acta Neurol Belg*. 2021;121(2):437–442. doi: 10.1007/s13760-019-01200-9. - [DOI](#) - [PubMed](#)

95. Pitman R.K., Rasmusson A.M., Koenen K.C., Shin L.M., Orr S.P., Gilbertson M.W., Milad M.R., Liberzon I. Biological Studies of Post-Traumatic Stress Disorder. *Nat. Rev. Neurosci.* 2012;13:769–787. doi: 10.1038/nrn3339. - [DOI](#) - [PMC](#) - [PubMed](#)
96. Garton A.L., Sisti J.A., Gupta V., Christophe B.R., Connolly E.S. Poststroke Post-Traumatic Stress Disorder. *Stroke.* 2017;48:507–512. doi: 10.1161/STROKEAHA.116.015234. Corrected in *Stroke* 2017, 48, e96. - [DOI](#) - [PMC](#) - [PubMed](#)
97. Zohar J, et al. . 2011. High dose hydrocortisone immediately after trauma may alter the trajectory of PTSD: interplay between clinical and animal studies. *Eur Neuropsychopharmacol.* 21(11):796–809. doi: 10.1016/j.euroneuro.2011.06.001. - [DOI](#) - [PubMed](#)
98. Kiphuth IC, Utz KS, Noble AJ, Köhrmann M, Schenk T. Increased prevalence of posttraumatic stress disorder in patients after transient ischemic attack. *Stroke.* 2014;45(11):3360–3366. doi: 10.1161/strokeaha.113.004459. - [DOI](#) - [PubMed](#)
99. Jiang, C., Li, Z., Du, C., Zhang, X., Chen, Z., Luo, G., Wu, X., Wang, J., Cai, Y., Zhao, G., & Bai, H. (2022). Supportive psychological therapy can effectively treat post-stroke post-traumatic stress disorder at the early stage. *Frontiers in neuroscience*, 16, 1007571. doi: 10.3389/fnins.2022.1007571. - [DOI](#) - [PubMed](#)
100. Smart C. M. (2022). Eye movement desensitization and reprocessing for post-stroke post-traumatic stress disorder: Case report using the three-phase approach. *Brain injury*, 36(12-14), 1372–1381. doi: 10.1080/02699052.2022.2140833. - [DOI](#)
101. Tang W. K., Wang L., Tsoi K. K. F., Rutovic S., Kim J. S. (2022). Post-traumatic stress disorder after stroke: a systematic review. *Neurol. India* 70, 1887–1895. doi: 10.4103/0028-3886.359285. - [DOI](#) - [PubMed](#)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.