

# A Recombinant Protein XBB.1.5 RBD/Alum/CpG Vaccine Elicits High Neutralizing Antibody Titers against Omicron Subvariants of SARS-CoV-2

[Syamala Rani Thimmiraju](#) , Rakesh Adhikari , Maria J Villar , Jungsoon Lee , Zhuyun Liu , Rakhi Kundu , Yi-Lin Chen , [Suman Sharma](#) , Karm Ghei , Brian Keegan , [Leroy Versteeg](#) , [Portia M Gillespie](#) , Allan Ciciriello , [Nelufa Yesmin](#) , [Cristina Poveda](#) , Nestor Uzcategui , [Wen-Hsiang Chen](#) , [Jason T Kimata](#) , [Bin Zhan](#) , [Ulrich Strych](#) , [Maria Elena Bottazzi](#) , [Peter Hotez](#) , [Jeroen Pollet](#) \*

Posted Date: 7 September 2023

doi: 10.20944/preprints202309.0139.v2

Keywords: Immune escape; Vaccine efficacy; COVID-19; SARS-CoV-2



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# A Recombinant Protein XBB.1.5 RBD/Alum/CpG Vaccine Elicits High Neutralizing Antibody Titers against Omicron Subvariants of SARS-CoV-2

Syamala Rani Thimmiraju <sup>1,2</sup>, Rakesh Adhikari <sup>1,2</sup>, Maria Jose Villar <sup>1,2</sup>, Jungsoon Lee <sup>1,2</sup>, Zhuyun Liu <sup>1,2</sup>, Rakhi Kundu <sup>1,2</sup>, Yi-Lin Chen <sup>1,2</sup>, Suman Sharma <sup>3</sup>, Karm Ghei <sup>3</sup>, Brian Keegan <sup>1,2</sup>, Leroy Versteeg <sup>1,2</sup>, Portia M. Gillespie <sup>1,2</sup>, Allan Ciciriello <sup>1,2</sup>, Nelufa Y. Islam <sup>1,2</sup>, Cristina Poveda <sup>1,2</sup>, Nestor Uzcategui <sup>1,2</sup>, Wen-Hsiang Chen <sup>1,2</sup>, Jason T. Kimata <sup>3</sup>, Bin Zhan <sup>1,2</sup>, Ulrich Strych <sup>1,2</sup>, Maria Elena Bottazzi <sup>1,2,4,5</sup>, Peter J. Hotez <sup>1,2,4,5</sup> and Jeroen Pollet <sup>1,2</sup>

<sup>1</sup> Texas Children's Hospital Center for Vaccine Development, Houston, TX, USA

<sup>2</sup> Department of Pediatrics, National School of Tropical Medicine, Baylor College of Medicine, Houston, TX, USA

<sup>3</sup> Department of Molecular Virology and Microbiology, Baylor College of Medicine, Houston, TX, USA

<sup>4</sup> Department of Biology, Baylor University, Waco, TX, USA

<sup>5</sup> James A. Baker III Institute for Public Policy, Rice University, Houston, TX, USA

\* Correspondence: Jeroen.pollet@bcm.edu; Tel.: +1 832-824-0545

**Abstract:** (1) Background: We previously reported the development of a recombinant protein SARS-CoV-2 vaccine, consisting of the Receptor-Binding Domain (RBD) of the SARS-CoV-2 spike protein, adjuvanted with aluminum hydroxide (alum) and CpG oligonucleotides. In mice and non-human primates, our wild-type (WT) RBD vaccine induced high neutralizing antibody titers against the WT isolate of the virus, and, with partners in India and Indonesia it was later developed into two closely resembling human vaccines, Corbevax and Indovac. Here, we describe the development and characterization of a next-generation vaccine adapted to the recently emerging XBB variants of SARS-CoV-2. (2) Methods: We conducted preclinical studies in mice using a novel yeast-produced SARS-CoV-2 XBB.1.5 RBD subunit vaccine candidate formulated with alum and CpG. We examined the neutralization profile of sera obtained from mice vaccinated twice intramuscularly at a 21-day interval with the XBB.1.5-based RBD vaccine, against ancestral wild type (WT), Beta, Delta, BA.4, BQ.1.1, BA.2.75.2, XBB.1.16, and XBB.1.5 SARS-CoV-2 pseudoviruses. (3) Results: The XBB.1.5 RBD/CpG/alum vaccine induced robust antibody production in mice that demonstrated strong neutralization of the XBB.1.5 pseudovirus and multiple other Omicron pseudoviruses. However, regardless of high antibody cross-reactivity by ELISA, the anti-XBB.1.5 RBD antigen serum showed low neutralizing titers against the WT and Delta virus variants. (4) Conclusions: Whereas we observed modest cross-neutralization against Omicron subvariants by sera from mice vaccinated with the WT RBD/CpG/Alum vaccine or with the BA.4/5-based vaccine, sera raised against the XBB.1.5 RBD showed robust cross-neutralization. These findings underscore the opportunity for an updated vaccine formulation utilizing the XBB.1.5 RBD antigen.

**Keywords:** immune escape; vaccine efficacy; COVID-19; SARS-CoV-2

---

## 1. Introduction

Since the beginning of the COVID-19 pandemic, the SARS-CoV-2 virus has evolved very rapidly, giving rise to a series of mutations, generating multiple variants of concern (VOCs) like Beta, Delta, and, subsequently, Omicron [1]. Within the Omicron lineage, recombinant strains have emerged constantly starting from the earliest BA.1 strain to the latest XBB.1.5, XBB 1.16, and EG.5 strains. As of August 2023, these Omicron subvariants appear to have become dominant worldwide. XBB lineages have likely evolved from a recombination event among two BA.2 strains with a mutation at

S486P [2]. The additional F486P substitution in XBB.1.5 is believed to offer higher affinity to the human angiotensin-converting enzyme 2 (ACE2) receptor than seen with BQ.1, XBB, and XBB.1 [3]. As a consequence, immunization or prior exposure to the ancestral wild-type (WT) variant provides suboptimal protection through neutralizing antibodies [4]. Specifically, studies have shown that XBB and other Omicron strains such as BQ.1.1 have higher resistance to humoral immunity induced by vaccination or natural infection, than earlier strains like BA.2 and BA.5 [5–7]. These findings were confirmed with our recombinant protein vaccines, either RBD219-N1 or RBD203-N1, which encodes the receptor-binding domain (RBD) of the SARS-CoV-2 spike protein. In mice and non-human primates, such RBDs, adjuvanted with CpG oligonucleotides and aluminum hydroxide (Alum) induced high neutralizing antibodies (Abs) against SARS-CoV-2 (WT)[8–10]. Such antigens became central components of the Corbevax vaccine produced by Biological E in India and IndoVac produced by BioFarma in Indonesia and have been administered close to 100 million times [11].

While these vaccines still offer protection from severe disease, the cross-neutralizing titers against the Omicron strains are suboptimal. As XBB.1.5 is highly resistant to antiviral immunotherapy, the most efficient way to control the current wave is to update vaccine antigens to induce a more effective immunity [12]. Here, we show the development and testing of an XBB.1.5 RBD-based vaccine together with vaccines matching the Beta, Delta, and BA.4/5 variants. We used pseudovirus neutralization assays to determine the cross-protection elicited by these antigens against the ancestral strain (WT), and seven additional variants (Beta, Delta, BA.4, BQ.1.1, BA.2.75.2, XBB.1.16, and XBB.1.5).

## 2. Materials and Methods

### 2.1. Sources of recombinant RBD203-N1 proteins

SARS-CoV-2 RBD203-N1 protein was designed and produced as previously described [8,13], and it encompasses amino acid residues 332–533 of the SARS-CoV-2 spike protein. The sequence alignment among the RBD variants included in this study is shown in **Figure 1**.

WT	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNSASFSTFKCYGVSPTKLNDLC	-389
Delta	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNSASFSTFKCYGVSPTKLNDLC	-389
Beta	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNSASFSTFKCYGVSPTKLNDLC	-389
BQ.1.1	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNFAPFFAFKCYGVSPTKLNDLC	-389
BA.4/5	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNFAPFFAFKCYGVSPTKLNDLC	-389
BA.2.75.2	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNFAPFFAFKCYGVSPTKLNDLC	-389
XBB.1.5	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNFAPFFAFKCYGVSPTKLNDLC	-389
XBB.1.16	-332	ITNLCPPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNFAPFFAFKCYGVSPTKLNDLC	-389
		*****	
WT	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
Delta	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
Beta	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
BQ.1.1	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
BA.4/5	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
BA.2.75.2	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
XBB.1.5	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
XBB.1.16	-390	FTNVYADSFVIRGDEVQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGNNY	-451
		*****	
WT	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
Delta	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
Beta	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
BQ.1.1	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
BA.4/5	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
BA.2.75.2	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
XBB.1.5	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
XBB.1.16	-452	LYRLFRKSNLKPFFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVV	-511
		*****	
WT	-512	VLSFELLHAPATVCGPKKSTNL	-533
Delta	-512	VLSFELLHAPATVCGPKKSTNL	-533
Beta	-512	VLSFELLHAPATVCGPKKSTNL	-533
BQ.1.1	-512	VLSFELLHAPATVCGPKKSTNL	-533
BA.4/5	-512	VLSFELLHAPATVCGPKKSTNL	-533
BA.2.75.2	-512	VLSFELLHAPATVCGPKKSTNL	-533
XBB.1.5	-512	VLSFELLHAPATVCGPKKSTNL	-533
XBB.1.16	-512	VLSFELLHAPATVCGPKKSTNL	-533
		*****	

**Figure 1.** Amino acid sequence alignment of the eight RBD variants investigated in this study (alignment made with <https://www.ebi.ac.uk/Tools/msa/clustalo/>).

## 2.2. Production of the five variant RBD antigens in *Pichia pastoris* X-33

To generate the recombinant proteins in yeast, DNAs encoding the SARS-CoV-RBD203-N1 proteins for the WT, Beta, Delta, Omicron BA.4/5 (BA.4 RBD is identical to BA.5), and XBB.1.5 variants were codon-optimized based on yeast codon preference and cloned into the yeast expression vector pPICZαA. The recombinant plasmid DNAs were transformed into *P. pastoris* X33 following a process described previously [8,14]. The highest expressing clones for each RBD were used to make research seed stocks that were saved at -80 °C. Fermentation was carried out at the 5 or 1 L scale, respectively, as previously described [14]. Briefly, the seed stocks for each construct were used to inoculate a 0.5 L buffered minimal glycerol (BMG) medium and were grown at 30 °C and 250 rpm until an OD<sub>600</sub> of 5–10. This culture was used to inoculate sterile low-salt medium (LSM, pH 5.0) with PTM1 trace elements and d-Biotin. Cell expansion was continued at 30 °C with a dissolved oxygen (DO) set point of 30% until glycerol depletion. Then methanol was pumped in from 1 mL/L/h to 11 mL/L/h over a 6 h period; the pH was adjusted to 6.5. The methanol induction was maintained at 11 mL/L/h at 25 °C for 70 h, except for XBB.1.5-RBD which was induced for 48 h. After fermentation, the culture was harvested by centrifugation, filtered, and kept frozen at -80 °C until purification. The recombinant RBD protein was captured from the fermentation supernatant using a butyl Sepharose high-performance column (Cytiva) in the presence of ammonium sulfate salt at a concentration of 0.8 M (XBB.1.5-RBD) or 1.1 M (WT-RBD, Beta-RBD, Delta-RBD and BA.4/5-RBD) in HIC buffer (30 mM Tris-HCl pH 8.0). RBD protein was further purified using a Q Sepharose XL (QXL) column (Cytiva) in a negative mode in QXL buffer A (20 mM Tris-HCl pH 7.5) with 100 mM NaCl (WT-RBD and Beta-RBD), 50 mM NaCl (Delta-RBD) or 0 mM NaCl (XBB.1.5-RBD). For BA.4/5-RBD, a negative QXL chromatography was performed in QXL buffer B (20 mM Tris-HCl pH 8.4 and 10 mM NaCl) followed by dialysis in QXL buffer C (20 mM L-histidine and 100 mM NaCl, pH 6.0) for storage. All RBD proteins were aseptically filtered using a 0.22 μm filter and stored at -80 °C until usage. To evaluate these RBD proteins' biophysical characteristics and functionality, SDS-PAGE-based densitometry, dynamic light scattering, and ACE-2 binding assays were performed following the methods previously described [15]. SE-HPLC was performed by injecting 50 μg of RBD on an XBridge Premier Protein SEC Column (Waters, Cat# 186009959) connected with a corresponding guard column (Waters, Cat# 186009969). The protein was eluted with 1X TBS, pH 7.5 at a 0.5 mL/min flow rate. The Bio-Rad gel filtration standard (Bio-Rad, catalog# 1511901) was used as a control.

## 2.3. Vaccine formulations and preclinical study design

All animal experiments were performed under a Baylor College of Medicine Institutional Animal Care and Use Committee-approved protocol. Female BALB/c mice (N=8/group), aged 6–8 weeks old, were immunized twice intramuscularly at 21-day intervals with the antigens shown in [Table S1](#) and then euthanized two weeks after the second vaccination. Each dose of the SARS-CoV-2 RBD vaccine contained 7 μg RBD protein, 200 μg alum (*Alhydrogel*®, aluminum hydroxide, Catalog # AJV3012, Croda Inc., UK), and 20 μg CpG1826 (*Invivogen*, USA). The SARS-CoV-2 RBD proteins were prepared with 1xTBS buffer (20 mM Tris, 100 mM NaCl, pH 7.5). Before injection, alum and CpG 1826 was added, and the sample was vortexed for 3 s.

## 2.4. Serological antibody measurements by ELISA

To examine SARS-CoV-2 RBD-specific antibodies in the mouse sera, indirect ELISAs were conducted as published before [13]. Briefly, plates were coated with 0.2 μg/well of SARS-CoV-2 RBD proteins from different variants. Mouse sera samples were 3-fold diluted from 1:200 to 1: 11,809,800 in 0.1% BSA in assay buffer (0.05% Tween20 in 1x PBS). Assay controls included a 1:400 dilution of pooled naive mouse sera (negative control) 1:10,000 dilution of pooled high titer mouse (positive control) and assay buffer as blanks. 100 μL/well of 1:6000 goat anti-mouse IgG HRP, in assay buffer was added. After incubation, plates were washed five times, followed by adding of 100 μL/well of TMB substrate. Plates were incubated for 15 min at room temperature (RT) while protected from light. After incubation, the reaction was stopped by adding 100 μL/well 1 M HCl. The absorbance at

a wavelength of 450 nm was measured using a Biotek Epoch 2 spectrophotometer. For each sample, the titer was determined using a four parameter logistic regression curve of the absorbance values. The titer cutoff value: negative serum control + 3 x standard deviation of the negative serum control.

### 2.5. Pseudovirus assay for determination of neutralizing antibodies

To test for neutralizing antibodies, we prepared non-replicating lentiviral particles expressing the SARS-CoV-2 spike protein on their membrane and encoding luciferase as a reporter. Infection was quantified using in vitro-grown human 293 T-hACE2 cells based on luciferase expression. The pseudovirus production system included the luciferase-encoding reporter plasmid, pNL4-3. lucR-E-, a Gag/Pol-encoding packaging construct (p $\Delta$ 8.9), and the codon-optimized SARS-CoV-2 spike VOC-expressing plasmids (pcDNA3.1-CoV-2 S gene), based on clone p278-1 [13]. Pseudovirus-containing supernatants were recovered after 48 h and passed through a 0.45  $\mu$ m filter, and saved at  $-80^{\circ}\text{C}$  until used for neutralization studies.

Expression plasmids for SARS-CoV-2 spike variants of concern WT, Beta, and Delta were generated by site-directed mutagenesis or replacement of segments of the codon-optimized Wuhan SARS-CoV-2 spike expression clone, p278-1 with variant sequences as previously described [8]. Omicron spike variants BA.4, BA.2.75.2, BQ.1.1, and XBB.1.5 sequences were generated by changing codons of the p278-1 spike clone sequence to produce the consensus amino acid sequence of each variant. The variant spike sequences were synthesized with the 3' Flag-tag (*Genscript*) and inserted into the pcDNA3.1 expression vector. The list with the variant-specific mutations added to the spike protein variant clones can be found in the supplementary data. All sequence changes were confirmed by commercial DNA sequencing.

The pseudovirus assay was performed as described earlier [13]; briefly, 10  $\mu$ L of pseudovirus were incubated with serial dilutions of the serum samples for 1 h at 37  $^{\circ}\text{C}$ . Next, 100  $\mu$ L of sera-pseudovirus were added to 293 T-hACE2 cells in 96-well poly-D-lysine coated culture plates. Following 48 h of incubation in a 5%  $\text{CO}_2$  environment at 37  $^{\circ}\text{C}$ , the cells were lysed with 100  $\mu$ L of Promega Glo Lysis buffer for 15 min at room temperature. Finally, 50  $\mu$ L of the lysate was added to 50  $\mu$ L luciferase substrate (Promega Luciferase Assay System). The amount of luciferase was quantified by luminescence (relative luminescence units (RLU)), using the Luminometer (*Biosynergy* H4, BioTek). Sera from vaccinated mice were compared by their 50% inhibitory dilution (IC<sub>50</sub>), defined as the serum dilution at which the virus infection was reduced by 50% compared to the negative control (virus + cells). IC<sub>50</sub> values were calculated as described in [16]. Sets of IC<sub>50</sub> titers elicited by sera from mice vaccinated with the RBD vaccine against five pseudovirus variants were analyzed using GraphPad Prism 8.0 to rank the RBD vaccines according to Spearman's correlation coefficient.

## 3. Results


### 3.1. Expression, purification, and characterization of recombinant proteins of different variants of SARS-CoV-2 RBD203

After *P. pastoris* X-33 plasmid transformation, recombinant proteins of different variants of the SARS-CoV-2 RBD, including WT, Beta, Delta, Omicron BA.4/5 and Omicron XBB.1.5, were expressed by induction with methanol at 30  $^{\circ}\text{C}$  for 72 hours (WT, Beta, Delta, BA.4/5) or 48 hours (XBB.1.5) in either a 1 or a 5 L fermentation process. Following the published protocol for RBD203-N1 [14], all proteins were purified by a combination of a hydrophobic interaction and ion exchange chromatography. While all proteins were purified to more than 96% homogeneity, their yields differed (Table 1). In particular, the RBDs of the two Omicron strains, BA.4/5 and XBB.1.5 showed reduced yields. We are currently optimizing the purification conditions of these strains to prepare for technology transfer to manufacturing partners.

Table 1. Biophysical characterization of five recombinant RBD proteins.

RBD Variant	Purification Yield [RBD (mg) / fermentation supernatant (L)]	SDS-PAGE with Coomassie Blue staining (non- reduced)		Homogeneity by SE-HPLC (%)	Radius by dynamic light scattering (nm)
		MW (kDa)	Purity (%)		
WT-RBD	270.5	25.3	94.1	99.7	2.57±0.01
Beta-RBD	204.3	23.8	93.2	99.7	2.63±0.01
Delta-RBD	294.5	25.1	93.9	98.4	2.73±0.02
BA.4/5-RBD	46.2	24.7	94.2	96.4	2.58±0.02
XBB.1.5-RBD	78.0	25.1	93.2	99.4	2.71±0.03

When analyzed by SE-HPLC, all RBD proteins produced a single peak detected at UV280 nm, confirming the homogeneity of each preparation (Figure 2A). Moreover, the apparent molecular weight by SE-HPLC matched that observed by densitometry scanning of SDS-PAGE gels (Table 1).




**Figure 2.** A) SE-HPLC and B) ACE-2 binding ELISA were performed to evaluate the identity/homogeneity and the functionality of five recombinant RBD proteins.

The functionality of the purified RBD molecules was established in an ACE-2 binding assay (Figure 2B), where all proteins were shown to bind recombinant human ACE-2. Consistent with prior data published by Mannar *et al.* [17], the affinity of the Omicron RBDs seemed to be higher than the other RBD variants, while the WT-RBD possessed the lowest binding affinity to ACE-2.

### 3.2. Immune response induced by adjuvanted RBD vaccines in mice

Preclinical studies were performed in eight to nine-week-old female BALB/c mice, using yeast-produced SARS-CoV-2 recombinant RBD proteins, adjuvanted with aluminum hydroxide and CpG1826. Mice (n=8/group) were immunized twice intramuscularly on days 0 and 21 with a dose of 7 µg RBD, 200 µg alum, and 20 µg CpG. On day 35, the study was terminated, and sera were collected and evaluated for total IgG and neutralizing antibodies against pseudoviruses matching the Wild type (WT) SARS-CoV-2 isolate, Beta, Delta, as well as the Omicron variants BA.4 and XBB.1.5 and XBB.1.16 (Figure 3).






**Figure 3.** A) Total IgG titers of pooled sera from mice immunized with five different RBD vaccine antigens, B) Neutralizing antibody titers of five RBD antigens tested against the same five pseudovirus variants. The Y-axis shows IC50 values of the neutralizing antibody titer on a log 10 scale (Figure S1), while the X-axis represents five RBD sera panels, each showing the neutralization titers against five pseudoviruses.

We found that the total IgG antibody titers elicited in mouse sera by all five (WT, Beta, Delta, BA.4/5, and XBB.1.5) RBDs were high when tested with immobilized antigens in an ELISA. (**Figure 3A**). However, when sera were tested for neutralizing antibodies in a pseudovirus assay, we observed that cross-protection of WT RBD-induced sera against Omicron BA.4/5 pseudovirus was minimal (**Figure 3B**), with a 73-fold reduction in the IC50 from 4174 against WT pseudovirus, to an IC50 of 57 against BA.4/5 pseudovirus. The titer against Omicron XBB.1.5 pseudovirus was even lower, with an IC50 of 29. Nab titers induced by Beta RBD and Delta RBD also failed to suggest protection against Omicron XBB.1.5 pseudovirus (**Figure 3B**). However, sera raised against the BA.4/5 RBD showed no protection against WT (IC50=20), Beta (IC50=193), Delta (IC50=20) pseudoviruses, and notably, the sera offered little protection against XBB.1.5 (IC50=40) pseudoviruses. These results contrast with the data of the mice vaccinated with the XBB.1.5 RBD vaccine, which induced high titers of neutralizing antibodies against the Omicron BA.4 and XBB.1.5 pseudovirus. However, the XBB.1.5 antigen sera showed low neutralizing antibody titers against the WT, Beta, and Delta pseudoviruses (**Figure 3B**).


In a second set of data (**Figure 4**), we tested the same sera against five of the Omicron pseudoviruses: BA.2.75.2, BQ.1.1., BA.4, XBB.1.5, and XBB.1.16. The sera of the mice vaccinated with XBB.1.5-RBD showed excellent cross-neutralization towards all tested omicron pseudoviruses. For most of the other sera, we observed a substantial lack of cross-neutralization against the Omicron variants.



**Figure 4.** Rader plot of IC50 values of neutralizing antibody against a panel of five omicron SARS-CoV-2 pseudoviruses (XBB.1.5, BA.4/5, XBB.1.16, BQ.1.1, BA.2.75.2) Sera are from mice that were immunized twice with one of five recombinant RBD vaccines adjuvanted with Alum+CpG.

Using a Spearman correlation on the IC50 values, we illustrate the number of different neutralizing epitopes comparing the RBD antigens of SARS-CoV variants. While we noticed a positive gradient of similarity (**Figure 5, Blue**) among the RBDs of three earlier variants of concern (WT, Beta, and Delta), We found a negative correlation coefficient value comparing WT, Beta, Delta RBDs to Omicron BA.4/5 and XBB.1.5 RBDs (**Figure 5, Red**). Likewise, Omicron RBD BA.4/5 and XBB.1.5 share many neutralizing epitopes, while no correlation was found with the earlier RBDs.





**Figure 5.** Non-parametric rank matrix of five variant RBD vaccines based on Spearman's rank correlation coefficients that were calculated based on the IC50 values against different SARS-CoV-2 pseudovirus variants.

#### 4. Discussion

In the fourth year of the pandemic, immune escape variants of SARS-CoV-2 continue to pose a significant threat to global health. Omicron variants such as BA.2.75.2, BQ.1.1, XBB.1.5, and XBB.1.16 are characterized by a large number of mutations with increased transmissibility, and more pronounced immune evasion. As these new variants emerge, intra-VOC recombination events are commonly observed, leading to an ever-increasing cascade of sub-lineages. Current data indicates that the XBB.1.5 strain is highly resistant to monoclonal antibodies and convalescent plasma treatment [7,18]. XBB subvariants high transmissibility, and its high number of mutations might have contributed to immune evasion. Thus, the vaccines that only targeted the ancestral virus are not as successful in raising neutralizing antibodies against the BA.5 or XBB subvariants. In addition, bivalent boosters that contain both ancestral and BA.4/5 antigens are ineffective against Omicron-derived next-generation subvariants, including XBB.1.5 [19]. Therefore, the broader immune response generated by our XBB.1.5 antigen against additional Omicron subvariants is encouraging for current circulating strains and potential future variants.

In this study, we compared the efficacy of various vaccines based on the RBD of the SARS-CoV-2 spike protein, with the goal of better characterizing cross-neutralization. We observed minimal heterologous cross-neutralization against the XBB.1.5 pseudovirus with sera generated with prior versions of viral RBD. Even immunizations with the more recent BA.4/5 RBD generated only partial cross-neutralization of XBB.1.5 pseudovirus, mirroring results published for breakthrough human infections of BA.4 that did not generate adequate antibodies that protect against XBB subvariants[19]. Our results also re-emphasized that, as compared to the cross-neutralization among the earlier three variants (like WT, Beta, and Delta), Omicron variants have deviated enough to successfully escape neutralization by parental vaccines. This was exemplified by the neutralization data of WT, Beta, and Delta RBDs against both BA 4 and XBB.1.5 pseudoviruses in this study. The IgG antibody binding data from ELISAs throws light on the pattern of almost complete binding of our ancestral RBDs (WT, Beta, and Delta) to all the five proteins, including BA. 4/5 and XBB 1.5, still showing low functional neutralization titers against BA.4/5 and XBB variant pseudoviruses.

In addition to the reported immune imprinting caused by previous vaccinations [19], the antigenic diversities caused by mutations on spike proteins [20] between a variety of Omicron sub-variants may further dampen the neutralization process and reduce the immune response against future variants. This is especially true in the case of XBB sub-lineages where the neutralizing

antibodies elicited against XBB sub-variants were very low after BA.5 infection, indicating the evolution of the XBB sub-lineage further away from BA.4/5 [19]. We have observed weak cross-neutralization from BA.4/5 RBD generated sera against XBB.1.5 and XBB.1.16 subvariants, while the inverse combination of XBB.1.5 RBD antigen against BA.4/5 pseudovirus offered very strong neutralizing antibody titers. The current study shows the value of the XBB.1.5 RBD antigen with its increased number of mutations in its RBD as a better vaccine candidate with successful cross-neutralization capacity against all subvariants of Omicron such as BA.4/5, BA.2.75, BQ.1.1, XBB.1.16, and XBB.1.5. Such levels of neutralization against those same variants were not imparted by BA.4/5 RBD, underscoring the importance of developing the next generation of COVID-19 vaccines customized towards XBB subvariant sequences. Further experiments using an XBB.1.5 boost after vaccination with the WT and BA.4/5 RBDs will mimic the real-world scenario and yield valuable information about the cross-protection levels against current and potential future versions of XBB derivatives.

**Author Contributions:** Conceptualization: SRT, BK, AC, LV, PMG, BZ, US, MEB, PJH, and JP; methodology: SRT, RA, MJV, JL, ZL, RTK, YC, BK, LV, NYI, CP, NU, WHC, SS, KG, JTK, and BZ; formal analysis: SRT, RA, LV, CP, WHC, RK, JTK, BZ, US, MEB, PJH, and JP; writing—original draft preparation: SRT, JTK, US, and JP; writing—review and editing: SRT, RA, MJV, JL, ZL, RK, YC, BK, PMG, LV, NYI, CP, NU, WHC, SS, KG, JTK, BZ, MEB, PJH, and JP; visualization: SRT, YC, WHC, US, and JP; supervision: JL, ZL, CP, WHC, JTK, BZ, US, MEB, PJH, and JP; project administration: PMG and AC; funding acquisition: MEB and PJH. All authors have read and agreed to the published version of the manuscript.

**Supplementary Materials:** The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

**Funding:** This work was supported by the following funding sources: Robert J. Kleberg Jr. and Helen C. Kleberg Foundation, USA; Fifth Generation, Inc., USA; JPB Foundation, USA,); and Texas Children's Hospital Center for Vaccine Development Intramural Funds, USA. In addition, we gratefully acknowledge donations from various private individuals received in support of our vaccine development program.

**Data Availability Statement:** The data that support the findings of this study are available on request from the corresponding author.

**Acknowledgments:** We are grateful to Dr. Vincent Munster (NIAID) for providing the spike expression plasmid for SARS-CoV-2, and thank Dr. Paula Aulicino for many discussions on SARS-CoV-2 variants.

**Conflicts of Interest:** Several of the authors are co-inventors of a COVID-19 recombinant protein vaccine technology owned by Baylor College of Medicine (BCM) that was license by BCM non-exclusively and with no patent restrictions to several companies committed to advancing vaccines for low- and middle-income countries. The co-inventors have no involvement in license negotiations conducted by BCM. Similar to other research universities, a long-standing BCM policy provides its faculty and staff, who make discoveries that result in a commercial license, a share of any royalty income.

## References

1. A. M. Carabelli *et al.*, "SARS-CoV-2 variant biology: immune escape, transmission and fitness," *Nat Rev Microbiol*, vol. 21, no. 3, Art. no. 3, Mar. 2023, doi: 10.1038/s41579-022-00841-7.
2. N. K. Channabasappa, A. K. Niranjana, and T. B. Emran, "SARS-CoV-2 variant omicron XBB.1.5: challenges and prospects – correspondence," *Int J Surg*, vol. 109, no. 4, pp. 1054–1055, Mar. 2023, doi: 10.1097/JS9.0000000000000276.
3. V. Gerardi, M. A. Rohaim, R. F. E. Naggari, M. O. Atasoy, and M. Munir, "Deep Structural Analysis of Myriads of Omicron Sub-Variants Revealed Hotspot for Vaccine Escape Immunity," *Vaccines (Basel)*, vol. 11, no. 3, p. 668, Mar. 2023, doi: 10.3390/vaccines11030668.
4. P. Qu *et al.*, "Enhanced evasion of neutralizing antibody response by Omicron XBB.1.5, CH.1.1, and CA.3.1 variants," *Cell Rep*, vol. 42, no. 5, p. 112443, May 2023, doi: 10.1016/j.celrep.2023.112443.
5. C. He *et al.*, "A recombinant spike-XBB.1.5 protein vaccine induces broad-spectrum immune responses against XBB.1.5-included Omicron variants of SARS-CoV-2," *MedComm*, vol. 4, no. 3, p. e263, 2023, doi: 10.1002/mco2.263.

6. D. Li *et al.*, “Neutralization of BQ.1, BQ.1.1, and XBB with RBD-Dimer Vaccines,” *N Engl J Med*, vol. 388, no. 12, pp. 1142–1145, Mar. 2023, doi: 10.1056/NEJMc2216233.
7. M. Imai *et al.*, “Efficacy of Antiviral Agents against Omicron Subvariants BQ.1.1 and XBB,” *N Engl J Med*, vol. 388, no. 1, pp. 89–91, Jan. 2023, doi: 10.1056/NEJMc2214302.
8. J. Pollet *et al.*, “Receptor-binding domain recombinant protein on alum-CpG induces broad protection against SARS-CoV-2 variants of concern,” *Vaccine*, vol. 40, no. 26, pp. 3655–3663, Jun. 2022, doi: 10.1016/j.vaccine.2022.05.007.
9. J. Pollet *et al.*, “SARS-CoV-2 RBD219-N1C1: A yeast-expressed SARS-CoV-2 recombinant receptor-binding domain candidate vaccine stimulates virus neutralizing antibodies and T-cell immunity in mice,” *Hum Vaccin Immunother*, vol. 17, no. 8, pp. 2356–2366, Aug. 2021, doi: 10.1080/21645515.2021.1901545.
10. M. Pino *et al.*, “A yeast-expressed RBD-based SARS-CoV-2 vaccine formulated with 3M-052-alum adjuvant promotes protective efficacy in non-human primates,” *Sci Immunol*, vol. 6, no. 61, p. eabh3634, doi: 10.1126/sciimmunol.abh3634.
11. P. J. Hotez *et al.*, “From concept to delivery: a yeast-expressed recombinant protein-based COVID-19 vaccine technology suitable for global access,” *Expert Rev Vaccines*, vol. 22, no. 1, pp. 495–500, 2023, doi: 10.1080/14760584.2023.2217917.
12. “Statement on the antigen composition of COVID-19 vaccines.” <https://www.who.int/news/item/18-05-2023-statement-on-the-antigen-composition-of-covid-19-vaccines> (accessed Aug. 18, 2023).
13. J. Pollet *et al.*, “SARS-CoV-2 RBD219-N1C1: A yeast-expressed SARS-CoV-2 recombinant receptor-binding domain candidate vaccine stimulates virus neutralizing antibodies and T-cell immunity in mice,” *Human Vaccines & Immunotherapeutics*, vol. 17, no. 8, pp. 2356–2366, Aug. 2021, doi: 10.1080/21645515.2021.1901545.
14. J. Lee *et al.*, “Process development and scale-up optimization of the SARS-CoV-2 receptor binding domain-based vaccine candidate, RBD219-N1C1,” *Appl Microbiol Biotechnol*, vol. 105, no. 10, pp. 4153–4165, 2021, doi: 10.1007/s00253-021-11281-3.
15. W.-H. Chen *et al.*, “Yeast-expressed recombinant SARS-CoV-2 receptor binding domain RBD203-N1 as a COVID-19 protein vaccine candidate,” *Protein Expr Purif*, vol. 190, p. 106003, Feb. 2022, doi: 10.1016/j.pep.2021.106003.
16. J. Nie *et al.*, “Quantification of SARS-CoV-2 neutralizing antibody by a pseudotyped virus-based assay,” *Nat Protoc*, vol. 15, no. 11, Art. no. 11, Nov. 2020, doi: 10.1038/s41596-020-0394-5.
17. D. Mannar *et al.*, “SARS-CoV-2 Omicron variant: Antibody evasion and cryo-EM structure of spike protein-ACE2 complex,” *Science*, vol. 375, no. 6582, pp. 760–764, Feb. 2022, doi: 10.1126/science.abn7760.
18. R. Uraki *et al.*, “Humoral immune evasion of the omicron subvariants BQ.1.1 and XBB,” *The Lancet Infectious Diseases*, vol. 23, no. 1, pp. 30–32, Jan. 2023, doi: 10.1016/S1473-3099(22)00816-7.
19. J. Yang *et al.*, “Low levels of neutralizing antibodies against XBB Omicron subvariants after BA.5 infection,” *Sig Transduct Target Ther*, vol. 8, no. 1, Art. no. 1, Jun. 2023, doi: 10.1038/s41392-023-01495-4.
20. W. Wang *et al.*, “Antigenic cartography of well-characterized human sera shows SARS-CoV-2 neutralization differences based on infection and vaccination history,” *Cell Host Microbe*, vol. 30, no. 12, pp. 1745–1758.e7, Dec. 2022, doi: 10.1016/j.chom.2022.10.012.

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.