

Article

Not peer-reviewed version

---

# High Resolution Computed Tomography in Middle ear Cholesteatoma: How Much do We Need it?

---

[Eugen Horatiu Stefanescu](#) <sup>\*</sup>, [Nicolae Constantin Balica](#), [Sorin Bogdan Motoi](#) <sup>\*</sup>, [Laura Grigorita](#),  
[Madalina Georgescu](#) <sup>\*</sup>, Gheorghe Iovanescu

Posted Date: 5 September 2023

doi: 10.20944/preprints202309.0205.v1

Keywords: High resolution computed tomography; cholesteatoma surgery; very good correlation



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

## Article

# High Resolution Computed Tomography in Middle ear Cholesteatoma: How Much do We Need it?

Eugen Horatiu Stefanescu <sup>1,\*</sup>, Nicolae Constantin Balica <sup>1</sup>, Sorin Bogdan Motoi <sup>2,\*</sup>, Laura Grigorita <sup>3</sup>, Madalina Georgescu <sup>4,\*</sup>, Gheorghe Iovanescu <sup>1</sup>

<sup>1</sup> Department of Otolaryngology, Victor Babeș University of Medicine and Pharmacy, Timișoara, Romania; stefanescu@umft.ro; balica@umft.ro; giovanescu@umft.ro

<sup>2</sup> Department of Radiology and Medical Imaging, Victor Babeș University of Medicine and Pharmacy, Timișoara, Romania; motoi.sorin@umft.ro

<sup>3</sup> Department of Anatomy and Embriology, Victor Babeș University of Medicine and Pharmacy, Timișoara, Romania; grigorita.laura@umft.ro

<sup>4</sup> Discipline of General Surgery and Qualified Care in Surgical Specialties, Carol Davila University of Medicine and Pharmacy, București, Romania; madalina.georgescu@umfcd.ro

\* Correspondence: stefanescu@umft.ro; motoi.sorin@umft.ro; madalina.georgescu@umfcd.ro

**Abstract:** *Background and Objectives:* The diagnosis of cholesteatoma is usually clinic and the only efficient treatment is surgical. High Resolution Computed Tomography (HRCT) is not considered absolutely necessary for the management of an uncomplicated cholesteatoma but unsuspected situations from a clinical point of view can be discovered using the scans, warning the surgeon. Our objective is to compare HRCT scan information with intraoperative findings in patients with cholesteatoma and analyze the usefulness of a preoperative HRCT scan from a surgical point of view. *Materials and Methods:* This is a prospective descriptive study, conducted in the department of Otolaryngology, Victor Babes University of Medicine and Pharmacy Timisoara, Romania from May 2021 to April 2022. It was carried out on 46 patients with clinical diagnosis of cholesteatoma who were consequently operated in our department. All patients received full clinical and audiological examination. In all cases a HRCT scan was performed preoperatively, as mandatory investigation. Preoperative HRCT scans were analyzed and their findings were compared to the intraoperative notes. The two sets of observations were analyzed using standard statistical methods. *Results:* Extensive cholesteatoma was the commonest type of disease involving 46% of the patients, followed by pars flaccida cholesteatoma (35%) and pars tensa cholesteatoma (19%). Eroded scutum was the most frequent lesion involving 70% of the patients, followed by incus erosion (67%). Comparison of the HRCT and intraoperative findings revealed very good correlation for tegmen tympani erosion, sigmoid plate erosion, scutum and malleus erosion, and moderate to good correlation for lateral semicircular canal erosion, incus and stapes erosion and Fallopian canal erosion. *Conclusions:* HRCT is a valuable tool in preoperative assessment of cholesteatoma, helping in making surgical decisions. It can accurately predict the extent of disease and is helpful for detecting unapparent dangerous situations. However it is not very accurate in detecting Fallopian canal and stapes erosion.

**Keywords:** High resolution computed tomography; cholesteatoma surgery; very good correlation

## 1. Introduction

When discussing a topic we usually start with its definition. When the topic is cholesteatoma the problems already start there. And they do continue in almost every aspect, being quite difficult to name one where there are no controversies. The old definition of "skin in the wrong place" is more misleading than helpful. Cholesteatoma is a benign, cystic lesion involving the temporal bone, which is derived from an abnormal growth of keratinizing squamous epithelium [1]. It consists of an outer lining of stratified squamous epithelium, an inner keratin mass and an external peri-matrix that produces enzymes, destroying the surrounding tissues [2]. Middle ear cholesteatoma has a higher incidence in individuals younger than 50 years of age but can occur at any age [3]. Cholesteatoma in

children tends to behave in a different manner than in adults; it has a more aggressive pattern of growth and recurs more frequently.

The clinical features of cholesteatoma vary considerably, ranging from an asymptomatic disease to life-threatening complications [4]. Even if today the complication rate has decreased significantly the danger is still there and a warning sign of any kind is always helpful for the physician. The management of this disease is strictly surgical and continues to be a challenge for otolaryngologists, and sometimes quite a nightmare for both patient and surgeon.

The temporal bone has a very complicated architecture and contains delicate structures like the ossicular chain, inner ear, facial nerve, and even vital ones like the internal carotid artery. Cholesteatoma may damage all of them and it also may involve adjacent structures like dura mater, the temporal lobe of the brain, and sigmoid sinus leading to both extra-cranial and intra-cranial complications as local invasion is a characteristic of the disease. It is considered to be a benign lesion but given the complications encountered one cannot be so sure this is the right word.

There are also many anatomic variants of the structures in the middle ear and the very narrow space inside the temporal bone makes discovering the lesions and tackling the complications quite challenging, so any preoperative information is highly appreciated.

HRCT of the temporal bone is one of the imaging modalities used to evaluate the extent of the disease and the lesions produced by cholesteatoma in and out the temporal bone prior to surgery and it is considered to be accurate as its slice thickness is usually less than 1 mm. Using special algorithms HRCT scans offer an excellent resolution [5]. On HRCT scans, cholesteatoma appears as a soft-tissue mass in the middle ear cavity and mastoid with associated signs of surrounding bony erosion [3]. However, HRCT cannot accurately differentiate cholesteatoma from inflammatory/granulation tissue or scar tissue inside the middle ear [6]. The sinus tympani and facial recess are two hidden areas of the middle ear and their involvement by cholesteatoma can be detected intraoperatively, especially if using an endoscope, but can also be identified preoperatively on a HRCT scan.

In attic cholesteatoma, erosion of the scutum in the coronal view can be assessed on HRCT scans and it is a useful sign for early diagnosis [7]. Ossicular chain erosion is a common feature in cholesteatoma, which can result in more or less significant conductive hearing loss. It can vary from small erosion of the long process of the incus and incudo-stapedial joint, which is the most frequent situation, to complete destruction of the entire ossicular chain. The interposition of the cholesteatoma between the tympanic membrane and stapes, no matter if intact or if its superstructure is eroded, can improve the sound transmission to the inner ear and be deceiving regarding the patients actual hearing level. Other dangerous site of erosion is the dome of the lateral semicircular canal (LSC), just above the second genu of the facial nerve canal. Speaking of the facial nerve canal the most frequent portion that can be dehiscent or eroded in cases of cholesteatoma is the tympanic one, usually just above the oval window area [8]. These are some of the reasons why such preoperative "inside information" are useful for planning the surgical approach and this can be best achieved using HRCT. Early detection and treatment of such dangerous situations has considerably reduced the morbidity and mortality in cholesteatoma [9].

## 2. Materials and Methods

This is a prospective descriptive study, conducted in the department of Otolaryngology, Head and Neck Surgery of the Victor Babes University of Medicine and Pharmacy Timisoara, Romania from May 2021 to April 2022. The study protocol was approved by the Ethics Committee of Victor Babes University of Medicine and Pharmacy Timisoara. The present study included 46 patients with clinical diagnosis of chronic otitis media with cholesteatoma in which we performed tympanomastoid surgery, both open and closed techniques, with ossicular chain reconstruction. All patients received preoperative full clinical and audiological examination. Collected clinical data included information about tympanic cavity status (tympanic membrane, ossicles, and mucosa lesions), hearing and vestibular status and facial nerve function. All patients underwent preoperative HRCT of the temporal bone as part of the mandatory investigations. All surgeries were performed between 3 and 8 weeks after imaging by the same surgical team under the surgical microscope. We only used

the endoscope to assess the extension of the disease to sinus tympani and to check for the complete removal of the lesions. Surgical findings were recorded by the surgical team.

Interpretation of preoperative HRCT images were focused to define the following: location and extent of cholesteatoma; bony erosions of the following structures: scutum, tegmen tympani, facial nerve (FN) canal and inner ear (LSC, promontory); integrity of the ossicular chain. All HRCT scans were assessed by a single radiologist with over 15 years experience in head and neck imaging.

The operative findings were compared to the radiological findings, assessing the usefulness of a preoperative HRCT scan in describing the status of the middle ear structures in cases of cholesteatoma with or without complications. Sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) of HRCT scans, and Cohen's Kappa coefficient (to evaluate their agreement with the intraoperative findings) were calculated for the following features, considered to be of interest for surgery- scutum, tegmen tympani, sigmoid sinus plate, LSC, FN canal, malleus, incus and stapes. For statistical analysis we used Microsoft Excel 2013 and Statistical Package for Social Sciences (SPSS) version 26 (IBM Corp., Armonk, NY, USA) The kappa coefficient in the range 0–0.40, 0.41–0.75, and 0.76–1.0 indicates poor, moderate to good, and very good correlation, respectively.

We did not include in the study cases with any kind of previous surgery in the same ear, cholesteatoma recurrences and patients who were operated later than 8 weeks after performing imaging. Written informed consent was obtained from all patients prior to data collection.

#### Scan protocol

HRCT scans were obtained on GE Revolution EVO (General Electric, Boston, USA), a 128 slice

CT scanner. The patient was laying in the supine position. The topogram was made in the lateral incidence, which includes the area between the tip of the mastoid and the arcuate eminence of the temporal bone as the lower and upper scanning point, respectively; scanning was not performed with an inclined gantry.

Axial projections were obtained by serial 1 mm sections of the temporal bone with the plane along the line joining the infra-orbital rim and external auditory meatus, perpendicular to the table with  $20 \times 0.625$  collimation. The images were reconstructed using the high-resolution bone algorithm in the axial plane with 0.625 mm section thickness, and a field of view (FOV) of 100-110 mm, with a matrix size of  $512 \times 512$ . The effective amperage in the tube was modulated according to the patient's age- between 130 mAs (for a newborn) up to 400mAs (for an adult). The voltage in the tube was between 110-140 kV, usually 120 kV.

The acquisition was in the spiral mode, which ensures a better reformation in the coronal and sagittal plane. The reconstruction in the axial plane is the basic one and includes sections parallel to the lateral semicircular canal; simultaneously with the reconstruction in the three planes, a set of images in the axial plane with a thickness of 2 mm was also obtained, which helps to visualize the tissue structures within the cholesteatoma as a whole.

The administration of the contrast substance was not always indicated in cholesteatoma involving the middle ear, but it becomes mandatory in situations where there is a suspicion of evolution towards complication: abscesses, coalescent otomastoiditis and sigmoid sinus thrombosis.

### 3. Results

Chronic otitis media with cholesteatoma represents 41% of the chronic otitis media cases operated in our department in 2022. There were 46 patients enrolled in our study, including 6 children, ages 8 to 15 years old. The oldest patient was 73 years old. The mean age of our patients was  $37.3 \pm 10.8$  years. The largest number of cases, 19 (41%) were in the 31- 40 year old group, and among them 11 (57%) were male and 8 (43%) were female. There was generally a slight male preponderance in our study population- 24 (52%) male and 22 (48%) female patients, with a male: female ratio of 1.09:1. Extensive cholesteatoma was the most commonly encountered type, detected in 21 patients (46%), and followed by pars flaccida cholesteatoma encountered in 16 patients (35%) and pars tensa cholesteatoma in 9 patients (19%).

The erosion of the scutum was the most common lesion but only by a narrow margin, and was found in 32 patients (70%). The other destructive lesions were: eroded incus, mainly the long process, in 31 patients (67%), eroded malleus in 21 patients (46%), eroded stapes in 14 patients (30%), significantly thinned/ eroded tegmen tympani in 10 patients (22%), dehiscent/ eroded facial nerve canal in 8 cases (17%), lateral semicircular canal fistula in 4 cases (9%), and eroded sigmoid sinus plate in 3 cases (7%). The patients enroled in the study presented the following complications – 2 patients with facial nerve palsy (grade III and IV House- Brackmann, respectively) and 4 patients with vertigo. All the patients with vertigo were correctly assessed from an imaging point of view. There was an extra case described as having a lateral semicircular canal fistula on HRCT scans which wasn't confirmed during surgery. When analyzing alterations of tegmen tympani we considered both erosion and thinning together as its clinical and surgical significance is comparable.

Table 1 shows the sensitivity, specificity, positive predictive and negative predictive values and correlation level for the considered HRCT findings.

**Table 1.** Accuracy of preoperative HRCT in detecting cholesteatoma lesions.

HRCT Findings	Sensitivity	Specificity	PPV	NPV	Correlation
Tegmen tympani erosion/thinning	90.0	88.8	69.2	96.9	0.97
Sigmoid plate erosion	100	97.6	75.0	100	0.87
Malleus erosion	85.7	96.0	94.7	88.8	0.82
Scutum erosion	87.5	92.8	96.5	76.4	0.76
LSC erosion	75.0	97.6	75.0	97.6	0.68
Incus erosion	80.6	86.6	92.5	68.4	0.62
Stapes erosion	57.1	84.3	61.5	81.8	0.44
Facial canal dehiscence/erosion	50.0	92.1	57.0	89.7	0.42

The greatest correlation between imaging and surgery was in detection of the tegmen tympani erosion/thinning ( $\kappa=0.97$ ). We also obtained very good imaging-surgical correlation in detection of sigmoid sinus plate dehiscence ( $\kappa=0.87$ ), malleus erosion ( $\kappa=0.82$ ), and scutum erosion ( $\kappa=0.76$ ). We obtained moderate to good results in detecting lateral semicircular canal dehiscence ( $\kappa=0.68$ ), incus erosion ( $\kappa=0.62$ ), stapes erosion ( $\kappa=0.44$ ), and facial nerve canal dehiscence ( $\kappa=0.42$ ). There were no poor imaging-surgical correlation for any of the investigated lesions in our study. The poorest correlation was in facial nerve canal dehiscence/erosion ( $\kappa=0.42$ ) and stapes erosion ( $\kappa=0.44$ ). The best correlation was in cases with extensive and severe lesions, as it was expected preoperatively.

#### 4. Discussion

The ability of HRCT to detect cholesteatoma and its complication, prior to surgery, is well known. It is especially helpful in detecting minimal erosion of the ossicular chain and other dangerous area like tegmen tympani, lateral semicircular canal, Fallopian canal [10]. Such information are very useful when discussing to the patient and getting informed consent for surgery. It is still worth remembering that HRCT can not accurately differentiate cholesteatoma from granulation tissue, and cholesterol granuloma, which are also encountered in chronic suppurative otitis media. All our cases had a clinical diagnosis of cholesteatoma prior to imaging so this was not our main concern, but cholesteatoma was confirmed and its extension accurately described on imaging.

Age- In our study we found that the majority of patients were aged between 31 and 40 years of age, involving 19 (41%) cases. In other studies it was observed a much younger age prevalence of less than 30 years of age [11,12].

Gender prevalence- In the present study the majority of patients were male (52%), but only by a thin margin, with a male to female ratio of 1.09: 1. Kemppainen et al., also reported that cholesteatoma was more frequent in men under the age of 50 years.[13]. Other studies reported a much higher male predominance with a male: female ratio of 1.39:1 [14].

Location and extent of cholesteatoma- In our study extensive cholesteatoma was the most commonly encountered type, detected in 46% of patients followed by pars flaccida cholesteatoma in 35% of patients and pars tensa cholesteatoma present in 19% of patients. Same prevalence was observed in the study done by Gomma et al. where extensive cholesteatoma were detected in 35.7% of patients. Also commonly detected was pars flaccida type [12]. In other studies the most frequently encountered type was pars flaccida cholesteatoma: a study on 30 cholesteatoma cases showed lesions in the epitympanum in 73.3% of patients followed by mesotympanum [15], and in another one pars flaccida cholesteatoma was described in 50.7% of cases, followed by extensive cholesteatoma in 30.1% of patients [16]. Pars tensa cholesteatoma was the least encountered type in most of the studies.

Accurate assessing of the extension of the lesions may provide crucial information regarding the most suitable surgical approach considering the given anatomy of the middle ear and mastoid. It is notorious the controversy between the surgeons supporting either the open or the closed technique in the treatment of cholesteatoma. However the closed technique might be extremely difficult in some temporal bones and this can be foreseen on imaging. There are also situations in which an open technique is not required because of limited extension of cholesteatoma [17].

From a surgical point of view the accuracy in detecting lesions of the delicate structures of the middle ear presents great importance and this is why several aspects regarding this topic deserve some comments.

Scutum erosion- It is considered to be an early sign of pars flaccida cholesteatoma but it does not appear in all cases. In our study it was detected in 32 cases (70%) with a sensitivity of 87,5% and a specificity of 92,8%. The correlation was still very good but borderline ( $\kappa=0,76$ ). In other study the scutum was eroded in 55.5% of the cholesteatoma cases [16]. Rai et al. reported scutum erosion in 65% of the patients [18]. The PPV was very good in our study (96.5%). In other studies scutum erosion was accurately predicted in 91% of cases [19,20]. However Sunitha reported no imaging-surgical correlation for scutum erosion [21]. Suat Keskin believes that differences between imaging and intraoperative findings regarding scutum erosion may be due to inappropriate angles of the coronal sections [22]. From a surgical point of view the erosion of the scutum may influence the approach to the ossicular chain, especially to incus and malleus.

Sinus plate erosion- In our study, erosion of the sigmoid sinus plate was described on HRCT in 4 cases but intraoperatively only 3 cases had sinus plate erosion so only 75% of cases were accurately detected by HRCT with relatively lower sensitivity when compared to other studies. In a study by Dutta et al. HRCT sensitivity was 100%, and specificity was also 100% for sinus plate erosion [11]. C Shah reported sensitivity and specificity of HRCT in detecting sigmoid erosion to be 91.7% and 95.25% respectively [17]. Kanotra et al. reported the sensitivity, specificity, positive and negative predictive value to be 100% [23]. Sinus plate erosion is quite a rare occurrence and this can explain why there are such differences between studies. Knowing that that the sinus plate is eroded requires special attention and careful exploration of the region even if there are no clinical signs of sigmoid sinus pathology. It can also alert the surgeon and prevent from inadvertent opening of the sinus wall with massive bleeding and prolonged surgical time.

Tegmen tympani erosion/thinning- We evaluated erosion and significant thinning of the tegmen tympani together, but not all studies did so. Probably this is one reason why we obtained such good specificity and sensitivity compared to others. HRCT finding shows eroded/thinned tegmen tympani in 13 cases but intraoperatively it was present in 9 cases (19%) with sensitivity 90% and specificity 88,8%. Other study identified tegmen tympani erosion in 38% of the cases [16]. In a study done by Jamal et al. tegmen tympani erosion was seen 30% of the patients [24]. Kanotra et al. reported a sensitivity of 100% in detecting tegmen tympani erosion by HRCT which was higher than the current study [23]. A high specificity rate of 95% was reported by Gerami et al. [25]. A poor sensitivity rate of HRCT in detecting tegmen tympani erosion was reported by Jackler et al. [9] and O'Reilly et al. [26], while a moderate sensitivity rate was described by Vlastarakos et al. [27] and Chee and Tan [10]. From a surgical point of view it is an invaluable information for the dissection of the cholesteatoma matrix away from that area even in the absence of dura related complications. Probably the poor

imaging-surgical correlation for tegmen tympani erosion may be due to partial volume of both the tympanic cavity and cerebral soft tissue [22].

Lateral semicircular canal erosion- This structure may be eroded by cholesteatoma especially in its dome region on the medial wall of the epitympanum, close to the second genu of the facial nerve [3]. In our study we found 4 cases (9%) of LSC fistula. Other studies noted LSC fistula in just 4.7% of the cases and even lower (4% of the patients) [16,28]. In our study HRCT scans assessed the LSC fistula cases with a sensitivity of 75% and specificity of 97.6%. In a study done by Gaurano et al. on 64 patients there were 4 cases (6.3%) that had labyrinthine fistula found on HRCT, but only 3 (4.7%) were confirmed intraoperatively [19]. Usually HRCT accurately predicted lateral semicircular canal fistula with excellent sensitivity and good specificity. Mafee et al., Chee et al., and Rocher et al., reported HRCT to be 100% sensitive in detecting lateral semicircular canal fistula [3,10,20]. Knowing in advance about the suspicion and location of the fistula is very important for the surgeon as dissection of the matrix of the cholesteatoma in that area has to be done very carefully avoiding unnecessary manipulation and suction. The closure of the fistula is of utmost importance for the inner ear function.

FN Canal dehiscence/erosion- Fallopian canal dehiscence/erosion is a relatively common finding in cholesteatoma affecting the middle ear, usually occurring in the tympanic portion of the facial nerve canal.[8,29] We found facial nerve canal dehiscence/erosion in 8 patients (17%). In other study the facial canal dehiscence was observed in 34.9% of the cholesteatoma cases [16]. The study by Jamal et al. reported facial canal dehiscence in 30% of the patients [24]. The sensitivity and specificity obtained in our study were 50% and 92.1% respectively with a moderate to poor correlation between imaging and surgery ( $\kappa=0.42$ ). Other studies presented similar observations with a sensitivity ranging from 33.3% to 83% and specificity between 60% and 97%. It is well known that the Fallopian canal can be so thin, as to appear dehiscent on a CT scan [30]. It's good to remember that the Fallopian canal may be dehiscent even in a healthy ear. The detection of the facial nerve dehiscence/erosion on imaging should influence the decision and timing of surgery, being a good reason to operate earlier, and alert the surgeon when working in that area. One should always consider the proximity of the stapes and facial nerve, especially when dissecting cholesteatoma matrix off these structures.

Ossicular chain- It is usually involved by cholesteatoma of the middle ear even if not always eroded or discontinued. This situation presents great importance for the surgeon as he/she has to drill and dissect a lot around the ossicular chain. Touching the ossicular chain during drilling might generate severe SNHL, and should be avoided at any cost. We found ossicular chain erosion in 37 out of 46 patients (80%). Incus was the most eroded ossicle, found in 67% of the patients in our study. In another study incus was found to be eroded in 60.3% of the cases, followed by malleus in 58.7% and stapes in 47.6% of the patients [16]. Manik et al. also showed incus to be the most commonly affected ossicle in 70% of the patients, followed by malleus in 42% and stapes in 34% of the patients [31]. In our study the most accurate information was regarding the erosion of the malleus, the head of the malleus being the most affected site, with a sensitivity of 85.7%, and a specificity of 96%, ( $\kappa=0.82$ ). It was followed by erosion of the incus - sensitivity of 80.6 and specificity 86.6%, ( $\kappa=0.62$ ). Other authors observed comparable sensitivity of 87% and 85% respectively [11,18]. Stapes is the least eroded ossicle in most of the studies. However, the poorest imaging- surgical correlation was regarding stapes erosion. In our study sensitivity was 57.1% and specificity of 84.3%, ( $\kappa=0.44$ ), which is borderline between moderate and poor correlation. Other authors also reported poor radiologic- surgical correlation in stapes erosion and stated that the small size of the bone may be the cause of poor detection on HRCT scans [32,33]. However, if an ossicle is involved by the cholesteatoma it has to be removed no matter if eroded or not. The only exception is the stapes footplate, in which case the cholesteatoma matrix should be carefully dissected off the structure. Anything else can and should be removed so preoperative information regarding erosion of the malleus, incus and even stapes superstructure are not essential for the ossicular chain reconstruction. The only unknown factor is which type of prosthesis should be used for the reconstruction of the sound transmission mechanism and this depends on the status of stapes superstructure- present and functional or absent/ eroded. Either way the stapes footplate needs to be mobile. However you can

always use a total ossicular replacement prosthesis (TORP) even if the stapes superstructure is intact by bypassing it.

HRCT is still considered the best available imaging method to describe the cholesteatoma of the middle ear benefiting both the patient and the surgeon. Early diagnosis in cholesteatoma is also crucial and there are important findings that can alert the physician such as erosion of scutum.

This study demonstrated good to excellent correlation between temporal bone HRCT findings and intraoperative lesions, particularly in tegmen tympani erosion, sigmoid plate dehiscence, malleus erosion, and scutum erosion. Our findings contrast with other reports which showed an excellent radio-surgical correlation for the stapes ( $\kappa=0.94$ ), and semicircular canals ( $\kappa=0.80$ ), but poorer for the tegmen tympani erosion/thinning ( $\kappa=0.65$ ) and facial nerve canal ( $\kappa=0.3$ ) [10]. Rocher and colleagues revealed good to excellent correlation for the scutum erosion, and the lateral semicircular canal dehiscence ( $\kappa>0.75$ ), and good correlation for tegmen tympani erosion ( $\kappa=0.6$ ), but poor correlation for the facial nerve canal dehiscence ( $\kappa<0.4$ ) [20].

The very good correlation between the findings on HRCT scans and intra-operative lesions may determine a higher degree of suspicion for unexpected problems during surgery and may improve the success rate of cholesteatoma surgery. The limitations and errors in interpretation may be improved by newer imaging technology.

## 5. Conclusions

The results of the present study indicates that HRCT is a most valuable preoperative imaging modality to evaluate cholesteatoma of the middle ear, playing an important role and guiding the surgical management. HRCT scans can accurately predict the extent of the disease and are helpful in detecting erosion of tegmen tympani, lateral semicircular canal fistula, sigmoid sinus plate and ossicular chain erosion with considerably high sensitivity and specificity. Accurate information on cholesteatoma lesions that are offered to the surgeon by the means of HRCT allow for more limited procedures to be performed when eradicating the disease while preserving the function of the middle and inner ear. However, this imaging technique is not yet able to accurately distinguish between cholesteatoma and granulation tissue, and cannot accurately detect facial nerve dehiscency/erosion and stapes erosion.

**Author Contributions:** The authors have contributed to this article as follows: Conceptualization, E.H.S.; Methodology, E.H.S., N.C.B., S.B.M.; Software, L.G.; Validation, E.H.S., N.C.B., M.G., G.I.; Investigation, E.H.S., M.G., S.B.M.; Data curation, E.H.S., S.B.M. L.G.; Writing—E.H.S., L.G.; Writing—review and editing, E.H.S., M.G., S.B.M.; Supervision, N.C.B., G.I.; Project administration, N.C.B., G.I.; All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki, and approved by the Ethics Committee of the Victor Babeş University of Medicine and Pharmacy (57/09.09.2020).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** Data supporting reported results can be found in the archives of the Department of Otolaryngology and Department of Radiology and Medical Imaging of the Victor Babes University of Medicine and Pharmacy Timisoara.

**Acknowledgments:** Nothing to declare

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Oiszewska E, Wagner M, Bernal-Sprekelsen M, et al. Etiopathogenesis of cholesteatoma. *Eur Arch Otorhinolaryngol*. 2004;261(1):6–24.2.
2. Ashutosh R, Soumit M, Yati RD, Rezaul K: Pre-operative HRCT of middle ear pathology with particular reference to intra-operative findings in cholesteatoma. *Int J Acad Med Pharm* 2023; 5 (3); 2303-2307.

3. Mafee MF, Levin BC, Applebaum EL, Campos CF: Cholesteatoma of the middle ear and mastoid . Otolaryngol Clin North Am. 1988, 21:265-8.
4. Lan MY, Lien CF, Liao WH. Using high resolution computed tomography to evaluate middle ear cleft aeration of postoperative cholesteatoma ears. J Chin Med Assoc 2003;66:217-23.
5. Bagul M. High-resolution Computed Tomography Study of Temporal Bone Pathologies. Headache. 2016;32(1):26-66.
6. Jackler RK, Dillon WP, Schindler RA: Computed tomography in suppurative ear disease: a correlation of surgical and radiographic findings. Laryngoscope. 1984, 94:746-52.
7. Mohammadi Gh, Naderpour M, Mousaviagdas M. Ossicular Erosion in Patients Requiring Surgery for Cholesteatoma. Iranian Journal of Otorhinolaryngology. 2012; 24(68): 125-8.
8. Meyer TA, Strunker JR, Chester L, Lambert PR. Cholesteatoma. In: Bailey BJ, Johnson JT. Head and neck surgery otolaryngology. 4th edition; 2006, Philadelphia, Lippincott Williams and Wilkins. Vol (2): 2081-91.
9. Vivek R, Gunasekaran P, Sethurajan S, Adaikappan M. Evaluation of HRCT temporal bone and pathologies. Journal of Evolution of Medical and Dental Sciences. 2014;3(52):12118-27.
10. Chee NC, Tan TY: The value of preoperative high resolution CT scans in cholesteatoma surgery. Singapore Med J. 2001, 2:155-9.
11. Datta G, Mohan C, Mahajan M, Mendiratta V. Correlation of preoperative HRCT findings with surgical findings in unsafe CSOM. J Dental Med Sci. 2014;13(1):120-5.
12. Gomaa MA, Abdel Karim AR, Abdel Ghany HS, et al. Evaluation of temporal bone cholesteatoma and the correlation between high resolution computed tomography and surgical finding. Clin Med Insights Ear, Nose, Throat 2013;6:21-8.
13. Kemppainen H, Puhakka HJ, Sipila MM, Manninen MP, Karma PH. Epidemiology and etiology of middle ear cholesteatomaActaOtolaryngol, 119 (1) (1999), pp. 568-572.
14. Khavasi P et al. International Journal of Otorhinolaryngology and Head and Neck Surgery, March-April 2018, Vol 4, Issue 2, Page 413-17.
15. Jacob A, Sreedhar S, Choolakkaparambu A, Anwar S, Bashir, Nalakath K: Utility of high resolution computed tomography in pre-operative evaluation of cholesteatoma. Int J Otorhinolaryngol Head Neck Surg. 2020, 6:1278.
16. Uz Zaman S, Rangankar V, Muralinath K, et al. (March 01, 2022) Temporal Bone Cholesteatoma: Typical Findings and Evaluation of Diagnostic Utility on High Resolution Computed Tomography. Cureus 14(3): e22730. DOI 10.7759/cureus.22730
17. Shah C, Shah P, Shah S. Role of HRCT temporal bone in pre operative evaluation of cholesteatoma. 2014.
18. Rai T: Radiological study of the temporal bone in chronic otitis media: prospective study of 50 cases . Indian J Otol. 2014, 20:48.
19. Gaurano JL, Joharjy IA. Middle ear cholesteatoma: characteristic CT findings in 64 patients. Annals of Saudi medicine. 2004;24(6):442-7.
20. Rocher P, Carlier R, Attal P, Doyon D, Bobin S, editors. Contribution and role of the scanner in the preoperative evaluation of chronic otitis. Radiosurgical correlation apropos of 85 cases. Annales d'otolaryngologie et de chirurgie cervico faciale: bulletin de la Societe d'oto-laryngologie des hopitaux de Paris; 1995.
21. Sunita M, Sambandan A. Importance of Pre-Operative HRCT Temporal Bone in chronic suppurative otitis media. Odisha J Otorhinolaryngol Head Neck Surg. 2015;9(1):10-3.
22. Keskin S, Çetin H, Töre HG. The correlation of temporal bone CT with surgery findings in evaluation of chronic inflammatory diseases of the middle ear. European Journal of General Medicine. 2011;8(1).
23. Kanotra S, Gupta R, Gupta N, Sharma R, Gupta S, Kotwal S. Correlation of high-resolution computed tomography temporal bone findings with intra-operative findings in patients with cholesteatoma. Indian Journal of Otology. 2015;21(4):280.
24. Jamal S, Rahman A, Mohan C, Srivastava A: Is pre-operative HRCT temporal bone findings consistent with tympano-mastoid surgical findings. Int J Health Clin Res. 2020, 3:104-9.
25. Gerami H, Naghavi E, Wahabi-Moghadam M, Forghanparast K, Akbar MH. Comparison of preoperative computerized tomography scan imaging of temporal bone with the intra-operative findings in patients undergoing mastoidectomy. Saudi medical journal. 2009;30(1):104-8.
26. O'reilly B, Chevretton E, Wylie I, Thakkar C, Butler P, Sathanathan N, et al. The value of CT scanning in chronic suppurative otitis media. The Journal of Laryngology & Otology. 1991;105(12):990-4.

27. Vlastarakos PV, Kiprouli C, Pappas S, Xenelis J, Maragoudakis P, Troupis G, et al. CT scan versus surgery: how reliable is the preoperative radiological assessment in patients with chronic otitis media? European Archives of Oto-rhino-laryngology. 2012;269(1):81-6.
28. Dashottar S, Bucha A, Sinha S, Nema D: Preoperative temporal bone HRCT and intra-operative findings in middle ear cholesteatoma: a comparative study. Int J Otorhinolaryngol Head Neck Surg. 2018, 5:77.
29. Ozbek C, Tuna E, Ciftci O, Yazkan O, Ozdem C. Incidence of fallopian canal dehiscence at surgery for chronic otitis media. Eur Arch Otorhinolaryngol 2009; 266(3): 357-62.
30. Yetiser S, Tosun F, Kazkayasi M. Facial nerve paralysis due to chronic otitis media. OtolNeurotol 2002;23:580-8.
31. Manik S, Dabholkar Y, Bhalekar S, Velankar H, Chordia N, Saberwal A: Sensitivity and specificity of high resolution computed tomography (HRCT) of temporal bone in diagnosing cholesteatoma and its correlation with intraoperative findings. Indian J Otolaryngol Head Neck Surg. 2021, 73:25-9.
32. Sreedhar S, Pujary K, Agarwal AC, Balakrishnan R. Role of high-resolution computed tomography scan in the evaluation of cholesteatoma: A correlation of high-resolution computed tomography with intraoperative findings. Indian Journal of Otology. 2015;21(2):103.
33. Rogha M, Hashemi SM, Mokhtarinejad F, Eshaghian A, Dadgostar A. Comparison of Preoperative Temporal Bone CT with Intraoperative Findings in Patients with Cholesteatoma. Iranian journal of otorhinolaryngology. 2014;26(74):7-12.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.