

Review

Not peer-reviewed version

From CGRP to PACAP: Unraveling the Next Chapter in Migraine Treatment

[Masaru Tanaka](#) , [Ágnes Szabó](#) , Tamás Körtési , [Délia Szok](#) , [János Tajti](#) , [And László Vécsei](#) *

Posted Date: 7 September 2023

doi: [10.20944/preprints202309.0459.v1](https://doi.org/10.20944/preprints202309.0459.v1)

Keywords: migraine disorders; headache disorders; nociceptive pain; analgesics; calcitonin gene-related peptide; pituitary adenylate cyclase-activating polypeptide (PACAP); vasoactive intestinal peptide; adrenomedullin; neuropeptides; drug development



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

From CGRP to PACAP: Unraveling the Next Chapter in Migraine Treatment

Masaru Tanaka ¹, Ágnes Szabó ^{2,3} Tamás Körtési ⁴, Délia Szok ², János Tajti ² and László Vécsei ^{1,2,*}

¹ HUN-REN-SZTE Neuroscience Research Group, Hungarian Research Network, University of Szeged (HUN-REN-SZTE), Danube Neuroscience Research Laboratory, Tisza Lajos krt. 113, H-6725 Szeged, Hungary; tanaka.masaru.1@med.u-szeged.hu (M.T.); vecsei.laszlo@med.u-szeged.hu (L.V.)

² Department of Neurology, Albert Szent-Györgyi Medical School, University of Szeged, Semmelweis u. 6, H-6725 Szeged, Hungary; szabo.agnes.4@med.u-szeged.hu (Á.S.); szok.delia@med.u-szeged.hu (D.S.); tajti.janos@med.u-szeged.hu (J.T.)

³ Doctoral School of Clinical Medicine, University of Szeged, Korányi fasor 6, H-6720 Szeged, Hungary

⁴ Faculty of Health Sciences and Social Studies, University of Szeged, Temesvári krt. 31, H-6726 Szeged, Hungary; kortesi.tamas@med.u-szeged.hu


* Correspondence: vecsei.laszlo@med.u-szeged.hu (L.V.); Tel.: +36-62-545-351 (L.V.).

Abstract: Migraine is a neurovascular disorder that can be debilitating for individuals and society. Current research focuses on finding effective analgesics and management strategies for migraines by targeting specific receptors and neuropeptides. However, the efficacy of calcitonin gene-related peptide (CGRP) receptor antagonists and monoclonal antibodies (mAbs) in treating migraines can vary among patients, and tolerance may develop over time, reducing their effectiveness. To address the need for novel therapeutic targets, researchers are exploring the potential of another secretin family peptide, pituitary adenylate cyclase-activating polypeptide (PACAP), as a ground-breaking treatment avenue for migraine. Animal models have revealed how PACAP affects the trigeminal system, which is implicated in headache disorders. Clinical studies have demonstrated the significance of PACAP in migraine pathophysiology. Intriguingly, the pituitary adenylate cyclase-activating peptide 1 receptor mAb, AMG 301 showed no benefit for migraine prevention, while the PACAP ligand mAb, Lu AG09222 significantly reduced the number of monthly migraine days over placebo in a phase 2 clinical trial. In addition, another secretin family peptide vasoactive intestinal peptide is gaining interest as a potential new target. In light of recent advances in PACAP research, we emphasize the potential of PACAP as a promising target for migraine treatment. Recent findings highlight the importance of exploring PACAP as a member of the antimigraine armamentarium, especially for patients who do not respond to anti-CGRP therapies. By updating our knowledge on PACAP and its unique contribution to migraine pathophysiology, we can pave the way for reinforcing PACAP and other secretin peptides as a novel treatment approach for migraines.

Keywords: migraine disorders; headache disorders; nociceptive pain; analgesics; calcitonin gene-related peptide; pituitary adenylate cyclase-activating polypeptide (PACAP); vasoactive intestinal peptide; adrenomedullin; neuropeptides; drug development

1. Introduction

Migraine is a neurological condition that causes moderate to severe throbbing or pulsating pain on usually one side of the head, along with sensitivity to light, sound, smell, or touch, nausea, and/or vomiting [1–7]. The exact cause of migraines remains incompletely understood, but a combination of genetic, environmental, and lifestyle factors contribute to their development and occurrence within individuals [8]. Migraine triggers can exhibit significant variations among individuals and may encompass diverse factors such as stress, hormonal fluctuations, certain diets, and disturbances in sleep patterns [9]. Identifying and managing triggers can be crucial in preventing the onset of



migraine attacks and reducing their frequency, duration, and severity [10]. Migraines are complex neurological disorders that can significantly impact individuals' quality of life [11,12]. Comprehending the distinct stages, symptoms, triggers, and treatment options is fundamental for healthcare professionals and researchers, as it facilitates enhanced management and support for individuals affected by migraines [10].

Neuropeptides like calcitonin gene-related peptide (CGRP), pituitary adenylate cyclase-activating polypeptide (PACAP), vasoactive intestinal polypeptide (VIP), islet amyloid polypeptide (IAPP)/amylin, substance P, and adrenomedullin (ADM) have been linked to the cause of migraines [13–18]. Notably, the secretin family of peptides is a group of evolutionarily related peptide hormones that control the activity of G protein-coupled receptors (GPCR). This group includes CGRP, PACAP, ADM, and amylin, which share homology, receptor cross-reactivity, and similar biological actions that suggest they belong in this family (Table 1) [19]. These neuropeptides play various roles in migraine pathogenesis and contribute to our understanding of the underlying mechanisms of the disorder [20,21].




Figure 1. The amino acid sequence alignment analysis of the main secretin family peptides. Those amino acids with matching hues are identical amino acid sequences. The alignment similarity between peptides is displayed as a percentage next to the brackets.

CGRP and PACAP are two neuropeptides that have been extensively studied in the pathogenesis of migraine [22–24]. Both CGRP and PACAP are released in conjunction with migraine and cluster headache attacks and are potent vasodilators that can cause migraine-like attacks when infused into people. The activity of spinal trigeminal neurons is a sensitive measure of trigeminal activity [25,26]. In the spinal trigeminal nucleus caudalis (TNC), the central terminals of trigeminal afferents release CGRP. Experiments have shown that CGRP helps send pain signals, most likely through presynaptic action. Experiments show that when the trigeminovascular system (TS) is activated, the expression of both CGRP and PACAP goes up at the same time in the central part of the system. This affects the formation of mechanical hyperalgesia [27,28]. PACAP and CGRP share similar vasodilatory and nociceptive functions, but PACAP is likely to play a similar but distinct role as CGRP based on similarities and differences observed in both clinical and preclinical studies [29,30]. In rodent models, the PACAP pathway appears to be independent of the CGRP pathway, suggesting that CGRP and PACAP act in parallel ways that cause a migraine-like symptom [31]. In addition to CGRP and PACAP, other neuropeptides have been implicated in the pathogenesis of migraine (Table 1). In patients suffering from migraine, the concentration of CGRP in the peripheral blood is increased during migraine attacks compared with the interictal period [32]. A very similar observation has recently been made for PACAP as well, suggesting a potential biomarker function of this peptide in migraine [33]. The intravenous administration of the 38-amino acid form of PACAP (PACAP1–38) caused headaches and dilated blood vessels in both healthy participants and migraine sufferers, but only in migraine sufferers did it delay attacks that were similar to migraines [34–36]. Additionally, the infusion of 27-amino acid form of PACAP variant, PACAP1–27, triggers migraine attacks without aura [37].

VIP is a neuropeptide mainly found in the trigeminal nerve. VIP levels are altered in individuals experiencing migraines, with levels increasing during a migraine attack, leading to blood vessel dilation, which is considered an essential mechanism in the development of migraines [38]. VIP influences the release of other neurotransmitters implicated in migraines, including serotonin and CGRP [39,40]. VIP is also involved in regulating inflammatory responses and modulating the immune system [41]. VIP's immunomodulatory properties may play a role in reducing inflammation and suppressing the release of pain-inducing molecules. Furthermore, VIP has been implicated in regulating the sensitivity of neurons to pain signals, potentially influencing the intensity and frequency of migraine attacks [42].

A new class of drugs has been developed to treat and prevent migraines by targeting CGRP ligands and receptors. This class of drugs includes CGRP receptor antagonists such as atogepant, rimegepant, and ubrogepant, humanized CGRP ligand monoclonal antibodies (mAbs) such as eptinezumab, fremanezumab, and galcanezumab, and hu-man CGRP receptor mAb erenumab [43,44]. Triptans and lasmiditan, which are serotonergic pharmacons, are used for the treatment of acute attacks, while gepants are used for both treating acute attacks and their prevention, and CGRP receptor mAbs are used for prevention [45–47]. CGRP ligand-targeting mAbs are indicated for the prevention of both episodic and chronic migraines in adults.

CGRP receptor antagonists such as gepants and CGRP-targeting mAbs are promising treatments for migraines, but there are some downsides to consider [48,49]. These include low efficacy, tolerance rate, cost, side effects, and the need for further research to determine their long-term safety. While CGRP-targeting mAbs and CGRP receptor antagonists have shown efficacy in reducing the frequency of migraine attacks, they may not work for everyone and may not work as well as other preventive treatments. Some patients may develop a tolerance to these drugs over time, which can reduce their effectiveness [50,51]. Additionally, these drugs can be expensive, and insurance coverage may be limited [52]. While generally well tolerated, CGRP-targeting mAbs can cause side effects such as injection site reactions and constipation, while the gepants can cause nausea [53].

Humans typically experience migraines, but preclinical research using animal models leads clinical research by revealing the interaction of genetic and environmental factors as well as pathological alterations that contribute to neurological disorders and neuropsychiatric conditions, including migraine [54–65]. These models simulate disease conditions, allowing for the identification of pathogenic processes, the evaluation of symptoms and comorbidities, and the discovery of interventions, including pharmacotherapy [66–72]. The integration of preclinical and clinical research contributes to the creation of innovative therapeutics and personalized medicine [73–76]. This narrative review introduces the topic of migraine pathophysiology and the need for new therapeutic targets for migraine treatment, providing an overview of the current understanding of migraine pathophysiology and highlighting the potential of PACAP ligands and receptors as a novel target for migraine treatment. Emphasizing the importance of further research to fully elucidate the role of PACAP in migraine pathophysiology and to develop targeted therapies for individuals suffering from migraines, The review also discusses the data implicating the pituitary adenylate cyclase-activating peptide 1 (PAC1) receptor as a future drug target in migraine and examines several potential emerging therapeutic targets, including PACAP1-38, a specific form of PACAP. Furthermore, it explores the similarities between PACAP and VIP, the latter of which is involved in sleep regulation and circadian rhythm, suggesting that PACAP may be a key neuropeptide involved in migraines. Overall, we aim to provide a comprehensive overview of the current state of research on migraine pathophysiology and the potential of PACAP as a therapeutic target for migraine treatment.

2. Pituitary adenylate cyclase-activating peptide

PACAP is a multi-functional peptide that has therapeutic potential in a variety of pathophysiological conditions and represents a promising avenue for therapeutic intervention. PACAP is a neuropeptide that plays a crucial role in both neural and endocrine functions [76]. This peptide is widely distributed throughout the body and is involved in diverse physiological processes,

including circadian rhythm regulation, modulation of pain perception, immune system regulation, and stress response [77]. PACAP also has neuroprotective effects and has been shown to support nerve cell survival and regeneration in various neurological disorders [78]. GPCRs control the signaling pathways and cause the activation of adenylate cyclase (AC), the release of cyclic AMP, and the activation of protein kinase A (PKA) and calcium channels [79,80]. PACAP is a multi-functional peptide that has therapeutic potential in a variety of pathophysiological conditions and represents a promising avenue for therapeutic intervention [81].

2.1. Background

PACAP was found in ovine hypothalamic extracts in 1989. It is a 38-amino-acid peptide hormone that stimulates AC activity in the pituitary gland [82]. Subsequently, it was found to be widely distributed in the central and peripheral nervous systems, as well as in non-neuronal tissues, including the adrenal gland, pancreas, gut, and reproductive system [83]. PACAP exists in three biologically active forms: PACAP1–38, 6–38-amino acid form of PACAP (PACAP6-38), and PACAP1–27 [84]. PACAP-related peptide (PRP) is also a member of the PACAP family [85]. Radioimmunoassay demonstrated that PACAP1–38 levels were approximately 60 times greater than PACAP1–27 levels and 10 times greater than PRP levels [86].

Since its discovery, PACAP has been extensively studied for its potent neuroprotective effects against a diverse range of neurological disorders, including stroke, traumatic brain injury, Parkinson's disease, and Alzheimer's disease [87,88]. Recent findings suggest that PACAP may also play a key role in the regulation of immune cell function and cytokine production, highlighting its potential as a therapeutic target for immune-mediated diseases such as rheumatoid arthritis, multiple sclerosis, and asthma [89]. Furthermore, PACAP has been implicated in the regulation of energy metabolism, making it a promising therapeutic agent for the treatment of metabolic disorders such as obesity and diabetes [90]. Overall, the growing body of evidence on the multifunctional properties of PACAP highlights its potential as a novel therapeutic target for a wide range of diseases.

2.2. Receptor and Signaling Mechanisms of PACAP

PACAP plays an important role in a wide range of biological processes such as feeding behavior, stress response, neuroprotection, and regulation of neurotransmitter release. It activates three different GPCRs named PAC1, vasoactive intestinal peptide receptor (VPAC) 1, and VPAC2; these receptors are widely expressed in the central and peripheral nervous systems, endocrine systems, and immune systems [91]. The binding of PACAP to these receptors leads to the activation of multiple signaling mechanisms (Table 2) [92].

Table 2. The secretin family peptides, their receptors, and their binding affinity.

Peptides	Receptors
CGRP	CLR
PACAP1–38	>>PAC1, <VPAC1, <VPAC2
PACAP6-38	?
PACAP1–27	>PAC1, <VPAC1, <VPAC2
PRP	?
VIP	>VPAC1, >VPAC2, <PAC1

CGRP: calcitonin gene-related peptide; PACAP: pituitary adenylate cyclase-activating polypeptide; PRP: PACAP-related peptide; VIP: vasoactive intestinal peptide; CLR: calcitonin receptor-like receptor; PAC1: pituitary adenylate cyclase activating polypeptide type I; VPAC: vasoactive intestinal peptide receptor; ?: unknown.

Activation of the PAC1 receptor by PACAP leads to the activation of the adenylyl cyclase enzyme, which in turn leads to the production of cyclic adenosine monophosphate (cAMP) and the activation of PKA [93]. It also triggers the activation of phospholipase C, which leads to the

breakdown of phosphatidyl inositol 4,5-bisphosphate (PIP2) into inositol triphosphate and diacylglycerol (DAG), which activates protein kinase C (PKC) [94]. On the other hand, VPAC1 and VPAC2 receptor activation leads to AC enzyme activation, which leads to the generation of cAMP and the activation of PKA [95]. Also, PACAP signaling turns on calcium signaling, which causes intracellular calcium to be released and calcium/calmodulin-dependent kinase II to be activated [96]. PACAP signaling also activates the mitogen-activated protein kinase (MAPK), extracellular signal-regulated kinase (ERK), and jun N-terminal kinase signaling pathways [97]. These signaling mechanisms contribute to the diverse biological effects of PACAP on cellular functions. The regulation of PACAP gene expression is presented in Figure 1.




Figure 1. PACAP receptors signaling to ERK activation. AC, adenylate cyclase; ATP: adenosine monophosphate; cAMP: cyclic adenosine monophosphate; DAG: diacylglycerol; ERK, extracellular signal-regulated kinase; Gs and Gq: stimulatory G protein; MEK: mitogen-activated protein kinase kinase; PKA: protein kinase A; PKC: protein kinase C; PACAP: pituitary adenylate cyclase-activating polypeptide; PAC1: PACAP 1 receptor; PIP2: phosphatidylinositol bisphosphate; VPAC1: vasoactive intestinal peptide receptor type 1; VPAC2: vasoactive intestinal peptide receptor type 2.

PACAP and VIP are neuropeptides that interact specifically with three receptors (VPAC1, VPAC2, and PAC1) from the class II B GPCR family [98]. The similarities between PACAP and VIP in receptor and signaling mechanisms include: PACAP and VIP share nearly 70% amino acid sequence identity; PACAP binds with high affinity to all three receptors, while VIP binds with high affinity to VPAC1 and VPAC2 receptors and has a thousand-fold lower affinity for the PAC1 receptor compared to PACAP; both PACAP and VIP receptors are preferentially coupled to G_s, leading to activation of AC, subsequent cAMP production, and activation of PKA; PKA may in turn activate ERKs; PACAP and VIP receptor-mediated signaling pathways [99–102]. Due to the wide distribution of VIP and PACAP receptors in the body, potential therapeutic applications of drugs targeting these receptors, as well as expected unwanted side effects, are numerous [103]. Designing selective therapeutics targeting these receptors remains challenging due to their structural similarities.

2.3. Role of PACAP in migraine

PACAP has been strongly associated with the pathophysiology of migraine. PACAP is found in high levels in the trigeminal nerve, which is known to play a critical role in migraine. PACAP is known to increase the sensitivity of the trigeminal nerve, cause dilation of blood vessels in the brain, and trigger inflammation. All these biological effects have been implicated in the development of migraine attacks [104]. Several studies have been conducted to investigate the role of PACAP in migraine. One study showed that PACAP levels in the blood are significantly higher in migraine patients during an attack compared to headache-free controls [105]. This study suggests that PACAP could be used as a potential biomarker for migraine. Another study demonstrated that the venous infusion of PACAP into migraine patients resulted in the development of migraine-like attacks [106]. This finding strongly supports the hypothesis that PACAP plays a crucial role in the pathophysiology of migraine and suggests that blocking PACAP could be a potential therapeutic target for the treatment of migraine. The role of PACAP in migraine is well established, and there is strong evidence that this neuropeptide plays a crucial role in the development of migraine attacks. Further research is needed to better understand the mechanism of action of PACAP and to develop new pharmacological agents that target PACAP for the treatment of migraine.

Both CGRP and PACAP are multifunctional peptides with many roles in the nervous, cardiovascular, respiratory, gastrointestinal, and reproductive systems. They play a role in vasodilation, neurogenic inflammation, and nociception. While CGRP plays an integral role in migraine, PACAP is likely to play a similar but distinct role as CGRP based on similarities and differences observed in both clinical and preclinical studies [107]. In rodent models, the PACAP pathway appears to be independent of the CGRP pathway, suggesting that CGRP and PACAP act in parallel ways that cause a migraine-like symptom [108]. In migraine without aura, the first double-blinded placebo-controlled study reported that 33% of the patients developed delayed migraine attacks after CGRP administration [109]. The studies have identified the involvement of two endogenous neuropeptides, CGRP and PACAP, in the pathogenesis of migraine [110].

VIP has also been implicated in the pathophysiology of migraine [111]. The similarities between PACAP and VIP in their roles in migraine include: PACAP and VIP are released in conjunction with migraine and cluster headache attacks [112]; PACAP and VIP are potent vasodilators and can cause migraine-like attacks when infused into people [113]; A 2-hour infusion of VIP caused migraine attacks, indicating that VIP plays a significant role in the pathophysiology of migraines and intravenous administration of PACAP-38 caused headaches in all healthy subjects and migraine-like attacks in 58% of patients with a history of migraine without aura [15,35]; PACAP and VIP receptors are preferentially coupled to G_{αs}, leading to activation of AC, subsequent cAMP production, and activation of PKA [114]; PKA may in turn activate ERKs [115]; PACAP and VIP receptor-mediated signaling pathways are shown to share activities, including vasodilation, neurogenic inflammation, and nociception in rodents [116]; PACAP and VIP receptors provide a rich set of targets to complement and augment the current CGRP-based migraine therapeutics; VPAC1 receptors play a dominant role in PACAP-induced vasorelaxation in female mice [117]. Also, PG 99-465, a selective VPAC2 receptor antagonist that has been used in a number of physiological studies, has been shown to have a significant activity at VPAC1 and PAC1 receptors [118].

2.4. Preclinical Studies

In addition to in vitro systems, a variety of organisms are used in experimental medicine [119–121]. Understanding the effects of endogenous neuropeptides, neurohormones, and metabolites has advanced significantly thanks to the information gathered using laboratory animals [122–127]. Animal models are a crucial tool for bridging the knowledge gap between data- and hypothesis-driven benchwork and its application to clinical bedside management. PACAP has been extensively studied as a neuromodulator in the trigeminal nociceptive pathway [128]. Preclinical studies have shown that PACAP is involved in the transmission of pain signals from the periphery to the central nervous system and is therefore a potential target for the treatment of migraine and other headache disorders [129,130].

In animal models, PACAP has been shown to play a role in trigeminal sensitization, which is the process by which nociceptive signals become amplified and persistent, leading to chronic pain [131]. Studies have also found that PACAP is involved in the activation of inflammatory pathways in the trigeminal nerve, further contributing to pain and inflammation [132]. In addition, PACAP has been implicated in the regulation of blood flow to the brain, which may also play a role in headache pathophysiology [133]. In an experimental model of migraine, intraperitoneal administration of nitroglycerol caused marked photophobia, meningeal vasodilatation, and increased the number of c-fos-positive activated neurons in the TNC in wild-type mice but not in PACAP1-38-deficient mice [134]. In line with this, an increased concentration of PACAP1-38 was detected in the TNC after the activation of the TS in different animal models [135,136].

PAC1 receptor antagonists includes PACAP6-38, N-stearyl-[Nle17] neuropeptides-(6-11)/VIP-(7-28), deletion mutants of maxadilan (M65 and Max.d.4), MK-0893, PG 97-269, and synthesized small molecule acyl hydrazides [137]. PACAP6-38 has been used as a PAC1 receptor antagonist in many studies, but it has affinity for VPAC2 receptors. M65 has not been extensively used due to problems of availability, and others have not yet been extensively characterized [138,139]. Thus, preclinical studies suggest that concentrating on the PACAP signaling pathways in the trigeminal nociceptive system could be an effective strategy for discovering novel treatments for headache disorders. However, more research is needed to fully understand the mechanisms underlying PACAP's role in headache pathophysiology and to develop effective and safe PACAP-targeted therapies.

VIP plays a key role in sensory processing and the modulation of pain pathways in the trigeminal system. In preclinical studies, VIP has been shown to change the activity of nociceptive neurons in the trigeminal ganglion and make the TNC more sensitive, which can cause chronic pain or migraines [140]. Trigeminal sensory neurons release VIP in response to noxious stimuli, which can activate VIP receptors on nearby neurons and cause the release of a number of signaling molecules involved in pain amplification [141]. VIP-mediated sensitization of trigeminal neurons can lead to hyperexcitability and increased responsiveness to noxious stimuli, which may contribute to the development and maintenance of chronic pain or migraine [142]. Targeting VIP signaling pathways may therefore represent a promising approach for the development of novel therapies for chronic pain or migraine.

2.5. Clinical studies

Clinical studies have shown that patients with migraines exhibit higher levels of PACAP compared to control groups [143]. PACAP is a neuropeptide that is known to play a role in the activation of nociceptive pathways, contributing to the development of migraines. The high levels of PACAP in migraineurs have been associated with increased headache severity and frequency, and this has led to the exploration of PACAP as a therapeutic target for migraine treatment [144]. In migraineurs without aura, the development of PACAP1-38-evoked migraine-like attacks was independent of the severity of family load [35,145]. In the same study, 90 minutes after the injection, the levels of numerous migraine-related molecular markers were increased in the plasma of patients [146]. Magnetic resonance imaging angiography examinations revealed that PACAP1-38-induced headache was associated with prolonged vasodilatation of the middle meningeal artery (MMA) but not the middle cerebral artery (MCA). Sumatriptan, an antimigraine medication, was able to alleviate the headache, which mirrored the contraction of the MMA but not the MCA, indicating that PACAP1-38-induced headaches may originate from extracerebral arteries [147].

Other clinical studies show that PACAP plays a significant role in migraine pathophysiology, and targeting PACAP signaling may be a potential therapeutic strategy for migraine treatment. In terms of safety, PACAP has been generally well tolerated in clinical trials [148]. One study found that PACAP induces headache via sustained vasodilation and that targeting the PACAP pathway may be a promising approach for the treatment of migraine [149]. AMG 301, a mAb that targets the PAC1 receptor, was administered to patients with episodic or chronic migraines in a randomized, double-blind, placebo-controlled phase 2 study. There was no significant difference between the AMG 301 group and the placebo group, suggesting that AMG 301 was ineffective for migraine prevention

[150,151]. On the other hand, the PACAP ligand mAb, Lu AG09222, was shown to reduce the number of monthly migraine days from baseline to weeks 1–4 of treatment statistically significantly more than placebo [152,153]. Additionally, the mAb targeting the PAC1 receptor, LY3451838, is currently undergoing phase 2 clinical trials for adults with treatment-resistant migraine. This trial is in progress, and the results are not yet available [154]. Overall, the efficacy and safety of PACAP as a migraine treatment in clinical studies suggest that it is a promising option for patients with this debilitating condition. Further research is needed to fully understand the potential of PACAP as a treatment for migraine, but the current evidence is encouraging.

Table 3. Calcitonin gene-related peptide (CGRP) monoclonal antibodies under clinical trials.

ClinicalTrials.gov Identifier	Monoclonal antibody	Target	Status	Ref.
NCT03238781	AMG 301	receptor prevention	No benefit over placebo for migraine prevention	[150,151]
NCT05133323	Lu AG09222	ligand	No results posted	[152,153]
NCT04498910	LY3451838	receptor	No results posted	[154]

VIP infusion has been studied in the context of migraine, with a particular focus on its potential to provoke migraine attacks and its role in migraine pathophysiology. A phase 2 clinical trial investigated the effects of a long-lasting infusion of VIP on headache, cranial hemodynamics, and autonomic symptoms in episodic migraine patients without aura [155]. The study found that a two-hour infusion of VIP promoted long-lasting cranial vasodilation and delayed headaches in healthy volunteers, resembling the effect of migraine prophylaxis. However, other studies have suggested that VIP infusions may actually provoke migraine attacks. For example, a randomized clinical trial found that a two-hour infusion of VIP caused migraine attacks, suggesting an important role for VIP in migraine pathophysiology [15]. It remains unclear whether the lack of migraine induction can be attributed to the only transient vasodilatory response after a 20-minute infusion of VIP. Overall, the search results suggest that VIP infusion may have a role in migraine pathophysiology, but further research is needed to fully understand its effects and potential therapeutic applications.

3. Discussion

This review paper aims to provide insights into the roles of PACAP in migraine by comparing its actions with those of VIP. By analyzing existing studies, this paper hopes to shed light on the pathophysiology of migraine and pave the way towards more effective treatments. The ultimate goal of this review is to explore the potential of developing antimigraine drugs that target the PACAP pathways. Identifying and producing new ways to target the PACAP system may provide an alternative therapeutic option for migraine sufferers. The authors aim to consolidate the current evidence on the PACAP system's role in migraines and evaluate potential drug targets within the pathway, hoping to pave the way for more extensive research to develop new and effective antimigraine drugs that target the PACAP pathways.

The PACAP system presents a significant challenge when it comes to targeted therapies due to its pleiotropic roles in the body, both physiologically and pathologically. PACAP plays crucial roles in various aspects of the body, such as neural development, pain regulation, immune functions, and stress responses. These diverse roles make the PACAP system difficult to target effectively without affecting other physiological functions. Furthermore, PACAP signaling is often dysregulated in pathological conditions such as inflammatory disorders, neurodegenerative diseases, and cancers. Conversely, PACAP has been shown to have protective effects in certain diseases, such as ischemic stroke and Alzheimer's disease. Thus, finding a balance between targeting the PACAP system to treat diseases while preserving its physiological functions remains a significant challenge in the field of medicine.

The PACAP system has emerged as a potential target for the treatment of migraine, especially after the discovery of the role of CGRP and its receptors in the pathophysiology of migraine. PACAP,

a neuropeptide similar to CGRP, is highly expressed in the TS, which is the neural network that is responsible for migraine pain. PACAP receptors have been found to be co-localized with CGRP receptors in the TS, suggesting that the two systems could be acting in a synergistic manner to induce migraine pain. Therefore, targeting the PACAP system could provide an additional therapeutic approach for the treatment of migraine, and several drugs that inhibit PACAP or its receptors are currently under development.

The present review holds notable significance in shedding light on the critical role of PACAP in comparison with other neuropeptides like CGRP and VIP, which have been extensively studied as potential therapeutic targets for various neurological disorders. By thoroughly analyzing the preclinical studies, the review highlights the promising findings that suggest the potential translation of PACAP's therapeutic benefits from laboratory settings to clinical practice. The authors' critical evaluation and systematic compilation of the latest research on PACAP is bound to have a relevant impact on the scientific community and serve as a foundation for further clinical research. Ultimately, the knowledge and insights gained from this review will be instrumental in developing advanced treatments for a range of debilitating neurological conditions.

The review also highlights limitations and challenges in PACAP research, such as the complexity of its signaling mechanism, variations in its effects on different cell types, and the limited availability of specific antibodies against PACAP and its receptors. The high cost of producing PACAP analogs and the lack of standardized protocols for their synthesis and purification are also limitations. These challenges and limitations make it difficult to fully understand the mechanisms of PACAP action and to develop effective therapeutic interventions.

The development of PACAP-based therapeutics for migraines will focus on two main approaches: targeting PACAP ligands and receptors. Studies using animal models of migraines have demonstrated that blocking the PACAP receptor reduces migraine symptoms, while inhibiting PACAP signaling reduces pain sensitivity. Currently, clinical trials are underway to assess the safety and effectiveness of various PACAP-based drugs for migraines in humans. PACAP-based therapies may offer an alternative to current treatments by targeting the underlying mechanisms of the disorder and reducing the risk of side effects. In addition, the role of additional secretin family peptides, ADM, and amylin in the pathogenesis of migraine remains to be investigated. Further research in this area could lead to the development of better treatments for migraines.

The future direction of migraine research holds great promise for advancing our understanding of this complex neurological disorder. The combination of preclinical and clinical data, along with computational tools, has provided invaluable insights into various aspects of diseases, including neurological and psychiatric disorders [156–173]. The use of preclinical models and clinical studies has shed light on the underlying mechanisms of migraine. These studies have contributed to the identification of structural and functional changes in the brain that occur in neurological and psychiatric disorders, such as migraine attacks [174–180]. Understanding these changes is crucial for developing targeted treatments and improving diagnosis.

Migraine is not just a pain disorder, but it is also interrelated to emotional and cognitive domains [181]. Migraine is commonly linked with a broad range of psychiatric comorbidities, especially among subjects with migraine with aura or chronic migraine [182]. The comorbidity between neurological and psychiatric disorders likely suggests multiple causes, such as unidirectional causal explanations or shared environmental and/or genetic risk factors, communication with other parts of the body, and their interaction on multiple levels [183–197]. Emotional distress is commonly recognized as a migraine trigger, and being affected by psychiatric disorders is considered an independent modifiable factor of progression toward chronicification of migraine and a tendency to medication overuse [198]. Therefore, revealing the mechanisms of comorbidity between migraine and psychiatric disorders may lead to a clue to migraine prevention and management. Many biological and neural aspects related to the comorbidity still need to be clearly elucidated to better approach the real nature of the association between migraine and psychiatric disorders.

The integration of computational tools in migraine research has allowed for the testing and evaluation of potential treatments. These tools enable researchers to simulate the effects of different

interventions, including brain stimulation and assess their therapeutic efficacy [199–203]. This approach holds promise for the development of novel and more effective migraine treatments. Advanced imaging techniques have played a crucial role in migraine research. Neuroimaging studies have revealed structural and functional brain changes associated with migraine [204–211]. These imaging techniques provide valuable insights into the pathophysiology of the disorder and can help identify unique clinical cases. The use of human brain organoids in migraine research is an emerging area of study [209]. Brain organoids are three-dimensional models that mimic the structure and function of the human brain [213]. They can be used to investigate altered neuronal pathways, protein expression, and metabolic pathways associated with migraines [214–215]. This approach offers a unique opportunity to study the disease in a more physiologically relevant system.

4. Conclusion

PACAP is a neuropeptide that has been linked to the pathophysiology of primary headaches such as migraine. The release of PACAP is associated with migraine and cluster headache attacks, and it has been shown to be a potent vasodilator that dilates cranial arteries and causes migraines when infused into patients. Like CGRP, PACAP is located near sensory nerve fibers and has nociceptive functions. Both peptides are promising targets for migraine therapeutics, and growing evidence supports the involvement of the PACAP-related mechanisms in migraine. While CGRP and PACAP share similar functions, the PACAP pathway appears to be independent of the CGRP pathway, suggesting that they act in parallel ways to cause a migraine-like symptom. Therefore, understanding the role of PACAP in migraine pathogenesis could lead to new treatment options for this debilitating condition.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “Conceptualization, M.T. and L.V.; methodology, N/A; software, N/A; validation, N/A; formal analysis, N/A.; investigation, N/A; resources, N/A; data curation, N/A; writing—original draft preparation, M.T.; writing—review and editing, M.T., Á.S., T.K., D.S. J.T., and L.V.; visualization, Á.S. and T.K.; supervision, J.T. and L.V.; project administration, L.V.; funding acquisition, L.V. All authors have read and agreed to the published version of the manuscript.”

Funding: This work was supported by OTKA-138125-K, TUDFO/47138-1/2019-ITM and HUN-REN-SZTE Neuroscience Research Group, Hungarian Research Network, University of Szeged (HUN-REN-SZTE).

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

AC	adenylate cyclase
ADM	adrenomedullin
cAMP	cyclic adenosine monophosphate
CGRP	calcitonin gene-related peptide
DAG	diacylglycerol
ERK	extracellular signal-regulated kinase
GPCR	G protein-coupled receptors
IAPP	islet amyloid polypeptide/amylin
mAbs	monoclonal antibodies
MAPK	mitogen-activated protein kinase
MCA	middle cerebral artery
MEK:	mitogen-activated protein kinase kinase
PAC1	pituitary adenylate cyclase-activating Peptide 1
PACAP	pituitary adenylate cyclase-activating polypeptide
PACAP1–38	38-amino acid form of PACAP
PACAP1–27	27-amino acid form of PACAP

PACAP6-38	6-38-amino acid form of PACAP
PIP2	phosphatidyl inositol 4,5-bisphosphate
PKA	protein kinase A
PKC	activates protein kinase C
PRP	PACAP-related peptide
TNC	trigeminal nucleus caudalis
TS	trigeminovascular system
VPAC	vasoactive intestinal peptide receptor
VIP	vasoactive intestinal polypeptide

References

1. Pescador Ruschel, M.A.; De Jesus, O. Migraine Headache. In: StatPearl; StatPearls Publishing; Treasure Island, FL, USA, 2023. Available online from: <https://www.ncbi.nlm.nih.gov/books/NBK560787/> (accessed on 28 August 2023).
2. Amiri, P.; Kazeminasab, S.; Nejadghaderi, S.A.; Mohammadinasab, R.; Pourfathi, H.; Araj-Khodaei, M.; Sullman, M.J.M.; Kolahi, A.A.; Safiri, S. Migraine: A Review on Its History, Global Epidemiology, Risk Factors, and Comorbidities. *Front. Neurol.* **2022**, *12*, 800605. <https://doi.org/10.3389/fneur.2021.800605>
3. Ferrari, M.D., Goadsby, P.J.; Burstein, R.; Kurth, T.; Ayata, C.; Charles, A.; Ashina, M.; van den Maagdenberg, A.M.J.M.; Dodick, D.W. Migraine. *Nat Rev Dis Primers.* **2022**, *8*(1), 2. <https://doi.org/10.1038/s41572-021-00328-4>
4. Ashina, M. Migraine. *N. Engl. J. Med.* **2020**, *383*(19), 1866-1876. <https://doi.org/10.1056/NEJMra1915327>
5. Burstein, R.; Noseda, R.; Borsook, D. Migraine: multiple processes, complex pathophysiology. *J. Neurosci.* **2015**, *35*(17), 6619-6629. <https://doi.org/10.1523/JNEUROSCI.0373-15.2015>
6. Dodick, D.W. A Phase-by-Phase Review of Migraine Pathophysiology. *Headache* **2018**, *58*(Suppl 1), 4-16. <https://doi.org/10.1111/head.13300>
7. Schankin, C.J.; Viana, M.; Goadsby, P.J. Persistent and Repetitive Visual Disturbances in Migraine: A Review. *Headache* **2017**, *57*(1), 1-16. <https://doi.org/10.1111/head.12946>
8. Gasparini, C.F.; Sutherland, H.G.; Griffiths, L.R. Studies on the pathophysiology and genetic basis of migraine. *Curr. Genomics* **2013**, *14*(5), 300-315. <https://doi.org/10.2174/13892029113149990007>
9. Khan, J.; Asoom, L.I.A.; Sunni, A.A.; Rafique, N.; Latif, R.; Saif, S.A.; Almandil, N.B.; Almohazey, D.; AbdulAzeez, S.; Borgio, J.F. Genetics, pathophysiology, diagnosis, treatment, management, and prevention of migraine. *Biomed. Pharmacother.* **2021**, *139*, 111557. <https://doi.org/10.1016/j.biopha.2021.111557>
10. Buse, D.C.; Rupnow, M.F.; Lipton, R.B. Assessing and managing all aspects of migraine: migraine attacks, migraine-related functional impairment, common comorbidities, and quality of life. *Mayo Clin. Proc.* **2009**, *84*(5), 422-435. [https://doi.org/10.1016/S0025-6196\(11\)60561-2](https://doi.org/10.1016/S0025-6196(11)60561-2).
11. Gupta, J.; Gaurkar, S.S. Migraine: An Underestimated Neurological Condition Affecting Billions. *Cureus* **2022**, *14*(8), e28347. <https://doi.org/10.7759/cureus.28347>
12. Villar-Martinez, M.D.; Goadsby, P.J. Pathophysiology and Therapy of Associated Features of Migraine. *Cells* **2022**, *11*, 2767. <https://doi.org/10.3390/cells11172767>
13. Durham, P.L. Calcitonin gene-related peptide (CGRP) and migraine. *Headache* **2006**, *46* Suppl 1(Suppl 1), S3-8. <https://doi.org/10.1111/j.1526-4610.2006.00483.x>
14. Waschek, J.A.; Baca, S.M.; Akerman, S. PACAP and migraine headache: immunomodulation of neural circuits in autonomic ganglia and brain parenchyma. *J. Headache Pain* **2018**, *19*(1), 23. <https://doi.org/10.1186/s10194-018-0850-6>
15. Pellesi, L.; Al-Karagholi, M.A.; De Icco, R.; Coskun H, Elbahi FA, Lopez-Lopez, C.; Snellman, J.; Hannibal, J.; Amin, F.M.; Ashina, M. Effect of Vasoactive Intestinal Polypeptide on Development of Migraine Headaches: A Randomized Clinical Trial. *JAMA Netw. Open* **2021**, *4*(8), e2118543. <https://doi.org/10.1001/jamanetworkopen.2021.18543>
16. Ghanizada, H.; Al-Karagholi, M.A.; Walker, C.S.; Arngrim, N.; Rees, T.; Petersen, J.; Siow, A.; Mørch-Rasmussen, M.; Tan, S.; O'Carroll, S.J.; et al. Amylin Analog Pramlintide Induces Migraine-like Attacks in Patients. *Ann. Neurol.* **2021**, *89*(6), 1157-1171. <https://doi.org/10.1002/ana.26072>
17. May, A.; Goadsby, P.J. Substance P receptor antagonists in the therapy of migraine. *Expert Opin. Investig. Drugs* **2001**, *10*(4), 673-678. <https://doi.org/10.1517/13543784.10.4.673>
18. Petersen, K.A.; Birk, S.; Kitamura, K.; Olesen, J. Effect of adrenomedullin on the cerebral circulation: relevance to primary headache disorders. *Cephalgia* **2009**, *29*(1), 23-30. <https://doi.org/10.1111/j.1468-2982.2008.01695.x>

19. Hay, D.L.; Garelja, M.L.; Poyner, D.R.; Walker, C.S. Update on the pharmacology of calcitonin/CGRP family of peptides: IUPHAR Review 25. *Br. J. Pharmacol.* **2018**, *175*(1), 3-17. <https://doi.org/10.1111/bph.14075>
20. Poyner, D.R.; Hay, D.L. Secretin family (Class B) G protein-coupled receptors - from molecular to clinical perspectives. *Br. J. Pharmacol.* **2012**, *166*(1), 1-3. <https://doi.org/10.1111/j.1476-5381.2011.01810.x>
21. Edvinsson, L.; Grell, A.S.; Warfvinge, K. Expression of the CGRP Family of Neuropeptides and their Receptors in the Trigeminal Ganglion. *J. Mol. Neurosci.* **2020**, *70*(6), 930-944. <https://doi.org/10.1007/s12031-020-01493-z>
22. Dux, M.; Vogler, B.; Kuhn, A.; Mackenzie, K.D.; Stratton, J.; Messlinger, K. The Anti-CGRP Antibody Fremanezumab Lowers CGRP Release from Rat Dura Mater and Meningeal Blood Flow. *Cells* **2022**, *11*, 1768. <https://doi.org/10.3390/cells1111768>
23. Pavelic, A.R.; Wöber, C.; Riederer, F.; Zebenholzer, K. Monoclonal Antibodies against Calcitonin Gene-Related Peptide for Migraine Prophylaxis: A Systematic Review of Real-World Data. *Cells* **2023**, *12*, 143. <https://doi.org/10.3390/cells12010143>
24. Körtési, T.; Spekker, E.; Vécsei, L. Exploring the Tryptophan Metabolic Pathways in Migraine-Related Mechanisms. *Cells* **2022**, *11*, 3795. <https://doi.org/10.3390/cells11233795>
25. Iyengar, S.; Johnson, K.W.; Ossipov, M.H.; Aurora, S.K. CGRP and the Trigeminal System in Migraine. *Headache* **2019**, *59*(5), 659-681. <https://doi.org/10.1111/head.13529>
26. Edvinsson, L.; Tajti, J.; Szalárdy, L.; Vécsei, L. PACAP and its role in primary headaches. *J. Headache Pain* **2018**, *19*(1), 21. <https://doi.org/10.1186/s10194-018-0852-4>
27. Körtési, T.; Tuka, B.; Nyári, A.; Vécsei, L.; Tajti, J. The effect of orofacial complete Freund's adjuvant treatment on the expression of migraine-related molecules. *J. Headache Pain* **2019**, *20*(1), 43. <https://doi.org/10.1186/s10194-019-0999-7>
28. Russo, A.F. Calcitonin gene-related peptide (CGRP): a new target for migraine. *Annu. Rev. Pharmacol. Toxicol.* **2015**, *55*, 533-552. <https://doi.org/10.1146/annurev-pharmtox-010814-124701>
29. Kaiser, E.A.; Russo, A.F. CGRP and migraine: could PACAP play a role too? *Neuropeptides* **2013**, *47*(6), 451-461. <https://doi.org/10.1016/j.npep.2013.10.010>
30. Kubaras, A.; Mason, B.N.; Hing, B.; Wattiez, A.S.; Reis, A.S.; Sowers, L.P.; Moldovan Loomis, C.; Garcia-Martinez, L.F.; Russo, A.F. PACAP Induces Light Aversion in Mice by an Inheritable Mechanism Independent of CGRP. *J. Neurosci.* **2021**, *26*, 41(21), 4697-4715. <https://doi.org/10.1523/JNEUROSCI.2200-20.2021>
31. Russo, A.F. CGRP as a neuropeptide in migraine: lessons from mice. *Br. J. Clin Pharmacol.* **2015**, *80*(3), 403-414. <https://doi.org/10.1111/bcp.12686>
32. Goadsby, P.J.; Edvinsson, L.; Ekman, R. Release of vasoactive peptides in the extracerebral circulation of humans and the cat during activation of the trigeminovascular system. *Ann. Neurol.* **1988**, *23*(2), 193-196. <https://doi.org/10.1002/ana.410230214>
33. Tuka, B.; Helyes, Z.; Markovics, A.; Bagoly, T.; Szolcsányi, J.; Szabó, N.; Tóth, E.; Kincses, Z.T.; Vécsei, L.; Tajti, J. Alterations in PACAP-38-like immunoreactivity in the plasma during ictal and interictal periods of migraine patients. *Cephalgia* **2013**, *33*(13), 1085-1095. <https://doi.org/10.1177/0333102413483931>
34. Lassen, L.H.; Haderslev, P.A.; Jacobsen, V.B.; Iversen, H.K.; Sperling, B.; Olesen, J. CGRP may play a causative role in migraine. *Cephalgia* **2002**, *22*(1), 54-61. <https://doi.org/10.1046/j.1468-2982.2002.00310.x>
35. Schytz, H.W.; Birk, S.; Wiencke, T.; Kruuse, C.; Olesen, J.; Ashina, M. PACAP38 induces migraine-like attacks in patients with migraine without aura. *Brain* **2009**, *132*(Pt 1), 16-25. <https://doi.org/10.1093/brain/awn307>
36. Vollesen, A.L.H.; Amin, F.M.; Ashina, M. Targeted Pituitary Adenylate Cyclase-Activating Peptide Therapies for Migraine. *Neurotherapeutics* **2018**, *15*(2), 371-376. <https://doi.org/10.1007/s13311-017-0596-x>
37. Ghanizada, H.; Al-Karagholi, M.A.; Arngrim, N.; Olesen, J.; Ashina, M. PACAP27 induces migraine-like attacks in migraine patients. *Cephalgia* **2020**, *40*(1), 57-67. doi: 10.1177/0333102419864507
38. Cernuda-Morollón, E.; Martínez-Camblor, P.; Alvarez, R.; Larrosa, D.; Ramón, C.; Pascual, J. Increased VIP levels in peripheral blood outside migraine attacks as a potential biomarker of cranial parasympathetic activation in chronic migraine. *Cephalgia* **2015**, *35*(4), 310-316. <https://doi.org/10.1177/0333102414535111>
39. Hery, M.; Faudon, M.; Hery, F. Effect of vasoactive intestinal peptide on serotonin release in the suprachiasmatic area of the rat. Modulation by oestradiol. *Peptides* **1984**, *5*(2), 313-317. [https://doi.org/10.1016/0196-9781\(84\)90226-2](https://doi.org/10.1016/0196-9781(84)90226-2)
40. Pellesi, L.; Al-Karagholi, M.A.; De Icco, R.; Chaudhry, B.A.; Lopez, C.L.; Snellman, J.; Hannibal, J.; Amin, F.M.; Ashina, M. Plasma Levels of CGRP During a 2-h Infusion of VIP in Healthy Volunteers and Patients With Migraine: An Exploratory Study. *Front. Neurol.* **2022**, *13*, 871176. <https://doi.org/10.3389/fneur.2022.871176>
41. De la Fuente, M.; Delgado, M.; Gomariz, R.P. VIP modulation of immune cell functions. *Adv. Neuroimmunol.* **1996**, *6*(1), 75-91. [https://doi.org/10.1016/s0960-5428\(96\)00002-2](https://doi.org/10.1016/s0960-5428(96)00002-2)
42. Hoffmann, J.; Baca, S.M.; Akerman, S. Neurovascular mechanisms of migraine and cluster headache. *J. Cereb. Blood. Flow Metab.* **2019**, *39*(4), 573-594. <https://doi.org/10.1177/0271678X17733655>

43. Ocheretyaner, E.R.; Kofman, M.; Quattrocchi, E. Calcitonin gene-related peptide (CGRP) receptor antagonists for the acute treatment of migraines in adults. *Drugs Context.* **2022**, *11*, 2022-3-5. <https://doi.org/10.7573/dic.2022-3-5>
44. Wang, X.; Chen, Y.; Song, J.; You, C. Efficacy and Safety of Monoclonal Antibody Against Calcitonin Gene-Related Peptide or Its Receptor for Migraine: A Systematic Review and Network Meta-analysis. *Front. Pharmacol.* **2021**, *12*, 649143. <https://doi.org/10.3389/fphar.2021.649143>
45. Berger, A.A.; Winnick, A.; Popovsky, D.; Kaneb, A.; Berardino, K.; Kaye, A.M.; Cornett, E.M.; Kaye, A.D.; Viswanath, O. Urits I. Lasmiditan for the Treatment of Migraines With or Without Aura in Adults. *Psychopharmacol. Bull.* **2020**, *50*(4 Suppl 1), 163-188. PMID: 33633424; PMCID: PMC7901123.
46. Rissardo, J.P.; Caprara, A.L.F. Gepants for Acute and Preventive Migraine Treatment: A Narrative Review. *Brain Sci.* **2022**, *12*(12), 1612. <https://doi.org/10.3390/brainsci12121612>
47. Ibekwe, A.; Perras, C.; Mierzwiński-Urban, M. Monoclonal Antibodies to Prevent Migraine Headaches. In: CADTH Issues in Emerging Health Technologies. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, Ottawa, Canada, 2016. Available online from: <https://www.ncbi.nlm.nih.gov/books/NBK538376/> (accessed on 28 August 2023).
48. Nie, L.; Sun, K.; Gong, Z.; Li, H.; Quinn, J.P.; Wang, M. Src Family Kinases Facilitate the Crosstalk between CGRP and Cytokines in Sensitizing Trigeminal Ganglion via Transmitting CGRP Receptor/PKA Pathway. *Cells* **2022**, *11*, 3498. <https://doi.org/10.3390/cells11213498>
49. Greco, R.; Demartini, C.; Francavilla, M.; Zanaboni, A.M.; Tassorelli, C. Antagonism of CGRP Receptor: Central and Peripheral Mechanisms and Mediators in an Animal Model of Chronic Migraine. *Cells* **2022**, *11*, 3092. <https://doi.org/10.3390/cells11193092>
50. Mavridis, T.; Deligianni, C.I.; Karagiorgis, G.; Daponte, A.; Breza, M.; Mitsikostas, D.D. Monoclonal Antibodies Targeting CGRP: From Clinical Studies to Real-World Evidence—What Do We Know So Far? *Pharmaceuticals* **2021**, *14*, 700. <https://doi.org/10.3390/ph14070700>
51. Pellesi, L.; Guerzoni, S.; Pini, L.A. Spotlight on Anti-CGRP Monoclonal Antibodies in Migraine: The Clinical Evidence to Date. *Clin. Pharmacol. Drug Dev.* **2017**, *6*(6), 534-547. <https://doi.org/10.1002/cpdd.345>
52. Nguyen, J.L.; Munshi, K.; Peasah, S.K.; Swart, E.C.S.; Kohli, M.; Henderson, R.; Good, C.B. Trends in utilization and costs of migraine medications, 2017-2020. *J. Headache Pain* **2022**, *23*(1), 111. <https://doi.org/10.1186/s10194-022-01476-y>
53. Haanes, K.A.; Edvinsson, L.; Sams, A. Understanding side-effects of anti-CGRP and anti-CGRP receptor antibodies. *J. Headache Pain* **2020**, *21*, 26. <https://doi.org/10.1186/s10194-020-01097-3>
54. Telegdy, G.; Adamik, A.; Tanaka, M.; Schally, A.V. Effects of the LHRH antagonist Cetrorelix on affective and cognitive functions in rats. *Regul. Pept.* **2010**, *159*(1-3), 142-147. <https://doi.org/10.1016/j.regpep.2009.08.005>
55. Tanaka, M.; Schally, A.V.; Telegdy, G. Neurotransmission of the antidepressant-like effects of the growth hormone-releasing hormone antagonist MZ-4-71. *Behav. Brain Res.* **2012**, *228*(2), 388-91. <https://doi.org/10.1016/j.bbr.2011.12.022>
56. Tanaka, M.; Telegdy, G. Neurotransmissions of antidepressant-like effects of neuromedin U-23 in mice. *Behav. Brain Res.* **2014**, *259*, 196-199. <https://doi.org/10.1016/j.bbr.2013.11.005>
57. Tanaka, M.; Csabai, K.; Telegdy, G. Neurotransmissions of antidepressant-like effects of kisspeptin-13. *Regul. Pept.* **2013**, *180*, 1-4. <https://doi.org/10.1016/j.regpep.2012.08.017>
58. Telegdy, G.; Tanaka, M.; Schally, A.V. Effects of the growth hormone-releasing hormone (GH-RH) antagonist on brain functions in mice. *Behav. Brain Res.* **2011**, *224*(1), 155-158. <https://doi.org/10.1016/j.bbr.2011.05.036>
59. Rákosi, K.; Masaru, T.; Zarándia, M.; Telegdy, G.; Tóth, G.K. Short analogs and mimetics of human urocortin 3 display antidepressant effects in vivo. *Peptides* **2014**, *62*, 59-66. <https://doi.org/10.1016/j.peptides.2014.09.023>
60. Tran, K.N.; Nguyen, N.P.K.; Nguyen, L.T.H.; Shin, H.-M.; Yang, I.-J. Screening for Neuroprotective and Rapid Antidepressant-like Effects of 20 Essential Oils. *Biomedicines* **2023**, *11*, 1248. <https://doi.org/10.3390/biomedicines11051248>
61. Tanaka, M.; Kádár, K.; Tóth, G.; Telegdy, G. Antidepressant-like effects of urocortin 3 fragments. *Brain Res. Bull.* **2011**, *84*(6), 414-418. <https://doi.org/10.1016/j.brainresbull.2011.01.016>
62. BAliellas, D.E.M.; Barros, M.P.; Vardaris, C.V.; Guariroba, M.; Poppe, S.C.; Martins, M.F.; Pereira, Á.A.F.; Bondan, E.F. Propentofylline Improves Thiol-Based Antioxidant Defenses and Limits Lipid Peroxidation following Gliotoxic Injury in the Rat Brainstem. *Biomedicines* **2023**, *11*, 1652. <https://doi.org/10.3390/biomedicines11061652>
63. Montanari, M.; Imbriani, P.; Bonsi, P.; Martella, G.; Peppe, A. Beyond the Microbiota: Understanding the Role of the Enteric Nervous System in Parkinson's Disease from Mice to Human. *Biomedicines* **2023**, *11*, 1560. <https://doi.org/10.3390/biomedicines11061560>
64. Garifulin, R.; Davleeva, M.; Izmailov, A.; Fadeev, F.; Markosyan, V.; Shevchenko, R.; Minyazeva, I.; Minekayev, T.; Lavrov, I.; Islamov, R. Evaluation of the Autologous Genetically Enriched Leucoconcentrate

- on the Lumbar Spinal Cord Morpho-Functional Recovery in a Mini Pig with Thoracic Spine Contusion Injury. *Biomedicines* **2023**, *11*, 1331. <https://doi.org/10.3390/biomedicines11051331>
- 65. Bueno, C.R.d.S.; Tonin, M.C.C.; Buchaim, D.V.; Barraiva, B.; Ferreira Junior, R.S.; Santos, P.S.d.S.; Reis, C.H.B.; Pastori, C.M.; Pereira, E.d.S.B.M.; Nogueira, D.M.B.; et al. Morphofunctional Improvement of the Facial Nerve and Muscles with Repair Using Heterologous Fibrin Biopolymer and Photobiomodulation. *Pharmaceuticals* **2023**, *16*, 653. <https://doi.org/10.3390/ph16050653>
 - 66. Sojka, A.; Żarowski, M.; Steinborn, B.; Hedzelek, W.; Wiśniewska-Spychała, B.; Dorocka-Bobkowska, B. Temporomandibular disorders in adolescents with headache. *Adv. Clin. Exp. Med.* **2018**, *27*(2), 193–199. <https://doi.org/10.17219/acem/64945>
 - 67. Polyák, H.; Galla, Z.; Nánási, N.; Cseh, E.K.; Rajda, C.; Veres, G.; Spekker, E.; Szabó, Á.; Klivényi, P.; Tanaka, M.; et al. The Tryptophan-Kynurenone Metabolic System Is Suppressed in Cuprizone-Induced Model of Demyelination Simulating Progressive Multiple Sclerosis. *Biomedicines* **2023**, *11*, 945. <https://doi.org/10.3390/biomedicines11030945>
 - 68. Tanaka, M.; Szabó, Á.; Vécsei, L. Preclinical modeling in depression and anxiety: Current challenges and future research directions. *Adv. Clin. Exp. Med.* **2023**, *32*(5), 505–509. <https://doi.org/10.17219/acem/165944>
 - 69. Chu, P.-C.; Huang, C.-S.; Chang, P.-K.; Chen, R.-S.; Chen, K.-T.; Hsieh, T.-H.; Liu, H.-L. Weak Ultrasound Contributes to Neuromodulatory Effects in the Rat Motor Cortex. *Int. J. Mol. Sci.* **2023**, *24*, 2578. <https://doi.org/10.3390/ijms24032578>
 - 70. Gecse, K.; Édes, A.E.; Nagy, T.; Demeter, A.K.; Virág, D.; Király, M.; Dalmadi Kiss, B.; Ludányi, K.; Környei, Z.; Denes, A.; et al. Citalopram Neuroendocrine Challenge Shows Altered Tryptophan and Kynurenone Metabolism in Migraine. *Cells* **2022**, *11*, 2258. <https://doi.org/10.3390/cells11142258>
 - 71. Nasini, S.; Tidei, S.; Shkodra, A.; De Gregorio, D.; Cambiaghi, M.; Comai, S. Age-Related Effects of Exogenous Melatonin on Anxiety-like Behavior in C57/B6J Mice. *Biomedicines* **2023**, *11*, 1705. <https://doi.org/10.3390/biomedicines11061705>
 - 72. Chen, W.-C.; Wang, T.-S.; Chang, F.-Y.; Chen, P.-A.; Chen, Y.-C. Age, Dose, and Locomotion: Decoding Vulnerability to Ketamine in C57BL/6J and BALB/c Mice. *Biomedicines* **2023**, *11*, 1821. <https://doi.org/10.3390/biomedicines11071821>
 - 73. Statsenko, Y.; Habuza, T.; Smetanina, D.; Simiyu, G.L.; Meribout, S.; King, F.C.; Gelovani, J.G.; Das, K.M.; Gorkom, K.N.-V.; Zaręba, K.; et al. Unraveling Lifelong Brain Morphometric Dynamics: A Protocol for Systematic Review and Meta-Analysis in Healthy Neurodevelopment and Ageing. *Biomedicines* **2023**, *11*, 1999. <https://doi.org/10.3390/biomedicines11071999>
 - 74. Dang, J.; Tao, Q.; Niu, X.; Zhang, M.; Gao, X.; Yang, Z.; Yu, M.; Wang, W.; Han, S. et al. Meta-Analysis of Structural and Functional Brain Abnormalities in Cocaine Addiction. *Front. Psychiatry* **2022**, *13*, 927075. <https://doi.org/10.3389/fpsyg.2022.927075>.
 - 75. Balogh, L.; Tanaka, M.; Török, N.; Vécsei, L.; Taguchi, S. Crosstalk between Existential Phenomenological Psychotherapy and Neurological Sciences in Mood and Anxiety Disorders. *Biomedicines* **2021**, *9*, 340. <https://doi.org/10.3390/biomedicines904034>
 - 76. Rassler, B.; Blinowska, K.; Kaminski, M.; Pfurtscheller, G. Analysis of Respiratory Sinus Arrhythmia and Directed Information Flow between Brain and Body Indicate Different Management Strategies of fMRI-Related Anxiety. *Biomedicines* **2023**, *11*, 1028. <https://doi.org/10.3390/biomedicines11041028>
 - 77. Arimura, A. Perspectives on pituitary adenylate cyclase activating polypeptide (PACAP) in the neuroendocrine, endocrine, and nervous systems. *Jpn. J. Physiol.* **1998**, *48*(5), 301–331. <https://doi.org/10.2170/jjphysiol.48.301>
 - 78. Holland, P.R.; Barloese, M.; Fahrenkrug, J. PACAP in hypothalamic regulation of sleep and circadian rhythm: importance for headache. *J. Headache Pain* **2018**, *19*(1), 20. <https://doi.org/10.1186/s10194-018-0844-4>
 - 79. Maugeri, G.; D'Amico, A.G.; Musumeci, G.; Reglodi, D.; D'Agata, V. Effects of PACAP on Schwann Cells: Focus on Nerve Injury. *Int. J. Mol. Sci.* **2020**, *21*, 8233. <https://doi.org/10.3390/ijms21218233>
 - 80. Johnson, G.C.; May, V.; Parsons, R.L.; Hammack, S.E. Parallel signaling pathways of pituitary adenylate cyclase activating polypeptide (PACAP) regulate several intrinsic ion channels. *Ann. N.Y. Acad. Sci.* **2019**, *1455*(1), 105–112. <https://doi.org/10.1111/nyas.14116>
 - 81. Clement, A.; Guo, S.; Jansen-Olesen, I.; Christensen, S.L. ATP-Sensitive Potassium Channels in Migraine: Translational Findings and Therapeutic Potential. *Cells* **2022**, *11*, 2406. <https://doi.org/10.3390/cells11152406>
 - 82. Miyata, A.; Arimura, A.; Dahl, R.R.; Minamino, N.; Uehara, A.; Jiang, L.; Culler, M.D.; Coy, D.H. Isolation of a novel 38 residue-hypothalamic polypeptide which stimulates adenylate cyclase in pituitary cells. *Biochem. Biophys. Res. Commun.* **1989**, *164*, 567–574. [https://doi.org/10.1016/0006-291X\(89\)91757-9](https://doi.org/10.1016/0006-291X(89)91757-9)
 - 83. Denes, V.; Geck, P.; Mester, A.; Gabriel, R. Pituitary Adenylate Cyclase-Activating Polypeptide: 30 Years in Research Spotlight and 600 Million Years in Service. *J. Clin. Med.* **2019**, *8*, 1488. <https://doi.org/10.3390/jcm8091488>
 - 84. Hypophysis Adenylate Cyclase Activating Polypeptide. Handbook of Biologically Active Peptides (Second Edition), 2013. Available online from: <https://www.sciencedirect.com/topics/pharmacology-toxicology>

- and-pharmaceutical-science/hypophysis-adenylate-cyclase-activating-polypeptide (accessed on 28 August 2023).
85. Tam, J.K.; Lee, L.T.; Chow, B.K. PACAP-related peptide (PRP)--molecular evolution and potential functions. *Peptides* **2007**, *28*(9), 1920-1929. <https://doi.org/10.1016/j.peptides.2007.07.011>
 86. Köves, K.; Szabó, E.; Kántor, O.; Heinzlmann, A.; Szabó, F.; Csáki, Á. Current State of Understanding of the Role of PACAP in the Hypothalamo-Hypophyseal Gonadotropin Functions of Mammals. *Front. Endocrinol. (Lausanne)* **2020**, *11*, 88. <https://doi.org/10.3389/fendo.2020.00088>
 87. Lee, E.H.; Seo, S.R. Neuroprotective roles of pituitary adenylate cyclase-activating polypeptide in neurodegenerative diseases. *BMB Rep.* **2014**, *47*(7), 369-375. <https://doi.org/10.5483/bmbrep.2014.47.7.086>
 88. Sadanandan, N.; Cozene, B.; Park, Y.J.; Farooq, J.; Kingsbury, C.; Wang, Z.J.; Moscatello, A.; Saft, M.; Cho, J.; Gonzales-Portillo, B.; Borlongan, C.V. Pituitary Adenylate Cyclase-Activating Polypeptide: A Potent Therapeutic Agent in Oxidative Stress. *Antioxidants (Basel)* **2021**, *10*(3), 354. <https://doi.org/10.3390/antiox10030354>
 89. Waschek, J.A. VIP and PACAP: neuropeptide modulators of CNS inflammation, injury, and repair. *Br. J. Pharmacol.* **2013**, *169*(3), 512-523. <https://doi.org/10.1111/bph.12181>
 90. Xiao, X.; Qiu, P.; Gong, H.Z.; Chen, X.M.; Sun, Y.; Hong, A.; Ma, Y. PACAP ameliorates hepatic metabolism and inflammation through up-regulating FAIM in obesity. *J. Cell Mol. Med.* **2019**, *23*(9), 5970-5980. <https://doi.org/10.1111/jcmm.14453>
 91. Langer, I.; Jeandriens, J.; Couvineau, A.; Sanmukh, S.; Latek, D. Signal Transduction by VIP and PACAP Receptors. *Biomedicines* **2022**, *10*, 406. <https://doi.org/10.3390/biomedicines10020406>
 92. Fizanne, L.; Sigaudo-Roussel, D.; Saumet, J.L.; Fromy, B. Evidence for the involvement of VPAC1 and VPAC2 receptors in pressure-induced vasodilatation in rodents. *J. Physiol.* **2004**, *554*(Pt 2), 519-528. <https://doi.org/10.1113/jphysiol.2003.053835>
 93. Parsons RL, May V. PACAP-Induced PAC1 Receptor Internalization and Recruitment of Endosomal Signaling Regulate Cardiac Neuron Excitability. *J Mol Neurosci.* 2019 Jul;68(3):340-347. doi: 10.1007/s12031-018-1127-x.
 94. Bill, C.A.; Vines, C.M. Phospholipase C. *Adv. Exp. Med. Biol.* **2020**, *1131*, 215-242. https://doi.org/10.1007/978-3-030-12457-1_9
 95. Barloese, M.; Chitgar, M.; Hannibal, J.; Møller, S. Pituitary adenylate cyclase-activating peptide: Potential roles in the pathophysiology and complications of cirrhosis. *Liver Int.* **2020**, *40*(11), 2578-2589. <https://doi.org/10.1111/liv.14602>
 96. Makhinson, M.; Chotiner, J.K.; Watson, J.B.; O'Dell, T.J. Adenylyl cyclase activation modulates activity-dependent changes in synaptic strength and Ca²⁺/calmodulin-dependent kinase II autophosphorylation. *J. Neurosci.* **1999**, *19*(7), 2500-2510. <https://doi.org/10.1523/JNEUROSCI.19-07-02500.1999>
 97. Johnson, G.L.; Lapadat, R. Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science* **2002**, *298*(5600), 1911-1912. <https://doi.org/10.1126/science.1072682>
 98. Lu, J.; Piper, S.J.; Zhao, P.; Miller, L.J.; Wootten, D.; Sexton, P.M. Targeting VIP and PACAP Receptor Signaling: New Insights into Designing Drugs for the PACAP Subfamily of Receptors. *Int. J. Mol. Sci.* **2022**, *23*, 8069. <https://doi.org/10.3390/ijms23158069>
 99. Hirabayashi, T.; Nakamachi, T.; Shioda, S. Discovery of PACAP and its receptors in the brain. *J. Headache Pain*, **2018**, *19*, 28. <https://doi.org/10.1186/s10194-018-0855-1>
 100. Vasoactive Intestinal Polypeptide Receptor 1. Methods in Enzymology, 2013. Available online: <https://www.sciencedirect.com/topics/medicine-and-dentistry/vasoactive-intestinal-polypeptide-receptor-1> (accessed on 28 August 2023).
 101. Vasoactive Intestinal Polypeptide Receptor. Autonomic Neuroscience, 2007. Available online: <https://www.sciencedirect.com/topics/neuroscience/vasoactive-intestinal-polypeptide-receptor> (accessed on 28 August 2023).
 102. May, V.; Buttolph, T.R.; Girard, B.M.; Clason, T.A.; Parsons, R.L. PACAP-induced ERK activation in HEK cells expressing PAC1 receptors involves both receptor internalization and PKC signaling. *Am. J. Physiol. Cell Physiol.* **2014**, *306*(11), C1068-1079. <https://doi.org/10.1152/ajpcell.00001.2014>
 103. Hou, X.; Yang, D.; Yang, G.; Li, M.; Zhang, J.; Zhang, J.; Zhang, Y.; Liu, Y. Therapeutic potential of vasoactive intestinal peptide and its receptor VPAC2 in type 2 diabetes. *Front. Endocrinol. (Lausanne)* **2022**, *13*, 984198. <https://doi.org/10.3389/fendo.2022.984198>
 104. Sundrum, T.; Walker, C.S. Pituitary adenylate cyclase-activating polypeptide receptors in the trigeminovascular system: implications for migraine. *Br. J. Pharmacol.* **2018**, *175*(21), 4109-4120. <https://doi.org/10.1111/bph.14053>
 105. Liu, J.; Wang, G.; Dan, Y.; Liu, X. CGRP and PACAP-38 play an important role in diagnosing pediatric migraine. *J. Headache Pain* **2022**, *23*(1), 68. <https://doi.org/10.1186/s10194-022-01435-7>
 106. Schytz, H.W.; Olesen, J.; Ashina, M. The PACAP receptor: a novel target for migraine treatment. *Neurotherapeutics* **2010**, *7*(2), 191-196. <https://doi.org/10.1016/j.nurt.2010.02.003>

107. Kuburas, A.; Russo, A.F. Shared and independent roles of CGRP and PACAP in migraine pathophysiology. *J. Headache Pain* **2023**, *24*(1), 34. <https://doi.org/10.1186/s10194-023-01569-2>
108. Ernstsen, C.; Christensen, S.L.; Rasmussen, R.H.; Nielsen, B.S.; Jansen-Olesen, I.; Olesen, J.; Kristensen, D.M. The PACAP pathway is independent of CGRP in mouse models of migraine: possible new drug target? *Brain* **2022**, *145*(7), 2450–2460. <https://doi.org/10.1093/brain/awac040>
109. Christensen, C.E.; Ashina, M.; Amin, F.M. Calcitonin Gene-Related Peptide (CGRP) and Pituitary Adenylate Cyclase-Activating Polypeptide (PACAP) in Migraine Pathogenesis. *Pharmaceuticals* **2022**, *15*, 1189. <https://doi.org/10.3390/ph15101189>
110. Anapindi, K.D.B.; Yang, N.; Romanova, E.V.; Rubakhin, S.S.; Tipton, A.; Dripps, I.; Sheets, Z.; Sweedler, J.V.; Pradhan, A.A. PACAP and Other Neuropeptide Targets Link Chronic Migraine and Opioid-induced Hyperalgesia in Mouse Models. *Mol. Cell Proteomics* **2019**, *18*(12), 2447–2458. <https://doi.org/10.1074/mcp.RA119.001767>
111. Silvestro, M.; Iannone, L.F.; Orologio, I.; Tessitore, A.; Tedeschi, G.; Geppetti, P.; Russo, A. Migraine Treatment: Towards New Pharmacological Targets. *Int. J. Mol. Sci.* **2023**, *24*, 12268. <https://doi.org/10.3390/ijms241512268>
112. Pellesi, L.; Chaudhry, B.A.; Vollesen, A.L.H.; Snoer, A.H.; Baumann, K.; Skov, P.S.; Jensen, R.H.; Ashina, M. PACAP38- and VIP-induced cluster headache attacks are not associated with changes of plasma CGRP or markers of mast cell activation. *Cephalgia* **2022**, *42*(8), 687–695. <https://doi.org/10.1177/03331024211056248>
113. Rasmussen, N.B.; Deligianni, C.; Christensen, C.E.; Karlsson, W.K.; Al-Khadali, H.M.; Van de Casteele, T.; Granhall, C.; Amin, F.M.; Ashina, M. The effect of Lu AG09222 on PACAP38- and VIP-induced vasodilation, heart rate increase, and headache in healthy subjects: an interventional, randomized, double-blind, parallel-group, placebo-controlled study. *J. Headache Pain* **2023**, *24*(1), 60. <https://doi.org/10.1186/s10194-023-01599-w>
114. Adenylate Cyclase. Adenylate cyclases are enzymes that catalyze the conversion of ATP to cAMP and pyrophosphate. From: The Senses: A Comprehensive Reference, 2008. Available online <https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/adenylate-cyclase> (accessed on 29 August 2023).
115. Roberts, R.E. The extracellular signal-regulated kinase (ERK) pathway: a potential therapeutic target in hypertension. *J. Exp. Pharmacol.* **2012**, *4*, 77–83. <https://doi.org/10.2147/JEP.S28907>
116. Lund, A.M.; Hannibal, J. Localization of the neuropeptides pituitary adenylate cyclase-activating polypeptide, vasoactive intestinal peptide, and their receptors in the basal brain blood vessels and trigeminal ganglion of the mouse CNS; an immunohistochemical study. *Front. Neuroanat.* **2022**, *16*, 991403. <https://doi.org/10.3389/fnana.2022.991403>
117. Ivic, I.; Balasko, M.; Fulop, B.D.; Hashimoto, H.; Toth, G.; Tamas, A.; Juhasz, T.; Koller, A.; Reglodi, D.; Solymár, M. VPAC1 receptors play a dominant role in PACAP-induced vasorelaxation in female mice. *PLoS One* **2019**, *14*(1), e0211433. <https://doi.org/10.1371/journal.pone.0211433>
118. VIP and PACAP receptors - IUPHAR/BPS Guide to PHARMACOLOGY. <https://www.guidetopharmacology.org/GRAC/FamilyDisplayForward?familyId=67> (accessed on 29 August 2023)
119. Datki, Z.; Sinka, R. Translational biomedicine-oriented exploratory research on bioactive rotifer-specific biopolymers. *Adv. Clin. Exp. Med.* **2022**, *31*(9), 931–935. <https://doi.org/10.17219/acem/152430>
120. Palotai, M.; Telegdy, G.; Tanaka, M.; Bagosi, Z.; Jászberényi, M. Neuropeptide AF induces anxiety-like and antidepressant-like behavior in mice. *Behav. Brain Res.* **2014**, *274*, 264–269. <https://doi.org/10.1016/j.bbr.2014.08.007>
121. Lieb, A.; Thaler, G.; Fogli, B.; Trovato, O.; Posch, M.A.; Kaserer, T.; Zangrandi, L. Functional Characterization of Spinocerebellar Ataxia Associated Dynorphin A Mutant Peptides. *Biomedicines* **2021**, *9*, 1882. <https://doi.org/10.3390/biomedicines9121882>
122. Skobeleva, K.; Shalygin, A.; Mikhaylova, E.; Guzhova, I.; Ryazantseva, M.; Kaznacheyeva, E. The STIM1/2-Regulated Calcium Homeostasis Is Impaired in Hippocampal Neurons of the 5xFAD Mouse Model of Alzheimer's Disease. *Int. J. Mol. Sci.* **2022**, *23*, 14810. <https://doi.org/10.3390/ijms232314810>
123. Martos, D.; Tuka, B.; Tanaka, M.; Vécsei, L.; Telegdy, G. Memory Enhancement with Kynurenic Acid and Its Mechanisms in Neurotransmission. *Biomedicines* **2022**, *10*, 849. <https://doi.org/10.3390/biomedicines10040849>
124. Tanaka, M.; Szabó, Á.; Spekker, E.; Polyák, H.; Tóth, F.; Vécsei, L. Mitochondrial Impairment: A Common Motif in Neuropsychiatric Presentation? The Link to the Tryptophan–Kynurenone Metabolic System. *Cells* **2022**, *11*, 2607. <https://doi.org/10.3390/cells11162607>
125. Tanaka, M.; Bohár, Z.; Martos, D.; Telegdy, G.; Vécsei, L. Antidepressant-like effects of kynurenic acid in a modified forced swim test. *Pharmacol. Rep.* **2020**, *72*(2), 449–455. <https://doi.org/10.1007/s43440-020-00067-5>

126. Tanaka, M.; Telegdy, G. Involvement of adrenergic and serotonergic receptors in antidepressant-like effect of urocortin 3 in a modified forced swimming test in mice. *Brain Res. Bull.* **2008**, *77*(5), 301-305. <https://doi.org/10.1016/j.brainresbull.2008.08.012>
127. Tanaka, M.; Spekker, E.; Szabó, Á.; Polyák, H.; Vécsei, L. Modelling the neurodevelopmental pathogenesis in neuropsychiatric disorders. Bioactive kynurenes and their analogues as neuroprotective agents-in celebration of 80th birthday of Professor Peter Riederer. *J. Neural. Transm. (Vienna)* **2022**, *129*(5-6), 627-642. <https://doi.org/10.1007/s00702-022-02513-5>
128. Reducha, P.V.; Edvinsson, L.; Haanes, K.A. Could Experimental Inflammation Provide Better Understanding of Migraines? *Cells* **2022**, *11*, 2444. <https://doi.org/10.3390/cells11152444>
129. Ojala, J.; Tooke, K.; Hsiang, H.; Girard, B.M.; May, V.; Vizzard, M.A. PACAP/PAC1 Expression and Function in Micturition Pathways. *J. Mol. Neurosci.* **2019**, *68*(3), 357-367. <https://doi.org/10.1007/s12031-018-1170-7>
130. Tamas, A.; Reglodi, D.; Farkas, O.; Kovács, E.; Pal, J.; Povlishock, J.T.; Schwarcz, A.; Czeiter, E.; Szanto, Z.; Doczi, T.; et al. Effect of PACAP in central and peripheral nerve injuries. *Int. J. Mol. Sci.* **2012**, *13*(7), 8430-8448. <https://doi.org/10.3390/ijms13078430>
131. Zhang, L.; Zhou, Y.; Yang, L.; Wang, Y.; Xiao, Z. PACAP6-38 improves nitroglycerin-induced central sensitization by modulating synaptic plasticity at the trigeminal nucleus caudalis in a male rat model of chronic migraine. *J. Headache Pain* **2023**, *24*(1), 66. <https://doi.org/10.1186/s10194-023-01603-3>
132. Takács-Lovász K, Kun J, Aczél T, Urbán P, Gyenesi A, Bölcsei K, Szőke É, Helyes Z. PACAP-38 Induces Transcriptomic Changes in Rat Trigeminal Ganglion Cells Related to Neuroinflammation and Altered Mitochondrial Function Presumably via PAC1/VPAC2 Receptor-Independent Mechanism. *Int. J. Mol. Sci.* **2022**, *23*(4), 2120. <https://doi.org/10.3390/ijms23042120>
133. Frederiksen, S.D.; Haanes, K.A.; Warfvinge, K.; Edvinsson, L. Perivascular neurotransmitters: Regulation of cerebral blood flow and role in primary headaches. *J. Cereb. Blood Flow Metab.* **2019**, *39*(4), 610-632. <https://doi.org/10.1177/0271678X17747188>
134. Markovics, A.; Kormos, V.; Gaszner, B.; Lashgarara, A.; Szoke, E.; Sandor, K.; Szabadfi, K.; Tuka, B.; Tajti, J.; Szolcsányi, J.; et al. Pituitary adenylate cyclase-activating polypeptide plays a key role in nitroglycerol-induced trigeminovascular activation in mice. *Neurobiol. Dis.* **2012**, *45*(1), 633-644. <https://doi.org/10.1016/j.nbd.2011.10.010>
135. Edvinsson, L. PACAP and its receptors in migraine pathophysiology: Commentary on Walker et al., Br. J. Pharmacol. 171: 1521-1533. *Br. J. Pharmacol.* **2015**, *172*(19), 4782-4784. <https://doi.org/10.1111/bph.12772>
136. Saposnik, G.; Montalban, X.; Selchen, D.; Terzaghi, M.A.; Bakdache, F.; Montoya, A.; Fruns, M.; Caceres, F.; Oh, J. Therapeutic Inertia in Multiple Sclerosis Care: A Study of Canadian Neurologists. *Front. Neurol.* **2018**, *9*, 781. <https://doi.org/10.3389/fneur.2018.00781>
137. Harding, S.D.; Armstrong, J.F.; Faccenda, E.; Southan, C.; Alexander, S.P.H.; Davenport, A.P.; Pawson, A.J.; Spedding, M.; Davies, J.A.; NC-IUPHAR. (2021) The IUPHAR/BPS guide to PHARMACOLOGY in 2022: curating pharmacology for COVID-19, malaria and antibacterials. *Nucl. Acids Res.* **2022**, *50*(D1), D1282-D1294. <https://doi.org/10.1093/nar/gkab1010>
138. Beebe, X.; Darczak, D.; Davis-Taber, R.A.; Uchic, M.E.; Scott, V.E.; Jarvis, M.F.; Stewart, A.O. Discovery and SAR of hydrazide antagonists of the pituitary adenylate cyclase-activating polypeptide (PACAP) receptor type 1 (PAC1-R). *Bioorg. Med. Chem. Lett.* **2008**, *18*(6), 2162-2166. <https://doi.org/10.1016/j.bmcl.2008.01.052>
139. Laburthe, M.; Couvineau, A.; Tan, V. Class II G protein-coupled receptors for VIP and PACAP: structure, models of activation and pharmacology. *Peptides* **2007**, *28*(9), 1631-1639. <https://doi.org/10.1016/j.peptides.2007.04.026>
140. Spekker, E.; Tanaka M, Szabó Á, Vécsei L. Neurogenic Inflammation: The Participant in Migraine and Recent Advancements in Translational Research. *Biomedicines.* **2022**; *10*(1):76. <https://doi.org/10.3390/biomedicines10010076>
141. Pinho-Ribeiro, F.A.; Verri, W.A. Jr.; Chiu, I.M. Nociceptor Sensory Neuron-Immune Interactions in Pain and Inflammation. *Trends Immunol.* **2017**, *38*(1), 5-19. <https://doi.org/10.1016/j.it.2016.10.001>
142. Guo, S.; Vollesen, A.L.; Hansen, R.D.; Esserlind, A.L.; Amin, F.M.; Christensen, A.F.; Olesen, J.; Ashina, M. Part I: Pituitary adenylate cyclase-activating polypeptide-38 induced migraine-like attacks in patients with and without familial aggregation of migraine. *Cephalalgia.* **2017**, *37*(2), 125-135. <https://doi.org/10.1177/0333102416639516>
143. ClinicalTrials.gov; PACAP Induced Migraine Attacks in Patients With High and Low Genetic Load. Available online: <https://classic.clinicaltrials.gov/ct2/show/NCT02158221> (accessed on 29 August 2023).
144. Togha, M.; Ghorbani, Z.; Ramazi, S.; Zavvari, F.; Karimzadeh, F. Evaluation of Serum Levels of Transient Receptor Potential Cation Channel Subfamily V Member 1, Vasoactive Intestinal Polypeptide, and Pituitary Adenylate Cyclase-Activating Polypeptide in Chronic and Episodic Migraine: The Possible Role in Migraine Transformation. *Front. Neurol.* **2021**, *12*, 770980. <https://doi.org/10.3389/fneur.2021.770980>

145. Körtési, T.; Tuka, B.; Tajti, J.; Bagoly, T.; Fülop, F.; Helyes, Z.; Vécsei, L. Kynurenic Acid Inhibits the Electrical Stimulation Induced Elevated Pituitary Adenylate Cyclase-Activating Polypeptide Expression in the TNC. *Front. Neurol.* **2018**, *8*, 745. <https://doi.org/10.3389/fneur.2017.00745>
146. Guo, S.; Vollesen, A.L.; Hansen, Y.B.; Frandsen, E.; Andersen, M.R.; Amin, F.M.; Fahrenkrug, J.; Olesen, J.; Ashina, M. Part II: Biochemical changes after pituitary adenylate cyclase-activating polypeptide-38 infusion in migraine patients. *Cephalgia* **2017**, *37*(2), 136-147. <https://doi.org/10.1177/0333102416639517>
147. Amin, F.M.; Asghar, M.S.; Guo, S.; Hougaard, A.; Hansen, A.E.; Schytz, H.W.; van der Geest, R.J.; de Koning, P.J.; Larsson, H.B.; Olesen, J.; Ashina, M. Headache and prolonged dilatation of the middle meningeal artery by PACAP38 in healthy volunteers. *Cephalgia* **2012**, *32*(2), 140-149. <https://doi.org/10.1177/0333102411431333>
148. Maasz, G.; Zrinyi, Z.; Reglodi, D.; Petrovics, D.; Rivnyak, A.; Kiss, T.; Jungling, A.; Tamas, A.; Pirger, Z. Pituitary adenylate cyclase-activating polypeptide (PACAP) has a neuroprotective function in dopamine-based neurodegeneration in rat and snail parkinsonian models. *Dis. Model Mech.* **2017**, *10*(2), 127-139. <https://doi.org/10.1242/dmm.027185>
149. Rubio-Beltrán, E.; Correnti, E.; Deen, M.; Kamm, K.; Kelderman, T.; Papetti, L.; Vigneri, S.; MaassenVanDenBrink, A.; Edvinsson, L.; European Headache Federation School of Advanced Studies (EHF-SAS). PACAP38 and PAC₁ receptor blockade: a new target for headache? *J. Headache Pain* **2018**, *19*, 64. <https://doi.org/10.1186/s10194-018-0893-8>
150. Ashina, M.; Doležil, D.; Bonner, J.H.; Zhou, L.; Klatt, J.; Picard, H.; Mikol, D.D. A phase 2, randomized, double-blind, placebo-controlled trial of AMG 301, a pituitary adenylate cyclase-activating polypeptide PAC1 receptor monoclonal antibody for migraine prevention. *Cephalgia* **2021**, *41*(1), 33-44. <https://doi.org/10.1177/0333102420970889>
151. Study to Evaluate the Efficacy and Safety of AMG 301 in Migraine Prevention 2020. Available online: <https://ClinicalTrials.gov/show/NCT03238781> (accessed on 4 September 2023).
152. Lundbeck News Room: Lundbeck announced the start of a phase II clinical study to assess Lu AG09222 for migraine prevention. Available online: <https://newsroom.lundbeckus.com/news-release/2021/lundbeck-announced-start-of-phase-ii-clinical-study-for-migraine-prevention> (Accessed on 29/August 2023).
153. A Study With Lu AG09222 in Adults With Migraine Who Have Not Been Helped by Prior Preventive Treatments 2023. Available online: <https://classic.clinicaltrials.gov/ct2/show/NCT05133323> (accessed on 4 September 2023).
154. A Study of LY3451838 in Participants With Migraine 2023. Available online: <https://ClinicalTrials.gov/show/NCT04498910> (accessed on 4 September 2023).
155. ClinicalTrials.gov; The Effects of a Long-lasting Infusion of Vasoactive Intestinal Peptide (VIP) in Episodic Migraine Patients. Available online: <https://classic.clinicaltrials.gov/ct2/show/NCT04260035> (accessed on 29 August 2023).
156. Tanaka, M.; Szabó, Á.; Vécsei, L. Integrating Armchair, Bench, and Bedside Research for Behavioral Neurology and Neuropsychiatry: Editorial. *Biomedicines* **2022**, *10*, 2999. <https://doi.org/10.3390/biomedicines10122999>
157. Gaebler, A.J.; Finner-Prével, M.; Sudar, F.P.; Langer, F.H.; Keskin, F.; Gebel, A.; Zweerings, J.; Mathiak, K. The Interplay between Vitamin D, Exposure of Anticholinergic Antipsychotics and Cognition in Schizophrenia. *Biomedicines* **2022**, *10*, 1096. <https://doi.org/10.3390/biomedicines10051096>
158. Castillo-Mariqueo, L.; Giménez-Llort, L. Impact of Behavioral Assessment and Re-Test as Functional Trainings That Modify Survival, Anxiety and Functional Profile (Physical Endurance and Motor Learning) of Old Male and Female 3xTg-AD Mice and NTg Mice with Normal Aging. *Biomedicines* **2022**, *10*, 973. <https://doi.org/10.3390/biomedicines10050973>
159. Lee, E.C.; Hong, D.-Y.; Lee, D.-H.; Park, S.-W.; Lee, J.Y.; Jeong, J.H.; Kim, E.-Y.; Chung, H.-M.; Hong, K.-S.; Park, S.-P.; et al. Inflammation and Rho-Associated Protein Kinase-Induced Brain Changes in Vascular Dementia. *Biomedicines* **2022**, *10*, 446. <https://doi.org/10.3390/biomedicines10020446>
160. Simonato, M.; Dall'Acqua, S.; Zilli, C.; Sut, S.; Tenconi, R.; Gallo, N.; Sfriso, P.; Sartori, L.; Cavallin, F.; Fiocco, U.; et al. Tryptophan Metabolites, Cytokines, and Fatty Acid Binding Protein 2 in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Biomedicines* **2021**, *9*, 1724. <https://doi.org/10.3390/biomedicines9111724>
161. Smagin, D.A.; Kovalenko, I.L.; Galyamina, A.G.; Belozertseva, I.V.; Tamkovich, N.V.; Baranov, K.O.; Kudryavtseva, N.N. Chronic Lithium Treatment Affects Anxious Behaviors and the Expression of Serotonergic Genes in Midbrain Raphe Nuclei of Defeated Male Mice. *Biomedicines* **2021**, *9*, 1293. <https://doi.org/10.3390/biomedicines9101293>
162. Tanaka, M.; Tóth, F.; Polyák, H.; Szabó, Á.; Mándi, Y.; Vécsei, L. Immune Influencers in Action: Metabolites and Enzymes of the Tryptophan-Kynurenone Metabolic Pathway. *Biomedicines* **2021**, *9*, 734. <https://doi.org/10.3390/biomedicines9070734>

163. Tanaka, M.; Török, N.; Tóth, F.; Szabó, Á.; Vécsei, L. Co-Players in Chronic Pain: Neuroinflammation and the Tryptophan-Kynurenine Metabolic Pathway. *Biomedicines* **2021**, *9*, 897. <https://doi.org/10.3390/biomedicines9080897>
164. Vila-Merkle, H.; González-Martínez, A.; Campos-Jiménez, R.; Martínez-Ricós, J.; Teruel-Martí, V.; Blasco-Serra, A.; Lloret, A.; Celada, P.; Cervera-Ferri, A. The Oscillatory Profile Induced by the Anxiogenic Drug FG-7142 in the Amygdala–Hippocampal Network Is Reversed by Infralimbic Deep Brain Stimulation: Relevance for Mood Disorders. *Biomedicines* **2021**, *9*, 783. <https://doi.org/10.3390/biomedicines9070783>
165. Santana-Santana, M.; Bayascas, J.-R.; Giménez-Llort, L. Fine-Tuning the PI3K/Akt Signaling Pathway Intensity by Sex and Genotype-Load: Sex-Dependent Homozygotic Threshold for Somatic Growth but Feminization of Anxious Phenotype in Middle-Aged PDK1 K465E Knock-In and Heterozygous Mice. *Biomedicines* **2021**, *9*, 747. <https://doi.org/10.3390/biomedicines9070747>
166. Muntasant, A.; Giménez-Llort, L. Genotype Load Modulates Amyloid Burden and Anxiety-Like Patterns in Male 3xTg-AD Survivors despite Similar Neuro-Immunoendocrine, Synaptic and Cognitive Impairments. *Biomedicines* **2021**, *9*, 715. <https://doi.org/10.3390/biomedicines9070715>
167. Giménez-Llort, L.; Marin-Pardo, D.; Marazuela, P.; Hernández-Guillamón, M. Survival Bias and Crosstalk between Chronological and Behavioral Age: Age- and Genotype-Sensitivity Tests Define Behavioral Signatures in Middle-Aged, Old, and Long-Lived Mice with Normal and AD-Associated Aging. *Biomedicines* **2021**, *9*, 636. <https://doi.org/10.3390/biomedicines9060636>
168. Komatsu, H.; Watanabe, E.; Fukuchi, M. Psychiatric Neural Networks and Precision Therapeutics by Machine Learning. *Biomedicines* **2021**, *9*, 403. <https://doi.org/10.3390/biomedicines9040403>
169. Caruso, G.; Godos, J.; Castellano, S.; Micek, A.; Murabito, P.; Galvano, F.; Ferri, R.; Grossi, G.; Caraci, F. The Therapeutic Potential of Carnosine/Anserine Supplementation against Cognitive Decline: A Systematic Review with Meta-Analysis. *Biomedicines* **2021**, *9*, 253. <https://doi.org/10.3390/biomedicines9030253>
170. Correia, B.S.B.; Nani, J.V.; Waladares Ricardo, R.; Stanisic, D.; Costa, T.B.B.C.; Hayashi, M.A.F.; Tasic, L. Effects of Psychostimulants and Antipsychotics on Serum Lipids in an Animal Model for Schizophrenia. *Biomedicines* **2021**, *9*, 235. <https://doi.org/10.3390/biomedicines9030235>
171. Ikonnikova, A.; Anisimova, A.; Galkin, S.; Gunchenko, A.; Abdukhalikova, Z.; Filippova, M.; Surzhikov, S.; Selyaeva, L.; Shershov, V.; Zasedatelev, A.; et al. Genetic Association Study and Machine Learning to Investigate Differences in Platelet Reactivity in Patients with Acute Ischemic Stroke Treated with Aspirin. *Biomedicines* **2022**, *10*, 2564. <https://doi.org/10.3390/biomedicines10102564>
172. Fan, P.; Miranda, O.; Qi, X.; Kofler, J.; Sweet, R.A.; Wang, L. Unveiling the Enigma: Exploring Risk Factors and Mechanisms for Psychotic Symptoms in Alzheimer's Disease through Electronic Medical Records with Deep Learning Models. *Pharmaceuticals* **2023**, *16*, 911. <https://doi.org/10.3390/ph16070911>
173. Parolini, F.; Goethel, M.; Becker, K.; Fernandes, C.; Fernandes, R.J.; Ervilha, U.F.; Santos, R.; Vilas-Boas, J.P. Breaking Barriers: Artificial Intelligence Interpreting the Interplay between Mental Illness and Pain as Defined by the International Association for the Study of Pain. *Biomedicines* **2023**, *11*, 2042. <https://doi.org/10.3390/biomedicines11072042>
174. Tanaka, M.; Diano, M.; Battaglia, S. Editorial: Insights into structural and functional organization of the brain: evidence from neuroimaging and non-invasive brain stimulation techniques. *Front. Psychiatry* **2023**, *14*, 1225755. doi: 10.3389/fpsyg.2023.1225755
175. Simon, C.; Soga, T.; Ahemad, N.; Bhuvanendran, S.; Parhar, I. Kisspeptin-10 Rescues Cholinergic Differentiated SHSY-5Y Cells from α -Synuclein-Induced Toxicity In Vitro. *Int. J. Mol. Sci.* **2022**, *23*, 5193. <https://doi.org/10.3390/ijms23095193>
176. Okanda Nyatega, C.; Qiang, L.; Jajere Adamu, M.; Bello Kawa, H. Altered striatal functional connectivity and structural dysconnectivity in individuals with bipolar disorder: A resting state magnetic resonance imaging study. *Front. Psychiatry* **2022**, *13*, 1054380. <https://doi.org/10.3389/fpsyg.2022.1054380>
177. Liu, N.; Li, Y.; Hong, Y.; Huo, J.; Chang, T.; Wang, H.; Huang, Y.; Li, W.; Zhang, Y. Altered brain activities in mesocorticolimbic pathway in primary dysmenorrhea patients of long-term menstrual pain. *Front. Neurosci.* **2023**, *17*, 1098573. <https://doi.org/10.3389/fnins.2023.1098573>
178. Du, H.; Yang, B.; Wang, H.; Zeng, Y.; Xin, J.; Li, X. The non-linear correlation between the volume of cerebral white matter lesions and incidence of bipolar disorder: A secondary analysis of data from a cross-sectional study. *Front. Psychiatry* **2023**, *14*, 1149663. <https://doi.org/10.3389/fpsyg.2023.1149663>
179. Chen, Y.; Yu, R.; DeSouza, J.F.X.; Shen, Y.; Zhang, H.; Zhu, C.; Huang, P.; Wang, C. Differential responses from the left postcentral gyrus, right middle frontal gyrus, and precuneus to meal ingestion in patients with functional dyspepsia. *Front. Psychiatry* **2023**, *14*, 1184797. <https://doi.org/10.3389/fpsyg.2023.1184797>
180. Adamu, M.J.; Qiang, L.; Nyatega, C.O.; Younis, A.; Kawa, H.B.; Jabire, A.H.; Saminu, S. Unraveling the pathophysiology of schizophrenia: insights from structural magnetic resonance imaging studies. *Front. Psychiatry* **2023**, *14*, 1188603. <https://doi.org/10.3389/fpsyg.2023.1188603>
181. Vuralli, D.; Ayata, C.; Bolay, H. Cognitive dysfunction and migraine. *J. Headache Pain* **2018**, *19*(1), 109. <https://doi.org/10.1186/s10194-018-0933-4>

182. Minen, M.T.; Begasse De Dhaem, O.; Kroon Van Diest, A.; Powers, S.; Schwedt, T.J.; Lipton, R.; Silbersweig, D. Migraine and its psychiatric comorbidities. *J. Neurol. Neurosurg. Psychiatry* **2016**, *87*(7), 741-749. <https://doi.org/10.1136/jnnp-2015-312233>
183. Tanaka, M.; Chen, C. Editorial: Towards a mechanistic understanding of depression, anxiety, and their comorbidity: perspectives from cognitive neuroscience. *Front. Behav. Neurosci.* **2023**, *17*, 1268156. <https://doi.org/10.3389/fnbeh.2023.1268156>
184. Gonzalez-Escamilla, G.; Dörfel, D.; Becke, M.; Trefz, J.; Bonanno, G.A.; Groppa, S. Associating Flexible Regulation of Emotional Expression With Psychopathological Symptoms. *Front. Behav. Neurosci.* **2022**, *16*, 924305. <https://doi.org/10.3389/fnbeh.2022.924305>.
185. Battaglia, S.; Cardellicchio, P.; Di Fazio, C.; Nazzari, C.; Fracasso, A.; Borgomaneri, S. The Influence of Vicarious Fear-Learning in "Infecting" Reactive Action Inhibition. *Front. Behav. Neurosci.* **2022**, *16*, 946263. <https://doi.org/10.3389/fnbeh.2022.946263>
186. Battaglia, S.; Cardellicchio, P.; Di Fazio, C.; Nazzari, C.; Fracasso, A.; Borgomaneri, S. Stopping in (e)motion: Reactive action inhibition when facing valence-independent emotional stimuli. *Front Behav. Neurosci.* **2022**, *16*, 998714. <https://doi.org/10.3389/fnbeh.2022.998714>.
187. Ironside, M.; DeVille, D.C.; Kuplicki, R.T.; Burrows, K.P.; Smith, R.; Teed, A.R.; Paulus, M.P.; Khalsa, S.S. The unique face of comorbid anxiety and depression: increased interoceptive fearfulness and reactivity. *Front. Behav. Neurosci.* **2023**, *16*, 1083357. <https://doi.org/10.3389/fnbeh.2022.1083357>
188. Rajkumar, R.P. Comorbid depression and anxiety: Integration of insights from attachment theory and cognitive neuroscience, and their implications for research and treatment. *Front. Behav. Neurosci.* **2022**, *16*, 1104928. <https://doi.org/10.3389/fnbeh.2022.1104928>.
189. Vila-Merkle, H.; González-Martínez, A.; Campos-Jiménez, R.; Martínez-Ricós, J.; Teruel-Martí, V.; Lloret, A.; Blasco-Serra, A.; Cervera-Ferri, A. Sex differences in amygdalohippocampal oscillations and neuronal activation in a rodent anxiety model and in response to infralimbic deep brain stimulation. *Front. Behav. Neurosci.* **2023**, *17*, 1122163. <https://doi.org/10.3389/fnbeh.2023.1122163>
190. Panov, G.; Panova, P. Obsessive-compulsive symptoms in patient with schizophrenia: The influence of disorganized symptoms, duration of schizophrenia, and drug resistance. *Front. Psychiatry* **2023**, *14*, 1120974. <https://doi.org/10.3389/fpsyg.2023.1120974>
191. Hakamata, Y.; Hori, H.; Mizukami, S.; Izawa, S.; Yoshida, F.; Moriguchi, Y.; Hanakawa, T.; Inoue, Y.; Tagaya, H. Blunted diurnal interleukin-6 rhythm is associated with amygdala emotional hyporeactivity and depression: a modulating role of gene-stressor interactions. *Front. Psychiatry* **2023**, *14*, 1196235. <https://doi.org/10.3389/fpsyg.2023.1196235>.
192. Fraile-Ramos, J.; Garrit, A.; Reig-Vilallonga, J.; Giménez-Llort, L. Hepatic Oxi-Inflammation and Neophobia as Potential Liver-Brain Axis Targets for Alzheimer's Disease and Aging, with Strong Sensitivity to Sex, Isolation, and Obesity. *Cells* **2023**, *12*, 1517. <https://doi.org/10.3390/cells12111517>
193. Sobolewska-Nowak, J.; Wachowska, K.; Nowak, A.; Orzechowska, A.; Szulc, A.; Plaza, O.; Gałecki, P. Exploring the Heart-Mind Connection: Unraveling the Shared Pathways between Depression and Cardiovascular Diseases. *Biomedicines* **2023**, *11*, 1903. <https://doi.org/10.3390/biomedicines11071903>
194. Festa, F.; Medori, S.; Macrì, M. Move Your Body, Boost Your Brain: The Positive Impact of Physical Activity on Cognition across All Age Groups. *Biomedicines* **2023**, *11*, 1765. <https://doi.org/10.3390/biomedicines11061765>
195. Spekker, E.; Bohár, Z.; Fejes-Szabó, A.; Szűcs, M.; Vécsei, L.; Párdutz, Á. Estradiol Treatment Enhances Behavioral and Molecular Changes Induced by Repetitive Trigeminal Activation in a Rat Model of Migraine. *Biomedicines* **2022**, *10*, 3175. <https://doi.org/10.3390/biomedicines10123175>
196. Park, S.Y.; Lee, S.P.; Kim, D.; Kim, W.J. Gut Dysbiosis: A New Avenue for Stroke Prevention and Therapeutics. *Biomedicines* **2023**, *11*, 2352. <https://doi.org/10.3390/biomedicines11092352>
197. Montanari, M.; Imbriani, P.; Bonsi, P.; Martella, G.; Peppe, A. Beyond the Microbiota: Understanding the Role of the Enteric Nervous System in Parkinson's Disease from Mice to Human. *Biomedicines* **2023**, *11*, 1560. <https://doi.org/10.3390/biomedicines11061560>
198. Altamura, C.; Corbelli, I.; de Tommaso, M.; Di Lorenzo, C.; Di Lorenzo, G.; Di Renzo A, Filippi, M.; Jannini, T.B.; Messina, R.; Parisi, P.; et al. Pathophysiological Bases of Comorbidity in Migraine. *Front. Hum. Neurosci.* **2021**, *15*, 640574. <https://doi.org/10.3389/fnhum.2021.640574>
199. Schwedt, T.J.; Vargas, B. Neurostimulation for Treatment of Migraine and Cluster Headache. *Pain Med.* **2015**, *16*(9), 1827-1834. <https://doi.org/10.1111/pme.12792>
200. de Albuquerque, L.L.; Pantovic, M.; Clingo, M.; Fischer, K.; Jalene, S.; Landers, M.; Mari, Z.; Poston, B. A Single Application of Cerebellar Transcranial Direct Current Stimulation Fails to Enhance Motor Skill Acquisition in Parkinson's Disease: A Pilot Study. *Biomedicines* **2023**, *11*, 2219. <https://doi.org/10.3390/biomedicines11082219>
201. Senevirathne, D.K.L.; Mahboob, A.; Zhai, K.; Paul, P.; Kammen, A.; Lee, D.J.; Yousef, M.S.; Chaari, A. Deep Brain Stimulation beyond the Clinic: Navigating the Future of Parkinson's and Alzheimer's Disease Therapy. *Cells* **2023**, *12*, 1478. <https://doi.org/10.3390/cells12111478>

202. Chu, P.-C.; Huang, C.-S.; Chang, P.-K.; Chen, R.-S.; Chen, K.-T.; Hsieh, T.-H.; Liu, H.-L. Weak Ultrasound Contributes to Neuromodulatory Effects in the Rat Motor Cortex. *Int. J. Mol. Sci.* **2023**, *24*, 2578. <https://doi.org/10.3390/ijms24032578>
203. Adeel, M.; Chen, C.-C.; Lin, B.-S.; Chen, H.-C.; Liou, J.-C.; Li, Y.-T.; Peng, C.-W. Safety of Special Waveform of Transcranial Electrical Stimulation (TES): In Vivo Assessment. *Int. J. Mol. Sci.* **2022**, *23*, 6850. <https://doi.org/10.3390/ijms23126850>
204. Battaglia, S.; Schmidt, A.; Hassel, S.; Tanaka, M. Editorial: Case reports in neuroimaging and stimulation. *Front. Psychiatry* **2023**, *14*, 1264669. <https://doi.org/10.3389/fpsyg.2023.1264669>
205. Chang, C.H.; Wang, W.L.; Shieh, Y.H.; Peng, H.Y.; Ho, C.S.; Tsai, H.C. Case Report: Low-Frequency Repetitive Transcranial Magnetic Stimulation to Dorsolateral Prefrontal Cortex and Auditory Cortex in a Patient With Tinnitus and Depression. *Front. Psychiatry* **2022**, *13*, 847618. <https://doi.org/10.3389/fpsyg.2022.847618>
206. Zakia, H.; Iskandar, S. Case report: Depressive disorder with peripartum onset camouflages suspected intracranial tuberculoma. *Front. Psychiatry* **2022**, *13*, 932635. <https://doi.org/10.3389/fpsyg.2022.932635>
207. Nyatega, C.O.; Qiang, L.; Adamu, M.J.; Kawaawa, H.B. Gray matter, white matter and cerebrospinal fluid abnormalities in Parkinson's disease: A voxel-based morphometry study. *Front Psychiatry* **2022**, *13*, 1027907. <https://doi.org/10.3389/fpsyg.2022.1027907>
208. Rymaszewska, J.; Wieczorek, T.; Fila-Witecka, K.; Smarzewska, K.; Weiser, A.; Piotrowski, P.; Tabakow, P. Various neuromodulation methods including Deep Brain Stimulation of the medial forebrain bundle combined with psychopharmacotherapy of treatment-resistant depression-Case report. *Front. Psychiatry* **2023**, *13*, 1068054. <https://doi.org/10.3389/fpsyg.2022.1068054>
209. Liu, M.; Xie, X.; Xie, J.; Tian, S.; Du, X.; Feng, H.; Zhang, H. Early-onset Alzheimer's disease with depression as the first symptom: a case report with literature review. *Front. Psychiatry* **2023**, *14*, 1192562. <https://doi.org/10.3389/fpsyg.2023.1192562>
210. Kim, B.H.; Kim, S.H.; Han, C.; Jeong, H.G.; Lee, M.S.; Kim, J. Antidepressant-induced mania in panic disorder: a single-case study of clinical and functional connectivity characteristics. *Front. Psychiatry* **2023**, *14*, 1205126. <https://doi.org/10.3389/fpsyg.2023.1205126>
211. Zhou, J.; Cao, Y.; Deng, G.; Fang, J.; Qiu, C. Transient splenial lesion syndrome in bipolar-II disorder: a case report highlighting reversible brain changes during hypomanic episodes. *Front. Psychiatry* **2023**, *14*, 1219592. <https://doi.org/10.3389/fpsyg.2023.1219592>
212. Statsenko, Y.; Habuza, T.; Smetanina, D.; Simiyu, G.L.; Meribout, S.; King, F.C.; Gelovani, J.G.; Das, K.M.; Gorkom, K.N.-V.; Zaręba, K.; et al. Unraveling Lifelong Brain Morphometric Dynamics: A Protocol for Systematic Review and Meta-Analysis in Healthy Neurodevelopment and Ageing. *Biomedicines* **2023**, *11*, 1999. <https://doi.org/10.3390/biomedicines11071999>
213. Gazerani, P. Human Brain Organoids in Migraine Research: Pathogenesis and Drug Development. *Int. J. Mol. Sci.* **2023**, *24*, 3113. <https://doi.org/10.3390/ijms24043113>
214. Jalink, P.; Caiazzo, M. Brain Organoids: Filling the Need for a Human Model of Neurological Disorder. *Biology* **2021**, *10*, 740. <https://doi.org/10.3390/biology10080740>
215. Zhang, D.Y.; Song, H.; Ming, G.L. Modeling neurological disorders using brain organoids. *Semin. Cell Dev. Biol.* **2021**, *111*, 4–14. <https://doi.org/10.1016/j.semcd.2020.05.026>