

Article

Not peer-reviewed version

---

# Optimization of Live Prey Enrichment Media for Rearing Juvenile Short-Snouted Seahorse, *Hippocampus hippocampus*

---

[Jorge Palma](#)\*, Ricardo Lima, [José Pedro Andrade](#), [Maria João Lança](#)

Posted Date: 7 September 2023

doi: 10.20944/preprints202309.0490.v1

Keywords: nutrition; *Hippocampus hippocampus*; live feed enrichment; microalgae; essential fatty acids; copepods



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Article

# Optimization of Live Prey Enrichment Media for Rearing Juvenile Short-Snouted Seahorse, *Hippocampus hippocampus*

Jorge Palma<sup>1,\*</sup>, Ricardo Lima<sup>2</sup>, José Pedro Andrade<sup>1</sup> and Maria João Lança<sup>2</sup>

<sup>1</sup> Centre of Marine Sciences (CCMAR), University of the Algarve, Faro, Portugal; jpalma@ualg.pt (J.P.); jandrade@ualg.pt (J.P.A.)

<sup>2</sup> MED – Mediterranean Institute for Agriculture, Environment and Development / CHANGE – Global Change and Sustainability Institute, Universidade de Évora, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal & Departamento de Zootecnia, Escola de Ciências e Tecnologia, Universidade de Évora, Évora, Portugal; ricardomduartelima@gmail.com (R.L.); mjlanca@uevora.pt (M.J.L.)

\* Correspondence: jpalma@ualg.pt (J.P.)

**Abstract:** This study aimed to determine a nutritionally adequate feeding protocol for *Hippocampus hippocampus* juveniles. In the experimental trial, seahorses were fed copepods from 0-7 days post-parturition (DPP) and, from 8-28 DPP, four dietary treatments: (copepods (control diet), microalgae-enriched *Artemia* with a DHA/EPA ratio of 2:1, microalgae-enriched *Artemia* with a DHA/EPA ratio of 2:1 along with 5% copepods, and with 10% copepods) were tested. Results showed that juvenile seahorses fed on copepods attained a significantly higher ( $P < 0.05$ ) growth performance ( $5.1 \text{ mg d}^{-1}$ ) than fish-fed diets with 5% and 10% copepod inclusion, ( $3.09$  and  $3.07 \text{ mg d}^{-1}$ , respectively), and fish-fed enriched *Artemia* ( $1.83 \text{ mg d}^{-1}$ ) which performed poorly. Data suggests that feeding copepods during the first 7 DPP promotes maturation of the digestive tract of juvenile seahorses, and the addition of a limited amount of copepods to the diet improves *H. hippocampus* juvenile growth performance when compared to the use of *Artemia* as a single diet due to the improvement of the essential fatty acid profile in the diets.

**Keywords:** nutrition; *Hippocampus hippocampus*; live feed enrichment; microalgae; essential fatty acids; copepods

**Key Contribution:** The growth and survival of juvenile *H. hippocampus* is significantly enhanced when the initial diet consists exclusively of copepods; allowing for the maturation of the digestive tract. Later, even a small inclusion of 5% to 10% copepods in the dietary protocol contributes to an improved fatty acid profile due to copepods' high EFA content and translates into substantial benefits in terms of growth and survival.

## 1. Introduction

Due to their unique characteristics and morphological peculiarities, seahorses (*Hippocampus* sp.) have captured the interest of the community, thereby enhancing their cultural, scientific, educational, and economic value. As a result of this growing interest, seahorse exploitation, both for traditional medicines and the ornamental fish trade, has increased. This has led to a decline in natural populations [1] and an amplified focus on their cultivation worldwide [2].

Presently, seahorse species encounter conservation challenges, with habitat degradation and overfishing being the two primary causes of significant declines in most wild populations over the past decades. These issues have resulted in these species becoming an integrant part of the IUCN Red List and Appendix II of CITES. Therefore, seahorse aquaculture holds significant potential as a conservation tool for these species, but also as a growing business, owing to the high demand for these species and their substantial market values. Presently, it is known that at least 13 species of seahorses are commercially produced [2] at different commercial scales.

Currently, there are 42 identified seahorse species, including *Hippocampus hippocampus*, which is found in the northeastern Atlantic and Mediterranean Sea [3]. Although there is insufficient

information to assess the conservation status of this species, natural populations in Portugal have exhibited a declining trend. Additionally, within the Mediterranean Sea, the species is already considered as Near Threatened [4]. This situation has promoted research into various wild populations, often with a focus on their long-term conservation, as well as increased investigation into their captive production. Therefore, this remains a recent topic, and numerous gaps in knowledge regarding seahorse aquaculture, including *H. hippocampus*, persist [2]. To prevent, or at the very least, mitigate the problems stemming from the illegal capture of seahorses, aquaculture can provide animals for international trade, thereby promoting the recovery of wild stocks and ensuring the conservation of natural populations [5].

It is widely acknowledged that captive breeding contributes significantly to preventing species extinction [6], and in this regard, seahorse captive breeding also assumes a preventive role in terms of species extinction [7]. Given the limited current understanding of the dietary requirements of *H. hippocampus*, the utilization and development of diets that align with the nutritional needs of the juveniles hold paramount importance. *Artemia* and rotifers have historically been utilized as live prey for feeding seahorses [7]. However, both *Artemia* nauplii and rotifers have consistently demonstrated nutritional deficiencies as food sources for larval and juvenile marine fishes, largely due to their inadequate fatty acid profiles and low levels of free amino acids [8]. Through enrichment, these live prey organisms can supply essential nutrients to the target species, with *Artemia* often serving as a carrier of essential fatty acids (EFA) [9–13]. Feeding on *Artemia* or copepods has resulted in high rates of survival and growth during the rearing of multiple seahorse species [5,14–22].

Numerous studies have also confirmed the significance of fatty acids in seahorse growth by enhancing *Artemia* with various emulsions e.g., [23–25]. This fact is related with the basis of the trophic chains because marine and freshwater trophic chains are distinct in terms of fatty acid profile. Producers of marine food webs are unicellular algae rich in polyunsaturated fatty acids (PUFAs), mainly from the  $\omega 3$  family, while freshwater food chain producers have a higher content of PUFAs from the  $\omega 6$  family [26].

Because marine fish acquire substantial PUFA levels directly from their diet, their capacity to elongate and desaturate PUFAs has diminished, owing to a reduction in the associated enzymatic processes [27]. Considering this aspect, it can be asserted that the majority of marine species exhibit limited, inadequate, or in some instances, even non-existent abilities to elongate and desaturate linoleic acid (LA; C18:2 $\omega 6$ ) and  $\alpha$ -linolenic acid (ALA; C18:3 $\omega 3$ ) is low, insufficient or, in some cases, even null as they receive high amounts of these fatty acids directly from the diet [28,29]. In contrast, freshwater fish have higher levels of PUFAs with 18-carbon chains (i.e., AL, LA, and ALA) and possess substantial quantities of AA (C20:4  $\omega 6$ ), EPA (C20:5 $\omega 3$ ), and DHA (C22:6 $\omega 3$ ) resulting from metabolism [27,29,30].

When comparing the lipid composition of eggs between the vast majority of marine fish species and seahorses, it becomes evident that the former is characterized by having eggs with high concentrations of palmitic acid (PA, C16:0) (20.0%), docosahexaenoic acid (DHA: 17.4%), and eicosapentaenoic acid (EPA: 13.2%). Additionally, they contain, to a lesser extent, oleic acid (C18:1 $\omega 9$ ), stearic acid (C18:0), palmitoleic acid (C16:1 $\omega 7$ ), and vaccenic acid (C18:1 $\omega 7$ ) [29].

However, as mentioned earlier, seahorse juveniles have a low content of essential fatty acids (EFAs), namely EPA and DHA [31], which are typically associated with their survival and growth performance [32,33]. As marine fish, seahorses have high requirements for fatty acids from the  $\omega 3$  family, which must be supplied through their diets [23,34–36]. Furthermore, the prey sizes may also restrict the ingesting capacity of seahorses [37]. In their natural habitat, copepods are the preferred prey for most marine fish larvae [38], including seahorses. Multiple studies have indicated that the preference for copepods among marine fish larvae is attributed to their nutritional composition [39,40] and the provision of numerous enzymes that aid in the digestive process [41]. Consequently, in the realm of fish larval culturing, copepods are generally considered superior to rotifers and *Artemia* [42]. However, the utilization of copepods as larval fish feed is hindered by their limited availability and the challenges associated with intensive production [43].

Hence, as copepods are a more nutritionally suitable prey, *Artemia* cultivation is a more feasible option and, through enrichment, can serve as a "vehicle" to fulfil the nutritional requirements of seahorses through a process known as "bio-encapsulation". Conversely, advancements have been made in marine fish incubation techniques over recent years. Nonetheless, these improvements have

not fully addressed the mortality issues observed during the early developmental stages of certain marine juvenile fish species [7,44,45], which are often linked to feeding, feeding behavior, and periods of food scarcity [46–48].

Considering the above, the objective of this study was to address some of the existing gaps in understanding the dietary requirements of the short-snout seahorse, *H. hippocampus*, during its early life stages, through the utilization of different live feed (*Artemia*) dietary enrichments and the implementation of an appropriate copepod co-feeding protocol.

## 2. Materials and Methods

A total of 420 *H. hippocampus* juveniles obtained from a captive-bred broodstock kept at the Ramalhete research station, CCMAR – Universidade do Algarve, Portugal were used to conduct two different rearing trials. The broodstock fish were kept in one 250 L plastic tank assembled in a semi-open flow system with a water turnover of approximately 125 L/h per tank. Holdfast units were added to provide support for seahorses [49]. Individuals were daily fed a mix of live mysid shrimp (*Mesopodopsis slabberi* and *Leptomysis* sp.) in different proportions, depending on availability. After natural spawning, juvenile seahorses from a single brood were randomly collected and used in the trials just after release from the male's pouch. Juvenile seahorses were gently transferred to the rearing tanks using a 250 ml beaker and with utmost caution to avoid exposure to air.

In this study, two rearing trials were conducted. The first one, referred to as preliminary trial, was designed to test an enrichment protocol for seahorse's feeding regime. The second, the experimental trial, was developed using the information obtained in the preliminary trial, to create an optimized feeding protocol that will be later described. In both trials, the rearing tanks (10 L glass rectangular tanks) were supplied with water at a stable flow rate within a semi-closed recirculation system. Prior to being introduced into the rearing tanks, water was circulated from the reservoir tank through a UV sterilization unit. To ensure that dissolved oxygen was always kept close to saturation, water was vigorously aerated in a filtration reservoir (sump). To prevent bubbling, rearing tanks water inflow was made through a transparent plastic tube ending below the water surface. The water outflow structure was assembled in the corner of the tanks. It consisted of a black polystyrene tube covered at the water surface with a 150 µm diameter mesh to prevent live feed from being flushed from the tanks. To improve prey detection, lateral and back tank walls were covered with a black adhesive like [50], whilst the front wall remained uncovered for observation. Each of the rearing tanks had two artificial holdfasts adjusted to juvenile size. For illumination, 2 x 36 W fluorescent tubes were placed above the tanks. The light intensity at the water surface was around  $900 \pm 40$  lux, and a photoperiod of 16L: 8D (06:00 – 22:00 h), controlled by a timer. Water temperature and chemical parameters pH, salinity, and dissolved oxygen were recorded daily and maintained at  $20.4 \pm 0.3$  °C, 7.8,  $37.5 \pm 0.1$  and  $7.6 \pm 0.1$  mg/L, respectively. Seawater quality parameters were recorded biweekly, values of ammonia were always below detectable levels, values of nitrate < 0.3 mg/L and values of nitrite < 1.25 mg/L. The tanks were daily cleaned by siphoning to remove faeces, uneaten feed, and dead juvenile seahorses (accounted for mortality).

### 2.1. Live Feed Culture and Enrichment

AF *Artemia* cysts (Sanders®, Ogden, UT, USA) were hatched according to the procedures described by [51]. Phytobloom Prof *Isochrysis galbana* and Phytobloom Prof *Nannochloropsis* spp. pure powders from Necton's Phytobloom Green Formula (Necton, S.A, Olhão, PT) were used as enrichment according to the manufacturer specifications. In brief, each microalgae quantity was weighted individually, added to 250 ml of seawater, mixed and left to hydrate for 15 minutes. The mix was then poured into a blender and homogenized for 2 min. The *Artemia* (approx. 48 000 nauplii) were added into the enrichment cup in a volume not higher than 750 ml and the enrichment media was added. This procedure was repeated for each dietary treatment. *Artemia* nauplii were left to enrich for 24h at room temperature (20–22 °C), under continuous moderate aeration and illumination.

### 2.2. Copepod Harvesting

Copepods (*Acartia clausi*) were naturally produced in the outflow pond of the Ramalhete research station and collected daily using a 60 µm hand net. After acclimation to room temperature,



copepods were strained through a 150 µm sieve to remove large plankters and debris and were counted to the adequate feed ratio.

### 2.3. Feeding Trials

In the preliminary trial, the single use of enriched *Artemia* was tested. For this, three different enrichment media with different DHA/EPA ratios were tested using two microalgae in different proportions: *I. galbana*, rich in DHA, and *Nannochloropsis* sp., rich in EPA. The calculations of the DHA and EPA ratios were derived from the fatty acid (FA) profile provided by the manufacturers. A total of 180 juvenile *H. hippocampus* from the same brood were distributed across nine rearing tanks using a completely randomized design, with three replicate tanks allocated to each dietary treatment. 20 juvenile *H. hippocampus* were stocked in each replicate tank. The first dietary treatment (*Art2:1*), contained *Artemia* metanauplii enriched with 2:1 DHA/EPA ratio; the second (*Art1:1*), *Artemia* metanauplii enriched with 1:1 DHA/EPA ratio; and the third (*Art1:2*), *Artemia* metanauplii enriched with 1:2 DHA/EPA ratio. The experiment was designed to last 28 days.

In the experimental trial, based on the results of the preliminary trial, the minimum copepod (*A. clausi*) addition to sustain optimal growth and survival of juvenile *H. hippocampus* was evaluated and compared to copepod and *Artemia* diets. 240 juvenile *H. hippocampus* from a same brood were assembled into twelve rearing tanks according to a completely randomised design, with three replicate tanks assigned to each dietary treatment. 20 juvenile *H. hippocampus* were stocked in each replicate tanks at a density of 2 fish l<sup>-1</sup>. All groups were first fed natural copepods from 0 – 7 days post-parturition (DPP), and only on the 8th day the transition to each of the following dietary treatments was made. The first dietary treatment (*Cop*), consisted in daily captured copepods (control diet); the second dietary treatment (*Art<sub>DHA/EPA</sub>*), of 2:1 DHA/EPA ratio enriched *Artemia* metanauplii used in the previous trial (dietary treatment *Art2:1*); the third (*Art<sub>DHA/EPA5%</sub>*), a mixture of 95% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 5% natural copepods; and the fourth dietary treatment (*Art<sub>DHA/EPA10%</sub>*), a mixture of 90% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 10% natural copepods. The experiment lasted for 28 days.

### 2.4. Sampling, Data Collection and Statistical Analysis

To obtain the accurate weight and length information at 0 DPP, due to the small size and to avoid morphological distortions, ten juveniles were immediately and rapidly sacrificed with an excess of anaesthetic (2-phenoxyethanol solution, 0.40 mg L<sup>-1</sup>). Individuals were placed in the anaesthetic solution for at least 20 minutes and removed no less than 10 minutes after ventilation stopped. Fish were then weighted using a high precision scale and measurements (total length as the sum of the head, trunk, and tail length) were done using a binocular microscope associated with an image analysis software (DinoCapture 2.0 (AnMo Electronics Corporation, Sanchong, New Taipei City, Taiwan)). Later, experimental fish were sampled at 14 and 28 DPP for growth performance analysis. To determine the total length, the sum of the head length and the height of the fish were considered. Head length was measured as the distance from the rostral zone of the snout (maxillary) to the midpoint of the cleithral ring. The height of the seahorse was measured by the length between the cleithral ring to the end of the tail, considering it in extension (Annex I), as suggested by [52]. This procedure was related to the adoption of a methodology that would inflict the least possible stress on seahorses, which is more advantageous than the traditional three-measurement method proposed by [53]. Mortality was monitored daily.

Seahorse's wet weight and total length, as well as mortality, were used to calculate: Survival ( $S$ ) =  $((N_i - M_{ij}) / N_i) \times 100$ , where  $N_i$  is the initial number of seahorses placed on each tank on day 1 and  $M_{ij}$  the total number of surviving individuals at the end of the trial; Condition Factor ( $CF$ ) =  $(WW / L^3) \times 100$ , where  $WW$  is the wet weight (g) and  $L$  the total length (cm); Mean Length Gain ( $LG$ ) (cm/fish) =  $(L_f / L_i) / t$ , where  $L_f$  is the final length (cm),  $L_i$  is the initial length (cm) and  $t$  is the number of experimental days; Mean Weight Gain ( $WG$ ) (mg/fish) =  $(W_f - W_i) / t$ , where  $W_f$  is the final wet weight (mg),  $W_i$  is the initial wet weight (mg) and  $t$  is the number of experimental days and the Thermal-unit Growth Coefficient ( $TGC$ ) (modified from [54] by [55]) =  $[(W_f^{1/3} - W_i^{1/3}) / \Sigma(T \times D)] \times 100$ , where  $W_f$  is the final seahorse wet weight (g),  $W_i$  is the initial wet weight (g),  $T$  is the water temperature (°C), and  $D$  is the number of days in each trial. Data processing and statistical analysis were conducted using the statistical analysis program package GraphPad Prism (version 8.4.2 for Windows,

GraphPad Software, San Diego, CA, USA). All data were tested for normality and homogeneity of variances. To study the effect of the diets on growth performance (Weight gain, Length gain, Thermal-unit Coefficient, Condition Factor, and Accumulative survival) between groups, a One-Way ANOVA, using Tukey's test for multiple comparisons was used.

### 2.5. Dietary Treatments Proximate Analysis and Fatty Acid Composition

To analyse the nutritional quality of prey and the dietary treatments used in the experimental trial, samples of natural copepods (*Cop*), unenriched *Artemia* (*Art<sub>Ref</sub>*) 24h enriched *Artemia* plus 5% copepods (*Art<sub>DHA/EPA5%</sub>*) and enriched *Artemia* plus 10% copepods (*Art<sub>DHA/EPA10%</sub>*) were collected. Even though unenriched *Artemia* (*Art<sub>Ref</sub>*) was not a dietary treatment tested in any of the groups, it was analysed along with all the regimes for comparison purposes and to determine the efficiency of the enrichment used and copepod addition.

The four dietary treatments and the unenriched *Artemia* were analyzed following the procedures outlined in [56]. In brief, the dry matter was analyzed by drying samples at 105 °C to constant weight; ash by incineration of samples at 550 °C for 5 h; crude protein ( $N \times 6.25$ ) using a Leco nitrogen analyzer (Leco Corporation, St. Joseph, USA); total lipid by extraction according to the method of [57], using accelerated solvent extraction (ASE) and lyophilized samples.; crude fiber by acid and basic digestion (Fibertec System M., 1020 Hot Extractor, Tecator) and gross energy content using an automatic oxygen bomb calorimeter (Parr 6400, U.S.A.) according to ISO 9831.

For the fatty acid analysis, to achieve the extract derivatization, total lipids were transferred to methylation tubes using 1.5 mL of  $CHCl_3/MeOH$  (3:2) solution and placed under a nitrogen stream to evaporate the transfer solvent and dry the extract for saponification and methylation. For methylation phase, the samples were cooled for approx. 1 min and 1 mL of  $BF_3$ -methanol was added, the tubes were then heated at 70°C for 10 min. Once methylation was complete, the solution was cooled to room temperature, 2 mL of distilled water and 2 mL of petroleum ether were added and stirred vigorously. The tubes were then placed in the fridge for 12 h, at 2°C. After separation of the phases was verified, each sample's supernatant (1.5 mL) was collected into 2 mL vials, which were hermetically sealed, labelled, and laced in the fridge until further use. Chromatographic analysis of methyl esters was performed as described by [58]. In brief, the analysis of the peaks and their respective MS was obtained by electronic impact at 70 eV, with a sweep of  $m/z = 40$  to 450 and analyzed using MSWS 8.2 software. The Fame standards used were 37-component FAME MIX and BAME MIX 26 (Supelco). Fatty acids were designated according to the nomenclature of the International Union of Pure and Applied Chemistry (IUPAC) for carbon chain length: number of double bonds and position of the double bond closest to the omega carbon.

### 2.6. Ethical Statement

CCMAR facilities and the research is certified to house and conduct experiments with live animals (Group-C licenses by the Direção Geral de Alimentação e Veterinária, Ministério da Agricultura, Florestas e Desenvolvimento Rural, Portugal). The experimental design of the present study was part of the Projects HIPPONUTRE (reference 1602-01-FMP-54) and HIPPOSAVE (reference MAR-01.04.02-FEAMP-0029), which obtained approval from the ethics committee of the Veterinary Medicines Directorate, Ministry of Agriculture, Rural Development and Fisheries, Portugal. The study adhered to the guidelines outlined by the European Union Council (86/609/EU) and the relevant Portuguese legislation concerning the use of laboratory animals.

## 3. Results

The proximate composition of each dietary treatment is presented in Table 1. The dry matter of *Cop* dietary treatment was considerably lower (13.1%) compared to the remaining diets (ranged between 29.9 and 33.2%). As for the gross protein, the *Art<sub>DHA/EPA10%</sub>* had the highest content (20 g/100g DM), followed by *Art<sub>DHA/EPA5%</sub>* (19.1 g/100g DM), *Art<sub>DHA/EPA</sub>* (17.9 g/100g DM), *Art<sub>Ref</sub>* (16.5g/100g DM) and *Cop* (10.9 g/100g DM). Concerning the total lipids content, *Cop* diet had the lowest value of total lipids, (4.9 g/100g DM), a value twofold lower than all other dietary treatments. Conversely, *Cop* diet had the higher gross energy content (503.9 Kcal/100g DM), followed by *Art<sub>DHA/EPA5%</sub>* (495.62 Kcal/100g DM), *Art<sub>DHA/EPA10%</sub>* (492.18 Kcal/100g DM), and *Art<sub>DHA/EPA</sub>* (489.84 Kcal/100g DM) (Table 1).

**Table 1.** Proximate composition of the four tested dietary treatments used (per 100g of Dry Matter (g/100g DM)). \* Although ArtRef was not directly tested is presented for comparison purposes, as it is the basis for the dietary treatments with enriched *Artemia* used. Cop - Natural copepods; ArtRef - Unenriched *Artemia* nauplii; ArtDHA/EPA - 2:1 DHA/EPA ratio enriched *Artemia* metanauplii; ArtDHA/EPA5% - 95% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 5% natural copepods; ArtDHA/EPA10% - 90% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 10% natural copepods.

Proximate Composition	Cop	Art <sub>Ref</sub> *	Art <sub>DHA/EPA</sub>	Art <sub>DHA/EPA</sub> 5%	Art <sub>DHA/EPA</sub> 10%
Moisture (%)	86,9	70,09	70,12	68,47	66,82
Dry Matter (%)	13,1	29,91	29,88	31,53	33,18
Gross Protein (g/100g DM)	10,91	16,5	17,93	19,1	20,04
Total Lipids (g/100g DM)	4,87	11,8	10,24	9,96	10,54
Gross Energy (Kcal/100g DM)	503,85	497,77	489,84	495,62	492,18

Regarding the presence of EFAs in the two microalgae used, *I. galbana* had the higher percentages of LA (C18:2 $\omega$ 6) (7.5%), ALA (C18:3 $\omega$ 3) (15.5%) and DHA (C22:6 $\omega$ 3) (16.5%), than *Nannochloropsis* sp. Conversely, although *Nannochloropsis* sp. did not contain any DHA in its profile, it had a higher concentration of ARA (C20:4 $\omega$ 6) (5.3%) and a substantially more EPA (C20:5 $\omega$ 3) (36.1%) (Table 2). Concerning fatty acid classes (SFA, MUFA, PUFA and HUFA) and families ( $\omega$ 3,  $\omega$ 6 and  $\omega$ 9), the biggest differences were in the percentage of HUFAs, EFAs and  $\omega$ 9 fatty acids. *Nannochloropsis* sp. presented higher amounts of HUFAs (42.3%) and EFAs (45.6%), while *I. galbana* showed a greater percentage in  $\omega$ 9 fatty acids, with a value of 15.2%. As for the ratios, *I. galbana* shows higher values for all the ratios presented, especially the very high DHA/EPA ratio (165) whereas *Nannochloropsis* sp. has a value of 0 for the same ratio (Table 2).

**Table 2.** Fatty acid profile of the two microalgae used as *Artemia* nauplii enrichment, in the diets, Art<sub>DHA/EPA</sub>, Art<sub>DHA/EPA</sub>5%, and Art<sub>DHA/EPA</sub>10%.


Fatty Acid	<i>Isochrysis galbana</i>	<i>Nannochloropsis</i> sp.
C18:2 $\omega$ 6 (LA)	7,5	4
C18:3 $\omega$ 3 (ALA)	15,5	0,2
C20:4 $\omega$ 6 (ARA)	0,2	5,3
C20:5 $\omega$ 3 (EPA)	0,1	36,1
C22:6 $\omega$ 3 (DHA)	16,5	-
$\Sigma$ SFA	25,5	26,2
$\Sigma$ MUFA	20,2	22,9
$\Sigma$ PUFA	50,1	48,1
$\Sigma$ HUFA	27,3	42,3
$\Sigma$ EFA	39,8	45,6
$\Sigma\omega$ 3	39,8	37,2
$\Sigma\omega$ 6	10,6	10,9
$\Sigma\omega$ 9	15,2	3,2
DHA/EPA	165	0
$\omega$ 3/ $\omega$ 6	3,75	3,41
PUFA/SFA	1,96	1,84

ALA - Alpha-Linolenic acid; ARA - Arachidonic acid; DHA - Docosahexaenoic acid; EFA - Essential Fatty Acid; EPA - Eicosapentaenoic acid; HUFA - Highly Unsaturated Fatty Acid; LA - Linoleic acid; MUFA - Monounsaturated Fatty Acid; PUFA - Polyunsaturated Fatty Acid; SFA - Saturated Fatty Acid.

The lipid profile of the control diet (Cop) and tested dietary treatments (Cop, Art<sub>DHA/EPA</sub>, Art<sub>DHA/EPA</sub>5% and Art<sub>DHA/EPA</sub>10%), and the additional Art<sub>Ref</sub>, for the most expressive fatty acids and respective sums of SFA, MUFA, PUFA, HUFA and EFA are detailed in Table 3. The Cop diet contained high values of all fatty acid classes, but MUFA. Cop and Art<sub>Ref</sub> diets also revealed the highest percentage values for PUFA and HUFA. Regarding PUFA, Cop diet showed a value of 31%, followed by Art<sub>Ref</sub> with 23.6%, and by the three diets including enriched *Artemia*. Dietary treatments with enriched *Artemia* had much higher values of MUFA ( $\approx$ 34%) and, consequently, lower PUFA and HUFA values ( $\approx$  20% and 10%, respectively).

The most relevant SFA were C12:0, C14:0, C16:0, C18:0, and C20:0. C16:0 was the SFA with the highest value in all tested diets, with Cop diet having 31.2%, followed by Art<sub>DHA/EPA</sub>10% (17.1%), Art<sub>DHA/EPA</sub>5% (16.9%), Art<sub>DHA/EPA</sub> (16%) diets (Table 3). The second most expressive SFA was C14:0, as for

C18:0, there were no major differences between dietary treatments. Amongst the PUFA, all EFAs (LA, ARA, ALA, EPA, and DHA) were present in the tested dietary treatments (Figure 1).



**Figure 1.** Essential fatty acids (EFA) present in each tested dietary treatment. **Cop** - Natural copepods; **Art<sub>Ref</sub>** - Unenriched *Artemia* nauplii; **Art<sub>DHA/EPA</sub>5%** - 95% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 5% natural copepods; **Art<sub>DHA/EPA</sub>10%** - 90% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 10% natural copepods. **LA** - Linoleic acid; **ALA** - Alpha-Linolenic acid; **ARA** - Arachidonic; **EPA** - Eicosapentaenoic acid; **DHA** - Docosahexaenoic acid.

Regarding LA and ARA, all diets containing *Artemia* (**Art<sub>DHA/EPA</sub>**, **Art<sub>DHA/EPA</sub>5%** and **Art<sub>DHA/EPA</sub>10%**) had about twice the amount of these fatty acids compared to the **Cop** control diet (Table 3). No relevant differences in the ALA amount ( $\approx 3\%$ ) were found among dietary treatments. Regarding EPA, **Cop** control diet had the highest value (10.48%), followed by **Art<sub>Ref</sub>**, and 8.26%, while the remaining samples have very similar amounts ( $\approx 6\%$ ). The **Cop** dietary treatment contained a high percentage of DHA (11.7%), compared to the remaining diets, with concentrations ranging between 0.2% and 0.6%. It is however noteworthy, that the low DHA concentrations in the **Art<sub>DHA/EPA</sub>5%** and **Art<sub>DHA/EPA</sub>10%** were present on the expenses of the 5 and 10% copepod inclusion, as in the **Art<sub>Ref</sub>** the DHA concentration was 0% (Table 3).


Figure 2. shows the distribution of fatty acid families by dietary treatment. The **Cop** control diet had the highest value of  $\omega 3$  fatty acids (27%), followed by the **Art<sub>Ref</sub>** (12.3%), **Art<sub>DHA/EPA</sub>5%** (10.5%), **Art<sub>DHA/EPA</sub>10%** (10.4%) and the **Art<sub>DHA/EPA</sub>** (10.2%). Concerning the  $\omega 6$ ,  $\omega 7$  and  $\omega 9$  fatty acid families, all diets but **Cop** revealed higher percentages. The diets with the highest  $\omega 6$  fatty acids were **Art<sub>DHA/EPA</sub>10%** (9.4%), **Art<sub>DHA/EPA</sub>** (7.9%) and **Art<sub>DHA/EPA</sub>5%** (7.9%). Regarding the  $\omega 7$  fatty acids, all regimes with enriched *Artemia* revealed to have high concentrations with similar values (approximately 22%) and **Cop** diet with half the concentration of the remaining diets (9.4%). Finally, for  $\omega 9$  fatty acids, it was observed that the **Art<sub>DHA/EPA</sub>** regime presented the highest value (20.7%), followed by **Art<sub>DHA/EPA</sub>5%** (19.9%) and **Art<sub>DHA/EPA</sub>10%** (19.2%). **Cop** diet presented the lowest value of  $\omega 9$  fatty acids, with 4.8%.

Some relevant ratios in terms of nutritional assessment of the tested diets are presented in Table 4. DHA/EPA and  $\omega 3/\omega 6$  ratios in **Cop** diet revealed the highest values (1.11 and 6, respectively). As for the DHA/EPA ratio in the **Art<sub>Ref</sub>** it was 0. Regarding the PUFA/SFA ratio, **Art<sub>Ref</sub>** presented the higher value (0.99), followed by **Art<sub>DHA/EPA</sub>10%** (0.78), **Art<sub>DHA/EPA</sub>** (0.75), **Art<sub>DHA/EPA</sub>5%** (0.74) and **Cop** (0.65) dietary treatments.



**Table 3.** Fatty acid profile of each dietary treatment (mean  $\pm$  standard deviation) expressed in percentage (%) relative to the total of fatty acids identified in the samples.

Fatty Acid	Cop	Art <sub>Ref</sub> *	Art <sub>DHA/EPA</sub>	Art <sub>DHA/EPA</sub> 5%	Art <sub>DHA/EPA</sub> 10%
C12:0	0,19 $\pm$ 0,01	0,16 $\pm$ 0,02	0,15 $\pm$ 0,02	0,18 $\pm$ 0,01	0,23 $\pm$ 0,07
C14:0	10,25 $\pm$ 0,19	3,06 $\pm$ 0,19	4,26 $\pm$ 0,16	4,25 $\pm$ 0,20	4,24 $\pm$ 0,15
C16:0	31,23 $\pm$ 0,40	15,69 $\pm$ 0,41	15,94 $\pm$ 0,88	16,91 $\pm$ 1,15	17,07 $\pm$ 0,47
C18:0	6,13 $\pm$ 0,28	4,84 $\pm$ 0,30	5,35 $\pm$ 0,20	5,19 $\pm$ 0,14	5,07 $\pm$ 0,29
C20:0	0,11 $\pm$ 0,00	0,11 $\pm$ 0,03	0,14 $\pm$ 0,00	0,14 $\pm$ 0,02	0,09 $\pm$ 0,03
$\Sigma$ SFA	47,90	23,87	25,84	26,65	26,70
C14:1 $\omega$ 5	-	0,33 $\pm$ 0,02	0,23 $\pm$ 0,01	0,21 $\pm$ 0,00	0,23 $\pm$ 0,02
C16:1 $\omega$ 7c	8,88 $\pm$ 0,20	13,79 $\pm$ 0,67	12,74 $\pm$ 0,46	13,06 $\pm$ 0,61	13,58 $\pm$ 0,47
C16:1 $\omega$ 7t	0,27 $\pm$ 0,07	0,47 $\pm$ 0,09	0,40 $\pm$ 0,04	0,36 $\pm$ 0,02	0,39 $\pm$ 0,02
C18:1 $\omega$ 9c	2,91 $\pm$ 0,04	6,07 $\pm$ 6,64	12,36 $\pm$ 0,92	12,21 $\pm$ 1,12	11,82 $\pm$ 1,64
C18:1 $\omega$ 9t	1,90 $\pm$ 0,09	8,08 $\pm$ 1,80	8,30 $\pm$ 1,30	7,73 $\pm$ 1,44	7,43 $\pm$ 1,29
C18:1 $\omega$ 7	0,29 $\pm$ 0,42	0,12 $\pm$ 0,06	0,10 $\pm$ 0,01	0,09 $\pm$ 0,07	0,07 $\pm$ 0,02
C20:1 $\omega$ 9c	-	0,34 $\pm$ 0,08	0,33 $\pm$ 0,03	0,33 $\pm$ 0,01	0,32 $\pm$ 0,02
$\Sigma$ MUFA	14,26	29,20	34,46	33,99	33,82
C16:2 $\omega$ 4	0,27 $\pm$ 0,13	1,04 $\pm$ 0,05	0,73 $\pm$ 0,45	0,71 $\pm$ 0,38	0,38 $\pm$ 0,22
C18:2 $\omega$ 6 (LA)	2,09 $\pm$ 0,04	5,05 $\pm$ 0,27	4,56 $\pm$ 0,02	4,49 $\pm$ 0,06	4,52 $\pm$ 0,13
C18:2 $\omega$ 4	-	0,72 $\pm$ 0,02	0,48 $\pm$ 0,03	0,46 $\pm$ 0,02	0,39 $\pm$ 0,08
C18:3 $\omega$ 6	0,34 $\pm$ 0,02	0,83 $\pm$ 0,13	0,63 $\pm$ 0,09	0,60 $\pm$ 0,06	1,56 $\pm$ 1,64
C18:3 $\omega$ 3 (ALA)	2,90 $\pm$ 0,02	3,23 $\pm$ 0,22	3,18 $\pm$ 0,10	3,07 $\pm$ 0,08	2,93 $\pm$ 0,18
C18:4 $\omega$ 3	1,50 $\pm$ 0,09	0,56 $\pm$ 0,04	1,14 $\pm$ 0,04	1,07 $\pm$ 0,03	0,85 $\pm$ 0,06
C20:3 $\omega$ 6	0,23 $\pm$ 0,02	0,24 $\pm$ 0,08	0,16 $\pm$ 0,05	0,17 $\pm$ 0,04	0,17 $\pm$ 0,08
C20:4 $\omega$ 6 (ARA)	1,28 $\pm$ 0,10	3,44 $\pm$ 0,17	2,55 $\pm$ 0,02	2,62 $\pm$ 0,01	3,14 $\pm$ 0,57
C20:4 $\omega$ 3	0,25 $\pm$ 0,08	0,24 $\pm$ 0,06	0,11 $\pm$ 0,05	0,14 $\pm$ 0,05	0,08 $\pm$ 0,04
C20:5 $\omega$ 3 (EPA)	10,48 $\pm$ 1,46	8,26 $\pm$ 0,30	5,58 $\pm$ 0,30	5,86 $\pm$ 0,46	5,94 $\pm$ 0,89
C22:6 $\omega$ 3 (DHA)	11,67 $\pm$ 0,79	-	0,23 $\pm$ 0,04	0,47 $\pm$ 0,09	0,58 $\pm$ 0,05
$\Sigma$ PUFA	31,01	23,60	19,35	19,63	20,78
$\Sigma$ HUFA	25,18	12,51	9,61	10,13	10,58
$\Sigma$ EFA	28,42	19,98	16,10	16,51	17,11
$\Sigma$ Others	6,83	23,33	20,35	19,73	18,70

**Figure 2.** - Characterization of the fatty acid families  $\omega$ 3,  $\omega$ 6,  $\omega$ 7 and  $\omega$ 9 in each of the four tested dietary treatments. **Cop** - Natural copepods; **Art<sub>Ref</sub>\*** - Unenriched *Artemia* nauplii; **Art<sub>DHA/EPA</sub>** - 95% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 5% natural copepods; - **Art<sub>DHA/EPA</sub>5%** - 90% 2:1 DHA/EPA ratio enriched *Artemia* metanauplii and 10% natural copepods.

**Table 4.** Main fatty acid ratios in each of the four tested dietary treatments.

Fatty Acid ratios	Cop	Art <sub>Ref</sub> <sup>a</sup>	Art <sub>DHA/EPA</sub>	Art <sub>DHA/EPA</sub> 5%	Art <sub>DHA/EPA</sub> 10%
DHA/EPA	1,11	0	0,04	0,08	0,1
ω3/ω6	6	1,29	1,29	1,34	1,1
PUFA/SFA	0,65	0,99	0,75	0,74	0,78

### 3.1. Growth Trials


In the preliminary trial, all dietary treatments (*Art*<sub>2/1</sub>, *Art*<sub>1/1</sub>, and *Art*<sub>1/2</sub>) resulted in 100% mortality by the 9th DPP, leading to the termination of the trial at that point. Regarding the subsequent experimental trial, comprehensive data regarding the survival and growth performance of juvenile *H. hippocampus* fed each of the four dietary treatments (*Cop*, *Art*<sub>DHA/EPA</sub>, *Art*<sub>DHA/EPA</sub>5%, and *Art*<sub>DHA/EPA</sub>10%) are presented in Table 5. It was observed that both the *Art*<sub>DHA/EPA</sub> and *Art*<sub>DHA/EPA</sub>5% groups exhibited similar survival rates, 88% and 87%, respectively. The *Art*<sub>DHA/EPA</sub>10% group followed with a survival rate of 77%, and the *Cop* group had the lowest survival rate at 72% (Table 5).

**Table 5.** - Survival and growth parameters of juvenile *Hippocampus hippocampus* fed each the four tested dietary treatments.

	<i>Cop</i>	<i>Art</i> <sub>DHA/EPA</sub>	<i>Art</i> <sub>DHA/EPA</sub> 5%	<i>Art</i> <sub>DHA/EPA</sub> 10%
Survival (%)	72	88	87	77
ML (cm) 0 DPP	1.1±0.1	1.1±0.1	1.1±0.1	1.1±0.1
ML (cm) 28 DPP	4.4±0.3 <sup>a</sup>	3.2±0.4 <sup>c</sup>	3.8±0.3 <sup>b</sup>	3.7±0.3 <sup>b</sup>
MW (mg) 0 DPP	2.4±0.4	2.4±0.5	2.4±0.6	2.4±0.7
MW (mg) 28 DPP	143.3±29.6 <sup>a</sup>	53.6±19.3 <sup>c</sup>	89±22.7 <sup>b</sup>	88.5±28.4 <sup>b</sup>
LG (cm/fish)	0.12 <sup>a</sup>	0.07 <sup>c</sup>	0.09 <sup>b</sup>	0.09 <sup>b</sup>
WG (mg/fish)	5.14 <sup>a</sup>	1.83 <sup>c</sup>	3.09 <sup>b</sup>	3.07 <sup>b</sup>
TGC	0.008 <sup>a</sup>	0.003 <sup>c</sup>	0.005 <sup>b</sup>	0.005 <sup>b</sup>
CF	0.17 <sup>a</sup>	0.16 <sup>a</sup>	0.17 <sup>a</sup>	0.17 <sup>a</sup>

### 3.2. Top of Form

At the end of the trial, juvenile *H. hippocampus* fed the *Cop* diet, grew significantly more ( $P < 0.05$ ) than all the remaining groups. *Art*<sub>DHA/EPA</sub>5% and *Art*<sub>DHA/EPA</sub>10% attained an intermediate growth performance, non-significantly different ( $P < 0.05$ ) among the two, and the group fed *Art*<sub>DHA/EPA</sub> grew significantly less ( $P < 0.05$ ) than all the remaining (Table 5 and Figure 3).

**Figure 3.** Mean weight (a) and mean length (b) gain of juvenile *Hippocampus hippocampus* fed each the four tested dietary treatments during the 28-day experiment.

This is corroborated by a *Cop* group WG of 5.1 mg d<sup>-1</sup>, and a final MW of 146.3 ± 29.6 mg at 28 DPP, the highest observed. The *Art*<sub>DHA/EPA</sub>5% and *Art*<sub>DHA/EPA</sub>10% groups showed identical WG (3.09 and 3.07 mg d<sup>-1</sup>, correspondingly) and a MW of 89 ± 22.7 mg and 88.5 ± 28.4 mg, respectively. The group fed with the *Art*<sub>DHA/EPA</sub> attained a WG of 1.8 mg d<sup>-1</sup> and a final MW of 53.6 ± 19.3 mg, showing the lowest growth performance.

Concordantly, the *Cop* group had the highest LG (0.12 cm d<sup>-1</sup>) averaging 4.4 ± 0.3 cm at the end of the experiment. *Art*<sub>DHA/EPA</sub>5% and *Art*<sub>DHA/EPA</sub>10% groups weren't significantly different ( $P > 0.05$ ), and

the LG was 0.09 cm d<sup>-1</sup>, resulting in 0.03 cm d<sup>-1</sup> less than the *Cop* group. The lowest LG and ML values were observed for the *Art<sub>DHA/EPA</sub>* regime group, with 0.07 cm d<sup>-1</sup>, and a ML of 3.2 ± 0.4 cm (Table 5).

Concerning the TGC, the highest value was observed in *Cop* group (0.008), followed by both *Art<sub>DHA/EPA5%</sub>* and *Art<sub>DHA/EPA10%</sub>* groups (0.005) and the *Art<sub>DHA/EPA</sub>* group (0.003) the lowest TGC value was observed. All groups attained similar, and non-significantly different ( $P > 0.05$ ) CF values at the end of the experiment (Table 5).

#### 4. Discussion

Juvenile seahorses require large amounts of good quality live feed [2] in which the fatty acids are a prevalent nutritional requirement. However, ensuring its appropriate provision remains one of the most significant challenges in successful seahorse mass farming. *Artemia* is a commonly used feed for seahorses due to its easy cultivation and commercial availability [14,16] but available *Artemia* enrichments are still inadequate for most seahorse species. *Artemia* nauplii has been identified as deficient in highly unsaturated fatty acids (HUFAs) [5,11], requiring HUFA enrichment media [14,25,59,60]. Conversely, copepods are regarded as an ideal diet for marine fish larvae, seahorse included, primarily due to their well-balanced fatty acid composition. However, copepods often cannot be cultured at high densities under controlled conditions as other live food options, because they need the use of larger volumes of water and larger culture vessels [61]. This, enriched *Artemia*-based diets cannot be ignored and can be used as an alternative or complement to an exclusive copepod-based diet if its enrichment is adequately tailored to the target species. Additionally, its utilization can also offer advantages by potentially reducing associated contaminants, as copepods frequently originate from wild populations and can introduce pathogens [62].

In the current study, all dietary treatments containing *Artemia* (*Art<sub>DHA/EPA</sub>*, *Art<sub>DHA/EPA5%</sub>*, and *Art<sub>DHA/EPA10%</sub>*) had higher levels of total protein and total lipids compared to the *Cop* control diet. This difference could be attributed to the greater moisture content in the *Cop* diet (86.9%) compared to that of the *Art<sub>Ref</sub>* (70.1%). However, it was the *Cop* diet that yielded a higher gross energy content. This same trend in gross energy content was observed when *Art<sub>DHA/EPA</sub>* was supplemented with 5% and 10% copepod inclusion, resulting in an increase in gross energy content in these dietary treatments. From a nutritional perspective, dietary lipids have been shown to play a critically important role in the early development of marine finfish larvae [29]. They serve as the primary energy source for larvae and are a vital source of highly unsaturated fatty acids (HUFA) and essential fatty acids (EFA), which are necessary for the development of new cellular structures. Additionally, they contribute to normal larval growth, morphogenesis, and bone formation [63]. Therefore, a deficiency in HUFA levels in prey can lead to detrimental effects on juvenile seahorse performance, including reduced feeding and swimming activity, as well as a delay in the settlement process, as observed in previous studies where *Artemia* was used, due to its incomplete assimilation [2,5]. Concordantly, a correlation between the type of prey enrichment and the occurrence of gas bubble disorder has also been observed. [17] reported that juveniles of *H. guttulatus* fed *Artemia* enriched with microalgae exhibited a significantly lower incidence of this disorder compared to those fed on *Artemia* enriched with commercial products. Similarly, it was noted that *H. guttulatus* juveniles fed DHA-Selco® enriched *Artemia* metanauplii displayed symptoms of gas bladder overinflation and intestinal disorders, unlike individuals fed solely on natural copepods [59].

In the present study, the selection of the two microalgae for *Artemia* enrichment was based on their DHA and EPA profiles. The analysis revealed that *I. galbana* had a higher proportion of C22:6ω3 (DHA) in its profile (16.5%), while *Nannochloropsis* sp. lacked any significant concentration of this essential fatty acid. On the contrary, *Nannochloropsis* sp. exhibited a substantial percentage of C20:5ω3 (EPA) (36.1%), while *I. galbana* displayed only a trace amount of this fatty acid. The presence of EPA in *Nannochloropsis* sp. notably influenced the HUFA content, leading to a 55% increase compared to that of *I. galbana*. Moreover, *I. galbana* was rich in ω9 HUFAs (15.2%), largely attributed to its high content of oleic acid (C18:1ω9) (15%), which represented 99% of ω9 HUFAs in this species. On its turn, the *Cop* diet had a higher PUFA percentage than *Art<sub>Ref</sub>* base diet. Notably, out of the PUFA content (31%) in the *Cop* diet, 79.9% comprised HUFA, indicating a significant presence of EPA and DHA in comparison to the *Art<sub>Ref</sub>* base diet. In contrast, only 53.4% of PUFA in the *Art<sub>Ref</sub>* diet constituted HUFA. Additionally, the analysis of both *Art<sub>Ref</sub>* and *Art<sub>DHA/EPA</sub>* diets unveiled that the enrichment process contributed to elevate the SFA and MUFA content. Conversely, the incorporation

of 5% or 10% copepods in combination with the enriched *Artemia* increased the PUFA and HUFA content.

Regarding the fatty acid families, the *Cop* diet showed high concentrations of  $\omega 3$  PUFA. Its fatty acid profile showed that EPA and DHA were the predominant polyunsaturated fatty acids. Furthermore, the substantial presence of C16:1 $\omega 7$ c (palmitoleic acid) in the *Artemia* profile contributes to the heightened  $\omega 7$  content in *Artemia*-based dietary treatments (*Art<sub>DHA/EPA</sub>*, *Art<sub>DHA/EPA</sub> 5%* and *Art<sub>DHA/EPA</sub> 10%*). Moreover, the enriched *Artemia* diets exhibited notably high levels of  $\omega 9$  fatty acids, which declined with the incorporation of copepods. This outcome is attributed to the elevated concentration of C18:1 $\omega 9$  (oleic acid) in the *I. galbana*, as mentioned earlier. The analysis of essential fatty acid profiles in the tested dietary treatments revealed the presence of all five essential fatty acids (LA, ALA, ARA, EPA, and DHA), standing out the content of EFA in the *Cop* diet ( $\Sigma$ EFA = 28.4%) (Table 3). The higher content of these fatty acids in the *Cop* diet could potentially justify a better lipid profile of this diet when compared to that of *Artemia* for juvenile seahorses, since high levels of these fatty acids are typically found in marine fish eggs [29]. In comparison, the EFA content in the *Art<sub>Ref</sub>* base diet was significantly lower than that of the *Cop* diet, which can be attributed to the instability of highly unsaturated fatty acids (HUFAs) and their metabolism in *Artemia* [31].

In the preliminary growth trial, the survival rate was 0% after 9 days of the experimental trial across all dietary treatments. This result suggests that the exclusive use of enriched *Artemia* diets since 0 DPP, even with differing DHA/EPA ratios, is not a viable alternative to copepods as a suitable diet for *H. hippocampus* juveniles. While these diets match the necessary fatty acid profile, the outcome agreed with the findings from other studies, in which, their authors linked this observation with the underdeveloped nature of a seahorse's digestive tract in the initial phase of its life cycle. Consequently, the correct assimilation of nutrients is hindered [7,64,65]. As *Artemia* can be more difficult to be digested by fish compared to other live prey such as copepods, when prey densities are high, continuous feeding reduces digestion time lowering the assimilation efficiency in the gut, possibly leading to an intact passage through the digestive tract. In some cases, preys might even remain alive upon passage [18], occasionally maintaining vitality [25]. In their natural habitat, newly born seahorses might exhibit preferences for specific prey items, making it advantageous to perform gut content analysis to assess prey preferences as they develop [66]. Since copepods constitute the primary prey for wild juvenile seahorses, it is expected that seahorses possess a greater capacity for digesting copepods compared to other prey. This capability could arise from an adaptive evolution of the gastrointestinal system and its secretions tailored for copepod digestion [60,67,68]. Moreover, the composition and permeability of the exoskeleton can also account for variations in digestive efficiency [69]. The exoskeleton of a copepod is comprised of a thin, easily breakable, segmented cuticle constructed from a protein matrix containing lipids and chitin rods. Conversely, the cuticle forming the exoskeleton of *Artemia* is thicker and more resistant to breaking [70]. This difference constitutes a significant barrier to the digestion and assimilation of the contents of specific live prey [65]. Additionally, the fact that *Artemia* is not a natural prey item for seahorses adds complexity. For instance, [18] discovered that *H. subelongatus* encountered difficulties in nutrient assimilation from the provided diet. [71] observed *Artemia* being excreted live during the first DPP when fed to juvenile *H. guttulatus*. Similarly, [7] found that *H. guttulatus* newborns exhibited a limited ability to efficiently digest *Artemia* until around 10 DPP, with significant improvement only after 15 DPP [65]. An additional explanation for the less effective use of *Artemia* as food could be attributed not only to the development of the seahorse's digestive system [7,64,72] but also to the size of their oral cavity [64,73], which may limit its capacity to accommodate larger prey. In terms of prey size, since both prey types (*Artemia* and copepods) were measured during the trials, and copepods were found to be almost twice the size of the *Artemia* used, this hypothesis is not plausible, and the current results only suggest that the maturation of the digestive system is the most probable underlying cause. This idea is reinforced by the results of [64], as their study did not establish a clear correlation between the mouth size of the juveniles and the size of the prey.

In contrast, the use of copepods as a preconditioning diet played a pivotal role in achieving the survival and growth of juvenile *H. hippocampus*. The preliminary use of a natural copepod diet appeared to facilitate the maturation of the digestive tract, subsequently enabling the ability to effectively digest other enriched crustacea diets with minimal mortality. Seahorses belong to the category of agastric teleost species, lacking a functional stomach. In these species, the primary site of



digestion is the intestine [74], which undergoes developmental changes during ontogeny, transitioning from a short, direct tube to an elongated, segmented duct. Within this context, the exocrine pancreas assumes the responsibility of synthesizing and releasing digestive enzymes, including proteases, glycosidases, and lipases, into the intestine. This biochemical process aids in the breakdown of dietary nutrients, transforming them into readily absorbable molecules [75].

Given the effectiveness of employing copepods as the primary diet for most finfish species, their inclusion, often co-fed with other diets to address the challenges of their exclusive use, has the potential to significantly enhance the growth and survival of juvenile seahorses e.g., [59,64]. Therefore, considering the unsatisfactory performance of *Artemia* in the preliminary trial, the second trial was conducted to test the beneficial use of limited copepod inclusion levels. During this trial, survival rates exhibited considerable fluctuation, yet the average survival rate across all dietary treatments reached 81%, with the lowest recorded at 72%. This promising outcome is noteworthy in comparison to findings from other studies [17,23,70]. [18] reported survival rates of >80% at 14 DPP for *H. subelongatus* juveniles fed copepods, compared to >40% for those fed enriched *Artemia*. In the present study, considering the appropriateness of a copepod diet, one would anticipate a higher survival rate among fish fed this diet. However, contrary to this expectation, the opposite was observed, as fish fed the *Cop* diet exhibited the lowest survival rate (72%) among the four tested groups. During the initial 7 days of the trial (0-7 DPP), all groups were provided with the same copepod diet, and only on the 8<sup>th</sup> day on the diet was changed to the respective dietary treatments. Therefore, the initial mortality observed within the first 7 DPP cannot be attributed to the diet. Similarly, previous studies achieved reasonably high rates of survival and growth for seahorse juveniles fed copepod-based diets [16,18,76].

Regarding the growth performance of fish fed various dietary treatments, it is noteworthy that the mean LG displayed greater homogeneity than the mean WG during the analysed time period. Furthermore, LG exhibited a more pronounced similarity among the groups. Once again, a recurring observation was made: smaller LG was associated with individuals fed *Art<sup>DHA/EPA</sup>*, whereas larger LG was evident among those fed on other dietary regimes, with the *Cop* diet resulting in the highest measurements. Consequently, juvenile seahorses fed the enriched *Artemia* diet (*Art<sup>DHA/EPA</sup>*) without any inclusion of copepods demonstrated the lowest MW gain at the conclusion of the experimental trial. As previously mentioned, this outcome likely resulted from the inadequate DHA/EPA ratio (0.04) identified in this particular diet, negatively impacting juvenile growth. Additionally, it was evident that when diets included either enriched *Artemia* co-fed with copepods (*Art<sup>DHA/EPA5%</sup>* and *Art<sup>DHA/EPA10%</sup>*) or were exclusively comprised of copepods (*Cop*), the MW gain increased significantly ( $P < 0.05$ ). This consequently yielded substantially higher mean WG values, thereby contributing to elevated thermal-unit growth coefficient (TGC) values. These findings can also be attributed to the heightened HUFA content observed in these diets compared to that of *Art<sup>DHA/EPA</sup>*. This observation agrees with [77], who demonstrated that optimal growth performance in *H. erectus* juveniles fed copepods was attributed to the abundant content of HUFAs.

In conclusion, the growth and survival of juvenile *H. hippocampus* are significantly enhanced when the initial diet consists exclusively of copepods, allowing for the maturation of the digestive tract. Subsequently, even a small inclusion of 5% to 10% copepods in the diet contributes to an improved fatty acid profile due to copepods' high EFA content. This dietary adjustment aligns with the species' requirements, translating into substantial benefits in terms of growth and survival.

**Author Contributions:** For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “Conceptualization, J.P.; methodology, J.P. and M.J.L.; formal analysis, R.L.; writing—original draft preparation, J.P. and R.L.; review and editing, M.J.L. and J.P.A.; supervision, J.P.; project administration, J.P.; funding acquisition, J.P. and J.P.A.. All authors have read and agreed to the published version of the manuscript.”

**Funding:** This study was funded by the projects HIPPOSAVE “Plano de ação para a recuperação das populações de cavalos marinhos da Ria Formosa” (Programa Operacional MAR2020, MAR-01.04.02-FEAMP-0029), and HIPPONUTRE “Cultivo do cavalo marinho de focinho comprido, *Hippocampus guttulatus*: Optimização zootécnica e avaliação de requisitos nutricionais” (Programa Operacional MAR2020-16-02-01-FMP-54), and received Portuguese national funds from FCT - Foundation for Science and Technology through projects UIDB/04326/2020, UIDP/04326/2020 and LA/P/0101/2020.

**Institutional Review Board Statement:** CCMAR facilities and the research is certified to house and conduct experiments with live animals (Group-C licenses by the Direção Geral de Alimentação e Veterinária, Ministério da Agricultura, Florestas e Desenvolvimento Rural, Portugal). The experimental design of the present study was part of the Projects HIPPONUTRE (reference 1602-01-FMP-54) and HIPPOSAVE (reference MAR-01.04.02-FEAMP-0029), which obtained approval from the ethics committee of the Veterinary Medicines Directorate, Ministry of Agriculture, Rural Development and Fisheries, Portugal. The study adhered to the guidelines outlined by the European Union Council (86/609/EU) and the relevant Portuguese legislation concerning the use of laboratory animals.

**Data Availability Statement:** The data used during the current study are available from the corresponding author upon reasonable request.

**Acknowledgments:** The authors would like to acknowledge André Jorge for his valuable contribution and assistance in the fatty acid laboratory analyses.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## References

1. Vincent, A.C.J. The international trade in seahorses. **1996**; TRAFFIC International; Cambridge, UK. ISBN 1 85850 098 2
2. Koldewey, H.J.; Martin-Smith, K.M. A global review of seahorse aquaculture. *Aquaculture* **2010**, *302*, 131-152. <https://doi.org/10.1016/j.aquaculture.2009.11.010>
3. IUCN. The IUCN Red List of Threatened Species. Version 2021.1. In: <http://www.iucnredlist.org>. Accessed in August 2023.
4. Pollom, R. 2016. *Hippocampus hippocampus* (Mediterranean assessment). *The IUCN Red List of Threatened Species* 2016: e.T10069A90866381.
5. Otero-Ferrer, F.; Molina, L.; Socorro, J.; Herrera, R. Live prey first feeding regimes for short-snouted seahorse *Hippocampus hippocampus* (Linnaeus, 1758) juveniles. *Aquacult. Res.* **2010**, *41*, 8–19. <https://doi.org/10.1111/j.1365-2109.2010.02505.x>
6. Leus, K. Captive breeding and conservation. *Zool. Middle East* **2011**, *54*(sup3), 151-158. <https://doi.org/10.1080/09397140.2011.10648906>
7. Olivotto, I.; Planas, M.; Simões, N.; Holt, G.J.; Avella, M.A.; Calado, R. Advances in breeding and rearing marine ornamentals. *J. World Aquac. Soc.* **2011**, *42*, 135-166. <https://doi.org/10.1111/j.1749-7345.2011.00453.x>
8. Rønnestad, I.; Thorsen, A.; Finn, R.N. Fish larval nutrition: a review of recent advances in the roles of amino acids. *Aquaculture* **1999**, *177*, 201–216. [https://doi.org/10.1016/S0044-8486\(99\)00082-4](https://doi.org/10.1016/S0044-8486(99)00082-4)
9. Barclay, W.; Zeller, S. Nutritional enrichment of n-3 and n-6 fatty acids in rotifers and *Artemia* nauplii by feeding spray-dried *Schizochytrium* sp. *J. World Aquac. Soc.* **1996**, *27*(3): 314-322. <https://doi.org/10.1111/j.1749-7345.1996.tb00614.x>
10. Furuita, H.; Konishi, K.; Takeuchi, T. Effect of different levels of eicosapentaenoic acid and docosahexaenoic acid in *Artemia* nauplii on growth, survival and salinity tolerances of larvae of the Japanese flounder, *Paralichthys olivaceus*. *Aquaculture* **1999**, *170*, 59 – 69. [https://doi.org/10.1016/S0044-8486\(98\)00386-X](https://doi.org/10.1016/S0044-8486(98)00386-X)
11. Sorgeloos, P.; Dhert, P.; Candreva, P. Use of the brine shrimp, *Artemia* spp., in marine fish larviculture. *Aquaculture* **2001**, *200*(1-2), 147-215. [https://doi.org/10.1016/S0044-8486\(01\)00698-6](https://doi.org/10.1016/S0044-8486(01)00698-6)
12. Watanabe, T. Nutritional quality of brine shrimp, *Artemia salina*, as a living feed from the viewpoint of essential fatty acids for fish. *Bull. Japan. Soc. Sci. Fish.* **1978**, *44*, 1115.
13. Watanabe, T. Lipid nutrition in fish. *Comp. Biochem. Physiol.* **1982**, *73B*, 3–15. NII Article ID: 20001329776
14. Job, S.; Do, H.; Meeuwig, J.; Hall, H. Culturing the oceanic seahorse, *Hippocampus kuda*. *Aquaculture* **2002**, *214*(1-4), 333-334. [https://doi.org/10.1016/S0044-8486\(02\)00063-7](https://doi.org/10.1016/S0044-8486(02)00063-7)
15. Martínez-Cárdenas L.; Purser G.J. Effect of tank colour on *Artemia* ingestion, growth and survival in cultured early juvenile pot-bellied seahorses (*Hippocampus abdominalis*). *Aquaculture* **2007**, *26*, 92–100. <https://doi.org/10.1016/j.aquaculture.2006.12.045>
16. Olivotto, I.; Avella, M. A.; Sampaolesi, G.; Piccinetti, C. C.; Ruiz, P. N.; Carnevali, O. Breeding and rearing the longsnout seahorse *Hippocampus reidi*: Rearing and feeding studies. *Aquaculture* **2008**, *283*, 92–96. <https://doi.org/10.1016/j.aquaculture.2008.06.018>

17. Palma, J.; Bureau, D. P.; Andrade, J. P. Effect of different *Artemia* enrichments and feeding protocol for rearing juvenile long snout seahorse, *Hippocampus guttulatus*. *Aquaculture* **2011**, 318(3–4), 439–443. <https://doi.org/10.1016/j.aquaculture.2011.05.035>
18. Payne, M.F.; Rippingale, R.J. Rearing West Australian seahorse, *Hippocampus subelongatus*, juveniles on copepod nauplii and enriched *Artemia*. *Aquaculture* **2000**, 188, 353–361. [https://doi.org/10.1016/S0044-8486\(00\)00349-5](https://doi.org/10.1016/S0044-8486(00)00349-5)
19. Sheng, J.; Lin, Q.; Chen, Q.; Gao, Y.; Shen, L.; Lu, J. Effects of food, temperature and light intensity on the feeding behavior of three-spot juvenile seahorses, *Hippocampus trimaculatus* Leach. *Aquaculture* **2006**, 256, 596–607. <https://doi.org/10.1016/j.aquaculture.2006.02.026>
20. Willadino, L.; Souza-Santos, L.P.; Mélo, R.C.S.; Brito, A.P.; Barros, N.C.S.; Araújo-Castro, C.M.V.; Galvão, D.B.; Gouveia, A.; Regis, C.G.; Cavalli, R.O. Ingestion rate, survival and growth of newly released seahorse *Hippocampus reidi* fed exclusively on cultured live food items. *Aquaculture* **2012**, 360, <https://doi.org/10.1016/j.aquaculture.2012.06.025>
21. Woods, C.M.C. Preliminary observations on breeding and rearing the seahorse *Hippocampus abdominalis* (Teleostei: Syngnathidae) in captivity. *N. Z. J. Mar. Freshw. Res.* **2000a**, 34, 475–485. <https://doi.org/10.1080/00288330.2000.9516950>
22. Zhang, D.; Yin, F.; Lin, J. Criteria for assessing juvenile quality of the lined seahorse, *Hippocampus erectus*. *Aquaculture* **2011**, 322, 255–258. <https://doi.org/10.1016/j.aquaculture.2011.10.008>
23. Chang, M.; Southgate, P. C. Effects of varying dietary fatty acid composition on growth and survival of seahorse, *Hippocampus* sp., juveniles. *Aquarium Sci. Conserv.* **2001**, 3(1), 205–214. <https://doi.org/10.1023/A:1011363807074>
24. Filleul, M.A. Optimizing growth of juveniles big bellied seahorse *Hippocampus abdominalis* Lesson. Unpublished B.Sc (Hons) thesis, University of Tasmania, Australia, **1996**.
25. Woods, C.M.C. Effects of varying *Artemia* enrichment on growth and survival of juvenile seahorses, *Hippocampus abdominalis*. *Aquaculture* **2003**, 220, 537–548. [https://doi.org/10.1016/S0044-8486\(02\)00639-7](https://doi.org/10.1016/S0044-8486(02)00639-7)
26. Sargent, J.R.; Bell, J.G.; Bell, M.V.; Henderson, R.J.; Tocher, D.R. Dietary origins and functions of long-chain (n-3) polyunsaturated fatty acids in marine fish. *J. Mar. Biotechnol.* **1995**, 3(1), 26–28. ISSN: 09412905
27. Tocher, D.R. Metabolism and functions of lipids and fatty acids in teleost fish. *Rev. Fish. Sci.* **2003**, 11(2), 107–118. <https://doi.org/10.1080/713610925>
28. Corraze, G. Lipid nutrition. In: *Nutrition and Feeding of Fish and Crustaceans*, Guillaume, J.; Kaushik, S.; Bergot, P.; Metailler, R. Eds. Springer-Praxis, UK, **2001**, pp. 111–129. ISBN 10: 1852332417; ISBN 13: 9781852332419
29. Sargent, J.R.; Tocher, D.R.; Bell, J.G. The lipids. In: *Fish Nutrition*, 3<sup>rd</sup> Ed.; Halver, J.E.; Hardy, R.W. Eds., Academic Press, U.S. **2002**, pp. 182–246. <http://www.elsevierdirect.com/imprint.jsp?iid=5>
30. Sargent, J.R.; McEvoy, L.A.; Bell, J.G. Requirements, presentation and sources of polyunsaturated fatty acids in marine fish larval feeds. *Aquaculture* **1997**, 155, 117–127. [https://doi.org/10.1016/S0044-8486\(97\)00122-1](https://doi.org/10.1016/S0044-8486(97)00122-1)
31. Faleiro, F.; Narciso, L. Lipid dynamics during early development of *Hippocampus guttulatus* seahorses: Searching for clues on fatty acid requirements. *Aquaculture*, **2010**, 307, 56–64. <https://doi.org/10.1016/j.aquaculture.2010.07.005>
32. Watanabe T.; Kiron V. Prospects in larval fish dietetics. *Aquaculture* **1994**, 124, 223–251. DOI:10.1016/0044-8486(94)90386-7
33. Watanabe, T.; Arakawa, T.; Takeuchi, T.; Satoh, S. Comparison between eicosapentaenoic and docosahexaenoic acids in terms of essential fatty acid efficiency in juvenile striped jack *Pseudocaranx dentex*. *Nippon Suisan Gakk.* **1989**, 55(11), 1989 – 1995. <https://doi.org/10.2331/SUISAN.55.1989>
34. Faleiro, F.; Narciso, L. Prey–predator dynamics in seahorses (*Hippocampus guttulatus*): deciphering fatty acid clues. *Aquacult. Res.* **2013**, 44, 618–633. <https://doi.org/10.1111/j.1365-2109.2011.03067.x>
35. Sargent, J.; Bell, G.; McEvoy, L.; Tocher, D.; Estevez, A. Recent developments in the essential fatty acid nutrition of fish. *Aquaculture*, **1999**, 177(14), 191–219. [https://doi.org/10.1016/S0044-8486\(99\)00083-6](https://doi.org/10.1016/S0044-8486(99)00083-6)
36. Webster C.D.; Lovell R.T. Response of striped bass larvae fed brine shrimp from different sources containing different fatty acid compositions. *Aquaculture* **1990**, 90, 49–62. [https://doi.org/10.1016/0044-8486\(90\)90282-R](https://doi.org/10.1016/0044-8486(90)90282-R)

37. Ostrowski, A.C.; Laidley, C.W. Application of marine food fish techniques in marine ornamental aquaculture: reproduction and larval first feeding. *Aquarium Sci. Conserv.* **2001**, *3*(1), 191-204. <https://doi.org/10.1023/A:1011349931035>
38. Holt, G.J. Research on culturing the early life stages of marine ornamental fish. In: *Marine ornamental species – Collection, culture & conservation*, Cato, J.C.; Brown, C.L. Eds. Iowa State Press, U.S; **2003**, pp. 251-254. <https://doi.org/10.1002/9780470752722.ch17>
39. Aragão, C.; Conceição, L.E.; Dinis, M.T.; Fyhn, H.J. Amino acid pools of rotifers and *Artemia* under different conditions: nutritional implications for fish larvae. *Aquaculture* **2004**, *234*, 429–445. <https://doi.org/10.1016/j.aquaculture.2004.01.025>
40. Hamre, K.; Srivastava, A.; Rønnestad, I.; Mangor-Jensen, A.; Stoss, J. Several micronutrients in the rotifer *Brachionus* sp. may not fulfil the nutritional requirements of marine fish larvae. *Aquacult. Nut.* **2008**, *14*(1), 51-60. <https://doi.org/10.1111/j.1365-2095.2007.00504.x>
41. Conceição, L.E.; Yúfera, M.; Makridis, P.; Morais, S.; Dinis, M.T. Live feeds for early stages of fish rearing. *Aquacult. Res.* **2010**, *41*(5), 613-640. <https://doi.org/10.1111/j.1365-2109.2009.02242.x>
42. Zeng, C.; Shao, L.; Ricketts, A.; Moorhead, J. The importance of copepods as live feed for larval rearing of the green mandarin fish *Synchiropus splendidus*. *Aquaculture* **2018**, *491*, 65-71. <https://doi.org/10.1016/j.aquaculture.2018.03.011>
43. Alajmi, F.; Zeng, C.; Jerry, D.R. Domestication as a novel approach for improving the cultivation of calanoid copepods: a case study with *Parvocalanus crassirostris*. *PLoS One* **2015**, *10*, e0133269. <https://doi.org/10.1371/journal.pone.0133269>
44. Kim, B.G.; Divakaran, S.; Brown, C.L.; Ostrowski, A. Comparative digestive enzyme ontogeny in two marine larval fishes, Pacific threadfin (*Polydactylus sexfilis*) and bluefin trevally (*Caranx melampygus*). *Fish Physiol. Biochem.* **2001**, *24*, 225-241. <https://doi.org/10.1023/A:1014054431627>
45. Pérez-Casanova, J.C.; Murray, H.M.; Gallant, J.W.; Ross, N.W.; Douglas, S.E.; Johnson, S.C. Development of the digestive capacity in larvae of haddock (*Melanogrammus aeglefinus*) and Atlantic cod (*Gadus morhua*). *Aquaculture* **2006**, *251*, 377-401. <https://doi.org/10.1016/j.aquaculture.2005.06.007>
46. Bolasina, S.; Pérez, A.; Yamashita, Y. Digestive enzymes activity during ontogenetic development and effect of starvation in Japanese flounder, *Paralichthys olivaceus*. *Aquaculture* **2006**, *252*, 503-515. <https://doi.org/10.1016/j.aquaculture.2005.07.015>
47. Furné, M.; García-Gallego, M.; Hidalgo, M.C.; Morales, A.E.; Domezain, A.; Domezain, J.; Sanz, A. Effect of starvation and refeeding on digestive enzyme activities in sturgeon (*Acipenser naccarii*) and trout (*Oncorhynchus mykiss*). *Comp. Biochem. Physiol. A* **2008**, *149*, 420-425. <https://doi.org/10.1016/j.cbpa.2008.02.002>
48. Lazo, J.P.; Mendoza, R.; Holt, G.J.; Aguilera, C.; Arnold, C.R. Characterization of digestive enzymes during larval development of red drum (*Sciaenops ocellatus*). *Aquaculture* **2007**, *265*, 194-205. <https://doi.org/10.1016/j.aquaculture.2007.01.043>
49. Palma, J.; Stockdale, J.; Correia, M.; Andrade, J.P. Growth and survival of adult long snout seahorse (*Hippocampus guttulatus*) using frozen diets. *Aquaculture* **2008**, *278*(1-4), 55–59. <https://doi.org/10.1016/j.aquaculture.2008.03.019>
50. Woods, C.M.C. Improving initial survival in cultured seahorses, *Hippocampus abdominalis* Leeson, 1827 (Teleostei: Syngnathidae). *Aquaculture* **2000b**, *190*, 377–388. [https://doi.org/10.1016/S0044-8486\(00\)00408-7](https://doi.org/10.1016/S0044-8486(00)00408-7)
51. Sorgeloos, P.; Lavens, P.; Léger, P.; Tackaert, W.; Versichele, D. *Manual for the culture of brine shrimp Artemia in aquaculture*. **1986**, University of Ghent, Ghent, Belgium. ISBN 92-5-103934-8
52. Palma, J.; Andrade, J.P.; Bureau, D.P. The impact of dietary supplementation with astaxanthin on egg quality and growth of long snout seahorse (*Hippocampus guttulatus*) juveniles. *Aquacult. Nut.* **2017**, *23*(2), <https://doi.org/10.1111/anu.12394>
53. Lourie, S.A.; Pritchard, J.C.; Casey, S.P.; Truong, S.K.; Hall, H.J.; Vincent, A.C.J. The taxonomy of Vietnam's exploited seahorses (family Syngnathidae). *Biol. J. Linn. Soc.* **1999**, *66*, 231-256. <https://doi.org/10.1111/j.1095-8312.1999.tb01886.x>
54. Iwama, G.K.; Tautz, A.F. A simple growth model for salmonids in hatcheries. *Can. J. Fish. Aquat. Sci.* **1981**, *38*, 649–656. <https://doi.org/10.1139/f81-087>
55. Cho, C.Y. Fish nutrition, feeds and feeding: with special emphasis on salmonid aquaculture. *Food Rev. Int.* **1990**, *6*, 333–357. <https://doi.org/10.1080/87559129009540876>



56. AOAC (Association of Official Analytical Chemists). *Official methods of analysis of AOAC International. Agricultural chemicals; contaminants drugs*. 16<sup>th</sup> edition; volume I. **1995**, Association of Official Analytical Chemists, Arlington, Virginia, USA.
57. Folch, J.; Lees, M.; Stanley, G.H.S. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* **1957**, 226(1), 497–509. [https://doi.org/10.1016/S0021-9258\(18\)64849-5](https://doi.org/10.1016/S0021-9258(18)64849-5)
58. Jorge, A.F.N. Determinação do perfil de ácidos gordos do músculo de achigã (*Micropterus salmoides*). **2018**. MsC Thesis; Universidade Nova de Lisboa; Lisboa.
59. Palma, J.; Bureau, D.; Andrade, J.P. The effect of diet on ontogenic development of the digestive tract in juvenile reared long snout seahorse *Hippocampus guttulatus*. *Fish Physiol. Biochem.* **2014**, 40(3), 739–775. <https://doi.org/10.1007/s10695-013-9881-8>
60. Schubert, P.; Vogt, L.; Eder, K.; Hauße, T.; Wilke, T. Effects of Feed Species and HUFA Composition on Survival and Growth of the Longsnout Seahorse (*Hippocampus reidi*). *Front. Mar. Sci.* **2016**, 3, 53. <https://doi.org/10.3389/fmars.2016.00053>
61. Gardner, T. The copepod/*Artemia* trade-off in the captive culture of *Hippocampus erectus*, a vulnerable species in New York State. 2004, MsC Thesis. Hofstra University; Long Island, New York, U.S.
62. Cohen, F.P.; Valenti, W.C. Opportunities and constraints for developing low-cost aquaculture of seahorses in mangrove estuaries. *Aquaculture* **2019**, 502, 121–127. <http://dx.doi.org/10.1016/j.aquaculture.2018.12.031>
63. Lall, S.P.; Lewis-McCrea, L.M. Role of nutrients in skeletal metabolism and pathology in fish—an overview. *Aquaculture* **2007**, 267(1–4), 3–19. <https://doi.org/10.1016/j.aquaculture.2007.02.053>
64. Blanco, A.; Planas, M. Mouth growth and prey selection in juveniles of the European long-snouted seahorse, *Hippocampus guttulatus*. *J. World Aquacult. Soc.* **2015**, 46, 596–607. <https://doi.org/10.1111/jwas.12240>
65. Blanco, A.; Planas, M.; Moyano, F.J. Ontogeny of digestive enzymatic capacities in juvenile seahorses *Hippocampus guttulatus* fed on different live diets. *Aquacult. Res.* **2016**, 47, 3558–3569. <https://doi.org/10.1111/are.12806>
66. Pham, N.; Lin, J. The effects of different feed enrichments on survivorship and growth of early juvenile long-snout seahorse, *Hippocampus reidi*. *J. World Aquacult. Soc.* **2013**, 44(3), <https://doi.org/10.1111/jwas.12037>
67. Castro, A.L.D.C.; Diniz, A.D.F.; Martins, I.Z.; Vendel, A.L.; Oliveira, T.P.R.D.; Rosa, I.M.D.L. Assessing diet composition of seahorses in the wild using a non-destructive method: *Hippocampus reidi* (Teleostei: Syngnathidae) as a study-case. *Neotrop. Ichthyol.* **2008**, 6(4), 637–644. <https://doi.org/10.1590/S1679-62252008000400012>
68. Tipton, K.; Bell, S. Foraging patterns of two syngnathid fishes: importance of harpacticoid copepods. *Mar. Ecol. Prog. Ser.* **1988**, 47, 31–113. <https://doi.org/10.3354/meps047031>
69. Bodin, N.; Delfosse, G.; Thi, T.; Thu, N.; Le, E.; Abboudi, T.; Larondelle, Y.; Rollin, X. Effects of fish size and diet adaptation on growth performances and nitrogen utilization of rainbow trout (*Oncorhynchus mykiss* W.) juveniles given diets based on free and/or protein-bound amino acids. *Aquaculture* **2012**, 356, 105–115. <https://doi.org/10.1016/j.aquaculture.2012.05.030>
70. Randazzo, B.; Rolla, L.; Ofelio, C.; Planas, M.; Gioacchini, G.; Vargas, A.; Giorgini, E.; Olivotto, I. The influence of diet on the early development of two seahorse species (*H. guttulatus* and *H. reidi*): Traditional and innovative approaches. *Aquaculture* **2018**, 490, 75–90. <https://doi.org/10.1016/j.aquaculture.2018.02.029>
71. Planas, M.; Quintas, P.; Chamorro, A.; Balcázar, J.L. Husbandry and rearing of the seahorse *Hippocampus guttulatus* (project *Hippocampus*). In: Proceedings of the World Aquaculture Society; World Aquaculture 2009, Veracruz (México), Sept 25–29, **2009**, p. 668.
72. Planas, M.; Blanco, A.; Chamorro, A.; Valladares, S.; Pintado, J. Temperature-induced changes of growth and survival in the early development of the seahorse *Hippocampus guttulatus*. *J. Exp. Mar. Biol. Ecol.* **2012**, 438, 154–162. <https://doi.org/10.1016/j.jembe.2012.10.003>
73. Manning, C.G.; Foster, S.J.; Vincent, A.C.J. A review of the diets and feeding behaviours of a family of biologically diverse marine fishes (Family Syngnathidae). *Rev. Fish Biol. Fisheries* **2019**, 1–25. <https://doi.org/10.1007/s11160-019-09549-z>
74. Rønnestad, I.; Yúfera, M.; Ueberschär, B.; Ribeiro, L.; Sæle, Ø.; Boglione, C. Feeding behaviour and digestive physiology in larval fish: current knowledge, and gaps and bottlenecks in research. *Rev. Aquac.* **2013**, 5, S59–S98. <https://doi.org/10.1111/raq.12010>

75. Zambonino-Infante, J.L.; Cahu, C.L. Ontogeny of the gastrointestinal tract of marine fish larvae. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* **2001**, *130*, 477-487. [https://doi.org/10.1016/S1532-0456\(01\)00274-5](https://doi.org/10.1016/S1532-0456(01)00274-5)
76. Celino, F.T.; Hilomen-Garcia, G.V.; del Norte-Campos, A.G. Feeding selectivity of the seahorse, *Hippocampus kuda* (Bleeker), juveniles under laboratory conditions. *Aquacult. Res.* **2012**, *43*(12), 1804-1815. <https://doi.org/10.1111/j.1365-2109.2011.02988.x>
77. Zhang, D.; Lin, T.; Liu, X. A comparison of growth, survival, and fatty acid composition of the lined seahorse, *Hippocampus erectus*, juveniles fed enriched *Artemia* and a calanoid copepod, *Schmackeria dubia*. *J. World Aquacult. Soc.* **2015**, *46*(6), 608-616. <http://dx.doi.org/10.1111/jwas.12233>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.