

Article

Not peer-reviewed version

On the Interpretation of Cosmic Acceleration

[Enrique Gaztanaga](#) *

Posted Date: 21 August 2024

doi: 10.20944/preprints202309.0873.v7

Keywords: Cosmology; Dark Energy; General Relativity; Black Holes; Cosmological Constant



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

On the Interpretation of Cosmic Acceleration

Enrique Gaztanaga 

¹ Institute of Cosmology & Gravitation, University of Portsmouth, Dennis Sciamia Building, Burnaby Road, Portsmouth PO1 3FX, UK; gaztanaga@gmail.com

² Institute of Space Sciences (ICE, CSIC), 08193 Barcelona, Spain

³ Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain

Abstract: In relativity, the newtonian concepts of velocity and acceleration are observer-dependent quantities that vary with the chosen frame of reference. It is well established that, in the comoving frame, cosmic expansion is currently accelerating; however, in the rest frame, this expansion is actually decelerating. In this paper, we explore the implications of this distinction. The traditional measure of cosmic acceleration, denoted by q , is derived from the comoving frame and describes the acceleration of the scale factor a for a 3D space-like homogeneous sphere. We introduce a new parameter, q_E , representing the acceleration experienced between observers within the light cone. By comparing q_E to the traditional q , using observational data from Type Ia supernovae (SN) and the radial clustering of galaxies and quasars (BAO)—including the latest results from DESI2024—our analysis demonstrates that q_E aligns more closely with this data. The core argument of the paper is that Λ —regardless of its origin—creates an Event Horizon that divides the manifold into causally disconnected regions, analogous to conditions inside a black hole's interior, thereby allowing for a rest frame perspective q_E in which cosmic expansion appears to be decelerating and the horizon acts like a friction term. Such horizon suggests that the universe cannot maintain homogeneity outside. The observed cosmological constant Λ can then be interpreted not as a driver of new dark energy or a modification of gravity, but as a boundary term exerting an attractive force, akin to a rubber band, resisting further expansion and preventing event horizon crossings. This interpretation calls for a reconsideration of current cosmological models and the assumptions underlying them.

Keywords: cosmology; dark energy; general relativity; black holes

1. Introduction

For over thirty years, cosmologists have accumulated compelling evidence that cosmic expansion is accelerating. More specifically, this acceleration appears to be dominated by a Cosmological Constant term, commonly denoted as Λ .

This Λ term can be understood in three distinct ways: (1) as a fundamental (or effective) modification of Einstein's classical General Relativity (GR), denoted Λ_F , (2) as an effective source term from a Dark Energy (DE) component, Λ_{DE} , or (3) as a boundary term, Λ_B . These three possible origins can be illustrated by writing the Einstein-Hilbert action for classical GR with the corresponding additional terms:

$$S = \int_{V_4} dV_4 \left[\frac{R}{16\pi G} - \frac{\Lambda_F}{8\pi G} + \mathcal{L}_{DE} \right] + \frac{1}{8\pi G} \oint_{\partial V_4} dV_3 K. \quad (1)$$

The first term in the volume integral represents the classical GR Lagrangian [1]. The second term corresponds to the fundamental cosmological constant Λ_F [2]. The third term is the DE or quintessence source term, given by $\mathcal{L}_{DE} = \frac{1}{2} \bar{\nabla}^2 \psi - V(\psi)$. For a comoving homogeneous perfect fluid with density ρ_{DE} and pressure p_{DE} in its ground state (where $V \gg \dot{\psi}^2$), this reduces to $\mathcal{L}_{DE} = p_{DE} = -\rho_{DE} = -\Lambda_{DE}/8\pi G$ (see also Eqs. B66-B68 in [3]). This illustrates how Λ_F and Λ_{DE} provide a totally degenerate interpretations for Λ .

The last integral represents the Gibbons-Hawking-York (GHY) boundary term [4–6], where K is the trace of the extrinsic curvature at the boundary $\partial V_4 = V_3$. As shown in [7], for a finite Friedman-Lemaître-Robertson-Walker (FLRW) metric with total mass-energy M_T , this term results in:

$$S_{GHY} = \frac{1}{8\pi G} \oint_{\partial V_4} dV_3 K = \int_{V_4} dV_4 \left[\frac{-2\Lambda_B}{16\pi G} \right], \quad (2)$$

where $\Lambda_B = \frac{3}{r_S^2}$ with $r_S = 2GM_T$. This provides a fundamentally different origin for Λ compared to Λ_F or Λ_{DE} . As noted in [8], the Λ_B boundary term cancels both Λ_F or Λ_{DE} potential contributions, solving the fine tune and coincidence problems [9].

Regardless of its origin, the Λ term is typically interpreted as a repulsive force between galaxies that counteracts gravity, driving the accelerated expansion of the universe. This phenomenon is often cited as one of the most profound challenges in contemporary physics and may offer a crucial observational pathway to understanding quantum gravity (e.g., see [10] and references therein).

Cosmic acceleration is typically measured using the adimensional coefficient q , defined as $(\ddot{a}/a)H^{-2}$, where $H \equiv \dot{a}/a$. If the universe follows an equation of state with $p = \omega\rho$, this leads to $q = -\frac{1}{2}(1 + 3\omega)$. For regular matter or radiation where $\omega > 0$, we'd expect deceleration in the expansion ($q < 0$) due to gravity. However, measurements from various sources, such as galaxy clustering, Type Ia supernovae, and CMB, consistently show an expansion asymptotically approaching $q \simeq 1$ or $\omega \simeq -1$ (e.g. see [11] and references therein for a review of more recent results, including weak gravitational lensing). This aligns well with a Cosmological Constant Λ , where H^2 approaches $H^2 = \Lambda/3 \equiv r_\Lambda^{-2}$ and q approaches 1. What does this mean?

The term Dark Energy (DE) was introduced by [12] to refer for any component with $\omega < -1/3$. However, there is no fundamental understanding of what DE is or why we measure a term with $\omega \simeq -1$. A natural candidate for DE is Λ , which is equivalent to $\omega = -1$ and can also be thought of as the ground state of a field (the DE), similar to the Inflaton but with a much smaller ($\simeq 10^{-50} - 10^{-100}$) energy scale. Λ can also be a fundamental constant in GR, but this has some other complications ([13–15]).

This paper critically examines the conventional concept of cosmic acceleration and proposes an alternative framework for understanding cosmic expansion dynamics. In section 2, we establish notation and derive standard definitions for cosmic expansion in the comoving frame. In Section 3, we demonstrate the dependence of these standard definitions on the observer's frame, highlighting the lack of covariance and potential for misinterpretation in the commonly used concept of cosmic acceleration. Sections 4 and 5 introduce an alternative definition for cosmic acceleration, which is grounded in the Λ event horizon. In Section 6, we compare both definitions to observational data, demonstrating that our proposed approach offers greater consistency with empirical observations.

The Appendix provides a detailed exposition of the correct method for defining 4D acceleration in relativity, based on the geodesic deviation equation. We also elaborate on the idea that Λ corresponds to a friction (attractive) force that decelerates cosmic events and revisit the Newtonian limit to show that Λ corresponds to an additional (attractive) Hooke's term to the inverse square gravitational law, envisioning a "rubber band Universe".

Finally, we conclude with a summary and discussion, emphasizing the significance of our findings for cosmological theory and observational practice, and suggesting avenues for further research and exploration in the field.

2. Cosmic Acceleration

Current observations of the cosmos seem consistent with General Relativity (GR) with a flat FLRW (Friedmann–Lemaître–Robertson–Walker) metric in comoving coordinates, corresponding to a homogeneous and isotropic space :

$$ds^2 = -d\tau^2 + a(\tau)^2 \left[d\chi^2 + \chi^2 d\Omega^2 \right], \quad (3)$$

where we use units of $c = 1$ and $a(\tau)$ is the scale factor. For a classical perfect fluid with matter and radiation density $\rho = \rho_m + \rho_R$, the solution to Einstein's field equations (called LCDM) is well known:

$$H^2 = \frac{8\pi G}{3}\rho + \frac{\Lambda}{3} \equiv H_0^2 \left[\Omega_m a^{-3} + \Omega_R a^{-4} + \Omega_\Lambda \right], \quad (4)$$

where $\Omega_X \equiv \frac{\rho_X}{\rho_c}$, where Ω_m and Ω_R represents the current ($a = 1$) matter and radiation density, $\rho_c \equiv \frac{3H_0^2}{8\pi G}$ and $\Omega_m + \Omega_R + \Omega_\Lambda = 1$. The cosmological constant (Λ) term corresponds to $\Omega_\Lambda = H_0^{-2}\Lambda/3 \simeq 0.7$ where $H_0 \simeq 70\text{Km/s/Mpc}$. Given $\Omega_m \simeq 0.3$ and $\Omega_R \simeq 4 \times 10^{-5}$ we can use the above equations to find $a(\tau)$.

Cosmic acceleration is usually defined as $\ddot{a}/a$, where the dot represents a derivative with respect to proper time τ at emission. A derivative over Equation (4) shows that:

$$q(z) \equiv \frac{\ddot{a}}{a} \frac{1}{H^2} = \left(\Omega_\Lambda - \frac{1}{2}\Omega_m a^{-3} - \Omega_R a^{-4} \right) \frac{H_0^2}{H^2}. \quad (5)$$

For $\Lambda = 0$ Equations (4)-(5) indicate that as time passes ($a \Rightarrow \infty$) we have that $H \Rightarrow 0$ and $q \Rightarrow -1/2$. This is because gravity opposes cosmic expansion and brings the expansion asymptotically to a halt. Including Λ brings the expansion to accelerate so that $H \Rightarrow r_\Lambda^{-1}$ and $q \Rightarrow 1$. This is illustrated as black continuous and dashed lines in Figure 1 for $\Omega_\Lambda = 0.7$. The effect of Λ is then interpreted as a mysterious new repulsive force (or Dark Energy) that opposes gravity.




Figure 1. Log of cosmic expansion rate (continuous lines) and acceleration (dashed lines) as a function of time (log scale factor a) for $\Omega_\Lambda = 0.7$. The black lines correspond to the usual interpretation in terms of 3D spatial coordinates: H and q . The blue lines show the corresponding results for the measurement in terms of 4D events: H_E and q_E . Without Λ both are equivalent. Gravity decelerates the expansion until it asymptotically brings it to a halt ($H \simeq H_E = 0$, $q \simeq q_E = -1/2$ with an EH: $R_{EH} \simeq \infty$). The effect of Λ according to the coordinate interpretation is to accelerate the expansion. While according to the proper distance R , it decelerates the expansion even further and brings it to an early halt: $H_E = 0$ and $q_E = -\infty$ at a finite horizon $R_{EH} = r_\Lambda$. This additional deceleration is caused by the friction term: $1/(HR_{EH})$ in Equation (15)-(16) (dashed-dot red line).

3. De-Sitter Phase

The FLRW metric with Λ , asymptotically tends to a constant: $H = H_\Lambda$ which corresponds to exponential inflation and de-Sitter metric, which can also be written as:

$$ds^2 = - \left(1 - r^2/r_\Lambda^2\right) dt^2 + \frac{dr^2}{1 - r^2/r_\Lambda^2} + r^2 d\Omega^2 \quad (6)$$

This form corresponds to a static 4D hypersphere of radius r_Λ . In this rest frame, events can only travel a finite distance $r \equiv a\chi < r_\Lambda \equiv 1/H_\Lambda$ within a static 3D surface of the imaginary 4D hypersphere.

This implies that there exists a frame duality, allowing us to equivalently describe de Sitter space either as static in proper or physical coordinates (t, r) , or as exponentially expanding in comoving coordinates (τ, χ) . In the static frame (t, r) , events are constrained within a limited region of the hypersphere, while in the expanding frame (τ, χ) , distances and coordinates evolve with time following an exponential expansion characterized by the de Sitter horizon r_Λ .

This frame duality can be understood as a Lorentz boost that results in both length contraction and time dilation. If define the coordinate $r = a\chi$ the radial velocity give us the Hubble law $\dot{r} = Hr$, leading to a Lorentz factor given by

$$\gamma = \frac{1}{\sqrt{1 - \dot{r}^2}} = \left(1 - \frac{r^2}{r_H^2}\right)^{-1/2}. \quad (7)$$

where $r_H = 1/H$. In the rest frame (t, r) , an observer sees the moving fluid element $ad\chi$ contracted by the Lorentz factor $1/\gamma$ in the radial direction and experiences a time dilation by γ : i.e. $dr = ad\chi/\gamma$ and $dt = \gamma d\tau$. More formally, we need to find a change of variables from comoving coordinates (τ, χ) in the FLRW metric of Equation (3) to rest frame deSitter coordinates (t, r) where $r = a(\tau)\chi$ (see [16]):

$$-d\tau^2 + a^2 d\chi^2 + a^2 \chi^2 d\Omega^2 = -(1 + 2\Psi)dt^2 + \frac{dr^2}{1 - r^2 H^2} + r^2 d\Omega^2, \quad (8)$$

which agrees with Equation (7) in [17] with:

$$(1 + 2\Psi) = \frac{a^2 \dot{a}^2}{1 - r^2 H^2} \left(\frac{d\tau}{d\sigma}\right)^2, \quad (9)$$

where $\sigma = \int_0^r \chi d\chi + \int_0^\tau \frac{dt}{a(t)\dot{a}(t)}$. This form reproduces the static deSitter metric Equation (6) when $H = H_\Lambda$. It also shows that t retains its time-like character as we cross inside $r = r_H = 1/H$. This is to be contrasted with the event horizon of the Schwarzschild metric, which requires a change of variables as we cross inside the horizon.

The spatial part of the light element in Equation (8) is illustrated in Figure 2. Geometrically it corresponds to the metric of a hypersphere of radius r_H that expands towards a constant radii $r_H \Rightarrow r_\Lambda$ which corresponds to an event horizon (see also Section 5 below and Appendix B in [16]). In the above rest (de-Sitter) frame, the FLRW background is asymptotically static, indicating no expansion or acceleration. While in the comoving frame there is cosmic acceleration ($q = 1$). This observation highlights that the concept of cosmic acceleration commonly used in cosmology critically depends on the chosen frame of reference.




Figure 2. The left panel shows a spatial representation of FLRW metric $ds^2 = (1 - r^2/r_H^2)^{-1}dr^2 + r^2d\theta^2$ in Equation (8) as a 2D metric $r^2 = x^2 + y^2$ (magenta plane) in polar coordinates (one angle is fixed) embedded in 3D flat space (where z in the plot corresponds to an extra dimension to illustrate the geometry). De-Sitter metric corresponds to the outer sphere $r_H = r_\Lambda$, while the FLRW metric is the blue sphere of radius r_H asymptotically expanding into r_Λ . The yellow/blue region shows the super/sub horizon causal regions. The right panel shows the same 2D (x, y) plane faced on, where we have also included (in red) the FLRW event Horizon R , discussed in Section 5 and Figure 4. Scales $r > R$ can never be reached from inside. Because $r > r_\Lambda$ is causally disconnected from the inside, this region should have the Schwarzschild metric with $r_s = r_\Lambda = 2GM$ if fully empty.

4. Event Acceleration

The interpretation of cosmic acceleration in Equation (5) is solely based on the definition for acceleration $\ddot{a}/a$ in Equation (5). We will show next, that such definition corresponds to events without a cause-and-effect connection and this lead us to the wrong picture of what is happening. We will then introduce a more physical alternative.

Consider the distance between two events corresponding to the light emission of a galaxy at (τ, χ) and the reception somewhere in its future (τ_1, χ_1) . The photon travels following an outgoing radial null geodesic $ds = 0$ which from Equation (3) implies $d\tau = a(\tau)d\chi$. This situation is depicted in Figure 3. We can define a 3D space-like distance d based in the comoving separation $\Delta\chi = \chi - \chi_1$:

$$d = a(\tau)\Delta\chi. \quad (10)$$

This is in fact the distance that corresponds to the acceleration given by $\ddot{a}/a$ in Equation (5), because $\dot{d}/d = \dot{a}/a$ and $\ddot{d}/d = \ddot{a}/a$, where the derivative is with respect to τ , the time at emission. Such distance corresponds to the distance between (τ, χ) and (τ, χ_1) , so that $d\tau = 0$. These events lack causal connection and are beyond observation. While using d isn't inherently incorrect, it involves extrapolating observed events (like luminosity distance) into non-observable realms. Essentially, d aligns with a non-local theory of Gravity or the Newtonian approximation, where action at a distance occurs with an infinite speed of light.




Figure 3. Comparison of different distances in the FLRW metric of Equation (3) between an observed event (at emission) at coordinates (τ, χ) and the corresponding null event (at reception) somewhere in its future (τ_1, χ_1) . The space-like distance $d = a(\tau)\Delta\chi$ in Equation (10), along the horizontal χ axis, is the one commonly used to define cosmic acceleration. It expands as $a(\tau)$, but is not causally connected. The distance $d_{LC} = c\Delta\tau$ in Equation (11), along the vertical τ axis, is the time-like distance traveled by light, but is independent of cosmic expansion $a(\tau)$. The event distance R in Equation (12) corresponds to the proper distance in the light-cone between the two events and is the one we should use to properly interpret cosmic expansion.

We could instead use the the distance traveled by the photon:

$$d_{LC} = \int_{\tau}^{\tau_1} a(\tau) d\chi = \int_{\tau}^{\tau_1} d\tau = \Delta\tau = (\tau_1 - \tau) \quad (11)$$

Note that we use units of $c = 1$, so that this should be read as $d_{LC} = c\Delta\tau$. But cosmic acceleration is zero $\ddot{d}_{LC} = 0$ for such distance because $\dot{d}_{LC} = -1$.

So the usual definition currently use by cosmologist, in Equation (10), corresponds to events that are space-like, i.e. at a fix comoving separation or fix cosmic time $d\tau = 0$. It only takes into account the change in the distance due to the expansion of the universe. To have a measurement of cosmic acceleration that is closer to actual observations, we need to use the distance between events that are causally connected, i.e. that not only takes into account how much the universe has expanded, but also how long it has taken for the two events to be causally connected.

To this end, we should use the proper future light-cone distance $R(\tau)$ obtained from $ds = 0$ in the FLRW of Equation (3) as $ds = 0 \Rightarrow d\chi = \int d\tau/a(\tau)$ and $R = ad\chi$ (see e.g. EquationA1 in [18]):

$$R(\tau, \tau_1) = a(\tau) \int_{\tau}^{\tau_1} d\chi = a(\tau) \int_{\tau}^{\tau_1} \frac{d\tau'}{a(\tau')} \quad (12)$$

Note that the term with the integral is not $\Delta\chi$, but it corresponds to the coordinate distance $d\chi$ travel by light between the two events including the effect of cosmic expansion. Thus, we argue that we should use R instead of d in Equation (10) as a measure of distance in cosmology to define cosmic acceleration and expansion rate. The difference between this 3 distances is illustrated in Figure 3.

Using R as a distance is equivalent to a simple change of coordinates in the FLRW metric of Equation (3), from comoving coordinates $d\chi$ to physical coordinates $dR = ad\chi$:

$$ds^2 = -d\tau^2 + dR^2 + R^2(\tau) d\Omega^2, \quad (13)$$

which is just Minkowski's metric in spherical coordinates with a radius $R = R(\tau)$.

We then have that: $\dot{R} = HR - 1$ and define the expansion rate between null events as:

$$H_E(\tau) \equiv \frac{\dot{R}}{R} = H \left(1 - \frac{1}{HR} \right) \quad (14)$$

where the additional term, $\frac{1}{HR}$ corresponds to a friction term. There is an ambiguity in this definition because R in Equation (12) depends also on the time τ_1 use to define R . To break this ambiguity we arbitrarily fix R to be the distance to $\tau_1 \Rightarrow \infty$ (which corresponds to a possible future Event Horizon):

$$H_E(\tau) = H \left(1 - \frac{1}{HR_{EH}} \right) \quad (15)$$

where $R_{EH} \equiv R[\tau, \tau_1 = \infty]$. The quantity $\frac{1}{HR_{EH}}$ corresponds to a friction term because it opposes the expansion. It just originates from change of frame. As we will see in next section, this choice implies that $\frac{1}{HR_{EH}}$ is zero unless $\Lambda \neq 0$. So this new invariant way to define cosmic expansion reproduces the standard definition when $\Lambda = 0$. But for $\Lambda > 0$ we have that the event expansion halts $H_E \Rightarrow 0$ (blue line in in Figure 1) due to the friction term (red line) for $a \gg 1$, while the standard Hubble rate definition approaches a constant $H \Rightarrow \sqrt{\Lambda/3} \equiv r_\Lambda^{-1}$ (black line). This might seem irrelevant at first look, but the resulting physical interpretation is quite different. In the standard definition, H , the expansion with Λ becomes asymptotically exponential (or inflationary expansion). While in our new definition, H_E , the expansion becomes static (as in the static de-Sitter metric).

The event acceleration can then be measured as:

$$q_E \equiv \frac{\ddot{R}}{R} \frac{1}{H_E^2} = \left(q - \frac{1}{HR_{EH}} \right) \left[1 - \frac{1}{HR_{EH}} \right]^{-2}. \quad (16)$$

The correct way to define a 4D acceleration in relativity is based on the geodesic deviation equation Equation (A1). The relation to q and q_E will be discussed in the Appendix.

As before, for $\Lambda = 0$ the friction term, $\frac{1}{HR_{EH}}$, makes little difference between q and q_E . For $\Lambda > 0$ the friction term asymptotically cancels the Λ term in $\frac{\ddot{a}}{a}$ (i.e. Equation (5)) so that $\frac{\ddot{R}}{R}$ is always negative, no matter how large is Λ ($HR_{EH} \Rightarrow 1$ and $q_E = -\infty$). The net effect of the Λ term is to bring the expansion of events to a faster stop ($H_E \Rightarrow 0$) than in the case with gravity alone. This is illustrated in Figure 1. The Λ term produces a faster deceleration (than with gravity alone). This corresponds to an attracting (and not repulsive) force, as explained in more detail in the Appendix.

5. Event Horizon

What is more relevant to understand the meaning of Λ is that the additional deceleration brings the expansion to a halt within a finite proper distance between the events, creating an Event Horizon (EH). The EH is the maximum distance that a photon emitted at time τ can travel following the outgoing radial null geodesic:

$$R_{EH}(a) = a \int_a^\infty \frac{da'}{H(a')a'^2} < \frac{1}{H[a = \infty]} = r_\Lambda \equiv \sqrt{3/\Lambda} \quad (17)$$

This is illustrated in Figure 4, which also demonstrates how inflation and the horizon problem (i.e., the observation that CMB measurements detect super-horizon $r > r_H$ frozen perturbations) occurred within R_{EH} . All the observable Universe (green line) is contained inside R_{EH} and we can therefore not measure anything outside.




Figure 4. Proper coordinate $r = a(\tau)\chi$ in units of c/H_0 as a function of cosmic time a (scale factor). The Hubble horizon $r_H = c/H$ (blue continuous line) is compared to the Λ future Event Horizon R_{EH} (red line) as defined in Equation (17). Larger radii (magenta shading) represent causally disconnected regions, while smaller ones (yellow shading), created during cosmic inflation, remain dynamically frozen. The full observable universe (dashed green line) encompasses both causal (blue shading) and frozen regions but is bounded by $r > R_{EH}$. Compared to Figure 2, which depicts a view at a fixed a using the same color coding. After inflation, r_H begins growing again, and by $a \approx 1$ (present epoch), both r_H and R_{EH} approach r_Λ (in black).

For $\Lambda = 0$ we have $R_{EH} = \infty$, so there is no EH. But for $\Lambda > 0$ we have that $R_{EH} \Rightarrow r_\Lambda$ (red line in Figure 1). We can then see that Λ corresponds to a causal horizon or boundary term. The analog force behaves like a rubber band between observed galaxies (null events) that prevents them crossing some maximum stretch (i.e. the EH). We can interpret such force as a boundary term that just emerges from the finite speed of light (see the Appendix).

The FLRW metric with Λ , asymptotically tends to the de-Sitter metric in Equation (6). This form corresponds to a static 4D hyper-sphere of radius r_Λ . So in this (rest) frame, events can only travel a finite distance $R < r_\Lambda$ within a static 3D surface of the imaginary 4D hyper-sphere. The region inside $r < r_\Lambda$ is causally disconnected from the outside. In the context of FLRW framework, this condition corresponds to $a < r_\Lambda/\chi$, where χ is a radial (space-like) distance. This condition implies that the expansion interpretation is valid only as long as $a < r_\Lambda/\chi$, indicating that it does not make sense for larger values where we cannot transition from $r < r_\Lambda$ to $r > r_\Lambda$. Essentially, beyond this threshold, the cosmological interpretation of expansion breaks down due to the causal disconnection imposed by the horizon defined at r_Λ .

As shown in Equation (8), this frame duality can be understood as a Lorentz boost. An observer in the rest frame, sees the moving fluid element $ad\chi$ contracted by the Lorentz factor γ . This duality is better understood using our new measures for the expansion rate H_E and cosmic deceleration q_E based on the distance between causal events.

6. Comparison to Data

We show next how to estimate the new measure of cosmic acceleration, q_E , using direct astrophysical observations. As an example consider the Supernovae Ia (SNIa) data as given by the 'Pantheon Sample' compilation ([19]) consisting on 1048 SNIas between $0.01 < z < 2.3$. Each SNIa provides a direct estimate of the luminosity distance $d_L(z)$ at a given measured redshift z . This corresponds to the comoving look-back distance:

$$\chi(z) = ad_L(z) = \int_0^z \frac{dz}{H(z)} \quad (18)$$

so that $\chi' \equiv d\chi/dz$ gives us directly $\chi' = r_H = H^{-1}$. The second derivative gives us the acceleration:

$$q = 1 + \frac{\chi''}{a\chi'} ; \quad q_E = \frac{q - \chi'/R_{EH}}{(1 - \chi'/R_{EH})^2}. \quad (19)$$

R_{EH} is given by the model prediction in Equation (17) (arbitrarily fixed at $\Omega_\Lambda = 0.85$ in both data and models). We adopt here the approach presented in [20], who used an empirical fit to the luminosity distance measurements, based on a third-order logarithmic polynomial:

$$\chi(a) = ad_L = a \left(x + Ax^2 + Bx^3 \right) H_0^{-1} \ln 10 \quad (20)$$

where $x = -\log_{10} a$. [20] find a good fit to: $A = 3.15 \pm 0.12$ and $B = 3.27 \pm 0.41$ to the full SNIa 'Pantheon Sample'. We use these values of A and B and its corresponding errors to estimate H , q and q_E using the above relations. Results for H and q are shown as shaded cyan regions in the left panel Figure 5. They are compared to the LCDM predictions in Equation (4) and (5).

There is a very good agreement in $H(z)$ for $\Omega_\Lambda \simeq 0.65$. At $z < 1$, the $q(z)$ estimates are also consistent with the $\Omega_\Lambda \simeq 0.65$ predictions. But the detailed $q(z)$ evolution with redshift in the SNIa data does not seem to follow any of the model predictions, specially for $z > 1$. The $q(z)$ estimates are too steep compare to the different models predictions. If we compare instead the q_E estimates (see right panel of Figure 5) we find a much better agreement with the model predictions. This seems to validate our q_E approach, but it is not clear from this comparison alone if this is caused by the fitting function use in Equation (20).

To test this further we use measurements of the radial BAO data to estimate q_E . Such measurements give us a direct estimate of $H(z)$ (as first demonstrated by [21]) so they have the advantage over SNIa that we only need to do a first order derivative, to estimate q or q_E :

$$q = 1 + \frac{1}{aH} \frac{dH}{dz} ; \quad q_E = \frac{q - r_H/R_{EH}}{(1 - r_H/R_{EH})^2}. \quad (21)$$

As an illustration, we utilize the $H(z)$ measurements presented in Table 2 of [22], which provide an expanded dataset including results up to 2023. This compilation of $H(z)$ values is shown as red points with 2σ error bars in the left panel of Figure 5, labeled as "<2023". Additionally, we incorporate four new data points from DESI2024 (LRG1, LRG2, LRG3+ELG1, and ELG2) derived from Table 18 of [23], using a sound horizon $r_o = 147$ Mpc from CMB measurements [24] to obtain $H(z)$, along with one data point from the Ly-alpha forest as described in Equation 7.3 of [25]. These are displayed as blue points with 2σ error bars in the left panel of Figure 5, labeled as "DESI2024".

The full compilation includes measurements from galaxy clustering ($z < 1.5$) and the Ly-alpha forest in quasars ($z > 2$). The combination of these distinct redshift ranges provides a robust measurement of $\frac{dH}{dz}$ at intermediate redshifts ($1 < z < 2$), where discrepancies in the supernova Type Ia data are observed when comparing the traditional q model and the q_E model (as discussed earlier). The radial BAO offers strong constraints on cosmic acceleration, independent of potential calibration errors in H_0 or sampling biases from small-area surveys. This level of precision is something

not yet achievable with current supernova datasets, but it will be exciting to see how upcoming wider and deeper surveys might address this issue in the near future.




Figure 5. Expansion rate $H(z)$ (upper left panel), cosmic acceleration q (lower left panel) and event acceleration q_E (right panel). Shaded areas correspond to a polynomial fit with 2σ region in a sample of SNIa (cyan) and radial BAO measurements (magenta). Dashed lines show the corresponding LCDM predictions for different values of Ω_Λ as labeled.

We fit a quadratic polynomial with inverse variance weighting to the radial BAO data:

$$H(z) = H_0 + H_1 z + H_2 z^2. \quad (22)$$

We have checked that the results presented here are very similar if we use a cubic polynomial. In units of Km/s/Mpc, we find $H_0 \simeq 69 \pm 3$, $H_1 \simeq 37 \pm 6$ and $H_2 \simeq 14 \pm 2$, with strong covariance between the errors (the cross-correlation coefficient between H_1 and H_2 is -0.965). The value of H_0 is in good agreement with the Planck CMB fit [24] but in some tension with the SNIa local calibration: $H_0 = 73 \pm 1$ (see [26]). This corresponds to either a local calibration problem (in SNIa, in radial BAO or in both) or a tension in the Λ CDM model at different times or distances (see e.g. [27]). We ignore this normalization problem here and just focus on the evolution of H/H_0 to measure cosmic acceleration q or q_E (which are fairly independent of H_0).

In the right panel of Figure 5 we show (as shaded regions) the measurements for q_E given by combining Equation (20) with Equation (19) and Equation (22) with Equation (21). The measurements clearly favour models with large negative cosmic event acceleration $q_E < 0$, which supports our interpretation of Λ as a friction term.

Comparing left and right panels in Figure 5 we see that both q and q_E are roughly consistent with models with $\Omega_\Lambda \simeq 0.7$ (or $\Lambda \simeq 2H_0^2$) in good concordance with $H(z)$ in the upper left panel of Figure 5.

Even when the underlying model for q and q_E is the same, note how the measured q and q_E data have different tensions with the model predictions as a function of redshift. In particular, the radial BAO and SNIa data sets show inconsistencies among them for q around $1.5 < z < 2.5$. This is a well known tension (see e.g. Figure 17 in [28]). This tension disappears when we use the corresponding estimates for q_E . Thus, data is more consistent with the q_E than with the q description.

One would expect that a perfect realizations of the LCDM model in Equation (4) would produce consistent results in both q and q_E . But deviations from LCDM and systematic effects can produce tensions in data, specially if we use a parametrization, like q , which refers to events that we never

observe. The q and q_E parametrization of acceleration are more general than the particular LCDM model and the fact that data prefers q_E is an important indication. Data lives in the light-cone, which corresponds to q_E rather than q . At $z \simeq 2$ the difference between a light-cone and space-like separations is very significant and any discrepancies in the data or model will show more pronounced in the q modeling.

We conclude that the data shows some tensions with LCDM predictions (as indicated by q) but confirms that cosmic expansion is clearly decelerating (as indicated by q_E) so that events are trapped inside an Event Horizon (R_{EH}).

7. Discussion & Conclusion

In our exploration, we've demonstrated that the commonly interpreted Λ term, thought to drive cosmic acceleration (as discussed in §2), actually leads to a quicker cosmic deceleration of events compared to the influence of gravity alone (as explained in §4). This relates to the nature of the Event Horizon (EH, see §5) that results from an expansion dominated by Λ . It suggests that Λ might not be a new form of dark or vacuum energy Λ_{DE} ([13–15]) or a modification of gravity Λ_F , but rather a boundary or surface term Λ_B in the corresponding action (see Equation (1)).

The measured R_{EH} in our cosmic expansion exhibits behavior analogous to the interior of a Schwarzschild Black Hole (BH), particularly under the assumption of a nearly empty space beyond R_{EH} (see [16]). This analogy is illustrated in Figure 2 and Figure 4, which challenge the conventional interpretation of the FLRW metric. The FLRW model assumes that the background density ρ remains constant both inside and outside R_{EH} , despite the absence of causal connections between these regions.

The absence of causality suggests that the universe could be inhomogeneous outside R_{EH} while remaining homogeneous within R_{EH} , as indicated by cosmic expansion. This concept diverges from the traditional horizon problem, as described by [3] and [29], in which the standard cosmological model envisions a homogeneous and isotropic universe fragmented into approximately 10^{83} causally disconnected regions in the past [30].

Inflationary theory, initially proposed by [31] and further developed by [30], [32], and [33], addresses the problem of cosmological fragmentation. It posits a period of exponential expansion driven by a state of energy characterized by the ground state of a field that translates into an effective negative pressure (similar to $\mathcal{L}_{DE} = p_{DE} = -\rho_{DE}$ in Equation (1)). This foundational phase allows all scales to initially exit the Hubble horizon (as depicted in the yellow regions of Figure 2 and Figure 4), and to re-enter post-inflation. Despite its success, the origins of this inflationary period remain elusive, posing a significant mystery in theoretical physics. Furthermore, inflationary theory does not address the causality outside R_{EH} (see Figure 4), raising critical considerations for our understanding of the universe's expansion.

The Event Horizon R_{EH} measured with q_E (i.e. Equation (17), which is equivalent to the presence of Λ) also tell us that there is a finite mass M_T trapped within R_{EH} . If we assume that the space outside R_{EH} is relatively empty, such finite mass M_T provides the explanation for the observed R_{EH} and therefore for Λ : i.e. $2GM_T = \sqrt{3/\Lambda}$, as resulting from a boundary term (see Equation (1)). This Black Hole Universe (BHU) model provides a new and completely classical explanation for the cosmological constant Λ within GR. It explains why Λ is so small but not zero: because M_T is so large, but not infinite. Yet it also raises new fundamental questions: If our local universe has fallen inside its own gravitational radius $r_s = 2GM_T = \sqrt{3/\Lambda}$, why is our universe now expanding and not collapsing?

The BHU interpretation, where the expansion happens inside R_{EH} (and is therefore a local solution [16]), opens the way to a new conjecture for the origin for cosmic expansion. Instead of emerging from a singular Big Bang (a global solution), it could result from the cold collapse of a large and low density

(local) cloud into a Black Hole¹. This is illustrated in Figure 6, reproduced from [9]. Unlike traditional models that lead to a singularity, this model suggests that a Big Bang-like explosion—termed the Big Bounce—prevents such an outcome. This Big Bounce could be driven by neutron degeneracy pressure, which occurs when densely packed neutron matter reaches a ground state governed by the Pauli exclusion principle. It could also be the result of a similar ground state happening at higher energies, like in standard Cosmic Inflation.




Figure 6. Illustration of the formation of a BHU, where $R = R_{EH}$ is the Event Horizon. Color regions $r < R$ (dashed red line) are filled with a uniform FLRW metric (blue corresponds to $r < r_H$ and yellow $r > r_H$) with empty space outside. A cloud of radius R and mass M collapses under gravity. When it reaches $R = r_s = 2GM$ it becomes a BH. The collapse proceeds inside the BH (where a frozen layer $r > r_H$ appears in yellow) until it bounces producing an expansion (the hot Big Bang). The event horizon r_s behaves like a cosmological constant with $\Lambda = 3/r_s^2$ so that the expansion freezes before it reaches back to $r_s = r_\Lambda \equiv 1/H_\Lambda$. The bottom panels shows the evolution of R and r_H during collapse and expansion in log units of $1/H_0$. Structure in-between R and r_H is frozen and seeds structure formation in our Universe, which could include smaller BHUs, CMB, stars and galaxies, and replaces the role of Inflation in Λ CDM (from [9]).

Mirroring the dynamics predicted by cosmic inflation, a ground state acts like a relativistic fluid with negative pressure in a closed cloud ($k > 0$). The combination of this two ingredients (positive curvature and positive background acceleration), not only halts the gravitational collapse but also catalyzes a fast rebound (exponential expansion that erases the original curvature), initiating the expansive phase of a flat Big Bang. The expansion drives the system away from the ground state and returns to a regular radiation and latter matter domination phases. This expansion will eventually be stopped by another quasi-deSitter phase, this time caused by the finite mass of the system $R_{EH} \rightarrow 2GM_T = \sqrt{3/\Lambda}$. Crucially, this quantum mechanism (Pauli exclusion principle) violates the strong energy condition $\rho + 3p \geq 0$ (but not the weak one $\rho + p \geq 0$) in Classical General Relativity

¹ Such collapse originates from a small initial over-density within a flat background, which can be modeled as a local FLRW closed curvature $k = 1/\chi_*^2 > 0$ solution, where the curvature radius χ_* is the comoving radius of the initial cloud. This important detail was overlooked in [7,9], who assumed a flat ($k = 0$) collapse. The equivalent case with curvature is presented in §12.5.1 of [34], with the only difference with respect to [7,9] of replacing χ_* by $\sin \chi_*$. The total relativistic mass M_T of the collapsing cloud is then given by $2GM_T = r_\Lambda = H_0^2 \sin^3 \chi_*$, which relates to both Λ (as a boundary term) and the initial curvature radius $k = 1/\chi_*^2$. As the collapse approaches the almost singular ground state, the curvature increases as $H^2 \simeq k/a^2$, which together with positive background acceleration (from the degenerate ground state) will enable the bounce to happen. After the bounce, the inflationary expansion erases the curvature term, while r_Λ will eventually dominate the bouncing expansion.

(GR) within a closed metric ($k > 0$) and sidesteps the singularity GR theorems proposed by Hawking and Penrose (e.g. [35]), thus presenting a novel solution to a pivotal issue in cosmological theory.

This idea is further validated by the observed large scale cut-off in the scale-invariance spectrum of metric perturbations, as observed in the CMB sky (see [36]). Such cut-off is measured to be 66 degrees, which corresponds to the R_{EH} radius at recombination (see dotted horizontal line in Figure 4) projected in the CMB sky. A recent research ([37]) has further revealed that several large-scale persistent CMB temperature anomalies originate from parity asymmetry $\mathcal{P}$. A groundbreaking explanation posits that the microscopic laws of quantum physics adhere to $\mathcal{PT}$ symmetry in a way that preserves causality and is promoted to curved spacetimes. Cosmic evolution disrupts the $\mathcal{T}$ symmetry, resulting in the observed $\mathcal{P}$ asymmetry. This idea was originally applied to inflationary quantum fluctuations but can be equally applied to the BHU Big Bounce picture above, as they are both defined by a period of quasi-deSitter expansion.

Additionally, we can conjecture from this notion that the interior dynamics of any other BH (e.g. stellar, binary or galactic) could also result from a similar BHU solution: a classical and non-singular, FLRW expanding interior (that becomes asymptotically deSitter, i.e. static in the rest frame). The mass (equivalent to Λ) boundary term in the BHU can then be interpreted as the actual physical mechanism that prevents anything from escaping the BH interior: i.e. it prevents the inside out crossing of the BH event horizon $r_S = 2GM = r_\Lambda$, which asymptotically results from $R_{EH} \rightarrow r_\Lambda$.

That the measured Λ term is fixed by the total mass M_T of our universe is in good agreement with the physical interpretation presented here that R_{EH} , in the rest frame, corresponds to a friction (attractive) force that decelerates cosmic events. In the Appendix we elaborate in this idea and revisit the Newtonian limit to show that Λ corresponds to an additional (attractive) Hooke's term to the inverse square gravitational law.

Funding: This work was partially supported by grants from Spain MCIN/AEI/10.13039/501100011033 grants PGC2018-102021-B-I00, PID2021-128989NB-I00 and Unidad de Excelencia María de Maeztu CEX2020-001058-M and from European Union funding LACEGAL 734374 and EWC 776247. IEEC is funded by Generalitat de Catalunya.

Data Availability Statement: No new data is presented.

Acknowledgments: I like to thank K. Sravan Kumar, Benjamin Camacho-Quevedo, Pablo Fosalba, Elizabeth Gonzalez and Pablo Renard for comments to the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Newtonian and Hookeon Limits

When we talk about classical forces we are making an analogy to Newton's law to gain some intuition on the physical problem. This is why we study next the role of Λ in the non-relativistic limit. Consider the geodesic acceleration $g^\mu = (g^0, g^i) = (g^0, \vec{g})$ defined from the geodesic deviation equation (see [34]):

$$g^\mu \equiv \frac{D^2 v^\mu}{D\tau^2} = R^\mu_{\alpha\beta\gamma} u^\alpha u^\beta v^\gamma, \quad (\text{A1})$$

where v^μ is the separation vector between neighbouring geodesics and u^α is the tangent vector to the geodesic. For an observer following the trajectory of the geodesic $u^\alpha = (1, 0)$ and $g^\alpha = (0, \vec{g})$:

$$g^i = R^i_{00\gamma} v^\gamma. \quad (\text{A2})$$

and we can choose the separation vector v^μ to be the spatial coordinate. The spatial divergence of $\vec{g}$ is then:

$$\vec{\nabla} \cdot \vec{g} = R^0_0 = -4\pi G(\bar{\rho} + 3\bar{p}) + \Lambda = 3\frac{\ddot{a}}{a}. \quad (\text{A3})$$

This equation is always valid for a comoving observer (see Equation 6.105 in [34]). Newtonian gravity is reproduced for the case of non-relativistic matter ($\bar{p}/c^2 \simeq 0$). The gravitational force (without Λ) is always attractive for $\bar{p} = 0$ (because $\bar{p} > 0$ and therefore $\vec{\nabla} \cdot \vec{g} < 0$) but it can be repulsive when

$\bar{p} < -\bar{\rho}/3$. For example, in the case of pure vacuum energy with $\Lambda = 0$, we have $p_{vac} = -\rho_{vac}$ and a repulsive gravitational force $\vec{\nabla}\vec{g} = +8\pi G\rho_{vac}$. The covariant version of Equation (A3) is the relativistic version of Poisson's Equation (see also [8,38]):

$$\nabla_{\mu}g^{\mu} = R_0^0 = -4\pi G(\bar{\rho} + 3\bar{p}) + \Lambda = 3\frac{\ddot{a}}{a}. \quad (\text{A4})$$

The solution to these equations is given by an integral over the usual propagators or retarded Green functions which account for causality.

This is also the Raychaudhuri equation for a shear-free, non-rotating fluid where $\Theta = \nabla_{\nu}u^{\nu}$ and u^{ν} is the 4-velocity:

$$\nabla_{\mu}g^{\mu} = \frac{d\Theta}{d\tau} + \frac{1}{3}\Theta^2 = R_{\mu\nu}u^{\mu}u^{\nu} = -4\pi G(\rho + 3p) + \Lambda \quad (\text{A5})$$

The above equation is purely geometric: it describes the evolution in proper time τ of the dilatation coefficient Θ of a bundle of nearby geodesics. Note that without Λ , the acceleration is always negative unless $p < -1/3\rho$ which is what we call DE today. This is degenerate with the Λ term for constant $p = -\rho$, so we can argue that Λ is a particular case of DE (but it can also be interpreted as a modify gravity term).

In the non-relativistic limit we see from Equation (A3) that indeed $\ddot{a}/a > 0$ corresponds to a repulsive force that dominates at large distances. For point like source:

$$\vec{g}(\vec{r}) = -\frac{GM\hat{r}}{r^2} + \frac{\Lambda}{3}\vec{r} \quad (\text{A6})$$

and acceleration can only be caused by Λ (see also [38,39]). Note how the linear term has the wrong sign compare to Hooke's law. It actually makes little sense to take the strict non-relativistic limit in Cosmology because in that limit, photons from different times will reach us instantly as in Equation (10). To make sense of observations we need to take into account the intrinsically relativistic effect that the speed of propagation is finite ($c = 1$). This corresponds to an additional term to the covariant acceleration $\nabla_{\mu}g^{\mu}$ which results in Equation (16). So besides gravitational deceleration, there is also a friction term proportional to H , caused by the expansion itself:

$$\vec{\nabla}\vec{g} = -4\pi G(\bar{\rho} + 3\bar{p}) + \Lambda - 3\frac{H}{R_{EH}} \quad (\text{A7})$$

So that the corresponding point like source is:

$$\vec{g}(\vec{r}) = -\frac{GM\hat{r}}{r^2} - \left(\frac{H}{R_{EH}} - \frac{\Lambda}{3}\right)\vec{r} \quad (\text{A8})$$

The negative friction term is always larger than the positive Λ term and asymptotically cancels it. This changes the sign of our interpretation of the role of Λ in terms of classical forces. The additional term has now the standard sign of Hooke's law in the above equation, so the effect of the Λ term could just be interpreted as a rubber band like force that prevents the crossing of the EH. We could summarize this as: Λ accelerates the 3D coordinate spatial expansion in $a(\tau)$ and this causes an additional deceleration in the expansion of events which results in an EH or a trapped surface.

References

1. Hilbert, D. Die Grundlage der Physik. *Konigl. Gesell. d. Wiss. Göttingen, Math-Phys K* **1915**, 3, 395–407.
2. Einstein, A. Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie. *S.K. Preußischen Akademie der W.* **1917**, pp. 142–152.
3. Weinberg, S. *Cosmology*, Oxford University Press; 2008.

4. York, J.W. Role of Conformal Three-Geometry in the Dynamics of Gravitation. *Phy.Rev.Lett.* **1972**, *28*, 1082–1085.
5. Gibbons, G.W.; Hawking, S.W. Cosmological event horizons, thermodynamics, and particle creation. *PRD* **1977**, *15*, 2738–2751.
6. Hawking, S.W.; Horowitz, G.T. The gravitational Hamiltonian, action, entropy and surface terms. *Class Quantum Gravity* **1996**, *13*, 1487.
7. Gaztañaga, E. The mass of our observable Universe. *MNRAS* **2023**, *521*, L59–L63.
8. Gaztañaga, E. The cosmological constant as a zero action boundary. *MNRAS* **2021**, *502*, 436–444.
9. Gaztañaga, E. The Black Hole Universe, Part II. *Symmetry* **2022**, *14*, 1984. doi:10.3390/sym14101984.
10. de Boer, J.; Dittrich, B.; Eichhorn, A.; Giddings, S.B.; Gielen, S.; Liberati, S.; Livine, E.R.; Oriti, D.; Papadodimas, K.; Pereira, A.D.; Sakellariadou, M.; Surya, S.; Verlinde, H. Frontiers of Quantum Gravity: shared challenges, converging directions. *e-prints* **2022**, p. arXiv:2207.10618.
11. DES Collaboration. DES Year 3 results: Cosmological constraints from galaxy clustering and weak lensing. *PRD* **2022**, *105*, 023520.
12. Huterer, D.; Turner, M.S. Prospects for probing the dark energy via supernova distance measurements. *PRD* **1999**, *60*, 081301.
13. Weinberg, S. The cosmological constant problem. *Reviews of Modern Physics* **1989**, *61*, 1–23.
14. Carroll, S.M.; Press, W.H.; Turner, E.L. The cosmological constant. *ARAA* **1992**, *30*, 499–542.
15. Peebles, P.J.; Ratra, B. The cosmological constant and dark energy. *Reviews of Modern Physics* **2003**, *75*, 559–606.
16. Gaztañaga, E. The Black Hole Universe, Part I. *Symmetry* **2022**, *14*, 1849. doi:10.3390/sym14091849.
17. Lochan, K.; Rajeev, K.; Vikram, A.; Padmanabhan, T. Quantum correlators in Friedmann spacetimes: The omnipresent de Sitter spacetime and the invariant vacuum noise. *Phys. Rev. D* **2018**, *98*, 105015, [arXiv:gr-qc/1805.08800]. doi:10.1103/PhysRevD.98.105015.
18. Ellis, G.F.R.; Rothman, T. Lost horizons. *American Journal of Physics* **1993**, *61*, 883–893.
19. Scolnic, D.M.; Jones, D.O.; Rest, A.; Pan, Y.C.; Chornock, R.; Foley, R.J.; Huber, M.E.; Kessler, R.; Narayan, G.; Riess, A.G.; Rodney, S.; Berger, E.; Brout, D.J.; Challis, P.J.; Drouot, M.; Finkbeiner, D.; Lunnan, R.; Kirshner, R.P.; Sanders, N.E.; Schlafly, E.; Smartt, S.; Stubbs, C.W.; Tonry, J.; Wood-Vasey, W.M.; Foley, M.; Hand, J.; Johnson, E.; Burgett, W.S.; Chambers, K.C.; Draper, P.W.; Hodapp, K.W.; Kaiser, N.; Kudritzki, R.P.; Magnier, E.A.; Metcalfe, N.; Bresolin, F.; Gall, E.; Kotak, R.; McCrum, M.; Smith, K.W. The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample. *ApJ* **2018**, *859*, 101.
20. Liu, Y.; Cao, S.; Biesiada, M.; Lian, Y.; Liu, X.; Zhang, Y. Measuring the Speed of Light with Updated Hubble Diagram of High-redshift Standard Candles. *ApJ* **2023**, *949*, 57.
21. Gaztañaga, E.; Cabré, A.; Hui, L. Clustering of luminous red galaxies - IV. Baryon acoustic peak in the line-of-sight direction and a direct measurement of $H(z)$. *MNRAS* **2009**, *399*, 1663–1680.
22. Niu, J.; Chen, Y.; Zhang, T.J. Reconstruction of the dark energy scalar field potential by Gaussian process. *e-prints* **2023**, p. arXiv:2305.04752.
23. DESI Collaboration. DESI 2024 III: Baryon Acoustic Oscillations from Galaxies and Quasars. *arXiv e-prints* **2024**, p. arXiv:2404.03000, [arXiv:astro-ph.CO/2404.03000]. doi:10.48550/arXiv.2404.03000.
24. Planck Collaboration. Planck 2018 results. VI. Cosmological parameters. *A&A* **2020**, *641*, A6.
25. DESI Collaboration. DESI 2024 IV: Baryon Acoustic Oscillations from the Lyman Alpha Forest. *arXiv e-prints* **2024**, p. arXiv:2404.03001, [arXiv:astro-ph.CO/2404.03001]. doi:10.48550/arXiv.2404.03001.
26. Riess, A.G. The expansion of the Universe is faster than expected. *Nature Reviews Physics* **2019**, *2*, 10–12.
27. Abdalla, E.; et al. Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies. *JHEA* **2022**, *34*, 49–211.
28. Bautista, J.; et al. Measurement of baryon acoustic oscillation correlations at $z = 2.3$ with SDSS DR12 $\text{Ly}\alpha$ -Forests. *A&A* **2017**, *603*, A12.
29. Rindler, W. Visual horizons in world models. *MNRAS* **1956**, *116*, 662.
30. Guth, A.H. Inflationary universe: A possible solution to the horizon and flatness problems. *PRD* **1981**, *23*, 347–356.
31. Starobinskiĭ, A.A. Spectrum of relict gravitational radiation and the early state of the universe. *Soviet J. of Exp. and Th. Physics Letters* **1979**, *30*, 682.
32. Linde, A.D. A new inflationary universe scenario. *Physics Letters B* **1982**, *108*, 389–393.

33. Albrecht, A.; Steinhardt, P.J. Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking. *Phy.Rev.Lett.* **1982**, *48*, 1220–1223.
34. Padmanabhan, T. *Gravitation*, Cambridge U. Press; 2010.
35. Hawking, S.W.; Penrose, R.; Bondi, H. The singularities of gravitational collapse and cosmology. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences* **1970**, *314*, 529–548.
36. Gaztañaga, E.; Camacho-Quevedo, B. What moves the heavens above? *Physics Letters B* **2022**, *835*, 137468. doi:10.1016/j.physletb.2022.137468.
37. Gaztañaga, E.; Sravan Kumar, K. Finding origins of CMB anomalies in the inflationary quantum fluctuations. *JCAP* **2024**, p. in Press. doi:10.48550/arXiv.2401.08288.
38. Gaztañaga, E. The size of our causal Universe. *MNRAS* **2020**, *494*, 2766–2772.
39. Calder, L.; Lahav, O. Dark energy: back to Newton? *Astronomy & Geophysics* **2008**, *49*, 1.13–1.18.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.