

Article

Not peer-reviewed version

---

# Development of qPCR and LAMP Assays for the Surveillance and Diagnosis of Herpes B Virus Infection

---

Murasaki Amano , Krittiga Sapkanarak , Wipaporn Thbthimthong , Suthirote Meesawat , Taratorn Kemthong , Nutchanat Suttisan , [Haruka Abe](#) , Suchinda Malaivijitnond , [Jiro Yasuda](#) \*

Posted Date: 26 September 2023

doi: [10.20944/preprints202309.1757.v1](https://doi.org/10.20944/preprints202309.1757.v1)

Keywords: Herpes B virus; zoonosis; LAMP; surveillance; cynomolgus macaque



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

## Article

# Development of qPCR and LAMP Assays for the Surveillance and Diagnosis of Herpes B Virus Infection

Murasaki Amano <sup>1, 2, 3</sup>, Krittiga Sapkanarak <sup>4, 5</sup>, Wipaporn Thbthimthong <sup>4</sup>, Suthirote Meesawat <sup>4, 5</sup>, Taratorn Kemthong <sup>4, 5</sup>, Nutchanat Suttisan <sup>4</sup>, Haruka Abe <sup>1, 6</sup>, Suchinda Malaivijitnond <sup>4, 5</sup> and Jiro Yasuda <sup>1, 2, 3, \*</sup>

<sup>1</sup> Department of Emerging Infectious Diseases, National Research Center for the Control and Prevention of Infectious Diseases (CCPID), Nagasaki University, Nagasaki 852-8523, Japan; bb55420102@ms.nagasaki-u.ac.jp (M.A.); abeh@nagasaki-u.ac.jp (H.A.)

<sup>2</sup> Department of Viral Emerging Infectious Diseases, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki 852-8523, Japan

<sup>3</sup> Department of Emerging Infectious Diseases, Institute of Tropical Medicine, Nagasaki University, Nagasaki 852-8523, Japan.

<sup>4</sup> National Primate Research Center of Thailand, Chulalongkorn University, Saraburi 18110, Thailand; Krittiga.S@chula.ac.th (K.S.); Wipaporn.T@chula.ac.th (W.T.); Suthirote.m@gmail.com (S.M.); Taratorn.K@chula.ac.th (T.K.); Nutchanatbell@gmail.com (N.S.); Suchinda.m@chula.ac.th (S.M.)

<sup>5</sup> Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

<sup>6</sup> Vietnam Research Station, Institute of Tropical Medicine (NEKKEN), Nagasaki University, Nagasaki 852-8523, Japan

\* Correspondence: j-yasuda@nagasaki-u.ac.jp .

**Abstract:** Herpes B virus (BV) is a zoonotic virus which can be transmitted from macaques to humans, which is often associated with high mortality rates. Because macaques often exhibit asymptomatic infections, individuals who come into contact with these animals face unexpected risks of BV infections. Serological test is widely performed to investigate BV infections. However, the assay's sensitivity and specificity appeared to be inadequate, and it does not necessarily indicate ongoing viral shedding. Here, we developed LAMP and qPCR assays aiming to detect BVs with high sensitivity and specificity in various macaque species and validated them using oral swab samples collected from 97 wild cynomolgus macaques living in Thailand. Our LAMP and qPCR assays detected more than 50 and 10 copies of the target sequences per reaction, respectively. The LAMP assay could detect BV within 25 min, indicating its advantages for the rapid detection of BV. Collectively, our findings indicated that both assays developed in this study exhibit advantages and usefulness for BV surveillance and diagnosis of BV infections in macaques. Furthermore, for the first time, we determined the partial genome sequences of BVs detected in cynomolgus macaques in Thailand. Phylogenetic analysis revealed the species-specific evolution of BV within macaques.

**Keywords:** Herpes B virus; zoonosis; LAMP; surveillance; cynomolgus macaque

## 1. Introduction

Herpes B virus (*Macacine alphaherpesvirus* 1; BV) belongs to the genus *Simplexvirus*, subfamily *Alphaherpesvirinae*, and is a zoonotic pathogen that causes a life-threatening disease in humans. The natural hosts of BV are macaques, and BV infection is usually asymptomatic in natural hosts. BV can be transmitted from infected macaques to humans via direct contact with body fluids, scratches, or bites, and results in fatal human central nervous system diseases, including encephalitis and encephalomyelitis [1]. Macaque monkeys are abundant in Southeast Asian countries. In Thailand, cynomolgus macaque or long-tailed macaque (*Macaca fascicularis*) and rhesus macaque (*Macaca mulatta*) are widely distributed and reside close to human residential areas [2], thereby posing a significant risk to public health. BV persists in trigeminal nerves and spinal ganglion of macaques throughout their lives. Viral shedding can potentially occur, especially in conjunction with



immunosuppression, breeding season, stress, and primary infection [3]. The virus can be secreted into the saliva, urine, and genitalia during the viral reactivation [4]. Previous study has shown that seropositivity was more than 50% in rhesus macaques, and 60% in cynomolgus macaques reared in laboratory animal facilities in Japan [5]. Moreover, 31 out of 38 (81.6%) wild adult rhesus macaques were BV seropositive in Bali [6]. Regardless of their habitats, the BV seroprevalence of adult macaques was higher than that of the young macaques [6]. Recently, the use of non-human primates for experiments has been gradually increasing every year for the evaluation of the efficacies of candidate drugs or vaccines. This is primarily due to the genetic, physiological, and immunological characteristics shared between humans and primates [7]. Therefore, workers who handle macaque monkeys are at a high risk of acquiring these viruses. Indeed, on June 30, 2021, a human infection case involving BV was reported in China. The patient eventually died two months after dissecting dead monkeys in a research institute [8], highlighting the importance of reliable BV detection before performing procedures with monkeys. Misdiagnosis and delay in treatment with patients may lead to fatal diseases in humans.

Thus, to evaluate the risk of BV transmission from macaques to humans in areas where humans and macaques live close, such as laboratory animal facilities, assays that can detect BV with high accuracy and sensitivity can be used for the surveillance and regular inspection of BV are required. At present, the standard assay used for the detection of BV involves viral isolation from clinical samples. However, a biosafety level (BSL)-3 or BSL-4 containment facility is required to propagate BV in cell lines, such as Vero cells and Hela cells [9] [10]. Moreover, it takes approximately one to seven days to isolate BV [11]. Although enzyme-linked immunosorbent assay (ELISA) has been commonly used in primate facilities as the serological method to detect the antibodies against BV, it can detect only a history of BV infection, but does not provide information on the shedding of the virus. Furthermore, detecting antibodies against BV in cases of recent infections is challenging, and there is a cross-reactivity between BV and other herpesviruses, including Herpes Simplex Virus (HSV) 1 and 2 [12]. Quantitative real-time PCR (qPCR) may be the most sensitive and accurate assay to diagnose BV infection. However, it has not gained widespread adoption as a surveillance assay, primarily due to the requirement for sophisticated instrumentation and stable electric power. In addition, previously reported primers for qPCR were designed based on the BV E2490 strain isolated from rhesus macaque [13], suggesting that it may be challenging to identify BV from other macaque species[14]. Therefore, a simple and rapid BV detection method is essential. One of the alternative methods to qPCR is the loop-mediated isothermal amplification (LAMP) assay, which amplifies the target DNA without thermal cycling steps. LAMP uses four to six primers, which increases the efficiency of target genome amplification [15]. We developed rapid and sensitive LAMP assays for the detection of SARS-CoV-2, ebolaviruses, Marburgvirus, Zika virus, Lassa virus, foot-and-mouth disease virus, bovine papular stomatitis virus [16–26]. Additionally, the LAMP assay is a simple technique with a battery-powered equipment, which doesn't require other advanced equipment, and can be made available in mobile clinics, as well as remote areas [18,19,27].

In this study, we developed the LAMP and qPCR assays, which can be employed to detect all BV strains identified so far. Both assays were validated using the samples collected from wild and captive cynomolgus macaques in Thailand.

## 2. Materials and Methods

### 2.1. Ethics statement

Sample collection from wild cynomolgus macaques was approved by the Department of National Parks, Wildlife, and Plant Conservation of Thailand and the Institutional Animal Care and Use Committee (IACUC) of the National Primate Research Center of Thailand-Chulalongkorn University (Protocol Review No. 2075007 and 2375008).

### 2.2. Sample collection and DNA extraction

Ninety-seven cynomolgus monkeys were captured from the central and southern areas of Thailand, and oral swab and blood samples were collected, and the animal health was checked. The oral cells and saliva were harvested from the inside of both cheeks, the base of the tongue and the soft palate. To minimize animal distress, anesthetics with 2-5 mg/kg body weight of Tiletamine HCl and zolazepam HCl (Virvac, France) were intramuscularly administered to all monkeys during specimen collection. All swabs were kept in a screw-cap tube containing a DNA/RNA shield (ZYMO Research, USA) to stabilize the DNAs until use. Following the manufacturer's protocol, whole DNA was extracted from swab samples using a QIAamp DNA Mini Kit (QIAGEN, Germany). The final elution volume was 150 µl.

### 2.3. Preparation of the BV DNA fragments

Synthesized DNAs based on the sequences of 248 bp of UL27 (1,850-2,097), 164 bp of gG (303 – 466), and 257 bp of UL29 (1,093 – 1,349) were purchased from Eurofins Genomics, Japan. Plasmids containing these BV sequences were constructed using a Zero Blunt TOPO PCR Cloning Kit (Invitrogen, USA). Plasmids were digested with *Eco*RI. DNA fragments were electrophoresed on agarose gels and purified using the QIAquick Gel Extraction Kit (QIAGEN). The purified DNAs were serially diluted with distilled water from 1×10<sup>6</sup> to 1×10<sup>0</sup> DNA copies/µl.

### 2.4. qPCR for validation of viral DNA quantity

The qPCR primer sequences are listed in Table 1. Primer sequences were based on the conserved sequences of the BV strains (accession No. KY628968-77, KY628979-85, and KJ566591). The TaqMan probe for BV detection was fluorescently labeled with 6-carboxyfluorescein (FAM) at the 5' ends and 6-carboxytetramethylrhodamine (TAMRA) at the 3' ends. All probes were synthesized by Eurofins Genomics (JAPAN). qPCR was performed using a QuantiStudio™ 5 qPCR system (Applied Biosystems, USA). The reaction mixture (10 µl) contained 5 µl of 2× TaqMan™ Fast Advanced Master Mix; 1 µl of 10× primer-probe mixture, including 0.9 µM forward and reverse primers and 0.2 µM probe, 3 µl of water, and 1 µl of extracted DNA. The reaction was conducted with a thermal cycle program of 50°C for 2 min, 95°C for 20 sec, followed by 40 cycles of 95°C for 1 sec, and 60°C for 20 sec. The cut-off values were set at the threshold cycle (C<sub>t</sub>) value of 40. To quantify the viral DNA, a standard curve was created with 10-fold serial dilutions of the purified BV DNA fragment possessing the qPCR target sequence. The assay was performed in duplicate for each sample.

**Table 1.** The sequences of qPCR and LAMP primers.

Primer set	Primer	Sequence (5'-3')
qPCR-gG	BV-qPCR-gG-323_Fw	TGGCCTACTACCGCGTGG
	BV-qPCR-gG-446_Rv	TGGTACGTGTTGGAGTAGCG
	BV-qPCR-gG-403_Probe	CCGCCCTCTCCGAGCACGTG
qPCR-UL29	BV-qPCR_UL29_Fw	TTGTCCGACTTCTGGTAGGC
	BV-qPCR_UL29_Rv	AAGGAGATGCGCGTCAAG
	BV-qPCR_UL29_Probe	CCGGCRAAGACCACCGCGTT
BV-gG_1st	BV-gG_Fw_1	GACCCCGCGTACTGCTAC
	BV-gG_Rv_1	CCCACCAGGATCTCGTAGTC
BV-gG_nested	BV-gG_Fw_2	GCCGAYGTCGACAGACAT
	BV-gG_Rv_1	CCCACCAGGATCTCGTAGTC
LAMP-UL27	LAMP_UL27_F3	CGCGCTACTTCACCTTC
	LAMP_UL27_B3	GGACCTCGGTGTAGTCAG
	LAMP_UL27_FIP	GTGCTGACCGTGGTCACGTCGCGGGCTAC
	LAMP_UL27_BIP	GTGTACTTTG
		CTCAACCTCACGATGCTCGAGGCCGCTGT
		CCTTGATCTCCT

---

LAMP_UL27_LF	GTGGGAGTACGCGTAATCCT
LAMP_UL27_LB	ACCACGAGTTCGTGCC

---

### 2.5. LAMP primer design

LAMP primers were designed based on the sequences of the conserved regions among the BV strains. The whole BV genome sequence was aligned using the MAFFT version 7 online software (<https://mafft.cbrc.jp/alignment/server/>) to identify conserved regions. Primer Explorer V5 software (Eiken, Japan; <http://primerexplorer.jp/>) was used to design LAMP primers. The LAMP method requires a set of six primers (F3, B3, FIP, BIP, LF, and LB) that recognize six different sequences in the target DNA.

### 2.6. LAMP reaction

LAMP assays were performed using a Genelyzer FIII real-time fluorescence detection platform (Canon Medical Systems, Japan), as previously described [16]. LAMP reaction mixture (25  $\mu$ l) included 15  $\mu$ l of Isothermal Master Mix (ISO004, Canon Medical Systems), 0.8  $\mu$ M each of FIP and BIP primer, 0.4  $\mu$ M each of F and B loop primers, 0.2  $\mu$ M each of F3 and B3, 2.5  $\mu$ l of water, and 5  $\mu$ L of each sample. The reaction was performed at 65°C for 40 min, followed by a dissociation analysis at 95-75°C with a temperature change rate of 0.1°C/s. Non-specific positive samples were excluded by comparing the melting temperature to that of the positive control. The sample was considered positive when its fluorescence intensity was > 20,000 within 30 min of amplification. The assay was performed in duplicate for each sample.

### 2.7. Sanger sequencing to confirm the conserved primer regions among BV strains

To examine whether the viral genome sequences of primer target regions are conserved, partial genomes, including the primer binding sites of BV-positive samples, were amplified by PCR using the primer sets shown in Table S2. Nested-PCR was performed using PrimeSTAR GXL DNA Polymerase (Takara Bio, Japan) under the following conditions: 30 cycles each of 10 sec at 98°C, 15 sec at 55°C, and 1min/kb at 68°C. All the PCR products were electrophoresed on agarose gels and purified from the gels using the QIAquick Gel Extraction Kit. The purified DNAs were processed using the BigDye Terminator v3.1 Cycle Sequencing kit (Thermo Fisher Scientific) and analyzed with the SeqStudio Genetic Analyzer. The collected sequences were identified using BLAST searches (<https://blast.ncbi.nlm.nih.gov>). The obtained sequences were deposited in the DNA Data Bank of Japan (DDBJ) database under the following accession numbers: LC777613-LC777618.

### 2.8. BV phylogenetic analysis

To characterize the BV strains, No. 8-14 and No. 8-17, identified in this study, the region of gJ (US5) to gD (US6) genes (720 or 723 bp), which is a hypervariable region of the BV genome, was amplified by PCR using the primer sets shown in Table S2. The PCR products were purified from the gels and then sequenced as described above. The phylogenetic analysis of the sequence data was performed using BioEdit version 7.2.5 and MEGA 7 software. A phylogenetic tree was constructed using the neighbor-joining method. The neighbor-joining tree was constructed at 1,000 bootstraps. The sequence data of No. 8-14 and No. 8-17 were deposited in DDBJ under the accession numbers LC777619 and LC777620, respectively.

## 3. Results

### 3.1. Development of a novel qPCR assay to detect BV


The qPCR method for detecting BV has been previously reported by Perelygina et al. [14]. The primer set for qPCR (hereafter called qPCR-gG) was designed based on the glycoprotein G (gG) region sequence of the BV strain E2490 isolated from the rhesus macaque (Macaca mulatta) (Table 1).

We analyzed the sequences of the binding region of the qPCR-gG primers in all the BV strains deposited in GenBank, including the strains isolated from other macaque species; KY628982 from Japanese macaque (*Macaca fuscata*), KY628968 from lion-tailed macaque (*Macaca silenus*), KY628985 bonnet macaque (*Macaca radiata*), and KY628969-70 from southern pig-tailed macaque (*Macaca nemestrina*), and found a nucleotide mismatch at the 3' end of the forward primer (G to C) and the probe (G to C) in the BV strains isolated from cynomolgus macaques (*Macaca fascicularis*) (Figure 1). The mismatches at the 3' end of primers plausibly affected the sensitivity of the assay. Therefore, to detect all the known BV strains with high sensitivity, we designed a new qPCR primer set targeting the UL29 region, which is conserved among all the BV strains (Figure 2). The mixed nucleotide probe was used in the qPCR, because a single nucleotide diversity (A or G) existed at the sixth nucleotide of the probe-binding region between the strains isolated from the cynomolgus and rhesus macaques.

We compared the sensitivities of qPCR-gG and the newly designed qPCR (qPCR-UL29), using plasmids containing the primer-binding sequences of the E2490 strain isolated from rhesus macaque or the E90-136 strain isolated from cynomolgus macaque. The detection limit of qPCR-gG was 10 copies or 100-1,000 copies for E2490 and for E90-136, respectively, whereas that of qPCR-UL29 was 10 copies for both E2490 and E90-136 (Table 2). Thus, qPCR-UL29 was more sensitive than qPCR-gG for the detection of E90-136, although both qPCR assays showed similar sensitivities for the detection of E2490. This suggests that qPCR-UL29 is more suitable for detecting a broad range of BV strains.



**Figure 1.** Sequence alignments of qPCR-gG primer target region of BV genome. The accession number of the genome sequence and the macaque origin of each BV strain are shown on the left.



**Figure 2.** Sequence alignments of qPCR-UL29 primer target region of BV genome. The accession number of the genome sequence and the macaque origin of each BV strain are shown on the left.

**Table 2.** Evaluation of qPCR-gG and qPCR-UL29 assays using DNA fragments based on either the E2490 or E90-136 strain.

DN A	E2490					E90-136					
	qPC R	qPCR-gG		qPCR-UL29		qPCR-gG	qPCR-UL29		qPCR-gG	qPCR-UL29	
No. of DNA copy	Ct value	Pos itiv e	Ct value	Pos itiv e	Ct value	Pos itiv e	Ct value	Pos itiv e	Ct value	Pos itiv e	
1x10 <sup>6</sup>	21. 16	21. 18	2/2	20. 68	20. 37	2/2	23. 16	23. 01	2/2	18. 99	18. 96
1x10 <sup>5</sup>	24. 65	24. 47	2/2	23. 33	22. 88	2/2	27. 49	27. 57	2/2	22. 38	22. 5
1x10 <sup>4</sup>	27. 91	27. 71	2/2	26. 33	26. 2	2/2	30. 48	30. 46	2/2	25. 77	26. 1
1x10 <sup>3</sup>	31. 21	30. 93	2/2	29. 76	29. 7	2/2	33. 28	33. 76	2/2	29. 36	30. 31
1x10 <sup>2</sup>	34. 89	34. 74	2/2	33. 12	33. 06	2/2	39. 3	-	1/2	33. 17	33. 3
1x10 <sup>1</sup>	37. 52	37. 12	2/2	38. 49	38. 3	2/2	-	-	0/2	38. 58	38. 3
1x10 <sup>0</sup>	-	-	0/2	-	-	0/2	-	-	0/2	-	-

### 3.2. Evaluation of the qPCR assay using monkey specimens

To further evaluate the qPCR assay for BV detection, 97 oral swab samples from the cynomolgus macaques were examined using qPCR-gG. Nineteen samples tested positive, which was further confirmed by employing a semi-nested PCR assay using an in-house primer set (Table 3). However, amplification of the target region was detected in only two samples, No. 8-14 and No. 8-17, among 19 qPCR-gG-positive samples (Figure S1), suggesting that the remaining 17 samples were false-positives detected via qPCR-gG. Subsequently, we evaluated the qPCR-UL29 assay using the same 97 oral specimens. The assay specifically detected BV in two samples, No. 8-14 and No. 8-17, which had shown positive results in both qPCR-gG and nested PCR (Table 3). The Ct values of No. 8-14 and No. 8-17 were 38.3 (1.25 copies) and 33.0 (15.95 copies) in qPCR-gG and 28.0 (12,748 copies) and 24.6 (177,970 copies) in qPCR-UL29, respectively (Table 3). This confirmed the fact that qPCR-UL29 exhibits high sensitivity and specificity in comparison to qPCR-gG.


**Table 3.** The results of qPCR-UL29 and nested PCR-gG for qPCR-gG positive samples.

Sample ID	qPCR-gG		qPCR-UL29		Nested PCR-gG
	Average of Ct value	Estimated DNA copy number	Average of Ct value	Estimated DNA copy number	
8-1	36.17	3.23	-	-	-
8-2	36.93	1.99	-	-	-
8-3	36.39	2.81	-	-	-
8-4	34.95	7.02	-	-	-
8-5	35.78	4.14	-	-	-
8-6	38.14	0.92	-	-	-
8-7	37	1.91	-	-	-
8-8	36.43	2.74	-	-	-
8-9	36.67	2.35	-	-	-
8-10	37.85	0.58	-	-	-
8-11	37.11	0.95	-	-	-
8-12	37.58	0.69	-	-	-
8-13	37.36	0.8	-	-	-
8-14	38.3	1.25	28	12,748	+
8-15	36.95	1.06	-	-	-
8-16	36.11	1.87	-	-	-
8-17	33	15.95	24.6	177,970	+
8-18	35.14	3.56	-	-	-
8-19	35.58	2.65	-	-	-

### 3.3. Development of the LAMP assay for BV

To establish a rapid and sensitive detection assay for BV, we developed the LAMP assay. LAMP primers were designed based on sequences within the UL27 region, which is conserved among BV strains. Among the primers, only the B3 primer had one nucleotide mismatch with the sequences of BV isolates other than rhesus macaque (Figure 3). However, this mismatch exhibited a minimal effect on the sensitivity because of its location near the 5'-terminus (Figure 3).

The LAMP assay was performed using DNA fragments containing primer-binding sequences of the E2490 strain isolated from rhesus macaque or E90-136 strain isolated from cynomolgus macaque. As shown in Table 4, the detection limit of the LAMP assay was 50 copies per reaction for both E2490 and E90-136. The assay detected more than 50 copies of the E2490 and E90-136 plasmid within 25 and 19 min, respectively, indicating that the LAMP assay exhibited rapidly and high sensitivity for the detection of BV.



**Figure 3.** Sequence alignments of LAMP primer target regions of BV genome. The accession number of the genome sequence and the macaque origin of each BV strain are shown on the left.

**Table 4.** Sensitivity of LAMP assay using DNA fragments based on either the E2490 or E90-136 strain.

DNA	E2490		E90-136	
	Amplification time (mm:ss)		Amplification time (mm:ss)	
DNA copies/reaction	Run 1	Run 2	Run 1	Run 2
5x10 <sup>6</sup>	6:46	6:31	8:16	7:46
5x10 <sup>5</sup>	7:16	7:16	9:31	9:01
5x10 <sup>4</sup>	12:01	8:16	10:31	10:16
5x10 <sup>3</sup>	15:16	9:31	12:31	12:16
5x10 <sup>2</sup>	15:16	11:16	19:16	16:16
5x10 <sup>1</sup>	24:16	21:31	17:46	18:16
5x10 <sup>0</sup>	-	-	-	-

#### 3.4. Validation of the LAMP assay using monkey specimens

To validate the LAMP assay, 97 oral swabs of wild cynomolgus macaques determined in the qPCR assays mentioned above were used (Table 5). Only two samples, No. 8-14 and No. 8-17, were

detected to be positive in the LAMP and qPCR-UL29 assays, indicating that the LAMP assay results were consistent with those of the qPCR-UL29 assay (Table 6). The average time for the detection of positivity was 13 min 8 sec and 9 min 54 sec for No.8-14 and No.8-17, respectively.

**Table 5.** Evaluation of the qPCR-UL29 and LAMP assays using monkey samples.

		LAMP assay	
		No. of positive samples	No. of negative samples
qPCR assay	No. of positive samples	2/97	0/97
	No. of negative samples	0/97	95/97

**Table 6.** The BV detection time of LAMP assay in positive samples.

Sample ID	LAMP-UL27		
	Amplification time (mm:ss)	Average (mm:ss)	
8-14	12:31	13:46	13:08
8-17	9:46	10:01	9:54

### 3.5. Sequence analyses for the primer binding sites

To investigate whether the primer sequences had mismatched sequences with the selected genes of the BV strains from the specimens of Thai cynomolgus macaques, we determined the sequences of the primer binding sites of No. 8-14 and No. 8-17. The qPCR-UL29 primers did not exhibit a mismatch in both No. 8-14 and No. 8-17 strains (Figure 2), while the LAMP primers possessed two nucleotide mismatches at the sixth position from the 5'-terminus of the F3 primer and fifth position from the 3'-terminus of the LF primer in both No. 8-14 and No. 8-17 strains (Figure 3). In addition, No. 8-14 strain exhibited another mismatch at the eighth position from the 3'-terminus of the B2 primer region. Therefore, the DNA fragment containing the sequence of the primer-binding region of No. 8-14 was prepared as described in the Materials and Methods section and used to evaluate the sensitivity of the LAMP assay. The detection limit was determined to be 50 copies (Table 7).


**Table 7.** The evaluation of LAMP sensitivity using fragmented DNA based on No.8-14 strain.

DNA	DNA based on No. 8-14	
	Amplification time (mm:ss)	
DNA copies/reaction	Run 1	Run 2
5x10 <sup>6</sup>	8:31	8:46
5x10 <sup>5</sup>	10:01	10:13
5x10 <sup>4</sup>	11:46	11:16
5x10 <sup>3</sup>	14:01	13:46
5x10 <sup>2</sup>	17:31	16:46
5x10 <sup>1</sup>	22:01	-
5x10 <sup>0</sup>	-	-

### 3.6. Phylogenetic analysis of BV strains isolated from Thai cynomolgus macaques

In this study, we detected BV in two Thai cynomolgus macaques. To date, genomic data on BV in macaques living in Southeast Asian countries have not been reported. Additionally, BV genomic data from cynomolgus macaques are limited, even globally. To characterize the BV strains detected in this study, we determined the sequences of gJ (US5) through gD (US6) genes (720 or 723 bp), which is a hypervariable region of the BV genome [28], and performed phylogenetic analysis. The phylogenetic tree showed three distinct lineages: rhesus and Japanese macaques (*M. mulatta* and *M. fuscata*), cynomolgus macaques (*M. fascicularis*), and pig-tailed macaques (*M. nemestrina*). BV strains derived from Thai cynomolgus macaques from our study clustered with the cynomolgus BV strain E90-136, as previously reported (accession No. AF082811; Figure 4). Among our cynomolgus

macaque's BV strains, No. 8-14 strain was slightly different from No. 8-17 strain which was clustered with BV strain E90-136. This might be because these BV strains were isolated from different subspecies of cynomolgus macaques; No. 8-14 strain was from common cynomolgus macaque (*M. fascicularis fascicularis*) and No. 8-17 strain was from Burmese cynomolgus macaque (*M. fascicularis aurea*). Unfortunately, we do not know the subspecies level of cynomolgus macaque infected with E90-136 strain. Noted, the BV genomes from cynomolgus macaques were entirely different from the larger clade of BV strains of rhesus and Japanese macaques.



**Figure 4.** Phylogenetic analysis for the BV strains from cynomolgus macaques in Thailand and other BV strains. LC777619 and LC777620 are the BVs identified in this study. The US5 (gJ) open reading frame (ORF) to US6 (gD) ORF sequence is amplified by PCR, and the product has been sequenced. The neighbor-joining tree is constructed at 1,000 bootstraps.

#### 4. Discussion

In this study, we developed the qPCR-UL29 assay that can efficiently detect BVs originating from both rhesus and cynomolgus monkeys, as well as the LAMP assay that can rapidly and specifically detect BVs. Both assays were practical for BV detection in macaque oral specimens. Although the multiplex LAMP assay which can detect both BV and Mpox virus simultaneously, was recently reported to demonstrate high specificity and sensitivity for the detection of BV, it was evaluated only using samples containing BV plasmid in BV negative monkey serum due to the limited availability of monkey specimens [29]. To the best of our knowledge, this is the first study to evaluate a LAMP assay for BV using monkey oral samples. The results of the LAMP assay were consistent with those of the qPCR assay, suggesting that the LAMP assay was a reliable method for BV detection.

The detection limit of the LAMP assay was 50 copies/reaction, whereas that of the qPCR-UL29 assay was 10 copies/reaction (Tables 2 and 4), indicating that the qPCR-UL29 assay exhibited a five-fold higher sensitivity than the LAMP assay. This may be due to the high GC content of the BV genome (74% G+C composition), which plausibly impact the efficiency of the LAMP amplification. In the complicated reaction steps of LAMP, GC-rich targets may cause unexpected dimers between the primers themselves or between the primers and amplicons. According to a previous study, viral shedding of HSV-2 frequently occurred at rates of  $10^{5.2}$  copies/ml ( $10^{2.2}$  copies/ $\mu$ l) and exhibited high infectivity [30,31]. Therefore, our LAMP assay displays adequate sensitivity to detect a transmissible number of BV. LAMP assay has several advantages, including simple experimental procedures, faster

detection rate than the qPCR, and the use of portable equipment [32], suggesting that the LAMP assay proves to be a valuable method for point-of-care diagnosis and field surveillance.

In this study, we identified two BV-positive samples, No. 8-14 and No. 8-17, in oral swab specimens from wild cynomolgus monkeys living in Thailand. The detection limit of the assay was 50 copies, which is comparable to a recent report [29], implying high sensitivity in the case of monkey specimens. Although the BV genome from cynomolgus macaques in Thailand had a sequence mismatch in the F3, LF, and BIP primer regions (Figure 3), the BV genome was appropriately detected in sample No. 8-14 as positive by our LAMP assay. A recent report showed that a sequence mismatch located near the 3' end of the FIP and BIP primers affects the sensitivity of the assay [33]. However, the sequence mismatch of primers had minimal effect on the sensitivity of our LAMP assay, since our primer set did not exhibit a mismatch near the 3' end of FIP and BIP.

Regarding the qPCR assay, the sequences of the qPCR-UL29 primers had no mismatch with the BV 8-14 and 8-17 strains identified in Thai cynomolgus macaques in this study and all the BV strains collected from at least seven macaque species of which the genome sequences have been registered in GenBank so far (Figure 2), suggesting that the qPCR-UL29 assay could be applied to most species of macaques.

BV is an important zoonotic disease that is transmitted from macaques to humans and shows a high mortality rate. In Southeast Asian countries, macaques live close to human residential areas in many towns. The transmission of BV from wild macaques in these towns and areas is a serious public health concern. Therefore, BV surveillance in macaques would be necessary in the context of "One Health."

Collectively, our findings indicated that the qPCR and LAMP assays developed in this study are useful for the diagnosis of BV for any macaque monkeys and the surveillance of BV in monkey populations.

Furthermore, we determined the partial genome sequences of two BV strains, No. 8-14 and No. 8-17, identified in Thai cynomolgus macaques. This is the first report of BV genome sequences from Thai cynomolgus macaque. Phylogenetic analysis showed that both the Thai strains belonged to the same clade as the E90-136 strain, which had previously been isolated from cynomolgus macaque reared in the primate facility in the US. Interestingly, the two BV strains (Nos. 8-14 and 8-17) were separated into different clades based on the host subspecies level (*M. fascicularis fascicularis* and *M. fascicularis aurea*). Recently, it was reported that the genetic characteristics, based on genetic markers of mtDNA, Y chromosome gene *SRY* and *TSPY*, whole genome sequences and autosomal SNPs, are very distinctive between these two subspecies [34–37]. This reflects that each clade of BVs has independently evolved in a specific macaque species/subspecies [38]. This was also confirmed by a big picture that *M. fascicularis*, *M. fuscata* and *M. mulatta* which belonged to the same species group of *fascicularis group* was separated from *M. nemestrina* which belonged to *silenus-sylvanus group* [39]. Thus, the susceptibility and the co-evolution of the host and BV's genetics should be taken into account.

**Supplementary Materials:** The following supporting information can be downloaded at: [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1); Figure S1: Nested PCR for gG-qPCR positive samples; Table S1: Sequences of the primers for conventional PCR and sequencing.

**Author Contributions:** Conceptualization, J.Y.; methodology, M.A., H.A., S.M., and J.Y.; validation, M.A., K.S., S.M., and J.Y.; formal analysis, M.A., K.S., W.T., H.A., S.M., and J.Y.; investigation, M.A., K.S., W.T., and J.Y.; resources, K.S., W.T., S.M., T.K., N.S., and S.M.; writing—original draft preparation, M.A.; writing—review and editing, H.A., S.M., and J.Y.; supervision, J.Y.; project administration, S.M. and J.Y.; funding acquisition, S.M. and J.Y. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by grants from the Japan Agency for Medical Research and Development (AMED) (JP23fm0208101 to JY) and the Japan Society for the Promotion of Science (JSPS)(JPJSCCB20220009 to JY), and the Research Fund Senior Scholar (RTA6280010 to SM).

**Institutional Review Board Statement:** The animal study protocol was approved by Animal Care and Use Committee (IACUC) of the National Primate Research Center of Thailand-Chulalongkorn University (protocol

review nos. 2075007 and 2375008) (Date of approval: 18<sup>th</sup> April 2020 and 28<sup>th</sup> Jun 2023) for studies involving animals.

**Data Availability Statement:** All available data are presented in the article.

**Acknowledgments:** We would like to thank all staff of the National Primate Research Center of Thailand, Chulalongkorn University for providing biological specimens and assistance. We are grateful to Ms. Miku Takano for the logistical management and Drs. Yasuteru Sakurai, Rokusuke Yoshikawa, and Takaaki Kinoshita, as well as all staff of Department of Emerging Infectious Disease, CCPID, for advice and discussions.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Jainkittivong, A.; & Langlais, R. P. Herpes B Virus Infection. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endodontology* **1998**, *85*, 399–403. [https://doi.org/10.1016/S1079-2104\(98\)90064-6](https://doi.org/10.1016/S1079-2104(98)90064-6).
2. Gumert, M. D. The Common Monkey of Southeast Asia: Long-Tailed Macaque Populations, Ethnophoresy, and Their Occurrence in Human Environments. In *Monkeys on the Edge: Ecology and Management of Long-Tailed Macaques and their Interface with Humans*; Fuentes, A., Jones-Engel, L., Gumert, M. D., Eds.; Cambridge Studies in Biological and Evolutionary Anthropology; Cambridge University Press: Cambridge, 2011; pp 3–44. <https://doi.org/10.1017/CBO9780511974434.003>.
3. Lee, M.-H.; Rostal, M. K.; Hughes, T.; Sitam, F.; Lee, C.-Y.; Japning, J.; Harden, M. E.; Griffiths, A.; Basir, M.; Wolfe, N. D.; Epstein, J. H.; & Daszak, P. Macacine Herpesvirus 1 in Long-Tailed Macaques, Malaysia, 2009–2011 - Volume 21, Number 7—July 2015 - Emerging Infectious Diseases Journal - CDC. **2015**, *21*, 1107–1113. <https://doi.org/10.3201/eid2107.140162>.
4. Estep, R. D.; Messaoudi, I.; & Wong, S. W. Simian Herpesviruses and Their Risk to Humans. *Vaccine* **2010**, *28*, B78–B84. <https://doi.org/10.1016/j.vaccine.2009.11.026>.
5. Sato, H.; Arikawa, J.; Furuya, M.; Kitoh, J.; Mannen, K.; Nishimune, Y.; Ohsawa, K.; Serikawa, T.; Shibahara, T.; Watanabe, Y.; Yagami, K.; Yamamoto, H.; & Yoshikawa, Y. Prevalence of Herpes B Virus Antibody in Nonhuman Primates Reared at the National University of Japan. *Exp. Anim.* **1998**, *47*, 199–202. <https://doi.org/10.1538/expanim.47.199>.
6. Engel, G. A.; Jones-Engel, L.; Schillaci, M. A.; Suaryana, K. G.; Putra, A.; Fuentes, A.; & Henkel, R. Human Exposure to Herpesvirus B-Seropositive Macaques, Bali, Indonesia. *Emerg. Infect. Dis.* **2002**, *8*, 789–795. <https://doi.org/10.3201/eid0808.010467>.
7. Grimm, D. Record number of monkeys being used in U.S. research. <https://www.science.org/content/article/record-number-monkeys-being-used-us-research> (accessed 2023-06-08).
8. Wang, W.; Qi, W.; Liu, J.; Du, H.; Zhao, L.; Zheng, Y.; Wang, G.; Pan, Y.; Huang, B.; Feng, Z.; Zhang, D.; Yang, P.; Han, J.; Wang, Q.; & Tan, W. First Human Infection Case of Monkey B Virus Identified in China, 2021. *China CDC Wkly.* **2021**, *3*, 632–633. <https://doi.org/10.46234/ccdcw2021.154>.
9. Sabin, A. B.; & Wright, A. M. ACUTE ASCENDING MYELITIS FOLLOWING A MONKEY BITE, WITH THE ISOLATION OF A VIRUS CAPABLE OF REPRODUCING THE DISEASE. *J. Exp. Med.* **1934**, *59*, 115–136.
10. Cohen, J. I.; Davenport, D. S.; Stewart, J. A.; Deitchman, S.; Hilliard, J. K.; Chapman, L. E.; & the B Virus Working Group. Recommendations for Prevention of and Therapy for Exposure to B Virus (*Cercopithecine Herpesvirus 1*). *Clin. Infect. Dis.* **2002**, *35*, 1191–1203. <https://doi.org/10.1086/344754>.
11. *Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis*; Arvin, A., Campadelli-Fiume, G., Mocarski, E., Moore, P. S., Roizman, B., Whitley, R., Yamanishi, K., Eds.; Cambridge University Press: Cambridge, 2007.
12. Fujima, A.; Ochiai, Y.; Saito, A.; Omori, Y.; Noda, A.; Kazuyama, Y.; Shoji, H.; Tanabayashi, K.; Ueda, F.; Yoshikawa, Y.; & Hondo, R. Discrimination of Antibody to Herpes B Virus from Antibody to Herpes Simplex Virus Types 1 and 2 in Human and Macaque Sera. *J. Clin. Microbiol.* **2008**, *46*, 56–61. <https://doi.org/10.1128/JCM.00342-07>.
13. Hilliard, J. K.; Eberle, R.; Lipper, S. L.; Munoz, R. M.; & Weiss, S. A. Herpesvirus Simiae (B Virus): Replication of the Virus and Identification of Viral Polypeptides in Infected Cells. *Arch. Virol.* **1987**, *93*, 185–198. <https://doi.org/10.1007/BF01310973>.
14. Perelygina, L.; Patrusheva, I.; Manes, N.; Wildes, M. J.; Krug, P.; & Hilliard, J. K. Quantitative Real-Time PCR for Detection of Monkey B Virus (*Cercopithecine Herpesvirus 1*) in Clinical Samples. *J. Virol. Methods* **2003**, *109*, 245–251. [https://doi.org/10.1016/S0166-0934\(03\)00078-8](https://doi.org/10.1016/S0166-0934(03)00078-8).
15. Notomi, T. Loop-Mediated Isothermal Amplification of DNA. *Nucleic Acids Res.* **2000**, *28*, 63e–663. <https://doi.org/10.1093/nar/28.12.e63>.

16. Yoshikawa, R.; Abe, H.; Igasaki, Y.; Negishi, S.; Goto, H.; & Yasuda, J. Development and Evaluation of a Rapid and Simple Diagnostic Assay for COVID-19 Based on Loop-Mediated Isothermal Amplification. *PLoS Negl. Trop. Dis.* **2020**, *14*, e0008855. <https://doi.org/10.1371/journal.pntd.0008855>.
17. Kurosaki, Y.; Takada, A.; Ebihara, H.; Grolla, A.; Kamo, N.; Feldmann, H.; Kawaoka, Y.; & Yasuda, J. Rapid and Simple Detection of Ebola Virus by Reverse Transcription-Loop-Mediated Isothermal Amplification. *J. Virol. Methods* **2007**, *141*, 78–83. <https://doi.org/10.1016/j.jviromet.2006.11.031>.
18. Kurosaki, Y.; Magassouba, N.; Oloniniyi, O. K.; Cherif, M. S.; Sakabe, S.; Takada, A.; Hirayama, K.; & Yasuda, J. Development and Evaluation of Reverse Transcription-Loop-Mediated Isothermal Amplification (RT-LAMP) Assay Coupled with a Portable Device for Rapid Diagnosis of Ebola Virus Disease in Guinea. *PLoS Negl. Trop. Dis.* **2016**, *10*, e0004472. <https://doi.org/10.1371/journal.pntd.0004472>.
19. Kurosaki, Y.; Magassouba, N.; Bah, H. A.; Soropogui, B.; Doré, A.; Kourouma, F.; Cherif, M. S.; Keita, S.; & Yasuda, J. Deployment of a Reverse Transcription Loop-Mediated Isothermal Amplification Test for Ebola Virus Surveillance in Remote Areas in Guinea. *J. Infect. Dis.* **2016**, *214*, S229–S233. <https://doi.org/10.1093/infdis/jiw255>.
20. Kurosaki, Y.; Grolla, A.; Fukuma, A.; Feldmann, H.; & Yasuda, J. Development and Evaluation of a Simple Assay for Marburg Virus Detection Using a Reverse Transcription-Loop-Mediated Isothermal Amplification Method. *J. Clin. Microbiol.* **2010**, *48*, 2330–2336. <https://doi.org/10.1128/JCM.01224-09>.
21. Kurosaki, Y.; Martins, D. B. G.; Kimura, M.; Catena, A. dos S.; Borba, M. A. C. S. M.; Mattos, S. da S.; Abe, H.; Yoshikawa, R.; de Lima Filho, J. L.; & Yasuda, J. Development and Evaluation of a Rapid Molecular Diagnostic Test for Zika Virus Infection by Reverse Transcription Loop-Mediated Isothermal Amplification. *Sci. Rep.* **2017**, *7*, 13503. <https://doi.org/10.1038/s41598-017-13836-9>.
22. Kurosaki, Y.; Okada, S.; Nakamae, S.; & Yasuda, J. A Loop-Mediated Isothermal Amplification Assay for Rapid and Sensitive Detection of Bovine Papular Stomatitis Virus. *J. Virol. Methods* **2016**, *238*, 42–47. <https://doi.org/10.1016/j.jviromet.2016.07.031>.
23. Fukuma, A.; Kurosaki, Y.; Morikawa, Y.; Grolla, A.; Feldmann, H.; & Yasuda, J. Rapid Detection of Lassa Virus by Reverse Transcription-Loop-Mediated Isothermal Amplification: Detection of Lassa Virus by RT-LAMP. *Microbiol. Immunol.* **2011**, *55*, 44–50. <https://doi.org/10.1111/j.1348-0421.2010.00286.x>.
24. Pemba, C. M.; Kurosaki, Y.; Yoshikawa, R.; Oloniniyi, O. K.; Urata, S.; Sueyoshi, M.; Zadeh, V. R.; Nwafor, I.; Iroeziindu, M. O.; Ajayi, N. A.; Chukwubike, C. M.; Chika-Igwensi, N. M.; Ndu, A. C.; Nwidi, D. U.; Maehira, Y.; Unigwe, U. S.; Ojide, C. K.; Onwe, E. O.; & Yasuda, J. Development of an RT-LAMP Assay for the Detection of Lassa Viruses in Southeast and South-Central Nigeria. *J. Virol. Methods* **2019**, *269*, 30–37. <https://doi.org/10.1016/j.jviromet.2019.04.010>.
25. Oloniniyi, O. K.; Kurosaki, Y.; Miyamoto, H.; Takada, A.; & Yasuda, J. Rapid Detection of All Known Ebolavirus Species by Reverse Transcription-Loop-Mediated Isothermal Amplification (RT-LAMP). *J. Virol. Methods* **2017**, *246*, 8–14. <https://doi.org/10.1016/j.jviromet.2017.03.011>.
26. Howson, E. L. A.; Kurosaki, Y.; Yasuda, J.; Takahashi, M.; Goto, H.; Gray, A. R.; Mioulet, V.; King, D. P.; & Fowler, V. L. Defining the Relative Performance of Isothermal Assays That Can Be Used for Rapid and Sensitive Detection of Foot-and-Mouth Disease Virus. *J. Virol. Methods* **2017**, *249*, 102–110. <https://doi.org/10.1016/j.jviromet.2017.08.013>.
27. Huang, S.; Do, J.; Mahalanabis, M.; Fan, A.; Zhao, L.; Jepeal, L.; Singh, S. K.; & Klapperich, C. M. Low Cost Extraction and Isothermal Amplification of DNA for Infectious Diarrhea Diagnosis. *PLOS ONE* **2013**, *8*, e60059. <https://doi.org/10.1371/journal.pone.0060059>.
28. Smith, A. L.; Black, D. H.; & Eberle, R. Molecular Evidence for Distinct Genotypes of Monkey B Virus (Herpesvirus Simiae) Which Are Related to the Macaque Host Species. *J. Virol.* **1998**, *72*, 9224–9232. <https://doi.org/10.1128/JVI.72.11.9224-9232.1998>.
29. Zeng, Y.; Zhao, Y.; Ren, X.; Zhou, X.; Zhang, C.; Wan, Z.; & Kuang, Y.-Q. Rapid Detection of Monkeypox Virus and Monkey B Virus by a Multiplex Loop-Mediated Isothermal Amplification Assay. *J. Infect.* **2023**, *86*, e114–e116. <https://doi.org/10.1016/j.jinf.2023.02.003>.
30. Schiffer, J. T.; Wald, A.; Selke, S.; Corey, L.; & Magaret, A. The Kinetics of Mucosal Herpes Simplex Virus-2 Infection in Humans: Evidence for Rapid Viral-Host Interactions. *J. Infect. Dis.* **2011**, *204*, 554–561. <https://doi.org/10.1093/infdis/jir314>.
31. Schiffer, J. T.; Mayer, B. T.; Fong, Y.; Swan, D. A.; & Wald, A. Herpes Simplex Virus-2 Transmission Probability Estimates Based on Quantity of Viral Shedding. *J. R. Soc. Interface* **2014**, *11*, 20140160. <https://doi.org/10.1098/rsif.2014.0160>.
32. Oliveira, B. B.; Veigas, B.; & Baptista, P. V. Isothermal Amplification of Nucleic Acids: The Race for the Next “Gold Standard.” *Front. Sens.* **2021**, *2*, 752600. <https://doi.org/10.3389/fsens.2021.752600>.
33. Tamanaha, E.; Zhang, Y.; & Tanner, N. A. Profiling RT-LAMP Tolerance of Sequence Variation for SARS-CoV-2 RNA Detection. *PLOS ONE* **2022**, *17*, e0259610. <https://doi.org/10.1371/journal.pone.0259610>.
34. Bunlungsup, S.; Imai, H.; Hamada, Y.; Matsudaira, K.; & Malaivijitnond, S. Mitochondrial DNA and Two Y-Chromosome Genes of Common Long-Tailed Macaques (*Macaca Fascicularis Fascicularis*) throughout Thailand and Vicinity. *Am. J. Primatol.* **2016**, *79*, e22596. <https://doi.org/10.1002/ajp.22596>.

35. Matsudaira, K.; Ishida, T.; Malaivijitnond, S.; & Reichard, U. H. Short Dispersal Distance of Males in a Wild White-Handed Gibbon ( *Hylobates Lar* ) Population. *Am. J. Phys. Anthropol.* **2018**, *167*, 61–71. <https://doi.org/10.1002/ajpa.23603>.
36. Osada, N.; Matsudaira, K.; Hamada, Y.; & Malaivijitnond, S. Testing Sex-Biased Admixture Origin of Macaque Species Using Autosomal and X-Chromosomal Genomic Sequences. *Genome Biol. Evol.* **2020**, *evaa209*. <https://doi.org/10.1093/gbe/evaa209>.
37. Phadphon, P.; Kanthaswamy, S.; Oldt, R. F.; Hamada, Y.; & Malaivijitnond, S. Population Structure of *Macaca Fascicularis Aurea*, and Their Genetic Relationships with *M. f. Fascicularis* and *M. Mulatta* Determined by 868 RADseq-Derived Autosomal SNPs—A Consideration for Biomedical Research. *J. Med. Primatol.* **2022**, *51*, 33–44. <https://doi.org/10.1111/jmp.12554>.
38. Kolb, A. W.; & Brandt, C. R. Genomic Nucleotide-Based Distance Analysis for Delimiting Old World Monkey Derived Herpes Simplex Virus Species. *BMC Genomics* **2020**, *21*, 436. <https://doi.org/10.1186/s12864-020-06847-w>.
39. Fooden, J. Provisional Classifications and Key to Living Species of Macaques (Primates: Macaca). *Folia Primatol. Int. J. Primatol.* **1976**, *25*, 225–236. <https://doi.org/10.1159/000155715>.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.