

Article

Not peer-reviewed version

In Vitro and In Vivo Evaluation for Antioxidant and Anti-Diabetic Properties of *Cyperus rotundus* L. Kombucha

[Ananya Dechakhamphu](#) , Nattapong Wongchum , Theeraphan Chumroenphat , Alongklod Tanomtong ,
[Somchai Pinlaor](#) , [Sirithon Siriamornpun](#) *

Posted Date: 19 October 2023

doi: 10.20944/preprints202310.1201.v1

Keywords: functional food; health benefit; herbal medicine; health promotion



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

In Vitro and In Vivo Evaluation for Antioxidant and Anti-Diabetic Properties of *Cyperus rotundus* L. Kombucha

Ananya Dechakhamphu ^{1,2}, Nattapong Wongchum ³, Theeraphan Chumroenphat ^{1,2}, Alongklod Tanomthong ⁴, Somchai Pinlaor ⁵ and Sirithon Siriamornpun ^{6,*}

¹ Thai Traditional Medicine Program, Faculty of Thai Traditional and Alternative Medicine, Ubon Ratchathani Rajabhat University, Ubonratchathani 34000 Thailand

² Aesthetic Sciences and Health Program, Faculty of Thai Traditional and Alternative Medicine, Ubon Ratchathani Rajabhat University, Ubonratchathani 34000 Thailand

³ Biology Program, Faculty of Science, Ubon Ratchathani Rajabhat University, Ubonratchathani 34000 Thailand

⁴ Biology Program, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand

⁵ Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand

⁶ Research Unit of Thai Food Innovation, Department of Food Technology and Nutrition, Mahasarakham University, Kantarawichai, Maha Sarakham 44150, Thailand

* Correspondence: sirithon.s@msu.ac.th

Abstract: *Cyperus rotundus* L. exhibits promising potential for the development of functional foods due to its documented pharmacological and biological activities. This study investigated the antioxidant and anti-diabetic properties of *C. rotundus* kombucha. The results demonstrated potent antioxidant activity with an IC₅₀ value of 76.7±9.6 µL/mL for the DPPH assay and 314.2±16.9 µL/mL for the ABTS assay. Additionally, the kombucha demonstrated alpha-glucosidase inhibitory with an IC₅₀ value of 142.7±5.2 µL/mL. This in vitro antioxidant potential was further validated in vivo using *Drosophila*. *Drosophila* fed a high-sugar diet and supplemented with pure kombucha revealed significant increases in DPPH and ABTS free radical scavenging activity. *Drosophila* on a high-sugar diet supplemented with varying kombucha concentrations manifested enhanced resistance to oxidative stresses induced by H₂O₂ and paraquat. Concurrently, there was a notable decline in lipid peroxidation levels. Additionally, significant upregulations in CAT, SOD1, and SOD2 activities were observed when the high-sugar diet was supplemented with kombucha. Furthermore, in vivo assessments using *Drosophila* demonstrated significant reductions in alpha-glucosidase activity when fed with kombucha. A comprehensive GC-MS and HPLC analysis of *C. rotundus* kombucha detected the presence of antioxidative and anti-glucosidase compounds. In conclusion, *C. rotundus* kombucha exhibits considerable antioxidant and anti-diabetic properties, demonstrating its potential as a beneficial beverage for health promotion.

Keywords: functional food; health benefit; herbal medicine; health promotion

1. Introduction

Cyperus rotundus L. from the Cyperaceae family, also known as purple nutsedge, is documented as a longevity remedy in ancient Thai manuscripts. Historically, this herb has been employed in traditional medicinal practices across China, India, Africa, Japan, and Arab nations to treat various ailments [1]. Numerous studies have demonstrated that the rhizomes and tubers of this plant contain a broad spectrum of medicinal properties. These include antioxidant [2–4], anti-diabetic [5,6], and anti-obesity effects [7]. They also have antiallergic [8], antimicrobial [9], and antidiarrheal capabilities [10]. Moreover, they offer cardioprotective [11], gastroprotective [12], and hepatoprotective benefits [13]. The plant also exhibits immunomodulatory [14], neuroprotective [15,16], and anticarcinogenic properties [17]. Additionally, it has been found to have antiarthritic [18–20], anti-inflammatory, anti-uropathogenic [21], anticonvulsant [22], and antidepressant effects [23]. Recently reports have revealed that the hydroethanolic extract of *C. rotundus* enhances sexual behavior and fitness in *Drosophila* [24]. The supplementation of *Drosophila* with this extract resulted in an increase in lifespan

and a decrease in oxidative stress-induced mortality [25]. In a *Drosophila* obesity model, supplementation with *C. rotundus* extract exhibited to extend lifespan in the presence of high-fat diet-induced mortality [26]. Given its demonstrated biological and pharmacological properties, this plant's rhizomes and tubers have the potential to be used in the development of functional foods and beverages. These products may offer viable alternatives for disease prevention and therapeutic interventions.

The suppression of alpha-glucosidase is essential for managing type-2 diabetes, a condition characterized by high blood glucose, insulin resistance, and inadequate insulin response. By decelerating the conversion of carbohydrates into simple sugars, alpha-glucosidase suppressants can reduce post-meal glucose increases, facilitating in maintaining of steady blood sugar levels and functioning as an effective diabetes treatment. In addition to their utility in the treatment of diabetes, alpha-glucosidase inhibitors have attracted attention for their therapeutic potential in treating a variety of diseases. Some findings indicate that a delayed carbohydrate digestion process might increase the feeling of fullness, potentially reducing overall calorie consumption and nurturing weight reduction. Currently, the majority of anti-glucosidase action research is conducted in in vitro settings, which may not replicate the complexity of living conditions. Animal research should be intensive in order to discover and validate promising drug candidates for the management of diabetes.

In scientific literature, kombucha is acknowledged for its putative health benefits, which range from improved digestive and immune function to cholesterol level modulation and detoxification processes [27,28]. Although empirical research addressing these assertions is still in its beginning, the beverage's rising popularity is influenced by the global trend toward natural and probiotics-enriched foods and beverages. Numerous in vitro studies have investigated the bioactivities of kombucha and its constituents, but the number of published in vivo studies is significantly lesser. To obtain a comprehensive understanding, it is essential to expand in vivo studies, conduct detailed bio-accessibility and bioavailability assessments, and precisely identify bioactive compounds. These initiatives will facilitate a thorough investigation of distinctive molecular functions.

Drosophila melanogaster or fruit flies is recognized as a model organism for investigating metabolic disorders. This organism has contributed to the development of diabetes models that replicate the complexities of type-2 diabetes in humans. A technique for inducing T2D symptoms in *Drosophila* via high-sugar diets has been developed [29]. Comparable to human physiology, *Drosophila* contains insulin-producing cells, insulin-like molecules, and an insulin-receptive mechanism [30]. In both the larval and adult stages of *Drosophila*, characteristics similar to insulin resistance, such as metabolic disorders and disrupted insulin signaling, can be replicated [31]. The established effects of high-sugar diets on *Drosophila* include high blood sugar levels, insulin resistance, increased fat accumulation, and decreased lifespan [32]. In addition, researchers have investigated the effect of botanical extracts on preventing metabolic disruptions in *Drosophila* caused by a high sucrose intake. Additionally, fruit flies contain the glucosidase enzyme, which is essential for sugar adaptation [33]. Therefore, *Drosophila* is a suitable model for evaluating the alpha-glucosidase-inhibitory properties of specific compounds.

In this study, we investigated the antioxidant and anti-diabetic properties of *C. rotundus* kombucha both in vitro and in vivo. Additionally, we analyzed its phytochemical composition. Our findings serve to bridge the gap between in vitro observations and in vivo conditions, potentially emphasizing the prospective role of *C. rotundus* kombucha as a functional beverage for disease prevention or intervention.

2. Materials and Methods

2.1. Chemicals

a,a-Diphenyl-b-picrylhydrazyl (DPPH), 2,20-azino-bis-(3-ethylbenzothiazoline- 6-sulfonic acid) diammonium salt (ABTS), Sodium phosphate monobasic, α -glucosidase, hydrogen peroxide, 1,1'-dimethyl-4,4'-bi-pyridinium dichloride (Paraquat dichloride hydrate), CAT assay kit, SOD assay kit,

and sodium phosphate dibasic were purchased from the Sigma Chemical Co. (St. Louis, MO). The SCOBY leavening agent was purchased from Master Lab Company (Thailand).

2.2. Plant materials and preparation of kombucha

C. rotundus rhizomes and tubers were collected from crop fields in the Thai province of Singburi. The Nattapong 01 voucher specimen was preserved at the Mahasarakham University Herbarium. The plant samples were washed and then dried for two days in a 50 °C hot air oven. After drying, the materials were pulverized to a fine powder. *C. rotundus* kombucha was prepared by boiling with 1% (w/v) *C. rotundus* powder for 10 minutes. After boiling, it was filtered through cheesecloth. Subsequently, 5% sugar (w/v) was added, and the mixture was boiled for an additional 5 minutes. The boiled solution was then transferred to a sterile container. Once boiled solution was cooled, a 10% SCOBY leavening agent (w/v) was added. The container was covered with cheesecloth and left to ferment at room temperature for ten days.

2.3. GC-MS analysis of *C. rotundus* kombucha

Phytochemical analysis of the *C. rotundus* kombucha was performed utilizing a GC-MS apparatus (7890B GC/5977B MSD, Agilent Technologies, Santa Clara, CA, USA) equipped with a 30 m HP-5MS capillary column, featuring a 250 mm diameter and a phase thickness of 0.25 mm. The kombucha sample was proportionally diluted with ethanol and introduced in a splitless mode. The oven started at 50 °C, maintained for 0.5 minutes, subsequently elevated to 250 °C at a rate of 10 °C per minute, and was sustained at this temperature for an additional 5 minutes. Helium (99.999%) served as the carrier gas with a regulated flow of 1.2 mL/min. The MS source temperature remained fixed at 230 °C, with a scan mode analysis spanning a mass range from 40 to 500.

2.4. HPLC analysis of *C. rotundus* kombucha

The evaluation of phenolic acids and flavonoids was conducted using High-Performance Liquid Chromatography (HPLC) from Agilent Technology, USA. The apparatus consisted of a protective guard column and an InertSustain® C18 column (250 mm x 4.6 mm i.d., 5 m; sourced from GL Sciences Inc., Tokyo, Japan). At wavelengths of 280 and 320 nm, a diode array detector was employed to detect phenolic acids and flavonoids. The use of external standards facilitated the identification and comparison of individual phenolic acids and flavonoids within the samples.

2.5. In vitro antioxidant activity

2.5.1. Scavenging DPPH free radical activity assay

The inhibitory effect on DPPH free radical activity of kombucha was evaluated by mixing 180 µL of kombucha solution with 20 µL of a 1 mM DPPH solution (solubilized in methanol). Reaction mixtures were incubated for 30 minutes at room temperature under no light. The absorbance at a wavelength of 517 nm was measured using an ELISA reader (Biochrome, England) to determine the inhibitory effect. Each sample was examined three replicates. The inhibitory activity was calculated using the following formula:

$$\text{Inhibition (\%)} = ((\text{Abs of control} - \text{Abs of sample}) / (\text{Abs of control})) \times 100$$

where "Abs of the control" refers to the DPPH solution's absorbance value without the sample.

The 50% inhibitory concentration (IC₅₀) was established from the regression equation derived from plotting the percentage of inhibition against kombucha concentrations (0-800 µL/mL, diluted with sterile water).

2.5.2. Scavenging ABTS free radical activity assay

The inhibitory effect of kombucha on ABTS free radical activity was assessed by mixing 20 µL of the kombucha solution with 180 µL of the ABTS• solution. The latter was prepared by combining

10 mL of 7 mM ABTS (in ultrapure water) with 10 mL of 2.45 mM ammonium persulphate and then incubating it overnight (12–16 h) at room temperature in the dark. The concentration of the ABTS radical (ABTS•) stock solution was verified at 734 nm. An ABTS• solution was prepared to achieve an absorption of approximately 0.700 at 734 nm. Reaction mixtures were incubated for 10 minutes at room temperature in the dark. The absorbance at a wavelength of 734 nm was measured using an ELISA reader (Biochrome, England) to determine the inhibitory effect. Each sample was examined three replicates. The inhibitory activity was calculated using the following formula:

$$\text{Inhibition (\%)} = ((\text{Abs of control} - \text{Abs of sample}) / (\text{Abs of control})) \times 100$$

where “Abs of the control” refers to the ABTS• solution’s absorbance value without the sample.

The 50% inhibitory concentration (IC₅₀) was established from the regression equation derived from plotting the percentage of inhibition against kombucha concentrations (0–800 uL/mL, diluted with sterile water).

2.6. *In vitro* anti- α -glucosidase activity

The inhibitory activity of the extracts against α -glucosidase was evaluated using a method described Pistia-Brueggeman and Hollingsworth (2001) [34], with slight modifications. Briefly, 50 μ L of α -glucosidase solution (0.5 U/mL in 50 mM potassium phosphate buffer, pH 7.4) was mixed with 50 μ L of either the kombucha solution or acarbose and incubated at room temperature for 10 minutes. Subsequently, 50 μ L of 5 mM p-nitrophenyl- α -D-glucopyranoside (dissolved in 50 mM potassium phosphate buffer, pH 7.4) was added and the mixture was further incubated for 10 minutes at room temperature. The absorbance was then measured at 405 nm using a spectrophotometer (Biochrome, UK). The 50% inhibitory concentration (IC₅₀) was established from the regression equation derived from plotting the percentage of inhibition against kombucha concentrations (0, 100, 200, 400, 600, and 800 uL/mL, diluted with sterile water).

2.7. *Drosophila* strain, culture conditions, and experimental design

The wild-type *D. melanogaster* Oregon-R-C strain was provided by the Department of Biology at Khon Kaen University. The flies were housed in a wheat cream medium enriched with yeast powder at 25.1±0.2 °C, 70–80% relative humidity, and subjected to a 12:12 light/dark cycle in a laboratory setting. Every two days, surviving flies were relocated to vials with fresh food.

For the preparation of the high sugar diet (HSD), 30 g of sucrose was dissolved in 100 mL of sterile water. In the case of the HSD+pure *C. rotundus* kombucha diet, 30 g of sucrose was dissolved in 100 mL of undiluted kombucha. For the HSD+400 uL/mL (40%) and HSD+100 uL/mL (10%) kombucha diets, kombucha was initially diluted to concentrations of 400 and 100 uL/mL, respectively, with sterile water. Thereafter, 30 g of sucrose was incorporated into each 100 mL of the diluted solutions.

Experiments were conducted using 5–7-day-old female flies, which were divided into four groups: HSD, HSD+ pure kombucha, HSD+400 (40%) uL/mL kombucha, and HSD+100 (10%) uL/mL kombucha. Each group consisted of 100 flies, with 20 flies per vial. The diets for each group were poured on the cotton sheet that placed in the bottom of the test vial (2.0 × 9.5 cm). Before treatment, flies were starved for 2 hours in an empty vial containing cotton cloth saturated with sterile water. After 2 hours of starvation, the flies were transferred to the test and fed for 4 hours. After the feeding period, the flies were transferred to an empty vial. The flies were euthanized with 5% CO₂ fumigation and then placed in the refrigerator at -20 °C for 10 minutes. After the flies were euthanized, they were transferred to a 1.5-mL microcentrifuge tube. The fly samples were washed three times with 1.0 mL of 50 mM potassium phosphate buffer. The samples were then homogenized using a pestle in 1.0 mL of the same buffer. Fly debris was removed from the homogenate by centrifugation at 10,000 rpm for 10 minutes at 4 °C. The supernatant was transferred to new 1.5 mL microcentrifuge tubes and kept on ice during preparation for measurement of antioxidant activity (DPPH and ABTS assays, CAT and SOD activities, and lipid peroxidation assay), α -glucosidase activity, and protein content.

The research protocol received approval from the Ethics Committee of Ubon Ratchathani Rajabhat University (Ethical Clearance No. AN63008).

2.8. *In vivo* antioxidant activity

2.8.1. Scavenging DPPH free radical activity assay

The inhibitory effect on DPPH free radical activity in *Drosophila* homogenate was evaluated by mixing 180 μ L of the supernatant with 20 μ L of a 1 mM DPPH solution (solubilized in methanol). Reaction mixtures were incubated for 30 minutes at room temperature under no light. The absorbance at a wavelength of 517 nm was measured using an ELISA reader (Biochrome, England) to determine the inhibitory effect. Each sample was examined three replicates. The inhibitory activity was calculated using the following formula:

$$\text{Inhibition (\%)} = ((\text{Abs of control} - \text{Abs of sample}) / (\text{Abs of control})) \times 100$$

where “Abs of the control” refers to the DPPH solution’s absorbance value without the sample.

The inhibitory activity was normalized to the protein content in the supernatant.

2.8.2. Scavenging ABTS free radical activity assay

The inhibitory effect on ABTS free radical activity in *Drosophila* homogenate was evaluated by mixing 20 μ L of the supernatant with 180 μ L of ABTS $^{\bullet}$ solution. Reaction mixtures were incubated for 10 minutes at room temperature in the dark. The absorbance at a wavelength of 734 nm was measured using an ELISA reader (Biochrome, England) to determine the inhibitory effect. Each sample was examined three replicates. The inhibitory activity was calculated using the following formula:

$$\text{Inhibition (\%)} = ((\text{Abs of control} - \text{Abs of sample}) / (\text{Abs of control})) \times 100$$

where “Abs of the control” refers to the ABTS $^{\bullet}$ solution’s absorbance value without the sample.

The inhibitory activity was normalized to the protein content in the supernatant.

2.8.3. SOD activity

SOD activity was determined using a Sigma-Aldrich reagent in accordance with the manufacturer’s instructions and the sample preparation procedure. Fruit flies (n = 100, 20/vial) were ground in 1 mL of cold buffer and then centrifuged (1,500 g for 5 minutes at 4 $^{\circ}$ C) to remove any remaining material. The supernatant (900 μ L) underwent another centrifugation (10,000 g for 15 min at 4 $^{\circ}$ C). This supernatant was then used to measure the activity of SOD1. To evaluate SOD2 activity, the resultant pellet was redissolved in 0.5 mL of cold buffer. Each test was performed three times.

2.8.4. CAT activity

The activity of CAT was evaluated using a kit from Sigma-Aldrich that measures the remaining hydrogen peroxide after the CAT reaction. In a summarized process, flies (n = 100, 20/vial) were ground in 1 mL of cold enzyme dilution buffer and then centrifuged (1,500 g for 5 min at 4 $^{\circ}$ C) to eliminate any waste. A portion of the supernatant (500 μ L) was then diluted in 20 mL of the assay buffer. It was combined with 25 μ L of hydrogen peroxide solution and, after 1 minute, the reaction was halted using 900 μ L of 15 mM sodium azide. Subsequently, a 10 μ L sample was introduced to a color reagent, which consisted of 0.25 mM 4-aminoantipyrine, 2 mM 3,5-dichloro-2-hydroxybenzenesulfinic acid, and peroxidase (0.5-1.5 U/mg). After incubating for 15 minutes at ambient temperature, the absorbance was recorded at 520 nm using a Biochrome instrument from the UK. This assay was repeated three times for accuracy.

2.8.5. Measurement of lipid peroxidation

Lipid peroxidation was assessed using a modified method from Zeb and Ullah (2016) [35], employing the TBAR technique and taking malondialdehyde as a reference. In a concise process, a 500 μ L portion of the fly homogenate was combined with 10 mL of TBA reagent. This mixture was then heated for 15 minutes, allowed to cool, and centrifuged at 3,000 rpm for 10 minutes to remove any sediment. The absorbance of the remaining supernatant was recorded at 532 nm using a Biochrome instrument based in the UK. The concentration of MDA was calculated based on a standard curve derived from 1,1,3,3-tetramethoxypropane.

2.8.6. H₂O₂ and paraquat challenge tests

To evaluate resistance to paraquat (superoxide anion) and hydrogen peroxide, a separate set of flies was used. Female flies, aged 5-7 days, were starved for 2 hours before being transferred to containers with filter paper saturated with either a 20 mM paraquat solution or a 10% hydrogen peroxide solution. These solutions were either unaltered or supplemented with *C. rotundus* kombucha at concentrations of pure, 400 μ L/mL, or 100 μ L/mL, and each was prepared using a 6% sucrose solution. The number of deceased flies was noted at four-hour intervals until no flies remained alive.

2.9. In vivo anti-alpha glucosidase activity

The glucosidase activity in *Drosophila* was evaluated by combining 50 μ L of supernatant with 50 μ L of 50 mM potassium phosphate buffer and 50 μ L of 5 mM p-nitrophenyl- α -D-glucopyranoside (dissolved in 50 mM potassium phosphate buffer). The reaction mixtures were incubated at ambient temperature for 30 minutes. The glucosidase activity was quantified at a wavelength of 405 nm using an ELISA reader (Biochrome, England). The standard curve for alpha-glucosidase was established by plotting the optical density at 405 nm against enzyme concentrations ranging from 0.03-0.5 U/mL. Each sample was evaluated in triplicate. The enzyme activities were normalized to the protein content in the supernatant.

2.10. Protein content assay

The protein content of the homogenate derived from the entire body was determined using the Bradford (1976) method and Merck's Bradford reagent according to the manufacturer's instructions. Bovine serum albumin was used as the standard, and concentrations were calculated in mg/mL.

2.11. Statistical analysis

The data were statistically analyzed using SPSS 20.0 (SPSS Inc., Chicago, USA). One-way ANOVA was used for mean comparisons. Kaplan-Meier analysis was used to determine the difference in survival curves. Statistical significance was defined as a $P < 0.05$. The significance levels for each experiment were denoted by the symbols * for $P < 0.05$ and ** for $P < 0.01$.

3. Results

3.1. GC-MS chemical profiles of *C. rotundus* kombucha

Table 1 shows the GC-MS chromatograms of *C. rotundus* kombucha. It has been documented that the following compounds in kombucha exhibit biological activity: 3,5-dimethylpyrazole, 4H-pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl, 2-(4-methyl-1H-1,2,3-triazol-1-yl)ethan-1-amine, 5-hydroxymethylfurfural, 10-azido-1-decanethiol, 3-deoxy-D-mannonic lactone, 1-beta-D-ribofuranosyl-3-[5-tetraazoly]-1,2,4-triazole, Nona-2,3-dienoic acid, ethyl ester, and stigmastrol.

The phenolic and flavonoid contents of *C. rotundus* kombucha, analyzed using HPLC, are presented in Figure 1. As demonstrated in Figure 1, chlorogenic acid and epicatechin were detected at the concentration of 6.04 and 1.07 μ g/g, respectively.

Table 1. The GC-MS profiling of *Cyperus rotundus* kombucha ferment.

No.	Name	Chemical formula	RT	Previously reported biological activity	Ref.
1	3,5-Dimethylpyrazole	C ₅ H ₈ N ₂	3.712	Hypoglycemic activity Anti-aging	[36,37]
2	Benzen-d5-amine	C ₆ H ₂ D ₅ N	4.008	Not reported	NA
3	5-.alpha.-Aminoethyltetrazole	C ₃ H ₇ N ₅	5.156	Not reported	NA
4	2,4-Dihydroxy-2,5-dimethyl-3(2H)-furan-3-one	C ₆ H ₈ O ₄	6.112	Flavourant	[38]
5	Xylopyranoside, methyl-4-azido-4-deoxy-, .beta.-L	C ₆ H ₁₁ N ₃ O ₄	6.475	Not reported	NA
6	Propylamine, N,N,2,2-tetramethyl-, N-oxide	C ₇ H ₁₇ N O	6.992	Not reported	NA
7	Furaneol	C ₆ H ₈ O ₃	7.546	Not reported	NA
8	4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl	C ₆ H ₈ O ₄	9.535	Antioxidant activity	[39]
9	Butyl-tert-butylisopropoxyborane	C ₁₁ H ₂₅ BO	10.119	Not reported	NA
10	1-Tetrazol-2-ylethanone	C ₃ H ₄ N ₄ O	10.224	Not reported	NA
11	2-(4-Methyl-1H-1,2,3-triazol-1-yl)ethan-1-amine	C ₅ H ₁₀ N ₄	10.587	Antioxidant, anti-proliferative, and anti-atherosclerotic activities	[40]
12	Lactic acid, 2-methyl-, monoanhydride with 1-butaneboronic acid, cyclic ester	C ₈ H ₁₅ BO ₃	10.606	Not reported	NA
13	1-Tetrazol-2-ylethanone	C ₃ H ₄ N ₄ O	10.903	Not reported	NA
14	Borinic acid, diethyl-	C ₄ H ₁₁ BO	10.684	Not reported	NA
15	5-Hydroxymethylfurfural	C ₆ H ₆ O ₃	11.294	Anticancer activity	[41]
16	10-Azido-1-decanethiol	C ₁₀ H ₂₁ N ₃ S	12.604	Anti-fungal activity	[42]
17	.beta.-D-Glucosyloxymethane	C ₈ H ₁₆ N ₂ O ₇	13.532	Not reported	NA
18	Ethanone, 1,1'-(2-ethyl-4,5-dimethyl-1,3,2-dioxaborolane-4,5-diyl)bis	C ₁₀ H ₁₇ BO ₄	14.144	Not reported	NA
19	1,3,2-Dioxaborolane, 2-ethyl-4-(3-oxiranylpropyl)-	C ₉ H ₁₇ BO ₃	15.330	Not reported	NA
20	Ethanone, 1-[5-(1,1-dimethylethyl)-2-ethyl-4-methyl-1,3,2-dioxaborolan-4-yl]-	C ₁₁ H ₂₁ BO ₃	18.036	Not reported	NA
21	3- Deoxy-D-mannonic lactone	C ₆ H ₁₀ O ₅	18.629	Antibacterial activity Antioxidant and alpha-glucosidase inhibitor activities	[43,44]
22	1-.beta.-d-Ribofuranosyl-3-[5-tetraazoly]-1,2,4-triazole	C ₈ H ₁₁ N ₇ O ₄	21.239	Analgesics, antipyretics, anti-convulsants, anti-inflammatory, immune modulatory activity	[45]
23	Nona-2,3-dienoic acid, ethyl ester	C ₁₁ H ₁₈ O ₂	24.949	Antiviral activity	[46]
24	Stigmasterol	C ₂₉ H ₄₈ O	29.530	Anti-osteoarthritic activity Anti-hypercholesterolemic activity Anti-tumor Antioxidant Antimutagenic activity Anti-inflammatory activity CNS activities	[47-55]

NA = Not available.




Figure 1. The HPLC chromatograms of *C. rotundus* kombucha reveal the presence of chlorogenic acid and epicatechin.

3.2. Effects of *C. rotundus* kombucha on *in vitro* and *in vivo* antioxidant activity

The antioxidant properties of *C. rotundus* kombucha were evaluated using two distinct assays: DPPH and ABTS. In the evaluation, the kombucha solution produced by *C. rotundus* exhibited significant antioxidant properties. For the DPPH assay, the IC_{50} was calculated to be 76.7 ± 9.6 μ L/mL (Figure 2A). The IC_{50} value for the kombucha solution in the ABTS assay was 314.2 ± 16.9 μ L/mL (Figure 2B). The antioxidant findings suggest that *C. rotundus* kombucha has significant antioxidant potential, as evidenced by its significant inhibitory activity at low concentrations. Consequently, consuming *C. rotundus* kombucha may provide the body with a beneficial source of antioxidants.

The antioxidant potential of *C. rotundus* kombucha, as observed *in vitro*, was further confirmed *in vivo* using *Drosophila* as a test model, employing DPPH and ABTS assays. As shown in Figure 2C, *Drosophila* consuming a high-sugar diet supplemented with 100% kombucha exhibited a 45.44% increase in DPPH free radical scavenging activity relative to those consuming a high-sugar diet alone ($P < 0.01$). When the same diet was supplemented with 40% or 10% *C. rotundus* kombucha, the scavenging efficiency increased by 36.35% and 30.74%, respectively, compared to the HSD group ($P < 0.05$). Similar to the DPPH results, *Drosophila* on a high-sugar diet supplemented with 100% kombucha demonstrated a 21.22% increase in the ABTS free radical scavenging capacity compared to the HSD group ($P < 0.01$). Adding 40% or 10% *C. rotundus* kombucha to the diet increased scavenging rates by 13.52% and 8.39%, respectively, compared to the HSD group ($P < 0.05$) (Figure 2D). The findings of this study indicate that *C. rotundus* kombucha has significant antioxidant properties when consumed. This suggests that *C. rotundus* kombucha may be a beneficial beverage for reducing oxidative stress caused by free radicals.




Figure 2. The effects of *C. rotundus* kombucha on both in vitro and in vivo antioxidant activities. (A) and (B) The % inhibition of DPPH and ABTS free radicals *in vitro*, respectively. (C) and (D) the mean \pm SD of % inhibition of DPPH and ABTS free radicals in *Drosophila* fed with a high sugar diet (HSD), HSD supplemented with 100% *C. rotundus* kombucha (HSD+CRK100), HSD supplemented with 40% *C. rotundus* kombucha (HSD+CRK40), and HSD supplemented with 10% *C. rotundus* kombucha (HSD+CRK10), respectively. The One-Way ANOVA test was employed to determine differences among groups. * indicates statistically significant differences compared to the control group at $P < 0.05$.

Figure 3A illustrates the effect of *C. rotundus* kombucha on the resistance of *Drosophila* on a high-sugar diet to H₂O₂-induced oxidative stress. When compared to the *Drosophila* group on a high sugar diet, those fed with pure kombucha on a high-sugar diet experienced a rise in their maximum and 50% survival rates by 13.23% and 12.52%, respectively ($P < 0.05$).

Figure 3B illustrates the effect of *C. rotundus* kombucha on the resistance of *Drosophila* fed a high-sugar diet to oxidative stress induced by paraquat. Compared to the group on a high-sugar diet alone, the *Drosophila* administered pure kombucha in addition to a high-sugar diet exhibited increases in their maximum and 50% survival rates by 21.74% and 23.67%, respectively ($P < 0.01$).

In this study, we evaluated the overall body lipid peroxidation (LPO) level to validate the antioxidant effects. As depicted in Figure 3C, *Drosophila* that consumed a high-sugar diet supplemented with pure kombucha demonstrated a notable reduction in LPO levels in comparison to the control group ($P < 0.05$).




Figure 3. The effect of *C. rotundus* kombucha supplementation on hydrogen peroxide (A) and paraquat (B) resistance and lipid peroxidation (C) in *Drosophila* fed a high sugar diet (HSD), HSD supplemented with 100% *C. rotundus* kombucha (HSD+CRK100), HSD supplemented with 40% *C. rotundus* kombucha (HSD+CRK40), and HSD supplemented with 10% *C. rotundus* kombucha (HSD+CRK10). Survival was analyzed using the Kaplan–Meier and log-rank tests. Differences were considered significant when the *P* value was less than 0.05.

The results shown in Figure 4A,B indicate that when the high-sugar diet was enriched with pure kombucha, there was a significant increase in CAT, SOD1, and SOD2 activities, compared to *Drosophila* that consumed only a high-sugar diet ($P < 0.05$).

The results shown in Figure 4A,B indicate that when the high-sugar diet was enriched with pure kombucha, there was a significant increase in CAT, SOD1, and SOD2 activities, compared to *Drosophila* that consumed only a high-sugar diet ($P < 0.05$).




Figure 4. CAT (A) and SOD1 and SOD2 (B) levels in *Drosophila* fed a high sugar diet (HSD), HSD supplemented with 100% *C. rotundus* kombucha (HSD+CRK100), HSD supplemented with 40% *C. rotundus* kombucha (HSD+CRK40), and HSD supplemented with 10% *C. rotundus* kombucha (HSD+CRK10). Data are presented as the mean \pm SD. The One-Way ANOVA test was employed to determine differences among groups. * indicates statistically significant differences compared to the HSD group at $P < 0.05$.

3.3. Effects of *C. rotundus* kombucha on in vitro and in vivo alpha-glucosidase inhibitory activity

The alpha-glucosidase inhibitory activity of the *C. rotundus* kombucha is shown in Figure 5A. The kombucha solution produced by *C. rotundus* showed inhibition potential with IC_{50} values of 142.7 ± 5.2 uL/mL. This result suggests that *C. rotundus* kombucha may function as an anti-diabetic beverage by inhibiting sugar absorption and, consequently, regulating blood glucose levels, thereby potentially preventing insulin resistance.

The in vitro observation of the inhibitory effect of *C. rotundus* kombucha on alpha-glucosidase activity was confirmed in vivo using a *Drosophila* model. As shown in Figure 5B, *Drosophila* fed a high-sugar diet supplemented with 100% kombucha demonstrated a 34.04% decrease in glucosidase activity compared to those fed a high-sugar diet alone ($P < 0.001$). When the high-sugar diet was

supplemented with either 40% or 10% *C. rotundus* kombucha, glucosidase activity decreased by 13.79% ($P<0.05$) and 11.60% ($P<0.05$), respectively, compared to *Drosophila* fed only the high-sugar diet. These results demonstrate in vitro and in vivo that *C. rotundus* kombucha significantly reduces glucosidase activity, indicating a possible anti-diabetic effect.




Figure 5. The effect of *C. rotundus* kombucha on the inhibition of α -glucosidase activity in vitro and in vivo. (A) The % inhibition of α -glucosidase activity in vitro. (B) α -Glucosidase activity in *Drosophila* fed a high sugar diet (HSD), HSD supplemented with 100% *C. rotundus* kombucha (HSD+CRK100), HSD supplemented with 40% *C. rotundus* kombucha (HSD+CRK40), and HSD supplemented with 10% *C. rotundus* kombucha (HSD+CRK10). The One-Way ANOVA test was employed to determine differences among groups. * and ** indicate statistically significant differences compared to the HSD group at $P<0.05$ and $P<0.01$, respectively.

4. Discussion

C. rotundus is a plant that has been used as medicine and food to promote health for a long time. Biological and pharmacological studies have been conducted both in vitro and in animals that demonstrate the beneficial effects of this plant. Therefore, the application of this plant in developing functional food to maintain health or prevent disease is very attractive. The purpose of this study was to process rhizomes and tubers into kombucha, a beverage that recently gained increasing recognition. To determine its health benefits, we investigated the antioxidant and anti-diabetic properties of *C. rotundus* kombucha both in vitro and in vivo. Our results revealed that *C. rotundus* kombucha exhibits potent antioxidant activity, particularly in scavenging DPPH and ABTS free radicals. Our findings suggest that *Drosophila* treated with kombucha are more resistant to oxidative stress induced by hydrogen peroxide and paraquat. This highlights the potent antioxidant properties of *C. rotundus* kombucha and indicates at the mechanism underlying its action.

Maintaining an appropriate balance between ROS production and elimination by various enzymatic and non-enzymatic agents is essential for cellular survival. In our research, we found that *Drosophila* treated with *C. rotundus* kombucha on a high-sugar diet exhibited enhanced activity of antioxidant enzymes, including SOD1, SOD2, and CAT, in comparison to flies on a high-sugar diet alone. This result is consistent with the findings of Wongchum et al. (2022) [26], who reported that the hydroethanolic extract from *C. rotundus* rhizome reduced paraquat-induced oxidative stress and increased the activity of antioxidant enzymes. Based on the chemical constituents found in the *C. rotundus* kombucha, Li et al. (2007) [56] showed that catechins from green tea can enhance the activity of antioxidant enzymes in *Drosophila*. The antioxidant capabilities of extracts might be attributed to the presence of phenols, flavonoids, and various other chemical constituents. GC-MS and HPLC analysis detected antioxidant-containing compounds, including 2-(4-Methyl-1H-1,2,3-triazol-1-yl)ethan-1-amine [40], 4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl [39], 3-Deoxy-D-mannonic lactone [44], stigmasterol [52], chlorogenic acid [57], and epicatechin [58]. The results of this study, combined with previous research, strongly suggest that the antioxidant potential of *C. rotundus* kombucha is linked to its antioxidant capacity and the activation of antioxidant enzymes.

A significant risk factor for the development of type-2 diabetes, which is characterized by elevated blood glucose levels, insulin resistance, and insufficient insulin secretion, is a high sugar intake. Reducing the activity of alpha-glucosidase is essential for managing and treating type-2 diabetes [59]. Similar to mammalian systems, *Drosophila* is a valuable model for investigating the anti-diabetic effects of natural substances [29]. Research has shown that a high-sugar diet leads to elevated blood sugar, insulin resistance, increased fat storage, and reduced life expectancy in *Drosophila* [32]. In addition, studies have revealed that fruit flies contain the glucosidase enzyme, which is essential for sugar absorption. In this study, we found that *C. rotundus* kombucha inhibited in vitro alpha-glucosidase activity with an IC_{50} of 142.7 ± 5.2 μ L/mL. Compared to other plants, the review by Kashtoh and Baek (2022) [60] reported that the IC_{50} values for glucosidase activity varied from 0.53 ± 0.014 μ g/mL to 1.873 ± 0.421 mg/mL. This suggests that *C. rotundus* kombucha is among the potent inhibitors. Our in vitro findings were verified in *Drosophila*; we observed a significant decrease in glucosidase activity in *Drosophila* fed a high-sugar diet supplemented with kombucha compared to those fed only a high-sugar diet. According to GC-MS and HPLC analysis, kombucha contains phytochemical compounds with anti-glucosidase properties. The evidence indicates that 3-Deoxy-D-mannoic lactone derived from *Distichochlamys citrea* effectively inhibits α -glucosidase activity both in vitro and in silico [44]. Studies have shown that plant-derived stigmasterol can inhibit alpha-glucosidase activity [61–63]. Wang et al. (2022) [64] reported that chlorogenic acid inhibited alpha-glucosidase activity, similar to acarbose. Therefore, based on our findings and prior research, we suggest that *C. rotundus* kombucha might act as a preventative measure against diabetes by limiting dietary sugar absorption. Nonetheless, further studies are essential to comprehensively understand the underlying mechanisms.

In conclusion, our study provides preliminary evidence supporting the potential of *C. rotundus* kombucha as an herbal remedy for enhancing antioxidant levels and preventing type-2 diabetes. In accordance with contemporary health demands and trends, we recommend further development of kombucha as a functional beverage intended to promote overall health and reduce the risk of chronic metabolic diseases.

Author Contributions: Conceptualization, conducted the experiments and wrote the manuscript, A.D. Conceptualization, Supervision, Reviewing and Editing: A.T. and S.P. Conducted the experiments: N.W. and T.C. Conceptualization, Methodology, Data curation, Writing—Reviewing and Editing and obtained funding: S.S. All the authors reviewed the manuscript.

Funding: This research was funded by Thailand Science Research and Innovation (TSRI) and Mahasarakham University.

Data Availability Statement: Data will be made available on request.

Acknowledgments: The authors thank the Faculty of Thai Traditional and Alternative Medicine, Ubon Ratchathani Rajabhat University and Research Unit of Thai Food Innovation, Department of Food Technology and Nutrition, Mahasarakham University, for providing instruments. We thank Mr. Chakkapong Thangthong, Plant Taxonomist, Ubon Ratchathani Rajabhat University, for his assistance with plant identification. We thank Assistant Professor Dr. Monthira Monthatong, Department of Biology, Faculty of Sciences, Khon Kaen University, Thailand, for providing the wild-type *Drosophila*.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Pirzada, A. M., Ali, H. H., Naeem, M., Latif, M., Bukhari, A. H., Tanveer, A. *Cyperus rotundus* L.: Traditional uses, phytochemistry, and pharmacological activities. *J. Ethnopharmacol.* 2015, 174, 540–560.
2. Yazdanparast, R., Ardestani, A. In vitro antioxidant and free radical scavenging activity of *Cyperus rotundus*. *J. Med. Food* 2007, 10, 667–674.
3. Hema, N., Avadhani, R., Ravishankar, B., Anupama, N. A comparative analysis of antioxidant potentials of aqueous and ethanolic extracts of *Cyperus rotundus* (L.). *Asian. J. Biomed. Pharm. Sci.* 2013, 3, 7–11.
4. Hu, Q. P., Cao, X. M., Hao, D. L., Zhang, L. L. Chemical composition, antioxidant, DNA damage protective, cytotoxic and antibacterial activities of *Cyperus rotundus* rhizomes essential oil against foodborne pathogens. *Sci. Rep.* 2017, 7, 45231.

5. Raut, N.A., Gaikwad, N.J. Antidiabetic activity of hydro-ethanolic extract of *Cyperus rotundus* in alloxan induced diabetes in rats. *Fitoterapia* 2006, 77, 585–588.
6. Tran, H. H., Nguyen, M. C., Le, H. T., Nguyen, T. L., Pham, T. B., Chau, V. M., Nguyen, H. N., Nguyen, T. D. Inhibitors of α -glucosidase and α -amylase from *Cyperus rotundus*. *Pharm. Biol.* 2014, 52, 74–77.
7. Athesh, K., Divakar, M., Brindha, P. Anti-obesity potential of *Cyperus rotundus* L. aqueous tuber extract in rats fed on high fat cafeteria diet. *Asian. J. Pharm. Clin. Res.* 2014, 7, 88–92.
8. Jin, J. H., Lee, D. U., Kim, Y. S., Kim, H. P. Anti-allergic activity of sesquiterpenes from the rhizomes of *Cyperus rotundus*. *Arch. Pharm. Res.* 2011, 34, 223–228.
9. Aeganathan, R., Rayar, A., Ilayaraja, S., Prabakaran, K., Manivannan, R. Anti-oxidant, antimicrobial evaluation and GC-MS analysis of *Cyperus rotundus* L. rhizomes chloroform fraction. *Am. J. Ethnomedicine* 2015, 2, 14–20.
10. Daswani, P.G., Brijesh, S., Tetali, P., Birdi, T.J. Studies on the activity of *Cyperus rotundus* Linn. tubers against infectious diarrhea. *Indian. J. Pharmacol.* 2011, 43, 340.
11. Nazish, J., Shoukat, A., 2012. Cardioprotective and antilipidemic potential of *Cyperus rotundus* in chemically induced cardiotoxicity. *Int. J. Agric. Biol.* 14, 989–992.
12. Güldür, M. E., Özgönül, A., Kilic, I. H., Söğüt, O., Ozaslan, M., Bitiren, M., Yalcin, M., Musa, D. Gastroprotective effect of *Cyperus rotundus* extract against gastric mucosal injury induced by ischemia and reperfusion in rats. *IJP-Int. J. Pharmacol.* 2010, 6, 104–110.
13. Parvez, M. K., Al-Dosari, M. S., Arbab, A. H., Niyazi, S. The in vitro and in vivo anti-hepatotoxic, anti-hepatitis B virus and hepatic CYP450 modulating potential of *Cyperus rotundus*. *Saudi. Pharm. J.* 2019, 27, 558–564.
14. Soumaya, K. J., Dhekra, M., Fadwa, C., Zied, G., Ilef, L., Kamel, G., Leila, C. G. Pharmacological, antioxidant, genotoxic studies and modulation of rat splenocyte functions by *Cyperus rotundus* extracts. *BMC Complement. Altern. Med.* 2013, 13, 1–11.
15. Mehdizadeh, M., Hashem Dabaghian, F., Shojaei, A., Molavi, N., Taslimi, Z., Shabani, R., Soleimani Asl, S. Protective effects of *Cyperus rotundus* extract on amyloid β -peptide (1-40)-induced memory impairment in male rats: A behavioral study. *Basic Clin. Neurosci.* 2017, 8, 249.
16. Sitalangka, C., Wattanathorn, J. Neuroprotective and cognitive-enhancing effects of the combined extract of *Cyperus rotundus* and *Zingiber officinale*. *BMC Complement. Altern. Med.* 2017, 17, 1–11.
17. Simorangkir, D., Masfria, M., Harahap, U., Satria, D. Activity anticancer n-hexane fraction of *Cyperus rotundus* L. Rhizome to breast cancer MCF-7 cell line. *Open Access Maced J. Med. Sci.* 2019, 7, 3904.
18. Biradar, S., Kangralkar, V. A., Mandavkar, Y., Thakur, M., Chougule, N. Antiinflammatory, antiarthritic, analgesic and anticonvulsant activity of *Cyperus* essential oils. *Int. J. Pharm. Pharm. Sci.* 2010, 2, 112–115.
19. Tsoyi, K., Jang, H. J., Lee, Y. S., Kim, Y. M., Kim, H. J., Seo, H. G., Lee, J. H., Kwak, J. H., Lee, D. U., Chang, K. C. (+)-Nootkatone and (+)-valencene from rhizomes of *Cyperus rotundus* increase survival rates in septic mice due to heme oxygenase-1 induction. *J. Ethnopharmacol.* 2011, 137, 1311–1317.
20. Rocha, F. G., Brandenburg, M. M., Pawloski, P. L., Soley, B. D. S., Costa, S. C. A., Meinerz, C. C., Baretta, I. P., Otuki, M. F., Cabrini, D. A. Preclinical study of the topical anti-inflammatory activity of *Cyperus rotundus* L. extract (Cyperaceae) in models of skin inflammation. *J. Ethnopharmacol.* 2020, 254, 112709.
21. Sharma, A., Verma, R., Ramteke, P. *Cyperus rotundus*: a potential novel source of therapeutic compound against urinary tract pathogens. *J. Herb. Med.* 2014, 4, 74–82.
22. Shivakumar, S. I., Suresh, H. M., Hallikeri, C. S., Hatapakki, B. C., Handiganur, J. S., Kuber, S., Shivakumar, B. Anticonvulsant effect of *Cyperus rotundus* Linn rhizomes in rats. *J. Nat. Remedies.* 2009, 9, 192–196.
23. Zhou, Z. L., Yin, W. Q., Yang, Y. M., He, C. H., Li, X. N., Zhou, C. P., Guo, H. New iridoid glycosides with antidepressant activity isolated from *Cyperus rotundus*. *Chem. Pharm. Bull (Tokyo)* 2016, 64, 73–77.
24. Wongchum, N., Dechakhamphu, A., Chaweerak, S., Pinlaor, S., Tanomtong, A. Evaluation of aqueous and ethanol extracts of *Cyperus rotundus* L. on sexual behaviours and reproductive fitness in *Drosophila melanogaster*. *Pharm. Sci. Asia.* 2021, 48, 516–522.
25. Wongchum, N., Dechakhamphu, A., Ma-ding, A., Khamphaeng, T., Pinlaor, S., Pinmongkhonkul, S., Tanomtong, A. The effects of *Cyperus rotundus* L. extracts on the longevity of *Drosophila melanogaster*. *South. Afr. J. Bot.* 2022, 148, 218–227.
26. Wongchum, N., Dechakhamphu, A., Panya, P., Pinlaor, S., Pinmongkhonkul, S., Tanomtong, A. Hydroethanolic *Cyperus rotundus* L. extract exhibits anti-obesity property and increases lifespan expectancy in *Drosophila melanogaster* fed a high-fat diet. *J. Herbm. Pharmacol.* 2022, 11, 296–304.

27. da Silva Júnior, J. C., Mafaldo, Í. M., de Lima Brito, I., de Magalhães Cordeiro, A. M. T. Kombucha: Formulation, chemical composition, and therapeutic potentialities. *Curr. Res. Food. Sci.* 2022, 5, 360–365.
28. Morales, D. Biological activities of kombucha beverages: The need of clinical evidence. *Trends. Food. Sci. Technol.* 2020, 105, 323–333.
29. Palanker Musselman, L., Fink, J. L., Narzinski, K., Ramachandran, P. V., Sukumar Hathiramani, S., Cagan, R. L., Baranski, T. J. A high-sugar diet produces obesity and insulin resistance in wild-type *Drosophila*. *Dis. Model. Mech.* 2011, 4, 842–849.
30. Partridge, L., Alic, N., Bjedov, I., Piper, M.D. Ageing in *Drosophila*: the role of the insulin/Igf and TOR signalling network. *Exp. Gerontol.* 2011, 46, 376–381.
31. Morris, S. N. S., Coogan, C., Chamseddin, K., Fernandez-Kim, S. O., Kolli, S., Keller, J. N., Bauer, J. H. Development of diet-induced insulin resistance in adult *Drosophila melanogaster*. *Biochim. Biophys. Acta. BBA-Mol. Basis. Dis.* 2012, 1822, 1230–1237.
32. Na, J., Musselman, L. P., Pendse, J., Baranski, T. J., Bodmer, R., Ocorr, K., Cagan, R. A *Drosophila* model of high sugar diet-induced cardiomyopathy. *PLoS. Genet.* 2013, 9, e1003175.
33. Tanimura, T., Kitamura, K., Fukuda, T., Kikuchi, T. Purification and partial characterization of three forms of α -glucosidase from the fruit fly *Drosophila melanogaster*. *J. Biochem (Tokyo)* 1979, 85, 123–130.
34. Pistia-Brueggeman, G., Hollingsworth, R.I. A preparation and screening strategy for glycosidase inhibitors. *Tetrahedron* 2001, 57, 8773–8778.
35. Zeb, A., Ullah, F. A simple spectrophotometric method for the determination of thiobarbituric acid reactive substances in fried fast foods. *J. Anal. Methods. Chem.* 2016, 2016, 9412767.
36. Gerritsen, G. C., Dulin, W. E. Effect of a new hypoglycemic agent, 3, 5-dimethylpyrazole, on carbohydrate and free fatty acid metabolism. *Diabetes* 1965, 14, 507–515.
37. Cavallini, G., Donati, A., Bergamini, E. Antiaging Therapy: A Novel Target for Antilipolytic Drugs. *Mini-Rev. Med. Chem.* 2014, 14, 551–556.
38. Chukwu, C., Omaka, O., Aja, P. Characterization of 2, 5-dimethyl-2, 4-dihydroxy-3 (2H) furanone, a flavourant principle from *Sysepalum dulcificum*. *Nat. Prod. Chem. Res.* 2017, 5, 2.
39. Dilek Tepe, H., Doyuk, F. Determination of phytochemical content by chromatographic methods and antioxidant capacity in methanolic extract of jujube (*Zizyphus jujuba* Mill.) and oleaster (*Elaeagnus a ngustifolia* L.). *Int. J. Fruit. Sci.* 2020, 20, S1876–S1890.
40. Priyaa, G.H., More, S.S. Antioxidant, anti-proliferative, and anti-atherosclerotic effect of phytochemicals isolated from *Trachyspermum ammi* with honey in RAW 264.7 and THP-1 cells. *Pharmacogn. Mag.* 2022, 18, 143–151.
41. Joel, O.O., Maharjan, R. Effects of 5-hydroxymethylfurfural isolated from *Cola hispida* on oral adenosquamous carcinoma and MDR *Staphylococcus aureus*. *JMPHTR.* 2021, 8, 1–7.
42. Doughari, J.H., Abraham, M. Antifungal activity of *Jatropha Curcas* Linn on *Candida albicans* and *Candida tropicalis* associated with neonatal and infantile infections in Yola, Nigeria. *Am. j. agric. biol. sci.* 2021, 16, 19–32.
43. Shobana, S., Vidhya, V., Ramya, M. Antibacterial activity of garlic varieties (ophioscordon and sativum) on enteric pathogens. *Curr. Res. J. Biol. Sci.* 2009, 1, 123–126.
44. Van Chen, T., Cuong, T. D., Quy, P. T., Bui, T. Q., Van Tuan, L., Van Hue, N., Triet, N. T., Ho, D. V., Bao, N. C., Nhung, N. T. A. Antioxidant activity and α -glucosidase inhibitability of *Distichochlamys citrea* MF Newman rhizome fractionated extracts: in vitro and in silico screenings. *Chem. Pap.* 2022, 76, 5655–5675.
45. Kakarla, L. Comparative Biochemical Studies on Indian Sedges *Cyperus scariosus* R.Br and *Cyperus rotundus* L. *Pharmacogn. J.* 2017, 8, 598–609.
46. Huang, X., Li, J., Hong, Y., Jiang, C., Wu, J., Wu, M., Sheng, R., Liu, H., Sun, J., Xin, Y., Su, W. Antiviral effects of the petroleum ether extract of *Tournefortia sibirica* L. against enterovirus 71 infection in vitro and in vivo. *Front. Pharmacol.* 2022, 13, 999798.
47. Gabay, O., Sanchez, C., Salvat, C., Chevy, F., Breton, M., Nourissat, G., Wolf, C., Jacques, C., Berenbaum, F. Stigmasterol: a phytosterol with potential anti-osteoarthritic properties. *Osteoarthritis. Cartilage* 2010, 18, 106–116.
48. Chandler, R.F., Hooper, S.N., Ismail, H.A. Antihypercholesterolemic studies with sterols: beta-sitosterol and stigmasterol. *J. Pharm. Sci.* 1979, 68, 245–247.
49. Batta, A. K., Xu, G., Honda, A., Miyazaki, T., Salen, G. Stigmasterol reduces plasma cholesterol levels and inhibits hepatic synthesis and intestinal absorption in the rat. *Metabolism* 2006, 55, 292–299.

50. Kasahara, Y., Kumaki, K., Katagiri, S., Yasukawa, K., Yamanouchi, S., Takido, M., Akihisa, T., Tamura, T. Carthami flos extract and its component, stigmasterol, inhibit tumour promotion in mouse skin two-stage carcinogenesis. *Phytother. Res.* 1994, 8, 327–331.
51. Gao, Z., Maloney, D.J., Dedkova, L.M., Hecht, S.M. Inhibitors of DNA polymerase β : Activity and mechanism. *Bioorg. Med. Chem.* 2008, 16, 4331–4340.
52. Panda, S., Jafri, M., Kar, A., Meheta, B. Thyroid inhibitory, antiperoxidative and hypoglycemic effects of stigmasterol isolated from *Butea monosperma*. *Fitoterapia* 2009, 80, 123–126.
53. Lim, J. C., Park, J. H., Budesinsky, M., Kasal, A., Han, Y. H., Koo, B. S., Lee, S. I., Lee, D. U. Antimutagenic constituents from the thorns of *Gleditsia sinensis*. *Chem. Biol. Pharm. Bull.* 2005, 53, 561–564.
54. Navarro, A., De Las Heras, B., Villar, A. Anti-inflammatory and immunomodulating properties of a sterol fraction from *Sideritis foetens* Clem. *Biol. Pharm. Bull.* 2001, 24, 470–473.
55. Garcia, M., Saenz, M., Gomez, M., Fernández, M. Topical antiinflammatory activity of phytosterols isolated from *Eryngium foetidum* on chronic and acute inflammation models. *Phytother. Res. Int. J. Devoted. Pharmacol. Toxicol. Eval. Nat. Prod. Deriv.* 1999, 13, 78–80.
56. Li, Y.M., Chan, H.Y.E., Huang, Y., Chen, Z.Y. Green tea catechins upregulate superoxide dismutase and catalase in fruit flies. *Mol. Nutr. Food. Res.* 2007, 51, 546–554.
57. Singh, A.K., Singla, R.K., Pandey, A.K. Chlorogenic acid: A dietary phenolic acid with promising pharmacotherapeutic potential. *Curr. Med. Chem.* 2023, 30, 3905–3926.
58. Sheng, Y., Sun, Y., Tang, Y., Yu, Y., Wang, J., Zheng, F., Li, Y., Sun, Y. Catechins: Protective mechanism of antioxidant stress in atherosclerosis. *Front. Pharmacol.* 2023, 14, 1144878.
59. Hossain, U., Das, A.K., Ghosh, S., Sil, P.C. An overview on the role of bioactive α -glucosidase inhibitors in ameliorating diabetic complications. *Food. Chem. Toxicol.* 2020, 145, 111738.
60. Kashtoh, H., Baek, K. H. Recent updates on phytoconstituent alpha-glucosidase inhibitors: An approach towards the treatment of type two diabetes. *Plants* 2022, 11, 2722.
61. Junejo, J. A., Zaman, K., Rudrapal, M., Celik, I., Attah, E. I. Antidiabetic bioactive compounds from *Tetrastigma angustifolia* (Roxb.) Deb and *Oxalis debilis* Kunth.: Validation of ethnomedicinal claim by in vitro and in silico studies. *South. Afr. J. Bot.* 2021, 143, 164–175.
62. Lolok, N., Sumiwi, S. A., Ramadhan, D. S. F., Levita, J., Sahidin, I. Molecular dynamics study of stigmasterol and beta-sitosterol of *Morinda citrifolia* L. towards α -amylase and α -glucosidase. *J. Biomol. Struct. Dyn.* 2023. 1–4.
63. Tasnuva, S. T., Qamar, U. A., Ghafoor, K., Sahena, F., Jahurul, M. H. A., Rukshana, A. H., Juliana, M. J., Al-Juhaimi, F. Y., Jalifah, L., Jalal, K. C. A., Ali, M. E., Zaidul, I. S. M. α -glucosidase inhibitors isolated from *Mimosa pudica* L. *Nat. Prod. Res.* 2019, 33, 1495–1499.
64. Wang, S., Li, Y., Huang, D., Chen, S., Xia, Y., Zhu, S. The inhibitory mechanism of chlorogenic acid and its acylated derivatives on α -amylase and α -glucosidase. *Food. Chem.* 2022, 372, 131334.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.