

Plasmonic-assisted water gas shift reaction of gold particles on TiO₂

K. W. Ahmed¹, K. Mudiyansele^{2#}, H. Idriss^{3*}

¹Department of Chemical Engineering, University of Waterloo, Waterloo, N2L 3G1, Ontario, Canada.

²Surface Science and Advanced Characterization, SABIC-CRD at King Abdullah University for Science and Technology (KAUST), Thuwal, 23955, Saudi Arabia

³Institute of Functional Interfaces, Karlsruhe Institute of Technology (KIT), 76344 Eggenstein-Leopoldshafen, Germany.

* hicham.idriss@kit.edu




Figure S1. Light flux as a function of wavelengths from 320 to 1100 nm (25% light intensity)




Figure S2. Light flux as a function of wavelengths from 320 to 1100 nm (40% light intensity)




Figure S3. Light flux as a function of wavelengths from 400 to 1100 nm (25% light intensity)




Figure S4. Light flux as a function of wavelengths from 320 to 620 nm (25% light intensity)




Figure S5. Light flux as a function of wavelengths from 400 to 660 nm (25% light intensity)




Figure S6. Light flux as a function of wavelengths from 610 to 700 nm (LED with band pass filter at 80% intensity)




Figure S7. Light flux as a function of wavelengths from 570 to 610 nm (band pass filter 40% light intensity)