

**Table S1. Filming locations and abundance of individuals.**

Locations: 1. Nieuwpoort 51.12848 N, 2.74802 E (Belgium); 2. Oostende, 51.22153 N (Belgium), 3. Baie de Saint-Brieuc, 48.52886 N, 2.73117 W (France); 4. Mazères 43.28701 N, 1.6333671 E, (France); 5. Saintes Maries de la Mer (France) 43.487854 N, 4.405823 E, 6. Arles 43.40801 N, 4.72683 E (France), 7. Arcachon 44.627503, 44.62750 W (France), 8. Séné, 47.6286 N, 2.7786 W (France), 9. Perros-Guirec, 48.815113 N, -3.439466 W (France), 10. Saint Cast-Le Guildo 48.633331 N 2.26667 W (France), 11. Fuente de Piedra 37.132954 N 4.740334 W (Spain). Abundance classes (number of individuals): 1: 1-5; 2: 6-25; 3: 26-125.

Species	Location	Abundance
<b>Suborder Charadri</b>		
<i>Charadrius hiaticula</i> L. 1758	1	2
<i>Haematopus ostralegus</i> L. 1758	2,3	2
<i>Himantopus himantopus</i> (L. 1758)	4,5	3
<i>Ochthodromus alexandrines</i> (L. 1758)	6	2
<i>Pluvialis squatarola</i> (L. 1758)	1	1
<i>Recurvirostra avosetta</i> L. 1758	5,6,7	3
<i>Thiornis dubius</i> (Scopoli 1786)	4	2
<i>Vanellus vanellus</i> (L. 1758)	8	2
<b>Suborder Scolopaci</b>		
<i>Actitis hypoleuca</i> (L. 1758)	3,4,6	2
<i>Arenaria interpres</i> (L. 1758)	2,9	2
<i>Calidris alba</i> (Pallas 1764)	2,10	2
<i>Calidris alpina</i> (L. 1758)	1,2,9	2
<i>Calidris maritima</i> (Brünnich 1764)	9	2
<i>Calidris pugnax</i> (L. 1758)	4	2
<i>Gallinago gallinago</i> (L. 1758)	4	2
<i>Limosa lapponica</i> (L. 1758)	3	1
<i>Limosa limosa</i> (L. 1758)	7,11	2
<i>Lymnocyptes minimus</i> (Brünnich 1764)	4	1
<i>Numenius arquata</i> (L. 1758)	1,3	2
<i>Numenius phaeopus</i> (L. 1758)	3	2
<i>Phalaropus fulicarius</i> (L. 1758)	2	1
<i>Tringa erythropus</i> (Pallas 1764)	4,5	3
<i>Tringa glareola</i> L. 1758	4	2
<i>Tringa nebularia</i> (Gunnerus 1767)	4,5	2
<i>Tringa ochropus</i> L. 1758	4	3
<i>Tringa totanus</i> (L. 1758)	1,4,9	3