

Communication

Not peer-reviewed version

Epitope Mapping of an Anti-CD44v4 Monoclonal Antibody (C₄₄Mab-108) using Enzyme-Linked Immunosorbent Assay

[Hiroyuki Suzuki](#)^{*}, Mayuki Tawara, Aoi Hirayama, [Nohara Goto](#), [Tomohiro Tanaka](#), Mika K Kaneko, [Yukinari Kato](#)^{*}

Posted Date: 3 November 2023

doi: 10.20944/preprints202311.0216.v1

Keywords: CD44; CD44 variant 4; monoclonal antibody; epitope; enzyme-linked immunosorbent assay



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Communication

Epitope Mapping of an Anti-CD44v4 Monoclonal Antibody (C₄₄Mab-108) Using Enzyme-Linked Immunosorbent Assay

Hiroyuki Suzuki, Mayuki Tawara, Aoi Hirayama, Nohara Goto, Tomohiro Tanaka, Mika K. Kaneko and Yukinari Kato

Department of Antibody Drug Development, Tohoku University Graduate School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai 980-8575, Miyagi, Japan

* Correspondence: hiroyuki.suzuki.b4@tohoku.ac.jp (H. S.); yukinari.kato.e6@tohoku.ac.jp (Y.K.); Tel.: +81-22-717-8207 (H.S., Y.K.)

Abstract: CD44 is a type I transmembrane glycoprotein, and possesses various isoforms which are largely classified into CD44 standard and CD44 variant (CD44v) isoforms. Some variant-encoded regions play critical roles in tumor progression. However, the function of CD44 variant 4 (CD44v4)-encoded region has not been fully understood. Using peptide immunization, we developed an anti-CD44v4 mAb, C₄₄Mab-108, which is useful for flow cytometry, western blotting, and immunohistochemistry. In this study, we determined the critical epitope of C₄₄Mab-108 by enzyme-linked immunosorbent assay (ELISA). We used the alanine (or glycine)-substituted peptides of the CD44v4-encoded region (amino acids 271–290 of human CD44v3–10), and found that C₄₄Mab-108 did not recognize the alanine-substituted peptides of D280A and W281A. Furthermore, these peptides could not inhibit the recognition of C₄₄Mab-108 in flow cytometry and immunohistochemistry. The results indicate that the critical binding epitope of C₄₄Mab-108 includes Asp280 and Trp281 of CD44v3–10.

Keywords: CD44; CD44 variant 4; monoclonal antibody; epitope; enzyme-linked immunosorbent assay

1. Introduction

CD44 has various isoforms, which are generated by the alternative splicing of CD44 pre-mRNA [1]. The mRNA of CD44 standard (CD44s) isoform is produced by constant region exons, including the first five (1 to 5) and the last five (16 to 20) [2]. The mRNAs of CD44 variant (CD44v) isoform are produced by the assembling of variant exons (v1–v10) with the constant region exons of CD44s [3]. CD44s and CD44v receive the post-translational modifications, including N- or O-glycosylation [4]. Both CD44s and CD44v can attach to hyaluronic acid, which is important for cellular adhesion, homing, and motility [5].

CD44v plays important roles in the tumor progression by specific functions of variant exon-encoded regions [6]. The heparin-binding growth factors are recruited to heparan sulphate modified in the v3-encoded region [7,8]. MET, a receptor tyrosine kinase for hepatocyte growth factor, associates with the v6-encoded region [9,10]. These functions are essential for the activation of growth factor signaling and tumor proliferation. However, the roles of CD44 variant 4 (CD44v4)-encoded region have not been investigated. Therefore, specific antibodies against CD44v4 are indispensable for basic research, tumor diagnosis, and therapy.

We have established anti-CD44 monoclonal antibodies (mAbs), which recognize the standard [11,12]- or each variant [13–20]-encoded region. All mAbs can be used for flow cytometry, western blotting, and immunohistochemistry, and are expected to contribute not only to basic research but also to pathogenic diagnosis. We also determined the critical epitopes of C₄₄Mab-5 [21] and C₄₄Mab-46 [22,23]. We previously established an anti-CD44v4 mAb, C₄₄Mab-108 (mouse IgG₁, kappa) using

the peptide immunization [19]. To clarify further characteristics of C₄₄Mab-108, we performed epitope mapping using enzyme-linked immunosorbent assay (ELISA).

2. Materials and Methods

2.1. Peptides

The CD44v4 peptide (₂₇₁-AFDHTKQNQDWTQWNPSHSN-₂₉₀) and 20 alanine (or glycine)-substituted peptides (Table 1) were synthesized by utilizing PEPscreen (Sigma-Aldrich Corp., St. Louis, MO). The number of amino acids (aa) is derived from human CD44v3-10 (accession No.: X66733).

2.2. ELISA

The CD44v4 peptides were immobilized on Nunc Maxisorp 96-well immunoplates (Thermo Fisher Scientific Inc., Waltham, MA) at a concentration of 10 µg/mL for 30 min at 37°C. After washing with phosphate-buffered saline (PBS) containing 0.05% Tween20 (PBST; Nacalai Tesque, Inc., Kyoto, Japan), wells were blocked with 1% bovine serum albumin (BSA)-containing PBST for 30 min at 37°C. The plates were incubated with 10 µg/mL of C₄₄Mab-108, followed by a peroxidase-conjugated anti-mouse immunoglobulins (1:2000 diluted; Agilent Technologies Inc., Santa Clara, CA). Enzymatic reactions were performed using the ELISA POD Substrate TMB Kit (Nacalai Tesque, Inc.). Optical density was measured at 655 nm using an iMark microplate reader (Bio-Rad Laboratories, Inc., Berkeley, CA).

2.3. Flow cytometry

The N-terminal PA16-tagged CD44v3-10-overexpressed Chinese hamster ovary-K1 (CHO/CD44v3-10) [19] cells were harvested after a brief exposure to 0.25% trypsin in 1 mM ethylenediaminetetraacetic acid (EDTA; Nacalai Tesque, Inc.) and washed with 0.1% BSA in PBS. C₄₄Mab-108 (10 µg/mL) was incubated with the CD44v4 peptides (10 µg/mL) for 30 min at 4°C. The cells were further treated with Alexa Fluor 488-conjugated anti-mouse IgG (1:2000). Fluorescence data were collected using the SA3800 Cell Analyzer (Sony Corp., Tokyo, Japan).

2.4. Immunohistochemical analysis

One formalin-fixed paraffin-embedded (FFPE) tissue of oral squamous cell carcinoma (OSCC) for peptide blocking assay was obtained from Tokyo Medical and Dental University [24]. The tissue slides were autoclaved in citrate buffer (pH 6.0; Nichirei Biosciences, Inc., Tokyo, Japan) for 20 min for antigen retrieval. After blocking with SuperBlock T20 (Thermo Fisher Scientific, Inc.), the sections were incubated with C₄₄Mab-108 (10 µg/mL) in the presence or absence of the CD44v4 peptides (10 µg/mL) and then treated with the EnVision+ Kit for mouse (Agilent Technologies Inc.) for 30 min. The color was developed using 3,3'-diaminobenzidine tetrahydrochloride (DAB; Agilent Technologies Inc.). Counterstaining was performed with hematoxylin (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan). Leica DMD108 (Leica Microsystems GmbH, Wetzlar, Germany) was used to examine the sections and obtain images.

3. Results

3.1. Epitope mapping of C₄₄Mab-108 with alanine (or glycine)-substituted CD44v4 peptide

We previously established an anti-CD44v4 mAb (C₄₄Mab-108) by peptide immunization of CD44v4 region (₂₇₃-DHTKQNQDWTQWNPSHSNP-₂₉₁) [19]. We confirmed that C₄₄Mab-108 recognizes with only the variant 4-encoded region peptide (aa 271-290), but not other regions of CD44v3-10 extracellular domain [19]. To identify the binding epitope of C₄₄Mab-108, we synthesized 20 alanine (or glycine)-substituted peptides of the CD44v4 (Table 1). C₄₄Mab-108 exhibited reaction with A271G, F272A, D273A, H274A, T275A, K276A, Q277A, N278A, Q279A, T282A, Q283A, W284A,

N285A, P286A, S287A, H288A, S289A, N290A, and wild-type (WT) (Figure 1A). In contrast, C₄₄Mab-108 did not react with D280A and W281A (Figure 1A). This result indicated that Asp280 and Trp281 are included in the critical epitope of C₄₄Mab-108. The results are summarized in Table 1. Figure 1B shows the schematic illustration of CD44s, CD44v3-10, and the critical aa (Asp280 and Trp281) recognized by C₄₄Mab-108.




Figure 1. Determination of the C₄₄Mab-108 epitope by ELISA using alanine (or glycine)-substituted CD44v4 peptides. (A) The alanine (or glycine)-substituted CD44v4 peptides in PBS or PBS alone were immobilized on immunoplates. The plates were incubated with C₄₄Mab-108 (10 µg/mL), followed by peroxidase-conjugated anti-mouse immunoglobulins. Error bars represent means ± SDs. N.C., negative control (PBS). (B) Schematic illustration of CD44s, CD44v3-10, and the C₄₄Mab-108 epitope. The C₄₄Mab-108 epitope includes Asp280 and Trp281 of CD44v3-10. The epitope is conserved in mouse and Chinese hamster (Cham), but not in rat.

Table 1. Identification of the C₄₄Mab-108 epitope using alanine (or glycine)-substituted CD44v4 peptides.

Peptides	Sequences	C ₄₄ Mab-108
WT (271-290)	AFDHTKQNQDWTQWNPSHSN	+++
A271G	GFDHTKQNQDWTQWNPSHSN	+++
F272A	AADHTKQNQDWTQWNPSHSN	+++
D273A	AFAHTKQNQDWTQWNPSHSN	+++
H274A	AFDATKQNQDWTQWNPSHSN	+++

T275A	AFDHAKQNQDWTQWNPSHSN	+++
K276A	AFDHTAQNQDWTQWNPSHSN	+++
Q277A	AFDHTKANQDWTQWNPSHSN	+++
N278A	AFDHTKQAQDWTQWNPSHSN	+++
Q279A	AFDHTKQNADWTQWNPSHSN	+++
D280A	AFDHTKQNQAWTQWNPSHSN	–
W281A	AFDHTKQNQDATQWNPSHSN	–
T282A	AFDHTKQNQDWAQWNPSHSN	+
Q283A	AFDHTKQNQDWTAWNPSHSN	+++
W284A	AFDHTKQNQDWTQANPSHSN	+
N285A	AFDHTKQNQDWTQWAPSHSN	+++
P286A	AFDHTKQNQDWTQWNASHSN	+++
S287A	AFDHTKQNQDWTQWNPAHSN	+++
H288A	AFDHTKQNQDWTQWNPSASN	+++
S289A	AFDHTKQNQDWTQWNPSHAN	+++
N290A	AFDHTKQNQDWTQWNPSHSA	+++

+++ , OD655≥0.6; ++ , 0.3≤OD655<0.6; + , 0.1≤OD655<0.3; – , OD655<0.1.

3.2. Flow cytometry using C44Mab-108 with alanine-substituted CD44v4 peptides

We next performed a peptide-blocking assay using flow cytometry to confirm the importance of the C44Mab-108 epitope. As shown in Figure 2, C44Mab-108 reacted with the CHO/CD44v3-10 cells. This reaction was completely neutralized by WT. In contrast, D280A and W281A did not block the reaction of C44Mab-108 with CHO/CD44v3-10. The result confirmed that Asp280 and Trp281 of CD44v3-10 are critical for detection by C44Mab-108 using flow cytometry.




Figure 2. Flow cytometry using C44Mab-108 and CD44v4 peptides. C44Mab-108 (10 µg/mL) plus WT, the alanine-substituted peptides (10 µg/ml), or control (0.1% DMSO in blocking buffer, –peptide) were reacted with CHO/CD44v3-10 cells for 30 min at 4°C, followed by treatment with Alexa Fluor 488-

conjugated anti-mouse IgG. The black line represents the control. WT, wild type; DMSO, dimethyl sulfoxide.

3.3. Immunohistochemistry using C₄₄Mab-108 with alanine-substituted CD44v4 peptides

We also performed a peptide-blocking assay using immunohistochemical analysis. As shown in Figure 3, C₄₄Mab-108 stained the FFPE section of OSCC, which was completely neutralized by WT. In contrast, D280A and W281A did not neutralize the reaction. These results were corresponding to that of Figure 2.




Figure 3. Immunohistochemistry using C₄₄Mab-108 and CD44v4 peptides. The FFPE sections of OSCC were incubated with C₄₄Mab-108 (10 µg/mL) plus WT, the alanine-substituted peptides (10 µg/mL), or control (0.1% DMSO in blocking buffer, -peptide), followed by that with the Envision + Kit. Scale bar = 100 µm. FFPE, formalin-fixed paraffin-embedded; OSCC, oral squamous cell carcinoma; WT, wild type; DMSO, dimethyl sulfoxide.

4. Discussion

In the present study, the critical epitope of C₄₄Mab-108 was determined as Asp280 and Trp281 in the CD44v4 region. Since the reactivity of C₄₄Mab-108 to T282A and W284A was also reduced (Figure 1A), Thr282 and Trp284 may contribute to the recognition partially. Figure 1B shows the homology of the v4 region among human, mouse, Chinese hamster, and rat sequences. The Asp280 and Trp281 are conserved in human, mouse, Chinese hamster, but not in rat. Furthermore, Thr282 and Trp284 are also conserved in human and mouse. Although the result suggests that C₄₄Mab-108 might recognize both human and mouse CD44v4, it did not react with mouse CD44v4 using flow cytometry (data not shown).

Since C₄₄Mab-108 was established by the peptide immunization, C₄₄Mab-108 can recognize the definite peptide structure of the variant 4-encoded region. In contrast, CD44 is predicted to carry 146 O-glycan sites in the variant region. Among them, 41 of these sites have already been experimentally confirmed [25]. Thr282 is also a confirmed O-glycan site [26]. Further studies are required to reveal whether the O-glycan at Thr282 affects the recognition by C₄₄Mab-108.

In our previous study, C₄₄Mab-108 could detect CD44v3-10-overexpressed cells such as CHO/CD44v3-10, but not detect endogenous CD44v4 in several cancer cell lines in flow cytometry [19]. In contrast, C₄₄Mab-108 could detect endogenous and membranous CD44v4 in immunohistochemistry [19]. These results suggest that the variant 4-encoded region is folded into the inside of protein in living cells, but exposed by antigen retrieval in immunohistochemistry. Recently, we provided a potential strategy for developing cancer-specific antibodies that target locally misfolded cell surface receptors such as human epidermal growth factor receptor 2 [27,28].

Further studies are needed to investigate whether C₄₄Mab-108 is involved in the recognition of specific CD44v4 type and/or specific condition of cells.

Funding: This research was supported in part by Japan Agency for Medical Research and Development (AMED) under Grant Numbers: JP23ama121008 (to Y.K.), JP23am0401013 (to Y.K.), 23bm1123027h0001 (to Y.K.), and JP23ck0106730 (to Y.K.), and by the Japan Society for the Promotion of Science (JSPS) Grants-in-Aid for Scientific Research (KAKENHI) grant nos. 22K06995 (to H.S.), 21K07168 (to M.K.K.), and 22K07224 (to Y.K.).

Conflicts of Interest: The authors have no conflict of interest.

References

1. Ponta, H.; Sherman, L.; Herrlich, P.A. CD44: from adhesion molecules to signalling regulators. *Nat Rev Mol Cell Biol* **2003**, *4*, 33-45, doi:10.1038/nrm1004.
2. Yan, Y.; Zuo, X.; Wei, D. Concise Review: Emerging Role of CD44 in Cancer Stem Cells: A Promising Biomarker and Therapeutic Target. *Stem Cells Transl Med* **2015**, *4*, 1033-1043, doi:10.5966/sctm.2015-0048.
3. Chen, C.; Zhao, S.; Karnad, A.; Freeman, J.W. The biology and role of CD44 in cancer progression: therapeutic implications. *J Hematol Oncol* **2018**, *11*, 64, doi:10.1186/s13045-018-0605-5.
4. Liao, C.; Wang, Q.; An, J.; Chen, J.; Li, X.; Long, Q.; Xiao, L.; Guan, X.; Liu, J. CD44 Glycosylation as a Therapeutic Target in Oncology. *Front Oncol* **2022**, *12*, 883831, doi:10.3389/fonc.2022.883831.
5. Mishra, M.N.; Chandavarkar, V.; Sharma, R.; Bhargava, D. Structure, function and role of CD44 in neoplasia. *J Oral Maxillofac Pathol* **2019**, *23*, 267-272, doi:10.4103/jomfp.JOMFP_246_18.
6. Zöller, M. CD44: can a cancer-initiating cell profit from an abundantly expressed molecule? *Nat Rev Cancer* **2011**, *11*, 254-267, doi:10.1038/nrc3023.
7. Bennett, K.L.; Jackson, D.G.; Simon, J.C.; Tanczos, E.; Peach, R.; Modrell, B.; Stamenkovic, I.; Plowman, G.; Aruffo, A. CD44 isoforms containing exon V3 are responsible for the presentation of heparin-binding growth factor. *J Cell Biol* **1995**, *128*, 687-698, doi:10.1083/jcb.128.4.687.
8. Jackson, D.G.; Bell, J.I.; Dickinson, R.; Timans, J.; Shields, J.; Whittle, N. Proteoglycan forms of the lymphocyte homing receptor CD44 are alternatively spliced variants containing the v3 exon. *J Cell Biol* **1995**, *128*, 673-685, doi:10.1083/jcb.128.4.673.
9. Matzke, A.; Sargsyan, V.; Holtmann, B.; Aramuni, G.; Asan, E.; Sendtner, M.; Pace, G.; Howells, N.; Zhang, W.; Ponta, H.; et al. Haploinsufficiency of c-Met in cd44^{-/-} mice identifies a collaboration of CD44 and c-Met in vivo. *Mol Cell Biol* **2007**, *27*, 8797-8806, doi:10.1128/mcb.01355-07.
10. Orian-Rousseau, V.; Chen, L.; Sleeman, J.P.; Herrlich, P.; Ponta, H. CD44 is required for two consecutive steps in HGF/c-Met signaling. *Genes Dev* **2002**, *16*, 3074-3086, doi:10.1101/gad.242602.
11. Yamada, S.; Itai, S.; Nakamura, T.; Yanaka, M.; Kaneko, M.K.; Kato, Y. Detection of high CD44 expression in oral cancers using the novel monoclonal antibody, C(44)Mab-5. *Biochem Biophys Rep* **2018**, *14*, 64-68, doi:10.1016/j.bbrep.2018.03.007.
12. Goto, N.; Suzuki, H.; Tanaka, T.; Asano, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Monoclonal Antibody for Multiple Applications against Esophageal Squamous Cell Carcinomas. *Int J Mol Sci* **2022**, *23*, doi:10.3390/ijms23105535.
13. Ejima, R.; Suzuki, H.; Tanaka, T.; Asano, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 6 Monoclonal Antibody C(44)Mab-9 for Multiple Applications against Colorectal Carcinomas. *Int J Mol Sci* **2023**, *24*, doi:10.3390/ijms24044007.
14. Ishikawa, K.; Suzuki, H.; Kaneko, M.K.; Kato, Y. Establishment of a Novel Anti-CD44 Variant 10 Monoclonal Antibody C(44)Mab-18 for Immunohistochemical Analysis against Oral Squamous Cell Carcinomas. *Curr Issues Mol Biol* **2023**, *45*, 5248-5262, doi:10.3390/cimb45070333.
15. Kudo, Y.; Suzuki, H.; Tanaka, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 5 Monoclonal Antibody C(44)Mab-3 for Multiple Applications against Pancreatic Carcinomas. *Antibodies (Basel)* **2023**, *12*, doi:10.3390/antib12020031.
16. Suzuki, H.; Goto, N.; Tanaka, T.; Ouchida, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 8 Monoclonal Antibody C(44)Mab-94 against Gastric Carcinomas. *Antibodies (Basel)* **2023**, *12*, doi:10.3390/antib12030045.
17. Suzuki, H.; Kitamura, K.; Goto, N.; Ishikawa, K.; Ouchida, T.; Tanaka, T.; Kaneko, M.K.; Kato, Y. A Novel Anti-CD44 Variant 3 Monoclonal Antibody C(44)Mab-6 Was Established for Multiple Applications. *Int J Mol Sci* **2023**, *24*, doi:10.3390/ijms24098411.
18. Suzuki, H.; Ozawa, K.; Tanaka, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 7/8 Monoclonal Antibody, C(44)Mab-34, for Multiple Applications against Oral Carcinomas. *Biomedicines* **2023**, *11*, doi:10.3390/biomedicines11041099.
19. Suzuki, H.; Tanaka, T.; Goto, N.; Kaneko, M.K.; Kato, Y. Development of a Novel Anti-CD44 Variant 4 Monoclonal Antibody C(44)Mab-108 for Immunohistochemistry. *Curr Issues Mol Biol* **2023**, *45*, 1875-1888, doi:10.3390/cimb45030121.

20. Tawara, M.; Suzuki, H.; Goto, N.; Tanaka, T.; Kaneko, M.K.; Kato, Y. A Novel Anti-CD44 Variant 9 Monoclonal Antibody C(44)Mab-1 Was Developed for Immunohistochemical Analyses against Colorectal Cancers. *Curr Issues Mol Biol* **2023**, *45*, 3658-3673, doi:10.3390/cimb45040238.
21. Asano, T.; Kaneko, M.K.; Kato, Y. Development of a Novel Epitope Mapping System: RIEDL Insertion for Epitope Mapping Method. *Monoclon Antib Immunodiagn Immunother* **2021**, *40*, 162-167, doi:10.1089/mab.2021.0023.
22. Asano, T.; Kaneko, M.K.; Takei, J.; Tateyama, N.; Kato, Y. Epitope Mapping of the Anti-CD44 Monoclonal Antibody (C(44)Mab-46) Using the REMAP Method. *Monoclon Antib Immunodiagn Immunother* **2021**, *40*, 156-161, doi:10.1089/mab.2021.0012.
23. Takei, J.; Asano, T.; Suzuki, H.; Kaneko, M.K.; Kato, Y. Epitope Mapping of the Anti-CD44 Monoclonal Antibody (C(44)Mab-46) Using Alanine-Scanning Mutagenesis and Surface Plasmon Resonance. *Monoclon Antib Immunodiagn Immunother* **2021**, *40*, 219-226, doi:10.1089/mab.2021.0028.
24. Itai, S.; Ohishi, T.; Kaneko, M.K.; Yamada, S.; Abe, S.; Nakamura, T.; Yanaka, M.; Chang, Y.W.; Ohba, S.I.; Nishioka, Y.; et al. Anti-podocalyxin antibody exerts antitumor effects via antibody-dependent cellular cytotoxicity in mouse xenograft models of oral squamous cell carcinoma. *Oncotarget* **2018**, *9*, 22480-22497, doi:10.18632/oncotarget.25132.
25. Steentoft, C.; Vakhrushev, S.Y.; Joshi, H.J.; Kong, Y.; Vester-Christensen, M.B.; Schjoldager, K.T.; Lavrsen, K.; Dabelsteen, S.; Pedersen, N.B.; Marcos-Silva, L.; et al. Precision mapping of the human O-GalNAc glycoproteome through SimpleCell technology. *Embo j* **2013**, *32*, 1478-1488, doi:10.1038/emboj.2013.79.
26. Mereiter, S.; Martins Á, M.; Gomes, C.; Balmaña, M.; Macedo, J.A.; Polom, K.; Roviello, F.; Magalhães, A.; Reis, C.A. O-glycan truncation enhances cancer-related functions of CD44 in gastric cancer. *FEBS Lett* **2019**, *593*, 1675-1689, doi:10.1002/1873-3468.13432.
27. Arimori, T.; Mihara, E.; Suzuki, H.; Ohishi, T.; Tanaka, T.; Kaneko, M.K.; Takagi, J.; Kato, Y. Locally misfolded HER2 expressed on cancer cells is a promising target for development of cancer-specific antibodies *Cell Press Community Review* **2023**, doi:dx.doi.org/10.2139/ssrn.4565236.
28. Suzuki, H.; Kaneko, M.K.; Kato, Y. A cancer-specific monoclonal antibody against HER2 for breast cancers *Preprint* **2023**, doi:10.20944/preprints202309.0906.v2.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.