

Review

Not peer-reviewed version

Cardiac Targeting Peptide: From Discovery to Applications

Daniella Sahagun and [Maliha Zahid](#) *

Posted Date: 21 November 2023

doi: 10.20944/preprints202311.1314.v1

Keywords: cell penetrating peptides; protein transduction domains; cardiac targeting peptide; heart failure; atrial fibrillation



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

Cardiac Targeting Peptide: From Discovery to Applications

Daniella Sahagun and Maliha Zahid *

Department of Cardiovascular Diseases, Mayo Clinic, Rochester, MN

* Correspondence: Guggenheim Gu9-01B; Mayo Clinic; 200 First Street SW; Rochester, MN, 55905; zahid.maliha@mayo.edu

Abstract: Despite significant strides in prevention, diagnosis, and treatment, cardiovascular diseases remain the number one cause of mortality in the United States, with rates climbing at an alarming rate in the developing world. Targeted delivery of therapeutics to the heart has been a lofty goal to achieve with strategies ranging from direct intra-cardiac or intra-pericardial delivery, intra-coronary infusion, to adenoviral, lentiviral, and adeno-associated viral vectors which have preference, if not complete cardio-selectivity, for cardiac tissue. Cell penetrating peptides (CPP) are 5-30 amino acid long peptides that are able to breach cell membrane barriers while carrying cargoes up to several times their size, in an intact functional form. Identified nearly three decades ago, the first of these CPPs came from the HIV coat protein transactivator of transcription. Although a highly efficient CPP, its clinical utility is limited by its robust ability to cross any cell membrane barrier, including crossing the blood-brain barrier and transducing neuronal tissue non-specifically. Several strategies have been utilized to identify cell or tissue specific CPPs, one of which is phage display. Using this latter technique, we identified a cardiomyocyte targeting peptide (CTP) more than a decade ago, a finding that has been corroborated by several independent labs across the world that have utilized CTP for myriad different purposes in pre-clinical animal models. The goal of this publication is to provide a comprehensive review of the identification, validation, and application of CTP, and outlining its potential in diagnostic and therapeutic applications especially in the field of targeted RNA interference.

Keywords: cell penetrating peptides; protein transduction domains; cardiac targeting peptide; heart failure; atrial fibrillation

Introduction

Despite major advances in diagnosis and treatment of cardiovascular disorders, they remain the number one killer in the developed world, with rates rising at an alarming rate in the developing world [1–4]. A major cardiovascular pathology is atrial fibrillation; the most common rhythm disorder in adults with an estimated prevalence of 46.3 million worldwide[5]. It affects 1-2% of adults in the United States, with an estimated 5.7 million patients[6]. Data from the Framingham heart study has documented an over threefold increase in its prevalence over the last 50 years[7]. These already high prevalence rates are further estimated to double by year 2050 due to an aging of the population, and increased prevalence of risk factors like coronary artery disease, hypertension, obesity, and congestive heart failure[7]. Superiority of rhythm over rate control in this disorder has been demonstrated in multiple randomized control trials[8–10]. Cardiomyocytes, the cells responsible for the contractility of the heart, are the seat of some common pathologies like heart failure with reduced ejection fraction due to myriad different reasons. Another type of heart failure, heart failure with preserved ejection fraction, results from not a “weak” but a “stiff” heart, and has been rising steadily as the proportion of all heart failure patients, due to the aging of the population, and increasing incidence of co-morbidities, like obesity, hypertension, and diabetes mellitus that lead to it. In stark contrast to heart failure with reduced ejection fraction, heart failure with preserved ejection fraction

is a recalcitrant disease with very few therapies shown to benefit it. Targeting the stiffened heart by changing the behaviour of the hypertrophied cardiomyocyte, hence addressing the underlying pathophysiology, would require the ability to successfully and specifically target the offending cardiomyocyte.

In our era of advanced cardiac diagnostics and multitude of therapeutics available to the clinician, targeted drug delivery is the next frontier[11]. It is increasingly being recognized that designing new drugs is not enough; they have to be delivered to a specific organ, and preferably to the cells of interest[12]. In the context of targeting the heart, the strategies have ranged from physical approaches ranging from direct microneedle injections into the myocardium[13,14], intra-coronary instillation[14], retrograde perfusion of the coronary sinus[15,16], and placement of hydrogels into the pericardial space[17–20]. On the vector side, viral vectors have ranged from adenovirus[21,22], lentivirus[23], adenoassociated viruses[24,25], or non-viral vectors comprising of exosomes[26], nanoparticles[27,28] or microbubbles targeted to the heart by destruction of these bubbles with ultrasound during their passage through the heart under direct visualization[29–31]. An alternate, non-viral, promising strategy is use of cell penetrating peptides (CPPs), also known as protein transduction domains[32].

First identified nearly three decades ago, CPPs are 5-30 amino acid long peptides that are able to cross cell membrane barriers while carrying cargoes several times their size in an intact, functional form. As so often happens in science, this was a serendipitous discovery. Two independent groups of scientists investigating the various functions of the HIV coat protein transactivator of transduction (Tat), an 86-amino acid protein, discovered the protein's ability to enter cells efficiently without the use of transfection reagents[33,34]. Soon to follow was the report that homeobox domain of antennapedia transcription factor of *Drosophila melanogaster* was able to enter neural cells in a receptor-independent manner and affect neural morphogenesis[35]. Mapping of the domains within Tat responsible for its transduction property led to the identification of a positively charged segment of the protein, only 11 amino acids long, rich in arginine and lysine that are two cationic amino acid residues (YGRKKRRQRRR)[36]. Subsequently, the ability to carry cargoes much larger than its size was demonstrated by fusing Tat to a beta-galactosidase reporter protein residue, which upon intra-peritoneal injection into wild-type mice led to robust transduction of multiple tissues, including heart, lung, liver, kidneys and brain tissue, at six hours post injection[37], high-lighting Tat's potential as a vector. However, this robust, ubiquitous uptake by myriad tissue, especially crossing of the blood-brain barrier, limited its clinical utility. Researchers have tackled this issue by using Tat locally instead of systemically, by intra-articular, intra-ocular or intra-tumoral injections. This approach is not feasible for the targeting of deeper organs such as the heart due to its invasiveness and resulting complications, which would limit repeat administrations.

Since the initial description of the first two CPPs, Tat and Antennapedia homeodomain, the family of CPPs has expanded exponentially, as has the number of publications related to them. Non-specific uptake of these peptides has been tackled by either use of localized delivery, or administration of peptides that are only activated for uptake in specific environments, like tumor micro-environments, or use of phage delivery to identify tissue specific CPPs. The concept of phage display was first proposed by G. P. Smith in 1985[38] in a single-author Science publication, where he demonstrated that foreign DNA could be inserted into filamentous phage gene III to create fusion protein that would display the inserted amino acids on the surface protein where it could be enriched over a 1000-fold. Since the initial report, combinatorial peptide libraries of various lengths utilizing different filamentous phage (M13, T7), displaying millions of different, randomized amino acid sequences have been utilized in phage display to identify peptides homing to various targets in vitro and in vivo. We utilized a combinatorial in vitro and in vivo phage display methodology[39] to identify a 12-amino acid long, non-naturally occurring peptide that specifically targets cardiomyocytes[40], which we termed *Cardiac Targeting Peptide* (CTP) and forms the subject of the rest of this review. Studies high-lighted in this review are peer-reviewed and span our work as well as work of other investigators from around the world published only in indexed medical journals available on PubMed.

Identification of Cardiac Targeting Peptide

We utilized a combinatorial in vitro and in vivo phage screening of a large, commercially available M13 phage display library to identify CTP, a cardiomyocyte-specific cell penetrating peptide[40]. The heart is a very vascular organ with only a small minority of the intravascularly injected phage being internalized, with the blood pool contaminating what is recovered from the cardiomyocytes themselves. To enrich for cardiac targeting peptides, we attempted to decrease non-specific phage by pre-screening in a rat cardiomyoblast, H9C2, cell line. Cells were plated in a 6-well plate and allowed to reach 70% confluency before being incubated with the phage library for 6 hours. At the end of the incubation period, cells were washed extensively, trypsinized, collected and lysed using freeze-thaw cycles. Phage recovered from the lysed cell was expanded, injected into a mouse and allowed to circulate for 24 hours. The rationale for prolonged circulation time was to allow for non-specific phage from the blood pool to clear out[41], allowing for only internalized phage to be recovered from heart tissue. In order to maximize viability of the internalized phage, mice were treated 24-hours before the injections and on day of the injection with Chloroquine to prevent acidification of lysosomes and resulting breakdown of internalized phage. Recovered phage from the heart was expanded and reinjected for subsequent cycles. After the 4th cycle, 10 plaques were picked and sent for sequencing, with 6 out of 10 bearing the identical sequence of APWHLSSQYSRT, a peptide that we later named CTP or cardiac targeting peptide. In vitro incubation of a number of different cell lines with fluorescently tagged CTP revealed a preferential uptake of the peptide by H9C2 cells[40]. Injecting mice with fluorescently tagged CTP showed robust uptake by heart in 30 minutes. Later biodistribution studies revealed peak uptake at 15 minutes (earliest time point tested) with almost complete disappearance of fluorescence by 6 hours[42].

Phage display had been utilized prior to above experiments in an in vitro display methodology to identify another, longer, cardiomyocyte homing peptide. McGuire and colleagues used an in vitro phage display approach to identify a 20-mer amino acid (WLSEAGPVVTVRALRGTGSW) which had partial sequence homology to tenascin-X, an extracellular matrix protein[43]. This peptide had a 180-fold preference for cardiomyocytes in vitro, the cell line used for phage display. They were able to show preferential isolation of this phage from heart tissue as compared to a random peptide bearing phage[43]. More recently, an independent group were able to engineer cardiosphere-derived cells to produce exosomes labeled with Lamp2b and WLSEAGPVVTVRALRGTGSW, which showed anti-apoptotic effects in vitro in primary mouse neonatal cardiomyocytes, and increased cardiac retention in vivo compared with non-labeled exosomes[44]. However, it is worth noting that all in vivo injections into mice were intramyocardial, and not systemic or intravenous, bringing into question the strength of the targeting as well as the clinical applicability of this approach for use in humans. This is an important point as intra-myocardial delivery is clinically feasible in a very limited number of patients such as those undergoing coronary artery bypass grafting or another open-heart procedure, precluding repeat therapies. In another application, L-arginine loaded gold nanoparticles modified with WLSEAGPVVTVRALRGTGSW exhibited increased cellular uptake by cardiomyocytes with resulting improved photoacoustic imaging in vitro and in vivo[45]. These nanoparticles were able to increase nitric oxide production, and decrease apoptosis, fibrosis and infarct size in a rat model of ischemia-reperfusion[45]. Hopefully, larger animal studies will be forthcoming and show similar efficacy, as this was a very recent publication.

Cardiac Targeting Peptide for Diagnostic Purposes

Coronary artery disease is the most common form of cardiovascular disorder. Atherosclerotic plaque buildup leads to progressive narrowing of coronary artery lumen. This is a chronic process, with symptoms developing only after a significant portion of the lumen (>50-70%) has been compromised. Symptoms, once they develop, range from chest pain, shortness of breath on exertion, to fatigue and decreased exercise tolerance, and are responsible for the most frequent reason for presentation to the emergency rooms in the US. These symptoms are investigated in patients by putting them through cardiac stress testing while continuously monitoring the heart with electrocardiograms (ECGs). The heart is stressed with exercise, the most physiological form of

stressor, or chemicals in patients who cannot exercise. Commonly, ECG alone has low sensitivity and specificity for detection of occlusive coronary artery disease, and hence is combined with an imaging modality, in the form of nuclear cardiac imaging or echocardiography using ultrasound. Myocardial perfusion imaging is most commonly done through using cardiac tracers like Thallium, or more recently Technetium 99m. Uptake of this tracer by liver and gut loops interferes with heart imaging, lowering the sensitivity of the examination. Targeting the radioisotopes to the heart would, in theory, improve cardiac uptake and decrease radio-isotope dose. CTP was tagged with Technetium 99m via a chelator, HYNIC (Figure 1), and showed almost exclusive uptake by the heart with predominantly renal excretion, when compared with Technetium Sestamibi, the formulation of Technetium 99m in wide-spread clinical use[42].




Figure 1. Chemical Structure of CTP-HYNIC Conjugate. An example of imaging application of CTP by synthesizing it with HYNIC conjugated to the N-terminus and labeled with ^{99m}Tc -HYNIC-CTP. Wild-type CD1 mice were injected with 1-5mCi of ^{99m}Tc Sestamibi or ^{99m}Tc -HYNIC-CTP, and single photon emission computed tomography (SPECT/CT) imaging performed⁴².

Cardiac PET imaging is a more recent addition to the clinician armamentarium of diagnostics. Compared to SPECT imaging, PET has a higher sensitivity (80.3% versus 68.7%) and similar specificity (63.8% versus 61.7%) and therefore overall superior accuracy in predicting occlusive coronary artery disease with significantly lower radiation exposure (by ~50%) than SPECT. Yet, it remains grossly under-utilized with data from 2019 indicating that of all nuclear cardiac stress tests, PET studies represented only 10% of them. The underlying reason for this is lack of widespread availability of cardiac PET tracers. The only FDA approved cardiac PET tracers in clinical use are ^{15}O water, ^{13}N ammonia, and ^{82}Rb rubidium^{13,14}. Availability of these tracers is limited by the need for an on-site (^{15}O water and ^{13}N ammonia) or nearby (^{13}N ammonia) cyclotron, or commitment to costly generators (^{82}Rb). Due to short half-lives ranging from 76sec for ^{82}Rb , to 2.1min for ^{15}O water and 10min for ^{13}N ammonia, their use in conjunction with treadmill exercise stress testing is either not possible (^{82}Rb and ^{15}O water) or not practical (^{13}N ammonia). Furthermore, the long positron range of ^{82}Rb makes image resolution suboptimal and its low extraction limits its spatial resolution. Of all stress modalities (exercise, dobutamine, dipyridamole, adenosine, regadenoson), exercise is the most physiological stressor, and provides important additive prognostic information, not available with other stressors.

With this background in mind, CTP could provide a solution. Our recent work showed that labeling of CTP with Gallium-68, another PET radioisotope, can be successfully performed using 1,4,7-Triazacyclononane-1,4,7-triacetic acid (NOTA) as a chelator, with preliminary animal studies showing promise. This conjugation was optimized to yield 98% radiolabeling efficiency at room temperature in ten minutes, and with >97% radiochemical purity, without the need of a heating block. The CTP-NOTA-Gallium 68 conjugate was taken up by heart tissue in as little as 5 minutes with isotope leaving the heart in ~30 minutes. At later time points, the gall bladder lit up on PET imaging,

with a significant portion being secreted by the kidneys (presented in abstract form at Society of Nuclear Medicine, Chicago, 2023).

Cardiac Targeting Peptide for Therapeutic Purposes

The first therapeutic application of CTP was studied by Avula and colleagues in order to perform cell-selective arrhythmia ablation[46]. They engineered a 8-arm pegylated nanoparticle with an average of 2.6 CTP molecules attached to the pegylated arms along with a photosensitizer chlorin e6. Injecting these nanoparticles into rats followed by laser illumination of the heart induced localized myocyte-specific ablation restoring normal rhythm with 85% efficiency. Of note, there was no damage to “innocent, bystander” cells like endothelial cells or myofibroblasts. There were also able to demonstrate this cardiomyocyte-specific ablation in sheep heart ex-vivo. This led to the formation of complete heart block at the ablated region leading to restoration of normal sinus rhythm[46].

Extracellular vesicles, or exosomes, are heterogeneous, membrane bound vesicles, ranging in size from nanometers to micrometers, and originate as extrusions from the endosome or plasma membrane. They are being actively explored by a number of researchers as delivery vehicles for cardiac and other applications. In a series of experiments, researchers from South Korea were able to engineer exosomes expressing CTP along with LampB, and showed ~15% increased uptake in vitro and in vivo compared to non-targeted (LampB but no CTP bearing) exosomes[47]. In another demonstration of this targeted delivery, exosomes labeled only with CTP were loaded with curcumin and miR-144-3p, and showed not only in vivo delivery to the murine heart, but also cardioprotection in a mouse infarct model[48]. In a separate application, Kim and colleagues engineered extracellular vesicles to express high-levels of CTP[49]. These vesicles were loaded with anti-RAGE (receptor for advanced glycation end products) siRNA. RAGE has been shown to contribute to inflammation in a number of cardiac pathologies that lead to myocarditis. Such labeled vesicles were injected intravenously into a rat model of myocarditis induced by immunization with cardiac myosin. On day 7 after induction of myocarditis, rats were treated with a single injection of non-targeted vesicles loaded with anti-RAGE siRNA or targeted vesicles labeled with CTP. Targeted vesicles significantly decrease RAGE, IL-6, and TNF- α levels in heart tissue. This decrease in multiple inflammatory markers was accompanied by a significant increase in left ventricular ejection fraction, and improvements in left ventricular end-systolic and end-diastolic dimensions[49]. Treatment also led to significant decrease in protein levels of RAGE (as expected), but also levels of IL-6, TNF- α , COX2 and proportion of phosphorylated p65, an activated component of the NF-Kappa-B pathway⁴⁹.

As stated earlier, heart failure with preserved ejection fraction is growing in incidence as well as a disease particularly recalcitrant to treatments by drugs and interventions that have been shown to be beneficial in heart failure with reduced ejection fraction. This is likely due to significant differences in underlying pathophysiology. One of the hallmarks of heart failure with preserved ejection fraction is a thick hypertrophied left ventricle with increased myocyte stiffness. To target this particular pathology at the cardiomyocyte level, Gallicano and colleagues worked to deliver microRNA106a to the cardiomyocyte. MicroRNAs are small non-coding, ~22 nucleotide long, non-coding RNAs, that silence gene expression via a post-translation modification of mRNAs. They target the 3'UTR of an mRNA through a 7-8 bases long seed sequence that marks the complex for degradation in the RISC or RNA-induced silencing complex. One of the targets of miRNA106a is calcium calmodulin kinase II δ that is upregulated in heart failure and responsible for the calcium mishandling intrinsic to these hypertrophied cardiomyocytes. In order to deliver miRNA106a specifically to hypertrophied cardiomyocytes, it was conjugated to the N-terminus of CTP via a disulfide linker. This linker was chosen due to its covalent nature and high likelihood of remaining stable in serum, as well as being broken down in the reducing intracellular environment, leading to release of cargo miRNA106a from its vector CTP, allowing it to bind to calcium calmodulin kinase II δ mRNA marking it for degradation in the RISC complex. Incubation of human left ventricular myocytes with fluorescently labeled CTP-miRNA106a conjugate led to robust uptake by the cells⁵⁰. This conjugate was selectively taken up by human cardiomyocyte cell line and not by HEK293 cells⁵⁰. Additionally, this conjugate was able to reverse angiotensin-phenylephrine induced hypertrophy of myocytes by decreasing expression of

both HDAC4 and calcium calmodulin kinase II δ , both targets of miRNA106a[50]. Studies on the effect of this conjugate in reversing cardiac hypertrophy and ameliorating heart failure in an in vivo mouse model are ongoing.

In our own work, we chose to utilize CTP to deliver amiodarone, a small molecule class III anti-arrhythmic. Amiodarone was approved by the FDA for use as an antiarrhythmic for ventricular tachycardia and fibrillation, two life-threatening rhythms, in December 1985. A frequent off-label use of it has been for atrial fibrillation, a common nuisance rhythm that puts patients at increased risk of congestive heart failure as well as strokes. In fact, as many as 20-25% of strokes in the US are secondary to clot formation in the heart resulting from atrial fibrillation. In spite of amiodarone being the most effective anti-arrhythmic, its use is limited due to its vast volume of distribution in the human body, prolonged half-life, with only a small fraction of it (<5%) going to the heart. There are also significant, and sometimes life-threatening, off-target toxicities associated with it. Our hypothesis was that by using CTP as a vector, we can direct amiodarone to the heart where it is needed to act, thereby reducing the total dose needed to see the same anti-arrhythmic effects and decreasing off-target uptake hence improving its safety profile. To that end, we conjugated amiodarone to the N-terminus of CTP via a disulfide bond[51]. Using high-pressure liquid chromatography, we showed that this conjugate is stable at 37°C for up to 22 days[51]. We tested this conjugate in vivo in guinea pigs that were injected with either vehicle, amiodarone (80mg/Kg for 7 days), CTP-amiodarone, or CTP, with the last two (CTP-amio and CTP) injected at 1/10th the molar dose of amiodarone, for 5 days[52]. At the end of the treatment period, guinea pigs were euthanized, hearts excised and perfused in a Langendorff apparatus. Voltage and calcium indicator dyes were injected and fluorescence from the epicardium of the heart split and focused onto two cameras to record cytosolic calcium currents and voltage conduction velocities. Amiodarone, as per expectations, slowed the heart rate, decreased calcium conduction and voltage conduction velocities, hence increasing the action potential durations. CTP-amiodarone also decreased calcium and voltage conduction velocities, similar to amiodarone. To our surprise CTP-amiodarone increased the heart rate, decreased action potential duration and decreased late calcium transients, effects that seem to be driven by CTP portion of the conjugate. In order to explore the underlying possible mechanisms for these observations, RNA was extracted from hearts, RNA sequencing performed, and results of CTP injected hearts compared to control (vehicle injections only) hearts. Two important calcium handling genes SERCA2a and DHPR were upregulated in CTP injected hearts, as were alpha adrenergic receptors. On the other hand, beta adrenergic receptors were downregulated, explaining the significant increase in heart rates. Most interestingly, several inflammatory genes in the NF-kappa B pathway, like TNF α , IL1 β , and COX2 were significantly downregulated in the CTP treated hearts[52]. These were unexpected findings as we had considered CTP to always be an inert vector. Whether CTP goes beyond being a vector and has anti-inflammatory effects and salutary effects on calcium handling, remains to be confirmed, and studies exploring these intriguing but preliminary findings are underway.

Challenges, Limitations and Future Studies

Since the report of its initial discovery in 2010[40], the literature surrounding CTP has grown with a number of laboratories independently verifying its cardiomyocyte-specific targeting abilities after systemic delivery. Not only is it a “cardiac” targeting peptide, it is cardiomyocyte-specific targeting peptide with possibility for cardiomyocyte physiology altering potential as demonstrated in work by Gallicano and colleagues⁵⁰. Additionally, the number of cargoes delivered using CTP continue to grow. Yet, many unanswered questions remain, the foremost being its mechanism of transduction that is instrumental in targeting cardiomyocytes. In spite of multiple efforts, that remains an enigma. Cardiomyocytes are constantly beating cells with cell-to-cell connections, sodium-potassium channels as well as calcium channels. However, these unique proteins form small pores meant for ions that are logarithmic degrees of magnitude smaller in size, compared with CTP with or without a cargo.

Another important question that requires further study is toxicity. Our own data, generated in our lab in a non good laboratory practices (non-GLP) environment, has shown no overt toxicities. However, our studies were not comprehensive or all encompassing, and only tested CTP by itself (and a chelator HYNIC added), without a cargo. Moreover, these toxicity studies were designed for micro-dose, diagnostic studies for humans application, and were hence single dose. To study the full genotoxicity profile of CTP and its cargo, multi-dose, longitudinal, longer term studies in at least two vertebrate species (of which only one can be a rodent animal model) need to be undertaken. Due to the need for a GLP environment, such studies are typically undertaken by independent, third-party, certified research organizations and are costly. Lastly, CTP is platform technology, a non-viral method of delivering diagnostic agents and therapeutics to the heart, with many potential cargoes that could be delivered. To fully realize its potential will require concerted, and costly, efforts from multiple independent investigators.

Conclusions

CTP belongs to the larger class of tissue-specific CPPs identified through phage display utilizing a commercially available phage display library. Uptake by the heart is in as little as 5-15 minutes making it ideal for time-sensitive conditions like myocardial infarction, also known as “heart attacks” in layman terms and ischemia-reperfusion injuries. During a myocardial infarction, cardiomyocytes are suffering necrosis due to a blocked artery with window of opportunity to intervene limited to 6 hours or less. This window is far shorter than the reach of viral, or plasmid vectors or direct intramyocardial injections. The ability of CTP to target cardiomyocytes with peak uptake in as little as 5-15 mins (depending on cargo size) adds to its utility as a vector. Furthermore, it is cardiomyocyte-specific, sparing other cells in the heart like myofibroblasts, endothelial cells or resident macrophages, an immensely valuable property in conditions like heart failure where cardiomyocytes are the seat of the pathology. Although many proof of principle studies in vitro and in vivo in small animal models have been carried out, studies in larger animals are lacking. Additionally, full toxicity studies of CTP with its cargo(s) of choice need to be carried out in order to allow human trials leading to clinical applications, and ultimately realization of its full potential.

Funding: DS and MZ are supported by NIH grant R01HL153407 awarded to MZ.

Conflicts of Interest: MZ holds a patent on use of cardiac targeting peptide for cardiac delivery along with Dr. Paul Robbins and University of Pittsburgh, PA.

References

1. Duggan JP, Peters AS, Trachiotis GD, Antevil JL. Epidemiology of Coronary Artery Disease. *Surg Clin North Am.* 2022;102:499-516. doi: 10.1016/j.suc.2022.01.007
2. Wang K, Shi X, Zhu Z, Hao X, Chen L, Cheng S, Foo RSY, Wang C. Mendelian randomization analysis of 37 clinical factors and coronary artery disease in East Asian and European populations. *Genome Med.* 2022;14:63. doi: 10.1186/s13073-022-01067-1
3. Writing Committee M, Lawton JS, Tamis-Holland JE, Bangalore S, Bates ER, Beckie TM, Bischoff JM, Bittl JA, Cohen MG, DiMaio JM, et al. 2021 ACC/AHA/SCAI Guideline for Coronary Artery Revascularization: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2022;79:e21-e129. doi: 10.1016/j.jacc.2021.09.006
4. Virani SS, Alonso A, Aparicio HJ, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Cheng S, Delling FN, et al. Heart Disease and Stroke Statistics-2021 Update: A Report From the American Heart Association. *Circulation.* 2021;143:e254-e743. doi: 10.1161/CIR.0000000000000950
5. Rahman F, Kwan GF, Benjamin EJ. Global epidemiology of atrial fibrillation. *Nat Rev Cardiol.* 2014;11:639-654. doi: 10.1038/nrcardio.2014.118
6. Turakhia MP, Shafrin J, Bognar K, Trocio J, Abdulsattar Y, Wiederkehr D, Goldman DP. Estimated prevalence of undiagnosed atrial fibrillation in the United States. *PLoS One.* 2018;13:e0195088. doi: 10.1371/journal.pone.0195088
7. Schnabel RB, Yin X, Gona P, Larson MG, Beiser AS, McManus DD, Newton-Cheh C, Lubitz SA, Magnani JW, Ellinor PT, et al. 50 year trends in atrial fibrillation prevalence, incidence, risk factors, and mortality in the Framingham Heart Study: a cohort study. *Lancet.* 2015;386:154-162. doi: 10.1016/S0140-6736(14)61774-8

8. Packer DL, Mark DB, Robb RA, Monahan KH, Bahnson TD, Poole JE, Noseworthy PA, Rosenberg YD, Jeffries N, Mitchell LB, et al. Effect of Catheter Ablation vs Antiarrhythmic Drug Therapy on Mortality, Stroke, Bleeding, and Cardiac Arrest Among Patients With Atrial Fibrillation: The CABANA Randomized Clinical Trial. *JAMA*. 2019;321:1261-1274. doi: 10.1001/jama.2019.0693
9. Willems S, Borof K, Brandes A, Breithardt G, Camm AJ, Crijns H, Eckardt L, Gessler N, Goette A, Haegeli LM, et al. Systematic, early rhythm control strategy for atrial fibrillation in patients with or without symptoms: the EAST-AFNET 4 trial. *Eur Heart J*. 2022;43:1219-1230. doi: 10.1093/eurheartj/ehab593
10. Vaduganathan M, Piccini JP, Camm AJ, Crijns H, Anker SD, Butler J, Stewart J, Bracer R, Albuquerque APA, Wieloch M, et al. Dronedronone for the treatment of atrial fibrillation with concomitant heart failure with preserved and mildly reduced ejection fraction: a post-hoc analysis of the ATHENA trial. *Eur J Heart Fail*. 2022;24:1094-1101. doi: 10.1002/ehf.2487
11. May M. Why drug delivery is the key to new medicines. *Nat Med*. 2022;28:1100-1102. doi: 10.1038/s41591-022-01826-y
12. May M. Why drug delivery is the key to new medicines. *Nature Medicine*. 2022;28:1100-1102. doi: 10.1038/s41591-022-01826-y
13. Wang X, Li Q, Hu Q, Suntharalingam P, From AH, Zhang J. Intra-myocardial injection of both growth factors and heart derived Sca-1+/CD31- cells attenuates post-MI LV remodeling more than does cell transplantation alone: neither intervention enhances functionally significant cardiomyocyte regeneration. *PLoS One*. 2014;9:e95247. doi: 10.1371/journal.pone.0095247
14. Vekstein AM, Wendell DC, DeLuca S, Yan R, Chen Y, Bishawi M, Devlin GW, Asokan A, Poss KD, Bowles DE, et al. Targeted Delivery for Cardiac Regeneration: Comparison of Intra-coronary Infusion and Intra-myocardial Injection in Porcine Hearts. *Front Cardiovasc Med*. 2022;9:833335. doi: 10.3389/fcvm.2022.833335
15. Tuma J, Fernandez-Vina R, Carrasco A, Castillo J, Cruz C, Carrillo A, Ercilla J, Yarleque C, Cunza J, Henry TD, et al. Safety and feasibility of percutaneous retrograde coronary sinus delivery of autologous bone marrow mononuclear cell transplantation in patients with chronic refractory angina. *J Transl Med*. 2011;9:183. doi: 10.1186/1479-5876-9-183
16. von Degenfeld G, Raake P, Kupatt C, Lebherz C, Hinkel R, Gildehaus FJ, Munzing W, Kranz A, Waltenberger J, Simoes M, et al. Selective pressure-regulated retroinfusion of fibroblast growth factor-2 into the coronary vein enhances regional myocardial blood flow and function in pigs with chronic myocardial ischemia. *J Am Coll Cardiol*. 2003;42:1120-1128. doi: 10.1016/s0735-1097(03)00915-x
17. Shvartz V, Kanametov T, Sokolskaya M, Petrosyan A, Le T, Bockeria O, Bockeria L. Local Use of Hydrogel with Amiodarone in Cardiac Surgery: Experiment and Translation to the Clinic. *Gels*. 2021;7. doi: 10.3390/gels7010029
18. Wang W, Mei YQ, Yuan XH, Feng XD. Clinical efficacy of epicardial application of drug-releasing hydrogels to prevent postoperative atrial fibrillation. *J Thorac Cardiovasc Surg*. 2016;151:80-85. doi: 10.1016/j.jtcvs.2015.06.061
19. Greenstein D, Beau J, Gottlieb G, Teller D, Kulik A. Topical amiodarone during cardiac surgery: Does epicardial application of amiodarone prevent postoperative atrial fibrillation? *J Thorac Cardiovasc Surg*. 2017;154:886-892. doi: 10.1016/j.jtcvs.2017.04.013
20. Garcia JR, Campbell PF, Kumar G, Langberg JJ, Cesar L, Deppen JN, Shin EY, Bhatia NK, Wang L, Xu K, et al. Minimally Invasive Delivery of Hydrogel-Encapsulated Amiodarone to the Epicardium Reduces Atrial Fibrillation. *Circ Arrhythm Electrophysiol*. 2018;11:e006408. doi: 10.1161/CIRCEP.118.006408
21. Stratford-Perricaudet LD, Makeh I, Perricaudet M, Briand P. Widespread long-term gene transfer to mouse skeletal muscles and heart. *J Clin Invest*. 1992;90:626-630. doi: 10.1172/JCI115902
22. Barr E, Carroll J, Kalynych AM, Tripathy SK, Kozarsky K, Wilson JM, Leiden JM. Efficient catheter-mediated gene transfer into the heart using replication-defective adenovirus. *Gene Ther*. 1994;1:51-58.
23. Fleury S, Simeoni E, Zuppinger C, Deglon N, von Segesser LK, Kappenberger L, Vassalli G. Multiply attenuated, self-inactivating lentiviral vectors efficiently deliver and express genes for extended periods of time in adult rat cardiomyocytes in vivo. *Circulation*. 2003;107:2375-2382. doi: 10.1161/01.CIR.0000065598.46411.EF
24. Kotchey NM, Adachi K, Zahid M, Inagaki K, Charan R, Parker RS, Nakai H. A potential role of distinctively delayed blood clearance of recombinant adeno-associated virus serotype 9 in robust cardiac transduction. *Mol Ther*. 2011;19:1079-1089. doi: 10.1038/mt.2011.3
25. Zincarelli C, Soltys S, Rengo G, Rabinowitz JE. Analysis of AAV serotypes 1-9 mediated gene expression and tropism in mice after systemic injection. *Mol Ther*. 2008;16:1073-1080. doi: 10.1038/mt.2008.76
26. Sun J, Shen H, Shao L, Teng X, Chen Y, Liu X, Yang Z, Shen Z. HIF-1 α overexpression in mesenchymal stem cell-derived exosomes mediates cardioprotection in myocardial infarction by enhanced angiogenesis. *Stem Cell Res Ther*. 2020;11:373. doi: 10.1186/s13287-020-01881-7
27. Wei J, Liu F, Lu Z, Fei Q, Ai Y, He PC, Shi H, Cui X, Su R, Klungland A, et al. Differential m(6)A, m(6)A(m), and m(1)A Demethylation Mediated by FTO in the Cell Nucleus and Cytoplasm. *Mol Cell*. 2018;71:973-985 e975. doi: 10.1016/j.molcel.2018.08.011

28. Zhang J, Ma A, Shang L. Conjugating Existing Clinical Drugs With Gold Nanoparticles for Better Treatment of Heart Diseases. *Front Physiol.* 2018;9:642. doi: 10.3389/fphys.2018.00642
29. Chen S, Shohet RV, Bekeredjian R, Frenkel P, Grayburn PA. Optimization of ultrasound parameters for cardiac gene delivery of adenoviral or plasmid deoxyribonucleic acid by ultrasound-targeted microbubble destruction. *J Am Coll Cardiol.* 2003;42:301-308. doi: 10.1016/s0735-1097(03)00627-2
30. Shohet RV, Chen S, Zhou YT, Wang Z, Meidell RS, Unger RH, Grayburn PA. Echocardiographic destruction of albumin microbubbles directs gene delivery to the myocardium. *Circulation.* 2000;101:2554-2556. doi: 10.1161/01.cir.101.22.2554
31. Guan L, Wang C, Yan X, Liu L, Li Y, Mu Y. A thrombolytic therapy using diagnostic ultrasound combined with RGDS-targeted microbubbles and urokinase in a rabbit model. *Sci Rep.* 2020;10:12511. doi: 10.1038/s41598-020-69202-9
32. Taylor RE, Zahid M. Cell Penetrating Peptides, Novel Vectors for Gene Therapy. *Pharmaceutics.* 2020;12. doi: 10.3390/pharmaceutics12030225
33. Frankel AD, Pabo CO. Cellular uptake of the tat protein from human immunodeficiency virus. *Cell.* 1988;55:1189-1193. doi: 10.1016/0092-8674(88)90263-2
34. Green M, Loewenstein PM. Autonomous functional domains of chemically synthesized human immunodeficiency virus tat trans-activator protein. *Cell.* 1988;55:1179-1188. doi: 10.1016/0092-8674(88)90262-0
35. Joliot A, Pernelle C, Deagostini-Bazin H, Prochiantz A. Antennapedia homeobox peptide regulates neural morphogenesis. *Proc Natl Acad Sci U S A.* 1991;88:1864-1868. doi: 10.1073/pnas.88.5.1864
36. Green M, Ishino M, Loewenstein PM. Mutational analysis of HIV-1 Tat minimal domain peptides: identification of trans-dominant mutants that suppress HIV-LTR-driven gene expression. *Cell.* 1989;58:215-223. doi: 10.1016/0092-8674(89)90417-0
37. Schwarze SR, Ho A, Vocero-Akbani A, Dowdy SF. In vivo protein transduction: delivery of a biologically active protein into the mouse. *Science.* 1999;285:1569-1572. doi: 10.1126/science.285.5433.1569
38. Smith GP. Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science.* 1985;228:1315-1317. doi: 10.1126/science.4001944
39. Zahid M, Robbins PD. Identification and characterization of tissue-specific protein transduction domains using peptide phage display. *Methods Mol Biol.* 2011;683:277-289. doi: 10.1007/978-1-60761-919-2_20
40. Zahid M, Phillips BE, Albers SM, Giannoukakis N, Watkins SC, Robbins PD. Identification of a cardiac specific protein transduction domain by in vivo biopanning using a M13 phage peptide display library in mice. *PLoS One.* 2010;5:e12252. doi: 10.1371/journal.pone.0012252
41. Molenaar TJ, Michon I, de Haas SA, van Berkel TJ, Kuiper J, Biessen EA. Uptake and processing of modified bacteriophage M13 in mice: implications for phage display. *Virology.* 2002;293:182-191. doi: 10.1006/viro.2001.1254
42. Zahid M, Feldman KS, Garcia-Borrero G, Feinstein TN, Pogodzinski N, Xu X, Yurko R, Czachowski M, Wu YL, Mason NS, et al. Cardiac Targeting Peptide, a Novel Cardiac Vector: Studies in Bio-Distribution, Imaging Application, and Mechanism of Transduction. *Biomolecules.* 2018;8. doi: 10.3390/biom8040147
43. McGuire MJ, Samli KN, Johnston SA, Brown KC. In vitro selection of a peptide with high selectivity for cardiomyocytes in vivo. *J Mol Biol.* 2004;342:171-182. doi: 10.1016/j.jmb.2004.06.029
44. Mentkowski KI, Lang JK. Exosomes Engineered to Express a Cardiomyocyte Binding Peptide Demonstrate Improved Cardiac Retention in Vivo. *Sci Rep.* 2019;9:10041. doi: 10.1038/s41598-019-46407-1
45. Wang Z, Yang N, Hou Y, Li Y, Yin C, Yang E, Cao H, Hu G, Xue J, Yang J, et al. L-Arginine-Loaded Gold Nanocages Ameliorate Myocardial Ischemia/Reperfusion Injury by Promoting Nitric Oxide Production and Maintaining Mitochondrial Function. *Adv Sci (Weinh).* 2023;10:e2302123. doi: 10.1002/advs.202302123
46. Avula UM, Yoon HK, Lee CH, Kaur K, Ramirez RJ, Takemoto Y, Ennis SR, Morady F, Herron T, Berenfeld O, et al. Cell-selective arrhythmia ablation for photomodulation of heart rhythm. *Sci Transl Med.* 2015;7:311ra172. doi: 10.1126/scitranslmed.aab3665
47. Kim H, Yun N, Mun D, Kang JY, Lee SH, Park H, Park H, Joung B. Cardiac-specific delivery by cardiac tissue-targeting peptide-expressing exosomes. *Biochem Biophys Res Commun.* 2018;499:803-808. doi: 10.1016/j.bbrc.2018.03.227
48. Kang JY, Kim H, Mun D, Yun N, Joung B. Co-delivery of curcumin and miRNA-144-3p using heart-targeted extracellular vesicles enhances the therapeutic efficacy for myocardial infarction. *J Control Release.* 2021;331:62-73. doi: 10.1016/j.jconrel.2021.01.018
49. Kim H, Mun D, Kang JY, Lee SH, Yun N, Joung B. Improved cardiac-specific delivery of RAGE siRNA within small extracellular vesicles engineered to express intense cardiac targeting peptide attenuates myocarditis. *Mol Ther Nucleic Acids.* 2021;24:1024-1032. doi: 10.1016/j.omtn.2021.04.018
50. Gallicano GI, Fu J, Mahapatra S, Sharma MVR, Dillon C, Deng C, Zahid M. Reversing Cardiac Hypertrophy at the Source Using a Cardiac Targeting Peptide Linked to miRNA106a: Targeting Genes That Cause Cardiac Hypertrophy. *Pharmaceutics (Basel).* 2022;15. doi: 10.3390/ph15070871

51. Yurko R, Islam K, Weber B, Salama G, Zahid M. Conjugation of amiodarone to a novel cardiomyocyte cell penetrating peptide for potential targeted delivery to the heart. *Front Chem.* 2023;11:1220573. doi: 10.3389/fchem.2023.1220573
52. Zahid M, Weber B, Yurko R, Islam K, Agrawal V, Lopuszynski J, Yagi H, Salama G. Cardiomyocyte-Targeting Peptide to Deliver Amiodarone. *Pharmaceutics.* 2023;15:2107.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.