

Case Report

Not peer-reviewed version

# VRAM flap for pelvic floor reconstruction after pelvic exenteration and abdominoperineal excision

[Ionut Flaviu Faur](#) , [Adelina Clim](#) , [Amadeus Dobrescu](#) \* , [Catalin Prodan](#) , Rami Hajjar , [Paul Pasca](#) , Marco Capitanio , [Ionel Nati](#) , [Cristi Tarta](#) , Alexandru Isaic , George Noditi , [Bogdan Totolici](#) , Ciprian Duta , Gabriel Lazar

Posted Date: 1 December 2023

doi: 10.20944/preprints202311.1995.v1

Keywords: vertical rectus abdominis muscle flap; perineal wound; pelvic exenteration; squamosal cell carcinoma of the uterine cervix; reconstruction



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Case Report*

# VRAM Flap for Pelvic Floor Reconstruction After Pelvic Exenteration and Abdominoperineal Excision

Ionut Flaviu Faur <sup>1,2</sup>, Adelina Clim <sup>3</sup>, Amadeus Dobrescu <sup>1,2,\*</sup>, Catalin Prodan <sup>1,2</sup>, Rami Hajjar <sup>1</sup>, Paul Pasca <sup>1</sup>, Marco Capitanio <sup>1</sup>, Cristi Tarta <sup>1,2</sup>, Alexandru Isaic <sup>1,2</sup>, George Noditi <sup>1,2</sup>, Ionel Nati <sup>4,5</sup>, Bogdan Totolici <sup>6,7</sup>, Ciprian Duta <sup>1,2</sup> and Gabriel Lazar <sup>4,5</sup>

<sup>1</sup> II nd Surgery Clinic, Timisoara Emergency County Hospital, Romania; dr.faurflaviuionut@gmail.com

<sup>2</sup> "Victor Babes" University of Medicine and Pharmacy Timisoara Romania

<sup>3</sup> II nd Obstetric and Gynecology Clinic "Dominic Stanca" 400123 Cluj-Napoca, Romania

<sup>4</sup> "Iuliu Hatieganu" University of Medicine and Pharmacy Cluj-Napoca, Romania

<sup>5</sup> Ist Clinic of Oncological Surgery, Oncological Institute "Prof Dr I Chiricuta" Cluj Napoca, Romania

<sup>6</sup> Ist Clinic of General Surgery, Arad County Emergency Clinical Hospital, Romania

<sup>7</sup> "Vasile Goldis" Western University of Arad, Romania

\* Correspondence: dobrescu.amadeus@umft.ro; Tel.: +40-71-577-091

**Abstract:** Due to the still large number of patients diagnosed with pelvic neoplasms (colorectal, gynecological, and urological) in advanced stages right from the initial diagnosis, surgery represents the mainstay of treatment, often implying wide, eventually multi-organ resections in order to achieve negative surgical margins. Perineal wound morbidity, particularly in extralevator abdominoperineal excision, leads to complications and local infection rates of up to 40%. Strategies to reduce postoperative wound complications are pursued to address this issue. The VRAM flap remains the gold standard for autologous reconstruction after pelvic oncological resection; it was initially designed for abdominal wall defects and later expanded for large pelvic tissue defects. The flap's application is based on its physical characteristics, including abundant tissue and a generous skin paddle, which effectively obliterates dead space after exenterations. The generous skin paddle offers good cosmetic and functional outcomes at the recipient site. This article describes the case of a patient histopathologically diagnosed with stage IIIA squamous cell carcinoma of the uterine cervix who received multimodal onco-surgical treatment. The surgical mainstay of this treatment is pelvic exenteration. Pelvic reconstruction after this major surgery was performed using a vertical flap with the rectus abdominis.

**Keywords:** vertical rectus abdominis muscle flap; perineal wound; pelvic exenteration; squamosal cell carcinoma of the uterine cervix; reconstruction

## 1. Introduction

Cancer still represents a major cause of morbidity and mortality worldwide, regardless of socioeconomic characteristics and/or demographics. According to the GLOBOCAN stats published for 2020, an estimated 19.3 million new cancer cases and almost 10.0 million cancer deaths occurred [1]. Malignant tumors arising from the pelvic region seem to particularly contribute to the numbers mentioned above. Pelvic malignancies death rates are decreasing due to increased screening rates and global sensibilization campaigns, reducing financial burdens and promoting better health outcomes. Nonetheless, patients often present with advanced stages due to lack of early and specific symptoms, reason why they do not seek medical attention until signs of extensive loco-regional involvement appear.

When such stages of advancement are reached, surgery represents the mainstay of treatment, often implying wide, eventually multi-organ resections in order to achieve negative surgical margins.

R0 resection represent the main goal of such procedures, as it has been proven to chiefly influence and consistently predict disease-free survival and postoperative recurrence rates. Pelvic exenteration and abdominoperineal resection are the main procedures for locally advanced malignancies in the anorectum, gynecological, or urological pelvic organs.

The main source of morbidity is the perineal wound and complications, with an estimated rate of local infection up to 40%. Flap reconstruction techniques for pelvic defects gain popularity due to their potential to address surgical challenges and reduce infection and dehiscence risks by appointing healthy, vascularized autologous tissue.

The aim of this paper is to reveal the benefits of a perineal reconstruction technique using a vertical rectus abdominis flap after pelvic exenteration. The present case is an example of reconstruction of the perineal anatomical region by means of a vertical rectus abdominis muscle flap, reducing the well-known complications of perineal lacerations after pelvic exenteration.

## 2. Materials and methods

### 2.1. Diagnostic and stadialisation

A 63-year-old female without other associated pathologies presented to Gynecology Department in April 2017 with metrorrhagia, faintness and dizziness.

Following the clinical and paraclinical examination, the diagnosis of menopausal metrorrhagia was established and a primary carcinoma of the cervix was suspected. Aside from the levels of red blood cell (RBC) and hemoglobin (HGB) of 4.01 and 7.1, respectively, the patient's laboratory results were normal.

A biopsy of the cervix was performed and the excised material has been sent for histopathological examination. Histopathological results: Squamous cell carcinoma of the cervix - G1 stage IIIA.

This CT scan of the Thorax, Abdomen and Pelvis (15.06.2017). showed rare solid nodules (1-2/ pulmonary field); the cervix enlarged in dimensions with transverse diameter of 5 cm, due to a proliferative process with an infiltrative appearance of approximately 4 cm in diameter, also extended to the level of uterine body. The tumor comes into contact with the posterior wall of the urinary bladder and the anterior wall of the rectum, which seems to be clearly defined. Infiltration of both parameters and minimal carcinomatous infiltration of pelvic fat showing fine solid septa and rare hyperdense micronodules. Rare infracentimeter lumboaortic adenopathies.

The MRI Abdomen+Pelvis (18.10.2022) showed a tumor formation developed at the level of the cervix, 7.8 cm circumferential, with bilateral parametrial invasion and lower extension at the level of the vagina and vulva on the right side, invasion of the levator ani muscle on the right side, with total invasion of the urethra along the entire path, and in contact with the anterior rectal wall without MRI signs of invasion, multiple bilateral enlarged inguinal lymph nodes up to 14 mm in size without secondary dissemination at the level of the scanned bone structures.

CT Thorax+Abdomen+Pelvis (31.10.2022) were performed due to the patient's enlarged inguinal lymph nodes. In the region of the cervix, a formation of approximately 55/40 mm diameter with a hypodense center and iodophilic, irregular periphery is evident. Without suspicious pelvic adenopathies, retroperitoneal lymph nodes of 6-7 mm and without signs of bone metastases.

The PET-CT scan (December 2022) shows the presence of a 75/45 mm FDG (SUVIbm= 7,9) hypercapturing tumor formation centered on the cervix with invasion in vagina up to perineal level was confirmed, without significant dimensional or metabolic active retroperitoneal, pelvic or inguinal adenopathy.

## 3. Results

### 3.1. Treatment plan

During the patient's hospitalization, hemostatic and analgesic treatment was administered (Alindor 500 mg, Adrenostazin 0.3mg/ml, Etamsylate 250 mg, Fitomenadion 10 mg/ml). The patient was discharged with the recommendation of a gynecological consultation in 2 weeks.

External Radiotherapy with Intensity Radiated Modulation Therapy (IMRT) for neoadjuvant purposes was proposed (06 – 07/2017). The patient was recommended to undergo radio-chemotherapy sessions with brachytherapy. In June and July 2017 radiotherapy was practiced for

neoadjuvant purposes up to a total dose of 42 Gy/ 21 fr.( which included pelvic nodules, uterus, both parameters and the upper third of the vagina). The chemotherapy was also proposed, but unfortunately, the patient refused it.

Cancellation of brachytherapy (17.07.2017). It should be mentioned that brachytherapy could no longer be performed due to hemorrhage (local hemostatic treatment with Gelaspon and intravenous Adrenostazin and Etamsylate was administered).

The patient is indicated for radical oncological surgery - pelvic exenteration.

### 3.2. *Expected outcome of the treatment plan*

Evolution without treatment (Disease progression without specialized therapy)

Without treatment, cervical cancer progresses more slowly or more rapidly towards loco-regional invasion (of regional lymph nodes, parametrium, nearby blood vessels and neighbouring organs) and systemic metastases (lung, bone, liver, distant lymph nodes).

Evolution with specialized treatment

This category details the evolution and possible complications following surgical, radiotherapy and chemotherapy treatment.

Total pelvic exenteration has been used as a salvage therapy. Candidates are those who have failed radiation therapy or primary surgical or combined treatment of the recurrence in the central pelvis and in cases of locally advanced primary pelvic tumors.

### 3.3. *Actual outcome*

In October 2022, the patient who was treated with exclusive irradiation in 2017, presents with pelvic recurrence with the invasion of the vagina, urethra and parameters bilaterally.

The clinical examination reveals the presence of a friable, hemorrhagic tumor formation at the level of the right hemivulva and vagina.

### 3.4. *Literature review*

A brief advanced research on PubMed about the topic - carried out using the MeSH terms "perineum" and "surgical flaps" - yielded a total of 14 results by filtering the level of the study to only meta-analysis and systematic review. Further on, studies addressing either primary benign or secondary pathology, such as local infection (either primary or secondary after previous pelvic surgery), benign proctology, hidradenitis suppurativa, were excluded, leaving a total pool of 6 contextually relevant studies – they are listed in Table 1. All of them seem to universally agree by proof of statistical evidence on how flap reconstruction procedures positively influence outcome and wound healing compared to primary closure techniques [16,17,19,20] even though the marked heterogeneity of the various study populations in terms of different types of flaps adopted does not allow for a better insight on the VRAM flap itself. While Johnstone et al. in 2017 managed to realize a statistical analysis that proved how the VRAM technique presents a lower incidence of perineal wound and flap complications [20], Esemé et al. in 2022 gathered data from 925 patients in a proportional meta-analysis strictly comparing the VRAM flap with the gracilis one, recording a higher incidence of donor site hernias [18]. Mean follow-up is not reported for all the studies mentioned above, while other patient data potentially crucial for the reliability of study outcomes are absent or heterogeneously expressed (e.g. mean age, prevalence of neoadjuvant treatment). Standardization probably represents the key factor for information reporting in order to establish some clear indications meant to aid the surgeon through the process of decision making in the best interest of the patient prognosis. A noteworthy insight has been already proposed by Yan et al into a meta-analysis that evaluate the results obtained through the application of Enhanced Recovery After Surgery (ERAS) protocols in the postoperative management of autologous flap-based reconstruction, proving a substantial benefit manifested by decreased length of hospital stay without observing an increase in the incidence of complications [23]. Despite that, the lack of a strong, statistically uniform body of data clearly supporting such protocols and measures for enhanced recovery in the field of




flap-based perineal reconstruction does not allow for their standard applications on a global scale; further research is therefore still warranted.

**Table 1.** Meta-Analysis and Systematic Reviews identified after advanced research on PubMed (MeSH terms “perineum”, “surgical flaps”) and exclusion.\* = expressed as a mean.

Reference	Study Design	Total No. of patients	Mean Age	Type of Cancer	Type of resection	Type of Flap	Duration of Follow-up	Conclusion
Devulapalli et al.16, 2016	Meta-Analysis	566	59.1	Rectal Cancer, Anal Cancer, Others	APE, PE	VRAM, Gracilis	24 months*	Flap reconstruction reduces wound morbidity after APE, PE compared to primary closure
Yang et al.17, 2019	Meta-Analysis	17.913	NS	Rectal Cancer	APE, ELAPE, PE	VRAM, Gracilis, Gluteal	17 – 50.4 months	Flap reconstruction reduces wound complication compared to primary closure
Eseme et al.18,2022	Proportional Meta-Analysis	925	50.3 - 67	Rectal, Anal, Vaginal, Others	APE, PE	VRAM, Gracilis	NS	Flaps are safe for vulvoperineal reconstruction; VRAM raises risk for donor site hernia
Buscail et al.19, 2021	Meta-Analysis	2180	NS	Rectal, Anal, Others	APE, PE	VRAM	NS	Flaps reduce incidence of wound complication after APE
Johnstone20, 2017	Systematic Review	649	NS	Rectal, Anal, Gynecological, Others	APE, PE	VRAM, Gracilis, Gluteal	NS	VRAM superior in terms of incidence of wound complication
Foster et al.21, 2012	Systematic Review	522 (one study NS)	NS	Rectal, Anal	ELAPE	VRAM, Gracilis, Gluteal,	10 – 38 months*	No significant difference in complications between flap and mesh repairs


3.5..... Ce scriem aici?



(a)



(b)




(c)




(d)



(e)



(f)



(g)



(h)



(i)



(j)



(k)

**Figure 1.** (a) - Completing abdominal surgical time and performing perineal time. Pre-incision measurement of the desired/ intended abdominal myocutaneous flap (16x8 cm). (b) - Preparation of the myocutaneous flap of the rectus abdominis muscle. (c;d;e) - Preparation of the myocutaneous flap of the rectus abdominis muscle. (f;g) - Appearance of the perineal wound before and after the application of the flap. (h;i) - En-bloc surgical resection specimen (j) - Final appearance of the perineal region after vertical abdominal rectus muscle flap reconstruction. (k) - Final aspect of the stoma and of the abdominal scar.

#### 4. Discussion

Malignant tumors arising from the pelvic region seem to particularly contribute to the numbers mentioned above. Declining death rates for pelvic malignancies such as colorectal cancer (CRC), prostate cancer, cervical cancer and other gynecological entities can be substantially ascribed to the increased rate of screening observed along the past years due to intensive campaigns of sensibilization deployed by governments around the world in order to buffer the financial burden that cancers as an entity exerts. Nonetheless, patients often present with advanced stages due to lack of early and specific symptoms, reason why they do not seek medical attention until signs of extensive loco-regional involvement appear.

Pelvic exenteration (PE), initially described by Brunschwig in 1948 in New York as a palliative procedure for recurrent carcinoma of the cervix [4], together with abdominoperineal resection (APR), firstly proposed by Miles in 1908 [5] but subsequently revised and modified [6], are the two main procedures performed nowadays for locally advanced malignancies arising from either the anorectum, gynecological or urological pelvic organs. What probably represent the main source of morbidity, especially in the case of extralevator abdominoperineal excision (ELAPE), is the perineal



wound and its complications, with an estimated rate of local infection up to 40%. Other complications are not uncommon: from early fluid collections, which eventually turn into abscesses, and wound dehiscence, to chronic fistulas and perineal hernias due to weakening of the pelvic floor; also, migration of bowel loops has been observed into the large remaining pelvic defect, leading to bowel movements disturbances and reduced patient's quality of life. Techniques using the omentum to suspend the bowel and ileal conduit out of the pelvis or fill it have reduced the incidence rate for abscesses and fistulas [7].

Due to such levels of perineal wound morbidity, coupled with the steadily increasing number of pelvic exenteration and ELAPE performed, different strategies have been pursued in order to try and lower the incidence of postoperative wound complications. Primary closure of the wound probably revealed itself as the riskier option available, especially considering the increasing prevalence of patients presenting for surgery after neoadjuvant radiotherapy, which significantly impacts pelvic tissues, making them prone to develop local ischemia, fibrosis and poor healing potential – some series report complication rates up to 57% [8]. Moreover, as already mentioned above, primary closure harbors a greater risk for bowel herniation into the pelvic space; this issue has also been addressed by the introduction of biological mesh repair's techniques that reduce tissue tension inside the wound: so far they have been widely employed around the world with satisfactory results and lower rates of perineal hernia compared to primary closure. Unfortunately, despite lowering tension, biological meshes do not bring benefits regarding other common points of concern encountered along the healing phase of the postoperative perineal wound, such as tendency to fluid collections with consequent early seromas and abscesses, and delayed healing due to the surrounding, poorly vascularized tissues affected by neoadjuvant radiotherapy [9].

Flap reconstruction techniques for pelvic defects gained growing popularity along the years, most likely because of their potential to aid facing the main surgical challenges presented by the rising prevalence of irradiated patients; in fact, by the apposition of new, healthy and vascularized autologous tissue, not only local tension levels decrease, but infection and dehiscence risk as well, due to the greater supply of oxygen, nutrients and inflammatory cells to the wound bed [10].

In terms of common practice, the current golden standard for autologous reconstruction after pelvic oncological resection remains the vertical rectus abdominis muscle (VRAM) flap, firstly described by Mathes and Botswick in 1977 for the repair of various defects of the abdominal wall, and further adapted later on for large pelvic tissue defects [11,12]. The rationale behind its use relies on some critical crucial factors: first, the physical characteristics of the flap itself, provided with abundant amounts of tissue forming a bulky, optimally vascularized mass well suited for obliterating the dead space left after large exenterations; second, the generous skin paddle provides good cosmetic and functional outcomes at the recipient site; last but not least, the anatomical features of the pedicle mobilized for this flap allow for a good range of movement and rotation to reach for distant defects [13,14].

Other options for flap closure are represented by inner thigh flaps, with particular attention reserved for the gracilis flap [15], requiring lower expertise for tailoring and preparation and supposedly harboring a lower risk for donor site complication, especially ventral hernias, compared to VRAM. The choice for the perfect technique for pelvic reconstruction fuels many debates worldwide in terms of comparison in between the various procedures and their individual indications and/or advantages, especially due to the various discrepancies in between findings published in the literature and the lack of large studies with low bias of interpretation.

## 5. Conclusions

The rectus abdominis muscle flap could eventually eclipse all other techniques for the reconstruction of the pelvic floor after abdominoperineal resection and pelvic exenteration. Multi-center studies are required in this context to collect a significant amount of data indicating and proving the utility of the rectus abdominis flap higher than other flap techniques already commonly used and discussed in the medical literature.



**Author Contributions:** Conceptualisation: Ionut Flaviu Faur and Amadeus Dobrescu. Methodology: Adelina Clim and Catalin Prodan. Software and validation: Rami Hajjar and Paul Pasca. Formal analysis and investigation: Marco Capitanio and Cristi Tarta. Resources and data curation: Alexandru Isaic and George Noditi. Writing: Ionel Nati. Project administration and supervision: Bogdan Totolici, Ciprian Duta and Gabriel Lazar.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the Helsinki ethics statement and approved by the Ethics Committee under the following number: 18906062022 - Application Number 10444.

**Informed Consent Statement:** Written informed consent has been obtained from the patients to publish this paper.

**Data Availability Statement:** MDPI research data politics.

**Acknowledgments:** We have nothing to declare in this section.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Sung, H, Ferlay, J, Siegel, RL, Laversanne, M, Soerjomataram, I, Jemal, A, Bray, F. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2021; 71: 209- 249. <https://doi.org/10.3322/caac.21660>
2. World Health Organization (WHO). Global Health Estimates 2020: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2019. WHO; 2020. Accessed December 11, 2020. <https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates/ghe-leading-causes-of-death>
3. PelvEx Collaboration. Surgical and survival outcomes following pelvic exenteration for locally advanced primary rectal cancer: results from an international collaboration. *Ann Surg.* 2019;269:315–21.
4. Brunschwig A. Complete excision of pelvic viscera for advanced carcinoma; a one-stage abdominoperineal operation with end colostomy and bilateral ureteral implantation into the colon above the colostomy. *Cancer.* 1948;1:177–83.
5. Miles W E. A method of performing abdominoperineal excision for carcinoma of the rectum and of the terminal portion of the pelvic colon. *Lancet.* 1908;2:1812–1813.
6. Beyond TME Consensus. Consensus statement on the multidisciplinary management of patients with recurrent and primary rectal cancer beyond total mesorectal excision planes. *Br J Surg.* 2013;100:1009–14
7. Valle G, Ferraris G. Use of the omentum to contain the intestines in pelvic exenteration. *Obstet Gynecol.* 1969 Jun;33(6):772-5.
8. Musters, G.D.; Buskens, C.J.; Bemelman, W.A.; Tanis, P.J. Perineal Wound Healing After Abdominoperineal Resection for Rectal Cancer. *Dis. Colon Rectum* 2014, 57, 1129–1139.
9. Foster, J.D., Tou, S., Curtis, N.J., Smart, N.J., Acheson, A., Maxwell-Armstrong, C., Watts, A., Singh, B. and Francis, N.K. (2018), Closure of the perineal defect after abdominoperineal excision for rectal adenocarcinoma – ACPGBI Position Statement. *Colorectal Dis*, 20: 5-23.
10. Stein, M. J., Karir, A., Ramji, M., Allen, M., Bain, J. R., Avram, R., Boushey, R., Auer, R., & Jarmuske, M. (2019). Surgical outcomes of VRAM versus gracilis flaps for the reconstruction of pelvic defects following oncologic resection. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 72(4), 565–571.
11. Mathes S, Bostwick J 3rd. A rectus abdominis myocutaneous flap to reconstruct abdominal wall defects. *Br J Plast Surg* 1977; 30: 282–283.
12. Shkula HS, Hughes LE. The rectus abdominis flap for perineal wounds. *Ann R Coll Surg Engl* 1984; 66: 337–339.
13. Nelson, R. A., & Butler, C. E. (2009). Surgical outcomes of VRAM versus thigh flaps for immediate reconstruction of pelvic and perineal cancer resection defects. *Plastic & Reconstructive Surgery*, 123(1), 175–183. <https://doi.org/10.1097/prs.0b013e3181904df7>
14. McMenamin, D. M., Clements, D., Edwards, T. J., Fitton, A. R., & Douie, W. J. P. (2011). Rectus Abdominis Myocutaneous flaps for perineal reconstruction: Modifications to the technique based on a large single-centre experience. *The Annals of The Royal College of Surgeons of England*, 93(5), 375–381. <https://doi.org/10.1308/003588411x572268>
15. Burke TW, Morris M, Roh MS, Levenback C, Gershenson DM. Perineal reconstruction using single gracilis myocutaneous flaps. *Gynecol Oncol.* 1995;57:221–225
16. Devulapalli, C., Jia Wei, A. T., DiBiagio, J. R., Baez, M. L., Baltodano, P. A., Seal, S. M., Sacks, J. M., Cooney, C. M., & Rosson, G. D. (2016). Primary versus flap closure of perineal defects following oncologic resection. *Plastic and Reconstructive Surgery*, 137(5), 1602–1613. <https://doi.org/10.1097/prs.0000000000002107>

17. Yang, X. Y., Wei, M. T., Yang, X. T., He, Y. Z., Hao, Y., Zhang, X. B., Deng, X. B., Wang, Z. Q., & Zhou, Z. Q. (2019). Primary myocutaneous flap closure of perineal defects following abdominoperineal resection for colorectal disease: A systematic review and meta-analysis. *Colorectal Disease*, 21(2), 138–155. <https://doi.org/10.1111/codi.14471>
18. Esemé, E. A., Scampa, M., Viscardi, J. A., Ebai, M., Kalbermatten, D. F., & Oranges, C. M. (2022). Surgical outcomes of Vram vs. Gracilis Flaps in vulvo-perineal reconstruction following oncologic resection: A proportional meta-analysis. *Cancers*, 14(17), 4300. <https://doi.org/10.3390/cancers14174300>
19. Buscail, E., Canivet, C., Shourick, J., Chantalat, E., Carrere, N., Duffas, J.-P., Philis, A., Berard, E., Buscail, L., Ghouti, L., & Chaput, B. (2021). Perineal wound closure following abdominoperineal resection and pelvic exenteration for cancer: A systematic review and meta-analysis. *Cancers*, 13(4), 721. <https://doi.org/10.3390/cancers13040721>
20. Johnstone, M. S. (2017). Vertical rectus abdominis myocutaneous versus alternative flaps for perineal repair after abdominoperineal excision of the rectum in the era of laparoscopic surgery. *Annals of Plastic Surgery*, 79(1), 101–106. <https://doi.org/10.1097/sap.0000000000001137>
21. Foster, J. D., Pathak, S., Smart, N. J., Branagan, G., Longman, R. J., Thomas, M.G., & Francis, N. (2012). Reconstruction of the perineum following extralevator abdominoperineal excision for carcinoma of the lower rectum: a systematic review. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland*, 14(9), 1052–1059.
22. Radwan, R. W., Tang, A. M., Harries, R. L., Davies, E. G., Drew, P., & Evans, M. D. (2021). Vertical rectus abdominis flap (VRAM) for perineal reconstruction following pelvic surgery: A systematic review. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 74(3), 523–529. <https://doi.org/10.1016/j.bjps.2020.10.100>
23. Tan, Y. Y., Liaw, F., Warner, R., Myers, S., & Ghanem, A. (2021). Enhanced Recovery Pathways for Flap-based reconstruction: Systematic Review and meta-analysis. *Aesthetic Plastic Surgery*, 45(5), 2096–2115. <https://doi.org/10.1007/s00266-021-02233-3>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.