

Short Note

Not peer-reviewed version

Brown-Séquard Syndrome: Understanding the Complexities of Spinal Cord Injury

[Dev Desai](#) , Shailkumar Desai , [Vismith Gami](#) *

Posted Date: 21 December 2023

doi: 10.20944/preprints202312.1595.v1

Keywords: Brown-Séquard syndrome; trauma; motor; sensory; autonomous



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Brown-Séquard Syndrome: Understanding the Complexities of Spinal Cord Injury

Dev Desai, Vismith Gami * and Shailekumar Desai

Smt. NHLMMC, Ahmedabad, Gujarat, India; devhdesai01@gmail.com; Shailekumar11@gmail.com

* Correspondence: Vismith.patel123@gmail.com

Abstract: Brown-Séquard Syndrome (BSS) is a rare spinal cord injury characterized by asymmetrical motor and sensory dysfunction. Trauma to the spinal cord, particularly the hemi section, is the primary cause. Diagnosis involves a thorough medical history, physical examination, and imaging studies. Treatment focuses on stabilization, rehabilitation, and potential emerging therapies. This article provides a comprehensive overview of BSS, including its causes, symptoms, diagnosis, and treatment.

Keywords: brown-séquard syndrome; trauma; motor; sensory; autonomous

Introduction

Spinal cord injuries can result in a wide range of neurological deficits, and one rare yet fascinating condition is Brown-Séquard Syndrome (BSS). Brown-Séquard Syndrome (BSS) causes an uncommon neurological condition that causes a unique collection of sensory, motor, and autonomic dysfunctions. This syndrome, which Charles-Édouard Brown-Séquard first identified around 1850, frequently results from spinal cord damage or other traumatic causes(1). This study of the literature intends an overview of the aetiology, clinical presentation, diagnostic techniques, treatment options, and prognosis of Brown-Séquard Syndrome.

Aetiology

The most frequent cause of Brown-Séquard Syndrome is trauma, which frequently results from piercing wounds like gunshots or stab wounds(2). Additional reasons include spinal cord tumours, degenerative illnesses, viral disorders, hematomas(3) and iatrogenic injuries sustained during surgical treatments. The affected side experiences varying degrees of motor paralysis and loss of proprioception (awareness of body position) due to damage to the corticospinal tract and dorsal columns, respectively.

Symptomatic Presentation

The symptoms of BSS depend on the level of the spinal cord injury. For instance, if the injury occurs in the cervical region, both arms may be affected, while a thoracic-level injury affects the lower limbs.(4) The ipsilateral motor paralysis, ipsilateral loss of proprioception, and contralateral loss of pain and temperature sensation below the site of the lesion make up the typical clinical trial of Brown-Séquard Syndrome(5). The spinal cord's physical configuration, in which proprioceptive and motor fibres ascend ipsilaterally and pain and temperature routes decussate and ascend contralaterally, results in this pattern(6) However, discrepancies in how BSS is presented have been noted, highlighting the need for individual patient assessment.

Diagnostic strategies

Brown-Séquard syndrome must be identified through a comprehensive clinical assessment. Imaging methods like computed tomography (CT) scans and magnetic resonance imaging (MRI) is essential for locating the exact position and size of the lesion in the spinal cord(7). Additional tests,

such as electromyography (EMG) and somatosensory evoked potentials (SSEP)(8), may be conducted to assess the integrity of neural pathways and provide further diagnostic confirmation. These tests help differentiate BSS from other spinal cord injuries and neurological conditions.

Management Techniques

A comprehensive approach is used to treat Brown-Séquard syndrome with the goal of reducing functional limitations and raising the quality of life(9). The main goals of acute interventions are to stabilise the patient, treat any injuries that are present, and guard against additional problems like respiratory compromise and deep vein thrombosis. Maximising functional recovery requires early rehabilitation, which includes physical and occupational therapy. Pharmacological interventions such as analgesics, muscle relaxants and spasticity management agents, may be utilized to alleviate symptoms and enhance mobility(10).

Rehabilitation plays a significant role in the recovery process for individuals with BSS. Physical therapy aims to restore strength, flexibility, and coordination in the affected limbs. Occupational therapy helps individuals regain independence in daily activities, while assistive devices and adaptive equipment can facilitate mobility and enhance the quality of life.

Pain management strategies may include medications, nerve blocks, or other interventional procedures tailored to the individual's needs. Psychological support is also essential to address the emotional and psychological impact of the syndrome.

Prognosis

Brown-Séquard Syndrome's prognosis is greatly influenced by the nature, severity, and extent of spinal cord damage. The result may also depend on the patient's age, general health, and the existence of comorbid conditions. While some people might recover significantly from their functional impairments, others would continue to have neurological deficiencies. Current studies into cutting-edge therapies, like stem cell therapy and neurodegenerative methods, show promise for enhancing long-term prognosis.

Conclusion

A distinctive pattern of spinal cord injury characterises the rare and complicated neurological illness known as Brown-Séquard Syndrome. For maximising patient outcomes, prompt diagnosis and a multidisciplinary therapy strategy are essential. Improvements in diagnostic procedures and therapy approaches offer hope for a better prognosis and functional recovery. Additional investigation is required to improve our comprehension of the underlying systems and create cutting-edge therapeutic approaches.

Funding and Sponsorship: None of the authors have a financial interest in any of the products, devices, or drugs mentioned in this manuscript.

Ethical Statement: Being a Short note, there were no ethical issues and IRB permission is not required.

Conflicts of interest: The authors declare no conflict of interest.

References

1. Rengachary SS, Colen C, Guthikonda M. CHARLES-ÉDOUARD BROWN-SÉQUARD. *Neurosurgery*. 2008 Apr;62(4):954–64.
2. Zhang XY, Yang YM. Scissors stab wound to the cervical spinal cord at the craniocervical junction. *The Spine Journal*. 2016 Jun;16(6):e403–6.
3. Kashyap S, Majeed G, Lawandy S. A rare case of Brown-Sequard syndrome caused by traumatic cervical epidural hematoma. *Surg Neurol Int*. 2018;9(1):213.
4. Shams S, Arain A. Brown-Sequard Syndrome. 2023.

5. Van Wittenberghe IC, Peterson DC. Corticospinal Tract Lesion. 2023.
6. Rodríguez-Quintero JH, Romero-Velez G, Pereira X, Kim PK. Traumatic Brown-Séquard syndrome: modern reminder of a neurological injury. *BMJ Case Rep.* 2020 Nov 30;13(11):e236131.
7. Dlouhy BJ, Dahdaleh NS, Howard MA. Radiographic and intraoperative imaging of a hemisection of the spinal cord resulting in a pure Brown-Séquard syndrome: case report and review of the literature. *J Neurosurg Sci.* 2013 Mar;57(1):81–6.
8. Titlic M, Milas I, Tonkic A, Jukic I, Kolic K. Diagnostics of incomplete Brown-Sequard syndrome caused by meningococcal myelitis. *Bratisl Lek Listy.* 2006;107(8):287–9.
9. Hachem LD, Ahuja CS, Fehlings MG. Assessment and management of acute spinal cord injury: From point of injury to rehabilitation. *J Spinal Cord Med.* 2017 Nov 2;40(6):665–75.
10. Wu YT, Ho CW, Chang ST, Chen LC. Brown-Séquard Syndrome Caused by Type III Odontoid Fracture. *Spine (Phila Pa 1976).* 2010 Jan;35(1):E27–30.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.