

Short Note

Not peer-reviewed version

Living on the Edge: Navigating the Complex Landscape of Borderline Personality Disorder

[Sharif Mohammed Sadat](#) , [Dev Desai](#) , Maria Eleni Malafi *

Posted Date: 26 December 2023

doi: 10.20944/preprints202312.1904.v1

Keywords: Borderline personality Disorder; instability; Impulsive; Stigma; Psychiatry



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Short Note

Living on the Edge: Navigating the Complex Landscape of Borderline Personality Disorder

Sharif Mohammed Sadat ¹, Dev Desai ² and Maria Eleni Malafi ^{3,*}

¹ Bangladesh Medical College, Dhaka, Bangladesh; sadat.bm33@gmail.com

² Smt. NHLMMC, Ahmedabad, India; devhdesai01@gmail.com

³ Medical School, Democritus University of Thrace, Alexandroupoli, Greece

* Correspondence: marilenamalafi@gmail.com

Abstract: Individuals with BPD grapple with unstable relationships, marked by extreme emotional swings and interpersonal conflicts. Emotional instability, a distorted self-image, and impulsivity further complicate matters, often leading to self-harm and suicide attempts as desperate coping mechanisms. Genetics and adverse childhood experiences contribute to BPD's development, shaping brain function and emotional regulation. Diagnosis involves meticulous assessments, paving the way for personalized treatments like dialectical behavior therapy and, in severe cases, hospitalization. Early intervention, professional support, and societal understanding are pivotal for positive outcomes, and challenging mental health stigma.

Keywords: borderline personality disorder; instability; impulsive; stigma; psychiatry

"It is as if my life were magically run by two electric currents: joyous positive and despairing negative - whichever is running at the moment dominates my life, floods it."

Sylvia Plath beautifully depicted the emotional whirlwind of Borderline Personality Disorder (BPD) in her journal through the metaphor of electric currents, which stands as a multifaceted conundrum that significantly impacts the daily functioning and overall well-being of individuals afflicted by it.

Symptoms

People with BPD may encounter various difficulties across multiple aspects of their lives. Relational difficulties arise due to extreme emotional highs and lows, leading to unstable connections with loved ones. This can result in rapidly shifting feelings towards others, oscillating between romanticization and devaluation. Interpersonal interactions become tumultuous, causing ongoing conflicts[1].

Emotional fluctuations are another hallmark of BPD, characterized by swift and intense changes in mood, including deep sadness or fury. This emotional rollercoaster significantly impacts daily functioning and can lead to difficulty making decisions and navigating personal identity issues. Moreover, people with BPD may struggle with a distorted sense of self, which influences thought patterns and fuels impulsive behavior[2].

Impulsiveness is a pervasive feature of BPD, displaying itself in behaviors like excessive spending, drug misuse, risky driving, and binge eating[3]. These actions act as maladaptive coping mechanisms to deal with the chaotic inner world caused by the disorder. Self-harm and suicide attempts are frequent among individuals with BPD, highlighting the importance of prompt therapeutic approaches[4,5].

Causes

Genetics play a significant role here; certain gene variations can increase the likelihood of developing BPD. However, it's essential to note that no single "borderline personality disorder gene"

has been identified. Rather, multiple genetic markers interact with each other and the environment to influence the emergence of symptoms[6,7].

The likelihood of having BPD might be considerably increased by adverse childhood experiences such as abuse, neglect, or trauma[8]. The structure and function of the brain are shaped throughout these formative years, which have an impact on later relationships, impulse control, and emotional regulation. Determining the potential effects of these events on your emotional, social, and cognitive development is crucial. This may be connected to issues with decision-making, memory formation, and emotion processing.

Diagnosis & Treatment:

Identifying this condition can be complex due to overlapping symptoms with other mental health issues. Healthcare providers use structured interviews and assessments to determine whether an individual exhibits signs of BPD and gauge its severity. Accurate diagnosis sets the stage for tailored treatment plans that address each person's distinctive challenges [9,10].

One particular psychotherapy that is commonly used to treating BPD is dialectical behaviour therapy. It gives people useful techniques to strengthen distress tolerance, increase emotional control, and improve interpersonal effectiveness. Hospitalisation might be necessary in cases when there are serious worries about self-harm or suicidal thoughts in order to ensure the individual's safety[11].

Early intervention, combined with professional mental health support and a strong network of social backups, plays a critical role in achieving favorable results. Raising public awareness and promoting an increased understanding of BPD can help combat stigma surrounding mental health issues, leading to more empathetic attitudes towards those coping with this condition.[12]

Conclusion

Navigating the complex landscape of Borderline Personality Disorder requires a nuanced understanding and a multifaceted approach to treatment. Recognizing the distinct difficulties encountered by individuals with BPD involves cultivating empathy and eliminating stigmatization. This creates an atmosphere conducive to early intervention and tailored therapy programs, such as psychotherapy and medication, which are crucial for favorable outcomes. As our society becomes more informed about BPD, we play a role in promoting a more compassionate grasp of mental wellness by highlighting the significance of empathy and backing for those struggling with this complex condition.

Funding: None of the authors have a financial interest in any of the products, devices, or drugs mentioned in this manuscript.

Institutional Review Board Statement: Being a Short note, there were no ethical issues and IRB permission is not required.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. P. Bozzatello, C. Garbarini, P. Rocca, and S. Bellino, "Borderline personality disorder: Risk factors and early detection," *Diagnostics*, vol. 11, no. 11, 2021, doi: 10.3390/diagnostics11112142.
2. R. S. Biskin and J. Paris, "Diagnosing borderline personality disorder," *CMAJ. Canadian Medical Association Journal*, vol. 184, no. 16, 2012. doi: 10.1503/cmaj.090618.
3. F. Kulacaoglu and S. Kose, "Borderline personality disorder (BPD): In the midst of vulnerability, chaos, and awe," *Brain Sciences*, vol. 8, no. 11, 2018. doi: 10.3390/brainsci8110201.
4. M. Oumaya, S. Friedman, A. Pham, T. Abou Abdallah, J.-D. Guelfi, and F. Rouillon, "[Borderline personality disorder, self-mutilation and suicide: literature review].," *Encephale*, vol. 34, no. 5, 2008.
5. P. Prada, N. Perroud, E. Rüfenacht, and R. Nicasro, "Strategies to deal with suicide and non-suicidal self-injury in borderline personality disorder, the case of DBT," *Frontiers in Psychology*, vol. 9, no. DEC. 2018. doi: 10.3389/fpsyg.2018.02595.

6. R. W. Carpenter, R. L. Tomko, T. J. Trull, and D. I. Boomsma, "Gene-environment studies and borderline personality disorder: A review," *Current Psychiatry Reports*, vol. 15, no. 1. 2013. doi: 10.1007/s11920-012-0336-1.
7. A. Bassir Nia, M. C. Eveleth, J. M. Gabbay, Y. J. Hassan, B. Zhang, and M. M. Perez-Rodriguez, "Past, present, and future of genetic research in borderline personality disorder," *Current Opinion in Psychology*, vol. 21. 2018. doi: 10.1016/j.copsyc.2017.09.002.
8. N. Cattane, R. Rossi, M. Lanfredi, and A. Cattaneo, "Borderline personality disorder and childhood trauma: Exploring the affected biological systems and mechanisms," *BMC Psychiatry*, vol. 17, no. 1, 2017, doi: 10.1186/s12888-017-1383-2.
9. O. J. Storebø *et al.*, "Psychological therapies for people with borderline personality disorder," *Cochrane Database of Systematic Reviews*, vol. 2018, no. 2, 2018, doi: 10.1002/14651858.CD012955.
10. J. M. May, T. M. Richardi, and K. S. Barth, "Dialectical behavior therapy as treatment for borderline personality disorder," *Mental Health Clinician*, vol. 6, no. 2, 2016, doi: 10.9740/mhc.2016.03.62.
11. A. A. Alabi, "Management of self-harm, suicidal ideation and suicide attempts," *South African Family Practice*, vol. 64, no. 1, 2022, doi: 10.4102/safp.v64i1.5496.
12. P. Klein, A. K. Fairweather, and S. Lawn, "Structural stigma and its impact on healthcare for borderline personality disorder: a scoping review," *Int J Ment Health Syst*, vol. 16, no. 1, 2022, doi: 10.1186/s13033-022-00558-3.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.