

Article

Not peer-reviewed version

---

# Particle Physics and Cosmology Intertwined

---

[Pran Nath](#) \*

Posted Date: 8 January 2024

doi: [10.20944/preprints202401.0557.v1](https://doi.org/10.20944/preprints202401.0557.v1)

Keywords: particle physics; cosmology



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

## Article

# Particle Physics and Cosmology Intertwined

Pran Nath

Department of Physics, Northeastern University, Boston, MA 02115-5000, USA; mailto:p.nath@northeastern.edu; p.nath@northeastern.edu

**Abstract:** While the standard model accurately describes data at the electroweak scale without inclusion of gravity, beyond the standard model physics is increasingly intertwined with gravitational phenomena and cosmology. Thus gravity mediated breaking of supersymmetry in supergravity models lead to sparticles masses, which are gravitational in origin, observable at TeV scales and testable at the LHC, and supergravity also provides candidate for dark matter, a possible framework for inflationary models and for models of dark energy. Further, extended supergravity models, and string and D-brane models contain hidden sectors some of which may be feebly coupled to the visible sector resulting in heat exchange between the visible and hidden sectors. Because of the couplings between the sectors both particle physics and cosmology are effected. The above implies that particle physics and cosmology are intrinsically intertwined in the resolution of essentially all of the cosmological phenomena such as dark matter and dark energy and in the resolution of cosmological puzzles such as Hubble tension and EDGES anomaly. Here we give a brief overview of the intertwining and implications for the discovery of sparticles, and the resolution of the cosmological anomalies and identification of dark matter and dark energy as major challenges for the coming decades.

**Keywords:** particle physics; cosmology

---

## 1. Introduction

This article is a contribution to Paul Frampton's 80th birthday volume marking his over five decades of contributions as a prolific researcher to theoretical physics. He is one of the few theoretical physicists who early on recognized no boundary between particle physics and cosmology and contributed freely to each in good measure. Prominent among his works in particle theory relate to physics beyond the standard model and anomaly cancellations in higher dimensions, and in cosmology on non-standard cosmological models and black hole physics. Since particle physics and cosmology are the two major areas of his work, this paper elaborates on the progressive intertwining of the fields of particle physics and cosmology, over the past several decades from the author's own perspective.

For a long period of time up to and including the period of the emergence of the standard model[1–7] and its tests, it was largely accepted that gravity could be ignored in phenomena related to particle physics. The contrary of course, was not true, as particle physics was already known to be central to a variety of astrophysical phenomena such as the Chandrasekhar limit [8], and the synthesis of elements in the work of B2FH[9] and Peebles [10]. For particle physics gravity became more relevant with the emergence of supersymmetry, supergravity and strings. Further, supergravity models in gravity mediated breaking of supersymmetry lead to soft terms which allow radiative breaking of the electroweak symmetry and predict sparticles observable at colliders. There is another aspect of supergravity and strings which has direct impact on particle physics. In extended supergravity, strings and D-brane models one finds hidden sectors which can couple feebly with the visible sector and affect particle physics phenomena observable at colliders and also have implications for cosmology as they can provide candidates for inflation, dark matter and dark energy. Thus, with the emergence of supergravity and strings deeper connection between particle physics and cosmology has emerged. Of course, one hopes that particles physics and cosmology are parts of strings and significant literature exists on particle physics-string connection (see, e.g., [11–15] and references therein) and on cosmology-string connection (see, e.g., [16–18] and references therein).

In this paper we will focus on the intertwining of particle physics and cosmology. As noted above this intertwining has occurred on two fronts: first, in supergravity models with gravity mediated breaking the sparticle spectra are direct evidence that gravitational interactions are at work even at the scale of electro-weak physics. Further, supergravity models with R-parity conservation lead to a candidate for dark matter, specifically a neutralino [19] which turns out to be the lightest supersymmetric particle most often in radiative breaking of the electroweak symmetry [20], which enters in simulations of cosmological evolution. At the same time supergravity provides models for the inflationary expansion of the universe. Second, also as noted above in extended supergravity and in string models, one finds hidden sectors some of which may be feebly coupled to the visible sector. Typically, the hidden sectors and the visible sector will have different temperatures but they have heat exchange which requires a synchronous evolution of the two sectors intertwining the two and affecting both particle physics and cosmology. The outline of rest of the paper is as follows. In section 2, we will discuss the implications of gravity mediated breaking of supergravity at low energy and in section 3 on the intertwining of the particle physics and cosmology via hidden sectors.

## 2. Gravitational imprint on particle physics at the electroweak scale

As noted above till the advent of sugra it was the prevalent view that gravity did not have much role in particle physics models. However, with the advent of supergravity grand unification[21,22] where supersymmetry is broken in the hidden sector and communicated to the visible sector by gravitational interactions, one finds that soft breaking terms are dependent on gravitational interactions [21,23,24]. Thus, the soft mass of scalars in the visible sector  $m_s \propto \kappa m^2$ , where  $\kappa = \sqrt{8\pi G_N}$  and  $G_N$  is Newton's constant, and  $m$  is an intermediate hidden sector mass. Here with  $m \sim 10^{10}$  and  $M_{Pl} = \kappa^{-1} = 2.43 \times 10^{18}$  GeV (in natural units:  $\hbar = c = 1$ ), one finds  $m_s$  to be of electroweak size. Since sparticle masses are controlled by the soft susy scale, the discovery of sparticles would be a signature indicating that gravity has a role in low energy physics. This would be very much in the spirit that the discovery of the  $W$  and the  $Z$  bosons are reflection of  $SU(2)_L \times U(1)_Y$  unification. It is notable that the soft terms are also responsible for generating spontaneous breaking of the electroweak symmetry[21,25]. An indication that some of the sparticles may be low lying comes from the  $g - 2$  data from Fermilab[26] which points to a deviation from the standard model prediction of about  $4\sigma$ . An attractive proposition is that the deviation arises from light sparticle exchange, specifically light charginos and light sleptons (see, e.g., [27–29] and the references therein), a deviation that was predicted quite a while ago[30]. However, a word of caution is in order in that the lattice analysis [31] for hadronic vacuum polarization contribution gives a smaller deviation from the standard model than the conventional result where the hadronic vacuum polarization contribution is computed using  $e^+e^- \rightarrow \pi^+\pi^-$  data. Thus further work is needed in reconciling the lattice analysis with the conventional result on the hadronic polarization contribution before drawing any definitive conclusions.

## 3. Hidden sectors intertwine particle physics and cosmology

As already noted in a variety of models beyond standard model physics, which include extended supergravity models, string models and extra dimension models, one has hidden sectors. While these sectors are neutral under the standard model gauge group they may interact with the visible sectors via feeble interactions. Such feeble interactions can occur via a variety of portals which include the Higgs portal[32], kinetic energy portal [33,34], Stueckelberg mass mixing portal[35,36], kinetic-mass-mixing portal[37], Stueckelberg-Higgs portal[38], as well as possible higher dimensional operators. The hidden sectors could be endowed with gauge fields, as well as with matter. At the reheat temperature the hidden sectors and the visible sector would in general lie in different heat baths. However, because of the feeble interactions between the sectors, there will be heat exchange between the visible and the hidden sectors and thus their thermal evolution will be correlated. The evolution of the relative temperatures of the two sectors then depends on the initial conditions, and specifically on the ratio  $\xi(T) = T_h/T$  at the reheat temperature, where  $T_h$  is the hidden sector temperature and  $T$  is the visible

sector temperature. The Boltzmann equations governing the evolution of the visible and the hidden sectors are coupled and involve the evolution equation for  $\xi(T)$ . Such an equation was derived in [39–41] and applied in a variety of settings in [42] consistent with all experimental constraints on hidden sector matter from terrestrial and astro-physical data [43]. It is found that hidden sectors can affect observable phenomena in the visible sector, such as the density of thermal relics. Hidden sectors provide candidates for dark matter and dark energy and help resolve cosmological anomalies intertwining particle physics and cosmological phenomena. We discuss some of these topics in further detail below.

Green-Schwarz [44] found that in the low energy limit of Type I strings the kinetic energy of 2-tensor  $B_{MN}$  of 10D supergravity multiplet has Yang-Mills and Lorentz group Chern-Simons terms (indicated by superscripts Y and L) so that  $\partial_{[P}B_{MN]} \rightarrow \partial_{[P}B_{MN]} + \omega_{PMN}^{(Y)} - \omega_{PMN}^{(L)}$ , where  $M, N, P$  are 10-dimensional indices. Inclusion of the Chern-Simons terms fully requires that one extend the 10D Sugra Lagrangian to order  $O(\kappa)^2$ . This was accomplished subsequent to Green-Schwarz work in [45] (for related works see [46–48]). Dimensional reduction to 4D with a vacuum expectation value for the internal gauge field strength,  $\langle F_{ij} \rangle \neq 0$  (where the indices are for the six-dimensional compact manifold), leads to  $\partial_\mu B_{ij} + A_\mu F_{ij} + \dots \sim \partial_\mu \sigma + m A_\mu$  ( $\mu$  in an index for four dimensional Minkowskian space-time) where the internal components  $B_{ij}$  give the pseudo-scalar  $\sigma$  and  $m$  arises from  $\langle F_{ij} \rangle$ , which is a topological quantity, related to the Chern numbers of the gauge bundle. Thus  $A_\mu$  and  $\sigma$  have a Stueckelberg coupling of the form  $A_\mu \partial^\mu \sigma$ . This provides the inspiration for building BSM models with the Stueckelberg mechanism [35,36,49–51]. Specifically, this allows the possibility of writing effective theories with gauge invariant mass terms. For the case of a single  $U(1)$  gauge field  $A_\mu$  one may write a gauge invariant mass term by letting  $A_\mu \rightarrow A_\mu + \frac{1}{m} \partial_\mu \sigma$  where the gauge transformations are defined so that  $\delta A_\mu = \partial_\mu \lambda$  and  $\delta \sigma = -m \lambda$ . In this case  $\sigma$ 's role is akin to that of the longitudinal component of a massive vector. The above technique also allows one to generate invariant mass mixing between two  $U(1)$  gauge fields. Thus consider two gauge groups  $U(1)_X$  and  $U(1)_Y$  with gauge fields  $A_\mu, B_\mu$  and an axionic field  $\sigma$ . In this case we can write a mass term  $(m_1 A_\mu + m_2 B_\mu + \partial_\mu \sigma)^2$  which is invariant under  $\delta_x A_\mu = \partial_\mu \lambda_x, \delta_x \sigma = -m_1 \lambda_x$  for  $U(1)_X$ , and  $\delta_y B_\mu = \partial_\mu \lambda_y, \delta_y \sigma = -m_2 \lambda_y$  for  $U(1)_Y$ . One of the interesting phenomena associated with effective gauge theories with gauge invariant mass terms is that they generate millicharges when coupled to matter fields [35,37,49,52]. We will return to this feature of the Stueckelberg mass mixing terms when we discuss the EDGES anomaly.

**Hubble tension:** Currently there exists a discrepancy between the measured value of the Hubble parameter  $H_0$  for low redshifts ( $z < 1$ ) and high redshifts ( $z > 1000$ ). Thus for ( $z < 1$ ) an analysis of data from Cepheids and SNIa gives [53]  $H_0 = (73.04 \pm 1.04)$  km/s/Mpc. On the other an analysis based on  $\Lambda$ CDM model the SH0ES Collaboration [53] using data from the Cosmic Microwave Background (CMB), Baryon Acoustic Oscillations (BAO), and from Big Bang Nucleosynthesis (BBN), determines the Hubble parameter at high  $z$  to be [54]  $H_0 = (67.4 \pm 0.5)$  km/s/Mpc. This indicates a  $5\sigma$  level tension between the low  $z$  and the high  $z$  measurements. There is a significant amount of literature attempting to resolve this puzzle at least partially and recent reviews include [55,56]. One simple approach is introducing extra relativistic degrees of freedom during the period of recombination which increases the magnitude of  $H_0$  which helps alleviate the tension. Models using this idea introduce extra particles such as the  $Z'$  of an extra  $U(1)$  gauge field which decays to neutrinos [57,58] or utilize other particles such as the majoron [59,60]. The inclusion of extra degrees of freedom, however, must be consistent with the BBN constraints which are sensitive to the addition of massless degrees of freedom. Thus the standard model prediction of  $N_{\text{eff}}^{\text{SM}} \simeq 3.046$  [61] is consistent with the synthesis of light elements and the introduction of new degrees of freedom must maintain this successful standard model prediction. The above indicates that the extra degrees of freedom should emerge only beyond the BBN time and in the time frame of the recombination epoch. It is to be noted that new degrees of freedom are also constrained by the CMB data as given by the Planck analysis [54].

A cosmologically consistent model based on the Stueckelberg extension of the SM with a hidden sector was proposed in [62] for alleviating the Hubble tension. The model is cosmologically consistent since the analysis is based on a consistent thermal evolution of the visible and the hidden sectors taking account of the thermal exchange between the two sectors. In addition to the dark fermions, and dark photon, the model also contains a massless pseudo scalar particle field  $\phi$  and a massive long-lived scalar field  $s$ . The fields  $\phi$  and  $s$  have interactions only in the dark sector with no interactions with the standard model fields. The decay of the scalar field occurs after BBN close to the recombination time via the decay  $s \rightarrow \phi\phi$  which provides the extra degrees of freedom needed to alleviate the Hubble tension. It should be noted that the full resolution of the Hubble tension would require going beyond providing new degrees of freedom and would involve a fit to all of the CMB data consistent with all cosmological and particle physics constraints. For some recent related work on Hubble tension see [63–68].

**EDGES anomaly:** The 21-cm line plays an important role in the analysis of physics during the dark ages and the cosmic dawn in the evolution of the early universe. The 21-cm line arises from the spin transition from the triplet state to the singlet state and vice-versa in the ground state of neutral hydrogen. The relative abundance of the triplet and the singlet states defines the spin temperature  $T_s$  (and  $T_B = T_s$ ) of the hydrogen gas and is given by  $n_1/n_0 = 3e^{-T_*/T_s}$ , where 3 is the ratio of the spin degrees of freedom for the triplet versus the singlet state,  $T_*$  is defined by  $\Delta E = kT_*$ , where  $\Delta E = 1420$  MHz is the energy difference at rest between the two spin states, and  $T_* = \frac{hc}{k\lambda_{21\text{cm}}}$  = 0.068K. EDGES (The Experiment to Detect the Global Epoch of Reionization Signature) reported an absorption profile centered at the frequency  $\nu = 78$  MHz in the sky-averaged spectrum. The quantity of interest is the brightness temperature  $T_{21}$  of the 21-cm line defined by  $T_{21}(z) = ((T_s - T_\gamma)/1 + z)(1 - e^{-\tau})$  where  $\tau$  is the optical depth for the transition. The analysis of Bowman et.al[69] finds<sup>1</sup> that at redshift  $z \sim 17$ ,  $T_{21} = -500^{+200}_{-500}$  mK at 99% C.L. On the other hand the analysis of [71] based on the  $\Lambda$ CDM model gives a  $T_{21}$  around  $-230$  mK, which shows that the EDGES result is a  $3.8\sigma$  deviation away from that of the standard cosmological paradigm.

The EDGES anomaly is not yet confirmed but pending its possible confirmation it is of interest to investigate what possible explanations there might be. In fact, several mechanisms have already been proposed to explain the  $3.8\sigma$  anomaly [72–88]. A list of some of the prominent possibilities consist of the following: (1) astrophysical phenomena such as radiation from stars and star remnants; (2) the CMB background radiation temperature is hotter than expected; (3) the baryons are cooler than what  $\Lambda$ CDM predicts; (4) modification of cosmological evolution: inclusion of dark energy such as Chaplin gas. Of the above, there appears to be a leaning towards baryon cooling and there is a substantial amount of work in this area following the earlier works of [89] and Barkana [78]. Specifically it was pointed out in [78] that the observed anomaly could be explained if the baryons were cooled down by roughly 3 K. Here one assumes a small percentage of DM ( $\sim 0.3\%$ ) is millicharged and baryons become cooler by Rutherford scattering from the colder dark matter. As mentioned earlier precisely such a possibility occurs via the Stueckelberg mass mixing if we assume one of the gauge fields  $U(1)_Y$  is the hyper charge gauge field while  $U(1)_X$  is a hidden sector field and the millicharge dark matter resides in the hidden sector while the rest of dark matter could be WIMPS. Within this framework a cosmologically consistent analysis of string inspired milli-charged model was proposed in[90] where a detailed fit to the data is possible consistent within a high scale model. For some recent work on EDGES anomaly see, [91–94].

**Inflation:** As is well known the problems associated with Big Bang such as flatness, horizon, and the monopole problem are resolved in inflationary models. In models of this type quantum fluctuations at horizon exit encode information regarding the characteristics of the inflationary model which can be extracted from the cosmic microwave background (CMB) radiation anisotropy [95–99].

<sup>1</sup> See, however, reference [70] on concerns regarding modeling of data.

In fact data from the Planck experiment [100–102] has already put stringent bounds on inflationary model eliminating some. A model which is especially attractive is the one based on an axionic field with a potential of the form [103,104]  $V(a) = \Lambda^4 \left(1 + \cos\left(\frac{a}{f}\right)\right)$  where  $a$  is the axion field and  $f$  is the axion decay constant. However, for the simple model above to hold the Planck data requires  $f > 10M_{Pl}$  which is undesirable since string theory indicates that  $f$  lie below  $M_{Pl}$  [105,106]. However, reduction of  $f$  turns out to be a non-trivial issue. Techniques used to resolve this issue include the alignment mechanism [107,108], n-flation, coherent enhancement [109] and models using shift symmetry, (for a review and more references see [110,111]).

We mention another inflation model which is based in an axion landscape with a  $U(1)$  symmetry[112]. This model involves  $m$  pairs of chiral fields and fields in each pair are oppositely charged under the same  $U(1)$  symmetry. Our nomenclature is such that we label the pseudo-scalar component of each field to be an axion and the corresponding real part to be a saxion. Since the model has only  $U(1)$  global symmetry, the breaking of the global symmetry leads to just one pseudo-Nambu-Goldstone-boson (PNGB) and the remaining pseudo-scalars are not PNGBs. Thus the superpotential of the model consists of a part which is invariant under the  $U(1)$  global symmetry and a  $U(1)$  symmetry breaking part which simulates instanton effects. The analysis of this work shows that the potential contains a fast roll-slow roll splitting mechanism which splits the axion potential into fast roll and slow roll parts where the fields entering fast roll are eliminated early on leaving the slow roll part which involves a single axion field which drives inflation. Here under the constraints of stabilized saxions, one finds inflation models with  $f < M_{Pl}$  consistent with Planck data. Similar results are found in the Dirac-Born-Infeld based models[113].

**Dark energy:** One of the most outstanding puzzles of both particle physics and of cosmology is dark energy which constitutes about 70% of the energy budget of the universe and is responsible for the accelerated expansion of the universe. Dark energy is characterized by negative pressure so that  $w$  defined by  $w = p/\rho$ , where  $p$  is the pressure and  $\rho$  the energy density for dark energy, must satisfy  $w < -1/3$ . The CMB and the baryon acoustic oscillation (BAO) data fits well with a cosmological constant  $\Lambda$  which corresponds to  $w = -1$ . Thus the Planck Collaboration [54] gives  $w = -1.03 \pm 0.03$  consistent with the cosmological constant. There are two puzzles connected with dark energy. First, the use of the cosmological constant appears artificial, and it is desirable to replace it by a dynamical field, i.e., a so-called quintessence field (for a review see [114]), which at late times can generate accelerated expansion similar to that given by  $\Lambda$ . The second problem relates to the very small size of the cosmological constant which is not automatically resolved by simply replacing  $\Lambda$  by a dynamical field. The extreme fine tuning needed in a particle physics model to get to the size of  $\Lambda$  requires a new idea such as vacuum selection in a landscape with a large number of possible allowed vacua[115], for instance those available in string theory. In any case, it is an example of the extreme intertwined nature of cosmology and particle physics. However, finding a quintessence solution that replaces  $\Lambda$  and is consistent all of the CMB data is itself progress. Regarding experimental measurement of  $w = -1.03 \pm 0.03$  if more accurate data in future gives  $w > -1$ , it would point to something like quintessence while  $w < -1$  would indicate phantom energy and an entirely new sector.

#### 4. Conclusion

In conclusion it is clear that particle physics and cosmology are deeply intertwined and models of physics beyond the standard model would in the future be increasingly constrained by particle physics experiments as well as by astrophysical data. We congratulate Paul for his notable contributions in the twin fields and wish him many productive years of contributions for the future.

**Acknowledgments:** PN is supported in part by the NSF Grant PHY-2209903.

## References

1. S. L. Glashow, Nucl. Phys. **22**, 579-588 (1961) doi:10.1016/0029-5582(61)90469-2
2. S. Weinberg, Phys. Rev. Lett. **19**, 1264-1266 (1967) doi:10.1103/PhysRevLett.19.1264
3. A. Salam, in N. Svartholm, ed., Elementary Particle Theory, Almqvist and Wiksell, (1969), p. 367.
4. G. 't Hooft, Nucl. Phys. B **35**, 167-188 (1971) doi:10.1016/0550-3213(71)90139-8
5. G. 't Hooft and M. J. G. Veltman, Nucl. Phys. B **44**, 189-213 (1972) doi:10.1016/0550-3213(72)90279-9
6. H. D. Politzer, Phys. Rev. Lett. **30**, 1346-1349 (1973) doi:10.1103/PhysRevLett.30.1346
7. D. J. Gross and F. Wilczek, Phys. Rev. Lett. **30**, 1343-1346 (1973) doi:10.1103/PhysRevLett.30.1343
8. S. Chandrasekhar, Astrophys. J. **74**, 81-82 (1931) doi:10.1086/143324
9. M. E. Burbidge, G. R. Burbidge, W. A. Fowler and F. Hoyle, Rev. Mod. Phys. **29**, 547-650 (1957) doi:10.1103/RevModPhys.29.547
10. P. J. E. Peebles, Astrophys. J. **146**, 542-552 (1966) doi:10.1086/148918
11. K. R. Dienes, Phys. Rept. **287**, 447-525 (1997) doi:10.1016/S0370-1573(97)00009-4 [arXiv:hep-th/9602045 [hep-th]].
12. P. Nath and P. Fileviez Perez, Phys. Rept. **441**, 191-317 (2007) doi:10.1016/j.physrep.2007.02.010 [arXiv:hep-ph/0601023 [hep-ph]].
13. T. Ibrahim and P. Nath, Rev. Mod. Phys. **80**, 577-631 (2008) doi:10.1103/RevModPhys.80.577 [arXiv:0705.2008 [hep-ph]].
14. S. Abel, K. R. Dienes and L. A. Nutricati, Phys. Rev. D **107**, no.12, 126019 (2023) doi:10.1103/PhysRevD.107.126019 [arXiv:2303.08534 [hep-th]].
15. S. Abel and K. R. Dienes, Phys. Rev. D **104**, no.12, 126032 (2021) doi:10.1103/PhysRevD.104.126032 [arXiv:2106.04622 [hep-th]].
16. S. Kachru, R. Kallosh, A. D. Linde and S. P. Trivedi, Phys. Rev. D **68**, 046005 (2003) doi:10.1103/PhysRevD.68.046005 [arXiv:hep-th/0301240 [hep-th]].
17. V. Balasubramanian, P. Berglund, J. P. Conlon and F. Quevedo, JHEP **03**, 007 (2005) doi:10.1088/1126-6708/2005/03/007 [arXiv:hep-th/0502058 [hep-th]].
18. J. Halverson, C. Long and P. Nath, Phys. Rev. D **96**, no.5, 056025 (2017) doi:10.1103/PhysRevD.96.056025 [arXiv:1703.07779 [hep-ph]].
19. H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983) [erratum: Phys. Rev. Lett. **103**, 099905 (2009)] doi:10.1103/PhysRevLett.50.1419
20. R. L. Arnowitt and P. Nath, Phys. Rev. Lett. **69**, 725-728 (1992) doi:10.1103/PhysRevLett.69.725
21. A. H. Chamseddine, R. L. Arnowitt and P. Nath, Phys. Rev. Lett. **49**, 970 (1982) doi:10.1103/PhysRevLett.49.970
22. P. Nath, R. L. Arnowitt and A. H. Chamseddine, Nucl. Phys. B **227**, 121-133 (1983) doi:10.1016/0550-3213(83)90145-1
23. R. Barbieri, S. Ferrara and C. A. Savoy, Phys. Lett. B **119**, 343 (1982) doi:10.1016/0370-2693(82)90685-2
24. L. J. Hall, J. D. Lykken and S. Weinberg, Phys. Rev. D **27**, 2359-2378 (1983) doi:10.1103/PhysRevD.27.2359
25. L. E. Ibanez and G. G. Ross, Phys. Lett. B **110**, 215-220 (1982) doi:10.1016/0370-2693(82)91239-4
26. D. P. Aguillard *et al.* [Muon g-2], Phys. Rev. Lett. **131**, no.16, 161802 (2023) doi:10.1103/PhysRevLett.131.161802 [arXiv:2308.06230 [hep-ex]].
27. A. Aboubrahim, M. Klasen and P. Nath, Phys. Rev. D **104**, no.3, 035039 (2021) doi:10.1103/PhysRevD.104.035039 [arXiv:2104.03839 [hep-ph]].
28. H. Baer, V. Barger and H. Serce, Phys. Lett. B **820**, 136480 (2021) doi:10.1016/j.physletb.2021.136480 [arXiv:2104.07597 [hep-ph]].
29. M. Chakraborti, S. Heinemeyer and I. Saha, Eur. Phys. J. C **81**, no.12, 1114 (2021) doi:10.1140/epjc/s10052-021-09900-4 [arXiv:2104.03287 [hep-ph]].
30. T. C. Yuan, R. L. Arnowitt, A. H. Chamseddine and P. Nath, Z. Phys. C **26**, 407 (1984) doi:10.1007/BF01452567
31. S. Borsanyi, Z. Fodor, J. N. Guenther, C. Hoelbling, S. D. Katz, L. Lellouch, T. Lippert, K. Miura, L. Parato and K. K. Szabo, *et al.* Nature **593**, no.7857, 51-55 (2021) doi:10.1038/s41586-021-03418-1 [arXiv:2002.12347 [hep-lat]].
32. B. Patt and F. Wilczek, [arXiv:hep-ph/0605188 [hep-ph]].
33. B. Holdom, Phys. Lett. B **166**, 196-198 (1986) doi:10.1016/0370-2693(86)91377-8

34. B. Holdom, Phys. Lett. B **259**, 329-334 (1991) doi:10.1016/0370-2693(91)90836-F

35. B. Kors and P. Nath, Phys. Lett. B **586**, 366-372 (2004) doi:10.1016/j.physletb.2004.02.051 [arXiv:hep-ph/0402047 [hep-ph]].

36. B. Kors and P. Nath, JHEP **12**, 005 (2004) doi:10.1088/1126-6708/2004/12/005 [arXiv:hep-ph/0406167 [hep-ph]].

37. D. Feldman, Z. Liu and P. Nath, Phys. Rev. D **75**, 115001 (2007) doi:10.1103/PhysRevD.75.115001 [arXiv:hep-ph/0702123 [hep-ph]].

38. M. Du, Z. Liu and P. Nath, Phys. Lett. B **834**, 137454 (2022) doi:10.1016/j.physletb.2022.137454 [arXiv:2204.09024 [hep-ph]].

39. A. Aboubrahim, W. Z. Feng, P. Nath and Z. Y. Wang, Phys. Rev. D **103**, no.7, 075014 (2021) doi:10.1103/PhysRevD.103.075014 [arXiv:2008.00529 [hep-ph]].

40. A. Aboubrahim, W. Z. Feng, P. Nath and Z. Y. Wang, [arXiv:2106.06494 [hep-ph]].

41. A. Aboubrahim and P. Nath, JHEP **09**, 084 (2022) doi:10.1007/JHEP09(2022)084 [arXiv:2205.07316 [hep-ph]].

42. J. Li and P. Nath, Phys. Rev. D **108**, no.11, 115008 (2023) doi:10.1103/PhysRevD.108.115008 [arXiv:2304.08454 [hep-ph]].

43. A. Aboubrahim, M. M. Altakach, M. Klasen, P. Nath and Z. Y. Wang, JHEP **03**, 182 (2023) doi:10.1007/JHEP03(2023)182 [arXiv:2212.01268 [hep-ph]].

44. M. B. Green and J. H. Schwarz, Phys. Lett. B **149**, 117-122 (1984) doi:10.1016/0370-2693(84)91565-X

45. A. H. Chamseddine and P. Nath, Phys. Rev. D **34**, 3769 (1986) doi:10.1103/PhysRevD.34.3769

46. L. J. Romans and N. P. Warner, Nucl. Phys. B **273**, 320-332 (1986) doi:10.1016/0550-3213(86)90249-X

47. H. Nishino and S. J. Gates, Jr., Phys. Lett. B **173**, 417-422 (1986) doi:10.1016/0370-2693(86)90407-7

48. E. Bergshoeff, A. Salam and E. Sezgin, Nucl. Phys. B **279**, 659-683 (1987) doi:10.1016/0550-3213(87)90015-0

49. B. Kors and P. Nath, JHEP **07**, 069 (2005) doi:10.1088/1126-6708/2005/07/069 [arXiv:hep-ph/0503208 [hep-ph]].

50. B. Kors and P. Nath, doi:10.1142/9789812701756\_0056 [arXiv:hep-ph/0411406 [hep-ph]].

51. D. Feldman, B. Kors and P. Nath, Phys. Rev. D **75**, 023503 (2007) doi:10.1103/PhysRevD.75.023503 [arXiv:hep-ph/0610133 [hep-ph]].

52. K. Cheung and T. C. Yuan, JHEP **03**, 120 (2007) doi:10.1088/1126-6708/2007/03/120 [arXiv:hep-ph/0701107 [hep-ph]].

53. A. G. Riess, W. Yuan, L. M. Macri, D. Scolnic, D. Brout, S. Casertano, D. O. Jones, Y. Murakami, L. Breuval and T. G. Brink, *et al.* Astrophys. J. Lett. **934**, no.1, L7 (2022) doi:10.3847/2041-8213/ac5c5b [arXiv:2112.04510 [astro-ph.CO]].

54. N. Aghanim *et al.* [Planck], Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].

55. E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess and J. Silk, Class. Quant. Grav. **38**, no.15, 153001 (2021) doi:10.1088/1361-6382/ac086d [arXiv:2103.01183 [astro-ph.CO]].

56. E. Abdalla, G. Franco Abellán, A. Aboubrahim, A. Agnello, O. Akarsu, Y. Akrami, G. Alestas, D. Aloni, L. Amendola and L. A. Anchordoqui, *et al.* JHEAp **34**, 49-211 (2022) doi:10.1016/j.jheap.2022.04.002 [arXiv:2203.06142 [astro-ph.CO]].

57. J. Gehrlein and M. Pierre, JHEP **02**, 068 (2020) doi:10.1007/JHEP02(2020)068 [arXiv:1912.06661 [hep-ph]].

58. M. Escudero, D. Hooper, G. Krnjaic and M. Pierre, JHEP **03**, 071 (2019) doi:10.1007/JHEP03(2019)071 [arXiv:1901.02010 [hep-ph]].

59. E. Fernandez-Martinez, M. Pierre, E. Pinsard and S. Rosauro-Alcaraz, Eur. Phys. J. C **81**, no.10, 954 (2021) doi:10.1140/epjc/s10052-021-09760-y [arXiv:2106.05298 [hep-ph]].

60. M. Escudero and S. J. Witte, Eur. Phys. J. C **81**, no.6, 515 (2021) doi:10.1140/epjc/s10052-021-09276-5 [arXiv:2103.03249 [hep-ph]].

61. G. Mangano, G. Miele, S. Pastor, T. Pinto, O. Pisanti and P. D. Serpico, Nucl. Phys. B **729**, 221-234 (2005) doi:10.1016/j.nuclphysb.2005.09.041 [arXiv:hep-ph/0506164 [hep-ph]].

62. A. Aboubrahim, M. Klasen and P. Nath, JCAP **04**, no.04, 042 (2022) doi:10.1088/1475-7516/2022/04/042 [arXiv:2202.04453 [astro-ph.CO]].

63. S. Vagnozzi, Universe **9**, no.9, 393 (2023) doi:10.3390/universe9090393 [arXiv:2308.16628 [astro-ph.CO]].

64. J. P. Hu and F. Y. Wang, Universe **9**, no.2, 94 (2023) doi:10.3390/universe9020094 [arXiv:2302.05709 [astro-ph.CO]].

65. G. A. C. Valdez, C. Quintanilla, M. A. García-Aspeitia, A. Hernández-Almada and V. Motta, *Eur. Phys. J. C* **83**, no.5, 442 (2023) doi:10.1140/epjc/s10052-023-11596-7 [arXiv:2301.07044 [astro-ph.CO]].
66. Z. Zhou, Y. Mu, G. Liu, L. Xu and J. Lu, *Phys. Rev. D* **107**, no.6, 063536 (2023) doi:10.1103/PhysRevD.107.063536 [arXiv:2210.06851 [astro-ph.CO]].
67. A. C. Sobotka, A. L. Erickcek and T. L. Smith, *Phys. Rev. D* **107**, no.2, 023525 (2023) doi:10.1103/PhysRevD.107.023525 [arXiv:2207.14308 [astro-ph.CO]].
68. W. F. Duan, S. P. Li, X. Q. Li and Y. D. Yang, *Chin. Phys. C* **47**, no.3, 033102 (2023) doi:10.1088/1674-1137/aca888 [arXiv:2111.05178 [hep-ph]].
69. J. D. Bowman, A. E. E. Rogers, R. A. Monsalve, T. J. Mozdzen and N. Mahesh, *Nature* **555**, no.7694, 67-70 (2018) doi:10.1038/nature25792 [arXiv:1810.05912 [astro-ph.CO]].
70. R. Hills, G. Kulkarni, P. D. Meerburg and E. Puchwein, *Nature* **564**, no.7736, E32-E34 (2018) doi:10.1038/s41586-018-0796-5 [arXiv:1805.01421 [astro-ph.CO]].
71. A. Cohen, A. Fialkov and R. Barkana, *Mon. Not. Roy. Astron. Soc.* **478**, no.2, 2193-2217 (2018) doi:10.1093/mnras/sty1094 [arXiv:1709.02122 [astro-ph.CO]].
72. J. B. Muñoz and A. Loeb, *Nature* **557**, no.7707, 684 (2018) doi:10.1038/s41586-018-0151-x [arXiv:1802.10094 [astro-ph.CO]].
73. J. B. Muñoz, C. Dvorkin and A. Loeb, *Phys. Rev. Lett.* **121**, no.12, 121301 (2018) doi:10.1103/PhysRevLett.121.121301 [arXiv:1804.01092 [astro-ph.CO]].
74. A. Halder and M. Pandey, *Mon. Not. Roy. Astron. Soc.* **508**, no.3, 3446-3454 (2021) doi:10.1093/mnras/stab2795 [arXiv:2101.05228 [astro-ph.CO]].
75. H. Liu, N. J. Outmezguine, D. Redigolo and T. Volansky, *Phys. Rev. D* **100**, no.12, 123011 (2019) doi:10.1103/PhysRevD.100.123011 [arXiv:1908.06986 [hep-ph]].
76. A. Berlin, D. Hooper, G. Krnjaic and S. D. McDermott, *Phys. Rev. Lett.* **121**, no.1, 011102 (2018) doi:10.1103/PhysRevLett.121.011102 [arXiv:1803.02804 [hep-ph]].
77. C. Feng and G. Holder, *Astrophys. J. Lett.* **858**, no.2, L17 (2018) doi:10.3847/2041-8213/aac0fe [arXiv:1802.07432 [astro-ph.CO]].
78. R. Barkana, *Nature* **555**, no.7694, 71-74 (2018) doi:10.1038/nature25791 [arXiv:1803.06698 [astro-ph.CO]].
79. R. Barkana, N. J. Outmezguine, D. Redigolo and T. Volansky, *Phys. Rev. D* **98**, no.10, 103005 (2018) doi:10.1103/PhysRevD.98.103005 [arXiv:1803.03091 [hep-ph]].
80. S. Fraser, A. Hektor, G. Hütsi, K. Kannike, C. Marzo, L. Marzola, C. Spethmann, A. Racioppi, M. Raidal and V. Vaskonen, *et al.* *Phys. Lett. B* **785**, 159-164 (2018) doi:10.1016/j.physletb.2018.08.035 [arXiv:1803.03245 [hep-ph]].
81. M. Pospelov, J. Pradler, J. T. Ruderman and A. Urbano, *Phys. Rev. Lett.* **121**, no.3, 031103 (2018) doi:10.1103/PhysRevLett.121.031103 [arXiv:1803.07048 [hep-ph]].
82. T. Moroi, K. Nakayama and Y. Tang, *Phys. Lett. B* **783**, 301-305 (2018) doi:10.1016/j.physletb.2018.07.002 [arXiv:1804.10378 [hep-ph]].
83. A. Fialkov and R. Barkana, *Mon. Not. Roy. Astron. Soc.* **486**, no.2, 1763-1773 (2019) doi:10.1093/mnras/stz873 [arXiv:1902.02438 [astro-ph.CO]].
84. K. Choi, H. Seong and S. Yun, *Phys. Rev. D* **102**, no.7, 075024 (2020) doi:10.1103/PhysRevD.102.075024 [arXiv:1911.00532 [hep-ph]].
85. C. Creque-Sarbinowski, L. Ji, E. D. Kovetz and M. Kamionkowski, *Phys. Rev. D* **100**, no.2, 023528 (2019) doi:10.1103/PhysRevD.100.023528 [arXiv:1903.09154 [astro-ph.CO]].
86. E. D. Kovetz, V. Poulin, V. Gluscevic, K. K. Boddy, R. Barkana and M. Kamionkowski, *Phys. Rev. D* **98**, no.10, 103529 (2018) doi:10.1103/PhysRevD.98.103529 [arXiv:1807.11482 [astro-ph.CO]].
87. K. Bondarenko, J. Pradler and A. Sokolenko, *Phys. Lett. B* **805**, 135420 (2020) doi:10.1016/j.physletb.2020.135420 [arXiv:2002.08942 [astro-ph.CO]].
88. G. Lanfranchi, M. Pospelov and P. Schuster, *Ann. Rev. Nucl. Part. Sci.* **71**, 279-313 (2021) doi:10.1146/annurev-nucl-102419-055056 [arXiv:2011.02157 [hep-ph]].
89. J. B. Muñoz, E. D. Kovetz and Y. Ali-Haïmoud, *Phys. Rev. D* **92**, no.8, 083528 (2015) doi:10.1103/PhysRevD.92.083528 [arXiv:1509.00029 [astro-ph.CO]].
90. A. Aboubrahim, P. Nath and Z. Y. Wang, *JHEP* **12**, 148 (2021) doi:10.1007/JHEP12(2021)148 [arXiv:2108.05819 [hep-ph]].

91. D. Paul, A. Dey, A. Dutta Banik and S. Pal, *JCAP* **11**, 015 (2023) doi:10.1088/1475-7516/2023/11/015 [arXiv:2308.04955 [astro-ph.CO]].
92. R. Barkana, A. Fialkov, H. Liu and N. J. Outmezguine, *Phys. Rev. D* **108**, no.6, 063503 (2023) doi:10.1103/PhysRevD.108.063503 [arXiv:2212.08082 [hep-ph]].
93. A. Halder, S. S. Pandey and A. S. Majumdar, *JCAP* **10**, 049 (2022) doi:10.1088/1475-7516/2022/10/049 [arXiv:2207.09177 [astro-ph.CO]].
94. S. Wu, S. Xu and S. Zheng, *Chin. Phys. C* **47**, no.9, 095101 (2023) doi:10.1088/1674-1137/ace17f [arXiv:2205.14876 [hep-ph]].
95. A. H. Guth, *Phys. Rev. D* **23**, 347-356 (1981) doi:10.1103/PhysRevD.23.347
96. A. A. Starobinsky, *Phys. Lett. B* **91**, 99-102 (1980) doi:10.1016/0370-2693(80)90670-X
97. A. D. Linde, *Phys. Lett. B* **108**, 389-393 (1982) doi:10.1016/0370-2693(82)91219-9
98. A. Albrecht and P. J. Steinhardt, *Phys. Rev. Lett.* **48**, 1220-1223 (1982) doi:10.1103/PhysRevLett.48.1220
99. A. D. Linde, *Phys. Lett. B* **129**, 177-181 (1983) doi:10.1016/0370-2693(83)90837-7
100. R. Adam *et al.* [Planck], *Astron. Astrophys.* **594**, A1 (2016) doi:10.1051/0004-6361/201527101 [arXiv:1502.01582 [astro-ph.CO]].
101. P. A. R. Ade *et al.* [Planck], *Astron. Astrophys.* **594**, A20 (2016) doi:10.1051/0004-6361/201525898 [arXiv:1502.02114 [astro-ph.CO]].
102. P. A. R. Ade *et al.* [BICEP2 and Keck Array], *Phys. Rev. Lett.* **116**, 031302 (2016) doi:10.1103/PhysRevLett.116.031302 [arXiv:1510.09217 [astro-ph.CO]].
103. K. Freese, J. A. Frieman and A. V. Olinto, *Phys. Rev. Lett.* **65**, 3233-3236 (1990) doi:10.1103/PhysRevLett.65.3233
104. F. C. Adams, J. R. Bond, K. Freese, J. A. Frieman and A. V. Olinto, *Phys. Rev. D* **47**, 426-455 (1993) doi:10.1103/PhysRevD.47.426 [arXiv:hep-ph/9207245 [hep-ph]].
105. T. Banks, M. Dine, P. J. Fox and E. Gorbatov, *JCAP* **06**, 001 (2003) doi:10.1088/1475-7516/2003/06/001 [arXiv:hep-th/0303252 [hep-th]].
106. P. Svrcek and E. Witten, *JHEP* **06**, 051 (2006) doi:10.1088/1126-6708/2006/06/051 [arXiv:hep-th/0605206 [hep-th]].
107. J. E. Kim, H. P. Nilles and M. Peloso, *JCAP* **01**, 005 (2005) doi:10.1088/1475-7516/2005/01/005 [arXiv:hep-ph/0409138 [hep-ph]].
108. C. Long, L. McAllister and P. McGuirk, *Phys. Rev. D* **90**, 023501 (2014) doi:10.1103/PhysRevD.90.023501 [arXiv:1404.7852 [hep-th]].
109. P. Nath and M. Piskunov, [arXiv:1906.02764 [hep-ph]].
110. E. Pajer and M. Peloso, *Class. Quant. Grav.* **30**, 214002 (2013) doi:10.1088/0264-9381/30/21/214002 [arXiv:1305.3557 [hep-th]].
111. D. J. E. Marsh, *Phys. Rept.* **643**, 1-79 (2016) doi:10.1016/j.physrep.2016.06.005 [arXiv:1510.07633 [astro-ph.CO]].
112. P. Nath and M. Piskunov, *JHEP* **03**, 121 (2018) doi:10.1007/JHEP03(2018)121 [arXiv:1712.01357 [hep-ph]].
113. P. Nath and M. Piskunov, *JHEP* **02**, 034 (2019) doi:10.1007/JHEP02(2019)034 [arXiv:1807.02549 [hep-ph]].
114. S. Tsujikawa, *Class. Quant. Grav.* **30**, 214003 (2013) doi:10.1088/0264-9381/30/21/214003 [arXiv:1304.1961 [gr-qc]].
115. S. Weinberg, *Rev. Mod. Phys.* **61**, 1-23 (1989) doi:10.1103/RevModPhys.61.1

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.