

# Supplementary Materials

## Antitrypanosomal constituents from *Brasenia schreberi* and *Nymphaea lotus* used in a single herbal preparation against sleeping sickness in Angola

Nina Vahekeni<sup>1,2\*</sup>, Théo Brillatz<sup>3</sup>, Marjan Rahmaty<sup>3</sup>, Monical Cal<sup>1,2</sup>, Sonja Keller-Maerki<sup>1,2</sup>, Romina Rocchetti<sup>1,2</sup>, Marcel Kaiser<sup>1,2</sup>, Sibylle Sax<sup>1,2</sup>, Kevin Mattli<sup>4</sup>, Evelyn Wolfram<sup>4</sup>, Laurence Marcourt<sup>3</sup>, Emerson F. Queiroz<sup>3</sup>, Jean-Luc Wolfender<sup>3</sup> and Pascal Mäser<sup>1,2</sup>

\*Corresponding author:

vahe@zhaw.ch; Tel.: +41 078 908 83 86

### Summary

**Table S1:** Overview of all the plants extracts, their preparation and antitrypanosomal activity.

**Figure S1:** ELSD chromatograms for the ethanolic extract of *B. schreberi* before and after VLC enrichment.

**Figure S2:** Growth inhibition activity (%) against *T. b. rhodesiense* of *B. schreberi* leave extracts at 20 and 10 µg/mL.

**Figure S3:** Separation of fraction F6 (chromatogram A), F10 (chromatogram B) and F12 (chromatogram C) of VLC methanolic extract of the leaves of *B. schreberi*.

**Figure S4:** <sup>1</sup>H NMR data and spectrum of compound 1 in CD<sub>3</sub>OD at 600 MHz.

**Figure S5:** NMR data and spectra of compound 2 in CD<sub>3</sub>OD at 600 MHz.

**Figure S6:** <sup>1</sup>H NMR data and spectrum of compound 3 in CD<sub>3</sub>OD at 600 MHz.

**Figure S7:** NMR data and spectra of compound 4 in CD<sub>3</sub>OD at 600 MHz.

**Figure S8:** NMR data and spectra of compound 5 in CD<sub>3</sub>OD at 600 MHz.

**Figure S9:** NMR data and spectra of compound 6 in DMSO-*d*<sub>6</sub> at 600 MHz.

**Figure S10:** NMR data and spectra of compound 7 in CD<sub>3</sub>OD at 600 MHz.

**Figure S11:** NMR data and spectra of compound 8 in CD<sub>3</sub>OD at 600 MHz.

**Figure S12:** UHPLC-HRMS chromatogram showing the presence of compounds 1 and 7 in the decoction of *N. lotus* and correspondence of their MS/MS spectra.

**Figure S13:** UHPLC-HRMS chromatogram showing the presence of compounds 1, 2, 5, 6 and 7 in the decoction of *B. schreberi*.

### References


**Table S2:** Overview of all the plants extracts, their preparation and antitrypanosomal activity.

Extract ID	Plant name	Plant part	Sample labeling	Solvent	Raw mass of dried plant material (g)	Drug/solvent ratio	Weight of dried extract (g)	Yield of extract (%)	Growth inhibition (%) at 20 µg/ml
1	<i>N. lotus &amp; B. schreberi (mixture)</i>	leaves	11	AqDec	10.18	1:20	1.703	16.7	97
2			12	EtOH80%	15.20	1:10	2.27	14.9	95
3			13	MeOH70%	14.8	1:8	1.95	13.2	96
4	<i>C. febrifuga</i>	trunk bark	46	AqDec	30.09	1:20	3.794	12.6	17
5			47	EtOH80%	15.1	1:10	1.88	12.5	26
6			4a	Hexane	100.04	1:3	0.075	0.1	38
7			4b	DCM		1:3	0.088	0.1	31
8			4c	MeOH		1:3	9.139	9.1	26
9			4d	H2O		1:3	1.479	1.5	17
10	<i>C. febrifuga</i>	root	58	AqDec	15.09	1:20	1.673	11.1	16
11			59	AqMac	15.1	1:20	1	6.6	18
12			510	EtOH80%	13.7	1:10	0.92	6.7	22
13			5a	Hexane	50.52	1:3	0.051	0.1	37
14			5b	DCM		1:3	0.080	0.2	37
15			5c	MeOH		1:3	4.576	9.1	31
16			5d	H2O		1:3	0.589	1.2	12
17	<i>C. febrifuga</i>	leaves	611	AqDec	15.08	1:20	3.749	24.9	22
18			612	EtOH80%	15.114	1:10	2.4	15.9	28
19			613	MeOH80%	15.1	1:10	2.95	19.5	26
20			6a	Hexane	100	1:3	1.247	1.2	85
21			6b	DCM		1:3	0.571	0.6	56
22			6c	MeOH		1:3	11.191	11.2	21
23			6d	H2O		1:3	2.656	2.7	12
24	<i>V. madiensis</i>	leaves	714	AqDec	15.0928	1:20	2.442	16.2	28
25			715	EtOH80%	15.2	1:10	1.82	12.0	33
26			716	MetT	15	1:20	2.656	17.7	29
27			717	MeOH	10.5	1:4.2	1.2	11.4	35
28			7a	Hexane	83.94	1:3	0.688	0.8	96
29			7b	DCM		1:3	0.746	0.9	62
30			7c	MeOH		1:3	12.191	14.5	22
31			7d	H2O		1:3	2.667	3.2	17
32	<i>B. owariensis</i>	leaves	818	AqDec	19.995	1:20	5.864	29.3	10
33			819	EtOH80%	20.3	1:10	2.681	13.2	23
34			820	MetT	15.1	1:20	3.509	23.2	11
35			8a	Hexane	100.13	1:3	1.007	1.0	96
36			8b	DCM		1:3	1.621	1.6	74
37			8c	MeOH		1:3	5.123	5.1	34
38			8d	H2O		1:3	8.353	8.3	22
39	<i>E. abyssinica</i>	trunk bark	1021	AqDec	30.259	1:20	5.539	18.3	61
40			1022	EtOH80%	50.19	1:10	3.4	6.8	70
41			10a	Hexane	100.18	1:8	0.404	0.4	75
42			10b	DCM		1:8	0.560	0.6	52
43			10c	MeOH		1:8	13.405	13.4	57
44			10d	H2O		1:8	0.579	0.6	13


45	<i>E. abyssinica</i>	root inside	11a23	AqDec	15.01	1:20	1.431	9.5	85
46			11a24	AqMac	15	1:20	2.1	14.0	103
47			11a25	EtOH80%	20.123	1:10	2.133	10.6	101
48			11a1	Hexane	100.08	1:8	2.608	2.6	58
49			11a2	DCM		1:8	0.167	0.2	33
50			11a3	MeOH		1:8	12.731	12.7	35
51			11a4	H2O		1:8	1.828	1.8	0
52	<i>E. abyssinica</i>	root bark	11b26	AqDec	20.329	1:20	4.349	21.4	88
53			11b27	AqMac	10.2	1:20	1.6	15.7	89
54			11b28	EtOH80%	15.121	1:10	2.249	14.9	98
55		root (all)	11b1	Hexane	100.08	1:4	0.395	0.4	20
56			11b2	DCM		1:4	0.317	0.3	53
57			11b3	MeOH		1:4	17.425	17.4	93
58			11b4	H2O		1:4	0.574	0.6	29
59	<i>M. charantia</i>	whole plant	1229	AqDec	15.015	1:20	2.822	18.8	16
60			1230	AqMac	5	1:20	0.97	19.4	16
61			1231	EtOH80%	24.2	1:10	2.514	10.4	27
62			1232	MeOH80%	9.9	1:10	1.27	12.8	22
63			12a	Hexane	40.81	1:3	0.152	0.4	58
64			12b	DCM		1:3	0.588	1.4	72
65			12c	MeOH		1:3	1.52	3.7	30
66	12d <sub>1</sub>	H2O	1:3	3.949		9.7	22		
67	<i>V. madiensis</i>	root	1333	AqDec	30.4	1:20	4.858	16.0	26
68			1334	EtOH80%	15.1	1:10	1.423	9.4	24
69			13a	Hexane	100.13	1:3	0.102	0.1	79
70			13b	DCM		1:3	0.069	0.1	50
71			13c	MeOH		1:3	7.818	7.8	7
72			13d	H2O		1:3	3.174	3.2	0
73	<i>S. latifolius</i>	root	1435	AqDec	15.136	1:20	2.357	15.6	18
74			1436	AqMac	5	1:20	0.47	9.4	16
75			1437	EtOH80%	29.8	1:10	1.922	6.4	18
76			1438	MeOH	10.1	1:10	0.64	6.3	32
77			14a	Hexane	52.08	1:3	1.911	3.7	18
78			14b	DCM		1:3	0.264	0.5	60
79			14c	MeOH		1:3	2.866	5.5	36
80			14d <sub>1</sub>	H2O		1:3	1.285	2.5	24
81	<i>P. schweinfurthii</i>	leaves	1539	AqDec	15.7	1:20	2.178	13.9	10
82			1540	AqMac	5.3	1:20	0.91	17.2	8
83			1541*	EtOH80%	30	1:10	2.208	7.4	15
84			15a	Hexane	50.06	1:5	0.234	0.5	41
85			15b	DCM		1:5	0.387	0.8	17
86			15c	MeOH		1:5	1.085	2.2	13
87	15d <sub>1</sub>	H2O	1:5	5.373		10.7	6		
88	<i>N. lotus</i>	whole plant	1642	AqDec	14.7	1:20	3.8	25.9	35
89			1643	MeOH70%	15.1	1:8	1.73	11.5	61
90			1644**	EtOH80%	15	1:10	2.196	14.6	58
91			16a	Hexane	21	1:3	0.616	2.9	98
92			16b	DCM		1:3	0.282	1.3	74
93			16c	MeOH		1:3	0.716	3.4	13
94			16d <sub>1</sub>	H2O		1:3	2.018	9.6	22
95	<i>B. schreberi</i>	whole plant	1745	AqDec	10.1	1:20	1.19	11.8	41
96			1746	EtOH80%	20.394	1:10	2.163	10.6	92
97			1747	MeOH70%	15	1:8	1.02	6.8	8
98			17a	Hexane	50.08	1:2	0.179	0.4	94
99			17b	DCM		1:2	0.096	0.2	79
100			17c	MeOH		1:2	1.954	3.9	76
101	17d <sub>1</sub>	H2O	1:2	1.585		3.2	16		

102	<i>B. schreberi</i>	stem (sub	1848	AqDec	10.51	1:20	1.232	11.7	20
103		aquatic	1849	EtOH80%	20.08	1:10	1.804	9.0	31
104			1850	MeOH70%	15.1	1:8	0.79	5.2	21
105			18a	Hexane	19.72	1:5	0.181	0.9	75
106			18b	DCM		1:5	0.079	0.4	70
107			18c	MeOH		1:5	1.645	8.3	23
108			18d <sub>1</sub>	H2O		1:5	1.225	6.2	27
109		<i>B. schreberi</i>	leaves	1951	AqDec	20	1:20	3.5	17.5
110	1952			EtOH80%	25.2	1:10	4.164	16.5	96
111	1953			MeOH70%	15.1	1:8	1.83	12.1	96
112	<i>N. lotus (bungo</i>	whole plant	2054	AqDec	15.1	1:20	3.9	25.8	32
113			2055	EtOH80%	19.806	1:10	2.31	11.7	46
114			2056	MeOH70%	14.96	1:8	2.07	13.8	51
115			20a	Hexane	20.12	1:5	0.404	2.0	96
116			20b	DCM		1:5	0.199	1.0	81
117			20c	MeOH		1:5	1.719	8.5	23
118			20d <sub>1</sub>	H2O		1:5	1.544	7.7	9
119			<i>N. lotus (Damba</i>	whole plant	2157	AqDec	15.1	1:20	3.5
120	2158*	EtOH80%			19.806	1:10	2.31	11.7	23
121	2159	MeOH70%			13.02	1:8	1.29	9.9	18
122	MIX	whole plant	Mix60	AqMac	5.08	1:20	2.33	15.4	6
		root							
		leaves							


**Figure S4:** ELSD chromatograms for the ethanolic extract of *B. schreberi* before and after VLC enrichment.



**Figure S5:** Growth inhibition activity (%) against *T. b. rhodesiense* of *B. schreberi* leave extracts at 20 and 10 µg/mL. The inhibitory activity of the crude ethanol 80% extract is compared to its enriched VLC fractions. VLC methanolic fraction (VLC\_MeOH) displayed most promising antitrypanosomal activity with a GI (%) of 84.6 at 10 µg/mL and 95.1% at 20 µg/mL. Legends for *B. schreberi* leave extracts: ethanol 80% crude extract (BS\_EE80), VLC aqueous fraction ((BS\_EE80\_VLC\_H2O), VLC methanolic fraction ((BS\_EE80\_VLC\_MeOH), VLC ethyl acetate fraction ((BS\_EE80\_VLC\_EtoAc).





**Figure S6:** Separation of fraction F6 (chromatogram A), F10 (chromatogram B) and F12 (chromatogram C) of VLC methanolic extract of the leaves of *B. schreberi*. F6 yielded methyl gallate (2) and 2,3,4,6 tetragalloyl-gluco-pyranoside (3). UV detector at 280 nm. Subfractionation of F10 yielded ethyl gallate (4) and 1,2,3,4,6 pentagalloyl- $\beta$ -gluco-pyranoside (5); UV detector at 280 nm. F12 yielded hypolaetin-7-*O*-glucoside (7); UV detector at 254 nm. P19\_EE80\_VLC\_MeOH = VLC ethanolic extract of the leaves of *B. schreberi*.



## NMR data


**Figure S4:**  $^1\text{H}$  NMR data and spectrum of compound **1** in  $\text{CD}_3\text{OD}$  at 600 MHz.


Gallic acid (**1**) (Kamatham *et al.*, 2015):  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 600 MHz)  $\delta$  7.04 (2H, s, H-2, H-6). HRESIMS  $m/z$  169.0136 [ $\text{M}-\text{H}$ ] $^-$  (calcd for  $\text{C}_7\text{H}_6\text{O}_5^-$ , 169.01370,  $\Delta = -3.7$  ppm).




**Figure S5:** NMR data and spectra of compound **2** in  $\text{CD}_3\text{OD}$  at 600 MHz.

Methyl gallate (**2**) (Kamatham *et al.*, 2015):  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 600 MHz)  $\delta$  (3H, s,  $\text{CH}_3$ -8), 7.04 (2H, s, H-2, H-6);  $^{13}\text{C}$  NMR ( $\text{CD}_3\text{OD}$ , 151 MHz)  $\delta$  51.9 ( $\text{CH}_3$ -8), 109.7 (C-2, C-6), 139.6 (C-4), 163.2 (C-3, C-5), 168.8 (C-7). HRESIMS  $m/z$  183.0302 [ $\text{M}-\text{H}$ ] $^-$  (calcd for  $\text{C}_8\text{H}_7\text{O}_5^-$ , 183.02935,  $\Delta = 1.8$  ppm).







<sup>1</sup>H NMR spectrum of compound 2 in CD<sub>3</sub>OD at 600 MHz.



COSY NMR spectrum of compound 2 in CD<sub>3</sub>OD.




Edited-HSQC NMR spectrum of compound 2 in CD<sub>3</sub>OD.



HMBC NMR spectrum of compound 2 in CD<sub>3</sub>OD.

**Figure S6:**  $^1\text{H}$  NMR data and spectrum of compound **3** in  $\text{CD}_3\text{OD}$  at 600 MHz.


2,3,4,6 tetragalloyl-glucopyranoside (**3**) (Tanaka *et al.*, 1983): Mixture of  $\alpha$  and  $\beta$ -glucopyranoside (1.0/0.7).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 600 MHz)  $\delta$  4.32 (2H, m,  $\alpha$ -H-6b,  $\beta$ -H-6b), 4.44 (1H, d,  $J = 11.1$  Hz,  $\alpha$ -H-6a), 4.50 (1H, dd,  $J = 12.3, 2.4$  Hz,  $\beta$ -H-6a), 5.08 (1H, d,  $J = 7.9$  Hz,  $\beta$ -H-1), 5.15 (1H, dd,  $J = 10.0, 3.5$  Hz,  $\alpha$ -H-2), 5.24 (1H, dd,  $J = 9.8, 7.9$  Hz,  $\beta$ -H-2), 5.52 (3H, m,  $\alpha$ -H-1,  $\alpha$ -H-4,  $\beta$ -H-4), 5.74 (1H, t,  $J = 9.8$  Hz,  $\beta$ -H-3), 6.01 (1H, t,  $J = 10.0$  Hz,  $\alpha$ -H-3), H-2'/H-6' of galloyl from  $\alpha$ -form: 6.91 (2H, s), 6.98 (2H, s), 7.02 (2H, s), 7.12 (2H, s), H-2'/H-6' of galloyl from  $\beta$ -form: 6.87 (2H, s), 6.96 (2H, s), 7.00 (2H, s), 7.11 (2H, s). HRESIMS  $m/z$  787.1049 and 787.1051 [ $\text{M}-\text{H}$ ] $^-$  (calcd for  $\text{C}_{34}\text{H}_{27}\text{O}_{22}$  $^-$ , 787.09940,  $\Delta = 6.3$  ppm).




$^1\text{H}$  NMR spectrum of compound **3** in  $\text{CD}_3\text{OD}$  at 600 MHz.

**Figure S7:** NMR data and spectra of compound **4** in CD<sub>3</sub>OD at 600 MHz.


Ethyl gallate (**4**) (Leela *et al.*, 2013): <sup>1</sup>H NMR (CD<sub>3</sub>OD, 600 MHz) δ 1.35 (3H, t, *J* = 7.1 Hz, CH<sub>3</sub>-9), 4.27 (2H, q, *J* = 7.1 Hz, H-8), 7.04 (2H, s, H-2, H-6); <sup>13</sup>C NMR (CD<sub>3</sub>OD, 151 MHz) δ 14.2 (CH<sub>3</sub>-9), 61.2 (C-8), 109.7 (C-2, C-6), 121.5 (C-1), 139.6 (C-4), 168.2 (C-7). HRESIMS *m/z* 197.0459 [M-H]<sup>-</sup> (calcd for C<sub>9</sub>H<sub>9</sub>O<sub>5</sub><sup>-</sup>, 197.04500, Δ = 1.9 ppm).



<sup>1</sup>H NMR spectrum of compound **4** in CD<sub>3</sub>OD at 600 MHz.




Edited-HSQC NMR spectrum of compound **4** in CD<sub>3</sub>OD.




HMBC NMR spectrum of compound **4** in CD<sub>3</sub>OD.

**Figure S8:** NMR data and spectra of compound **5** in CD<sub>3</sub>OD at 600 MHz.


1,2,3,4,6 pentagalloyl- $\beta$ -glucopyranoside (**5**) (Cui *et al.*, 2002): <sup>1</sup>H NMR (CD<sub>3</sub>OD, 600 MHz)  $\delta$  4.38 (1H, dd,  $J$  = 12.1, 4.4 Hz, H-6b), 4.41 (1H, m, H-5), 4.51 (1H, dd,  $J$  = 12.1, 1.6 Hz, H-6a), 5.59 (1H, dd,  $J$  = 9.7, 8.3 Hz, H-2), 5.62 (1H, t,  $J$  = 9.7 Hz, H-4), 5.90 (1H, t,  $J$  = 9.7 Hz, H-3), 6.24 (1H, d,  $J$  = 8.3 Hz, H-1), 6.90 (2H, s, H-2'-3, H-6'-3), 6.95 (2H, s, H-2'-2, H-6'-2), 6.98 (2H, s, H-2'-4, H-6'-4), 7.05 (2H, s, H-2'-1, H-6'-1), 7.11 (2H, s, H-2'-6, H-6'-6); <sup>13</sup>C NMR (CD<sub>3</sub>OD, 151 MHz)  $\delta$  63.1 (C-6), 69.8 (C-4), 72.2 (C-2), 74.1 (C-3), 74.4 (C-5), 93.8 (C-1), 110.3 (C-2'-6, C-6'-6), 110.4 (C-2'-3, C-6'-3), 110.4 (C-2'-2, C-6'-2), 110.5 (C-2'-4, C-6'-4), 110.6 (C-2'-1, C-6'-1), 140.0 (C-4'-6), 140.1 (C-4'-3), 140.3 (C-4'-4), 140.4 (C-4'-2), 140.8 (C-4'-1), 146.3 (C-3'-3, C-5'-3), 146.4 (C-3'-2, C-5'-2), 146.5 (C-3'-4, C-5'-4), 146.5 (C-3'-6, C-5'-6), 146.6 (C-3'-1, C-5'-1), 166.2 (C-7'-1), 166.9 (C-7'-4), 167.0 (C-7'-2), 167.3 (C-7'-3), 167.9 (C-7'-6). HRESIMS  $m/z$  939.1157 [M-H]<sup>-</sup> (calcd for C<sub>41</sub>H<sub>31</sub>O<sub>26</sub><sup>-</sup>, 939.11036,  $\Delta$  = 5.1 ppm), only observed as  $m/z$  469.0521 [M-H]<sup>2-</sup> in the decoction extract of *B. schreberi*.




<sup>1</sup>H NMR spectrum of compound **5** in CD<sub>3</sub>OD at 600 MHz.




COSY NMR spectrum of compound **5** in CD<sub>3</sub>OD.




<sup>13</sup>C-DEPTQ NMR spectrum of compound **5** in CD<sub>3</sub>OD at 151 MHz.



Edited-HSQC NMR spectrum of compound 5 in CD<sub>3</sub>OD.





HMBC NMR spectrum of compound 5 in CD<sub>3</sub>OD.




ROESY NMR spectrum of compound 5 in CD<sub>3</sub>OD.

**Figure S9:** NMR data and spectra of compound **6** in DMSO-*d*<sub>6</sub> at 600 MHz.


Gossypetin-7-*O*-β-glucopyranoside (**6**) (Yang *et al.*, 2003): <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 600 MHz) δ 3.18 (1H, t, *J* = 9.8, 8.8 Hz, H-4''), 3.31 (1H, t, *J* = 9.2, 8.8 Hz, H-3''), 3.36 (1H, dd, *J* = 9.2, 7.6 Hz, H-2''), 3.42 (1H, ddd, *J* = 9.8, 6.0, 2.1 Hz, H-5''), 3.48 (1H, dd, *J* = 11.8, 6.0 Hz, H-6''b), 3.73 (1H, dd, *J* = 11.8, 2.1 Hz, H-6''a), 4.91 (1H, d, *J* = 7.6 Hz, H-1''), 6.62 (1H, s, H-6), 6.90 (1H, d, *J* = 8.5 Hz, H-5'), 7.63 (1H, dd, *J* = 8.5, 2.2 Hz, H-6'), 7.76 (1H, d, *J* = 2.2 Hz, H-2'), 11.92 (1H, s, 5OH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>, 151 MHz) δ 60.7 (C-6''), 69.7 (C-4''), 73.3 (C-2''), 75.7 (C-3''), 77.3 (C-5''), 97.8 (C-6), 101.4 (C-1''), 104.7 (C-10), 115.3 (C-2'), 120.3 (C-6'), 122.1 (C-1'), 126.8 (C-8), 145.0 (C-3'), 147.9 (C-4'), 150.4 (C-7), 151.5 (C-5). HRESIMS *m/z* 479.0845 [M-H]<sup>-</sup> (calcd for C<sub>21</sub>H<sub>19</sub>O<sub>13</sub><sup>-</sup>, 479.08257, Δ = 2.9 ppm).



<sup>1</sup>H NMR spectrum of compound **6** in DMSO-*d*<sub>6</sub> at 600 MHz.





HMBC NMR spectrum of compound **6** in DMSO-*d*<sub>6</sub>.




ROESY NMR spectrum of compound **6** in DMSO-*d*<sub>6</sub>.

**Figure S10:** NMR data and spectra of compound **7** in CD<sub>3</sub>OD at 600 MHz.


Hypolaetin-7-*O*-glucoside (**7**) (Zapesochnaya et al., 1973): <sup>1</sup>H NMR (CD<sub>3</sub>OD, 600 MHz) δ 3.55 (4H, m, H-2'', H-3'', H-4'', H-5''), 3.75 (1H, dd, *J* = 12.2, 5.3 Hz, H-6''b), 3.92 (1H, dd, *J* = 12.2, 2.3 Hz, H-6''a), 5.00 (1H, d, *J* = 7.6 Hz, H-1''), 6.61 (1H, s, H-3), 6.70 (1H, s, H-6), 6.92 (1H, d, *J* = 8.2 Hz, H-5'), 7.50 (1H, dd, *J* = 8.2, 2.3 Hz, H-6'), 7.51 (1H, d, *J* = 2.3 Hz, H-2'); <sup>13</sup>C NMR (CD<sub>3</sub>OD, 151 MHz) δ 61.9 (C-6''), 71.0 (C-4''), 74.4 (C-2''), 77.2 (C-3''), 78.2 (C-5''), 100.0 (C-6), 102.7 (C-1''), 103.4 (C-3), 107.0 (C-10), 114.1 (C-2'), 116.4 (C-5'), 120.3 (C-6'), 123.4 (C-1'), 128.6 (C-8), 146.8 (C-3'), 150.8 (C-4'), 166.6 (C-2). HRESIMS *m/z* 463.0897 [M-H]<sup>-</sup> (calcd for C<sub>21</sub>H<sub>19</sub>O<sub>12</sub><sup>-</sup>, 463.08765, Δ = 3.3 ppm).




<sup>1</sup>H NMR spectrum of compound **7** in CD<sub>3</sub>OD at 600 MHz.




COSY NMR spectrum of compound 7 in CD<sub>3</sub>OD.



Edited-HSQC NMR spectrum of compound 7 in CD<sub>3</sub>OD.




HMBC NMR spectrum of compound 7 in CD<sub>3</sub>OD.




ROESY NMR spectrum of compound 7 in CD<sub>3</sub>OD.

**Figure S11:** NMR data and spectra of compound **8** in CD<sub>3</sub>OD at 600 MHz.


5-[(8Z,11Z,14Z)-heptadeca-8,11,14-trienyl] resorcinol (**8**) (Barrow and Capon, 1991): <sup>1</sup>H NMR (CD<sub>3</sub>OD, 600 MHz) δ 0.97 (3H, t, *J* = 7.5 Hz, CH<sub>3</sub>-25), 1.34 (8H, m, H-9, H-10, H-11, H-12), 1.57 (2H, m, H-8), 2.08 (4H, m, H-13, H-24), 2.44 (2H, m, H-7), 2.81 (4H, m, H-16, H-19), 5.35 (6H, m, H-14, H-15, H-17, H-18, H-22, H-23), 6.08 (1H, t, *J* = 2.2 Hz, H-4), 6.12 (2H, d, *J* = 2.2 Hz, H-2, H-6); <sup>13</sup>C NMR (CD<sub>3</sub>OD, 151 MHz) δ 14.6 (CH<sub>3</sub>-25), 21.5 (C-24), 26.4 (C-16), 26.5 (C-19), 28.2 (C-13), 30.3 (C-9), 30.3 (C-11), 30.5 (C-10), 30.7 (C-12), 32.4 (C-8), 37.0 (C-7), 100.9 (C-4), 107.9 (C-2, C-6), 128.2 (C-22), 128.8 (C-15), 129.2 (C-18), 129.2 (C-17), 131.1 (C-14), 132.7 (C-23), 146.3 (C-1), 159.3 (C-3, C-5). HRESIMS *m/z* 341.24808 [M-H]<sup>-</sup> (calcd for C<sub>23</sub>H<sub>33</sub>O<sub>2</sub><sup>-</sup>, 341.24806, Δ = 1.7 ppm).




<sup>1</sup>H NMR spectrum of compound **8** in CD<sub>3</sub>OD at 600 MHz.




COSY NMR spectrum of compound **8** in CD<sub>3</sub>OD.




<sup>13</sup>C-DEPTQ NMR spectrum of compound **8** in CD<sub>3</sub>OD at 151 MHz.



Edited-HSQC NMR spectrum of compound 8 in CD<sub>3</sub>OD.




HMBC NMR spectrum of compound 8 in CD<sub>3</sub>OD.




ROESY NMR spectrum of compound **8** in CD<sub>3</sub>OD.


**Figure S12:** UHPLC-HRMS chromatogram showing the presence of compounds **1** and **7** in the decoction of *N. lotus* and correspondence of their MS/MS spectra.




Chromatogram of the decoction of *N. lotus* highlighting compounds **1** and **7**.




MS/MS spectra of compound **1** in the decoction of *N. lotus*.



MS/MS spectra of compound 7 in the decoction of *N. lotus*.




MS/MS spectra of compound 1.




MS/MS spectra of compound 7.


Figure S13: UHPLC-HRMS chromatogram showing the presence of compounds 1, 2, 5, 6 and 7 in the decoction of *B. schreberi*.




Chromatogram of the decoction of *B. schreberi* highlighting compounds 1, 2, 5, 6 and 7.




MS/MS spectra of compound 1 in the decoction of *B. schreberi*.



MS/MS spectra of compound 2 in the decoction of *B. schreberi*.




MS/MS spectra of compound **5** in the decoction of *B. schreberi*, only observed as  $m/z$  469.0521  $[M-H]^-$ .



MS/MS spectra of compound **6** in the decoction of *B. schreberi*.

190711\_LP\_TB\_P19\_EaqD\_neg.raw#782 @1.54 MS2 (463.0888) c -, base peak: 300.0282 m/z (8.6E5)

Scan definition: FTMS - c ESI d Full ms2 463.0888@hcd30.00 [50.0000-490.0000]




■ Scan #782 ■ Peaks in 190711\_LP\_TB\_P19\_EaqD\_neg.raw chromatograms deconvoluted deisotoped

MS/MS spectra of compound 7 in the decoction of *B. schreberi*.

190606\_AR\_TB\_P19\_F06\_2\_01\_Neg.raw#498 @0.92 MS2 (183.0302) c -, base peak: 183.0302 m/z (6.9E7)

Scan definition: FTMS - c ESI d Full ms2 183.0302@hcd30.00 [50.0000-205.0000]




■ Scan #498 ■ Peaks in 190606\_AR\_TB\_P19\_F06\_2\_01\_Neg.raw chromatograms deconvoluted deisotoped

MS/MS spectra of compound 2.

190606\_AR\_TB\_P19\_F10\_2\_13\_Neg.raw#814 @1.51 MS2 (469.0529) c -, base peak: 169.0145 m/z (1.6E7)


Scan definition: FTMS - c ESI d Full ms2 469.0529@hcd30.00 [65.0000-975.0000]



MS/MS spectra of compound 5.

190711\_LP\_TB\_P19\_EaqD\_neg.raw#728 @1.43 MS2 (479.0836) c -, base peak: 317.0305 m/z (1.8E5)

Scan definition: FTMS - c ESI d Full ms2 479.0836@hcd30.00 [50.0000-505.0000]



MS/MS spectra of compound 6.

## References :

Barrow, R.A., Capon, R.J., Alkyl and alkenyl resorcinols from an Australian marine sponge, *Haliclona* sp. (Haplosclerida: haliclonidae). *Aust. J. Chem.* **1991** *44*, 1393-1405.

Cui, C.B., et al., Two new and four known polyphenolics obtained as new cell-cycle inhibitors from *Rubus aleaefolius* Poir. *J Asian Nat Prod Res* **2002** *4*, 243-252.

Kamatham, S., et al., Isolation and characterization of gallic acid and methyl gallate from the seed coats of *Givotia rottleriformis* Griff. and their anti-proliferative effect on human epidermoid carcinoma A431 cells. *Toxicol Rep* **2015** *2*, 520-529.

Leela, V.; Saraswathy, A. Isolation and characterization of phytoconstituents from *Acacia leucophloea* flowers (Roxb) wild. *Int Res J Pharm* **2013**, *4*, 107-109.

Tanaka, T., et al., Tannins and Related-Compounds .14. 7-O-Galloyl-(+)-Catechin and 3-O-Galloylprocyanidin-B-3 from *Sanguisorba officinalis*. *Phytochemistry* **1983** *22*, 2575-2578.

Yang, C., et al., Chemical constituents of *Pyrrosia petiolosa*. *J Asian Nat Prod Res* **2003** *5*, 143-150.

Zapesochnaya, G.; Pangarova, T. Hypolaetin 7-glucoside from *Caryopteris monolica*. *Chemistry of Natural Compounds* **1973**, *9*, 521-521.