

Article

Not peer-reviewed version

Larvicidal Potency of Some Selected Nigerian Plants against *Aedes aegypti*

[Eze Elijah Ajaegbu](#)*, Gloria Tochukwu Onah, [Adeniran John Ikuesan](#), Abdulsheed Momoh Bello

Posted Date: 18 January 2024

doi: 10.20944/preprints202401.1358.v1

Keywords: *Aedes aegypti*, dengue; diseases; arboviruses; toxicity; phytochemical



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Larvicidal Potency of Some Selected Nigerian Plants against *Aedes aegypti*

Eze E. Ajaegbu ^{1,*}, Gloria T. Onah ², Adeniran J. Ikuesan ³ and Abdurashed M. Bello ⁴

¹ Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, David Umahi Federal University of Health Sciences, Uburu Ebonyi State Nigeria; ajaegbuee@yahoo.com

² Department of Applied Sciences, Federal College of Dental Technology and Therapy, Trans-Ekulu, Enugu State, Nigeria; ikuesanjo@gmail.com

³ Science Laboratory Technology, Institute of Management and Technology, Enugu, Nigeria; onah.gloria@yahoo.com

⁴ Department of Applied Sciences, Federal College of Dental Technology and Therapy, Trans-Ekulu, Enugu State, Nigeria; abdurashedbello82@gmail.com

* Correspondence: ajaegbuee@yahoo.com, +2348069744845

Abstract: Many public health-related problems, such as dengue diseases, are caused by the vector *Aedes aegypti*. This paper's objective is to develop larvicidal activities against the larvae of *Ae. aegypti*. Standard methods were utilized for the collection, extraction and phytochemical screening of the plant parts and extracts. Different extract concentrations were tested against the larvae of *Ae. aegypti* mosquitoes. The different toxicities observed were due to changes in the compounds present in each of the plant extracts. Seven phytochemical constituents were detected for plant extracts. Nigerian medicinal plants could be used to curb the spread of the dengue vector, *Aedes aegypti*.

Keywords: *Aedes aegypti*; dengue; diseases; arboviruses; toxicity; phytochemical

1. Introduction

Arboviruses, which have for ages caused infectious diseases like malaria, dengue, Zika, etc., are linked to vector-borne illnesses as a public health problem. Arboviruses are a significant public health issue [1–3]. Arboviruses have expanded more widely as a result of variables like climate change, accelerating globalization, and the development of anthropogenic activities like travel. Indeed, it has been estimated to have impacted more than 60% of the global population and killed millions of people every year. Many underdeveloped and emerging countries' economies have increased and their costs have been constrained as a direct result of these tropical diseases [4–6]. With more than twenty-five (25) vaccine candidates in various phases of research, slow advancement has accelerated dramatically in the last ten years despite considerable obstacles in underdeveloped countries where a very effective yellow fever vaccine is available [7,8]. Recent viral outbreaks have drawn the attention of the entire globe and produced a severe public health issue, sparking intense concern over how to stop the spread of these fatal infections. The application of personal defense strategies to prepare for *Ae. aegypti* bites is crucial for preventing dengue and other arbovirus infections [9–11]. Synthetic pesticides have been shown to significantly lower the risk of vector-borne diseases. However, indiscriminate and ongoing use of pyrethroids and organophosphate insecticides has caused mosquito populations to become resistant to them, which has had unfavorable effects such as insecticide resistance in populations, mammalian toxicity, harm to non-target organisms, bioaccumulation, and environmental damage [12–14]. In order to effectively manage and stop the spread of vector-borne diseases, the current main strategies for reducing human-vector contact rely on the use of synthetic insecticides in the form of long-lasting insecticidal nets (LLINs), indoor residual spraying (IRS), and powerful antimalarial medications. For the past few years, these treatments have been effective in lowering the disease burden and mosquito vector populations in various African regions. But due to the overuse of pyrethroids, resistance has begun to appear in a

number of malaria-endemic regions of Africa and the rest of the world. It has been identified as a potential health risk to the population [15–17].

In general, plant-based pesticides are relatively non-toxic and particular to their intended targets. As a result, research has been done to create environmentally friendly herbicides with lower dangers. Consequently, the larvicidal, antifeedant, repellent, oviposition deterrent, growth regulating, and anti-vector actions of nearly 4,000 plant species were assessed as prospective insecticidal compounds [18,19]. The promise of replacing the use of artificial larvicides is signaled by the extraction of bioactive substances found in plants. In light of these issues, research has been concentrated on a number of objectives, including the identification and development of secondary metabolites insecticidal with efficient and safe therapies against arboviruses [20–22].

The available research shows that the potential larvicidal effects of several plants on mosquito vector behavior and reproductive fitness have not been thoroughly studied. It has been demonstrated that these secondary metabolites are effective against both the larvae and adults of numerous mosquito species [23,24]. Therefore, this study aimed at developing larvicidal activities with a certain number of Nigerian plant extracts against larvae of *Ae. aegypti*, the dengue vector.

2. Materials and Methods

2.1. Plant Collection

Some selected Nigerian plants were collected from the Institute of Management and Technology, Enugu State (IMT) in May 2022 and brought to the laboratory, which include: *Mentha pulegium* leaf (MUML); *Salvia rosmarinus* stem (MISS); *Salvia rosmarinus* leaf (MISL); *Nepeta cataria* leaf (MNCL); *Nepeta cataria* stem (MNCS); *Nepeta cataria* root (MNCR); *Ageratum conyzoides* stem (MCCS); *Psidium guajava* (MUGL); *Lantana camara* stem (MELS); *Lantana camara* leaf (MELL); *Mentha pulegium* stem (MUMS); *Cymbopogon citratus* leaf (MULL); *Ocimum gratissimum* stem (MCSS); *Mentha piperita* L. (peppermint) leaf (MOML); *Mentha piperita* L. (peppermint) stem (MOMS); *Ageratum houstonianum* leaf (MMML); *Melissa officinalis* root (MELR); *Azadirachta indica* (neem) leaf (MMNL); *Geranium* leaf (MMGL); *Azadirachta indica* (neem) stem (MONS). Following the previous protocol of Onah et al., 2022 [25] and Ajaegbu et al., 2022 [26], the different parts of these plants (leaves, flowers, roots, and stems) were collected, identified, cleaned, dried, powdered and prepared for extraction.

2.2. Preparation of the Nigerian Plant Extracts

The powders of different Nigerian plants were accurately and separately weighed, and 100g of each of the plant materials were extracted in 500 ml of methanol by a cold maceration process. The extraction proceeded for a period of two days at 10 hours per day with thorough shaking in the laboratory of the Chemistry unit, Department of Applied Sciences, Federal College of Dental Technology and Therapy, Trans-Ekulu, Enugu State. The extracts in suspension were filtered with Whatman filter paper. The crude extracts of different Nigerian plants were concentrated to dryness at 40–5°C using a rotary vacuum evaporator [27].

2.3. Rearing of Test Organism

The eggs of *Ae. aegypti*, were bought from the National Arbovirus and Vectors Research Centre, Enugu. The colony of *Ae. aegypti* was nurtured and maintained with tap water in the laboratory of the School of Preliminary Studies, Federal College of Dental Technology and Therapy, Trans-Ekulu, Enugu State. Mosquitoes were reared at (26 ± 3°C) of room temperature, 80 ± 4% relative humidity (RH), and 12:12 light/dark (L:D) under photoperiod cycles. Larvae of *Ae. aegypti* were fed upon a mixture of fish and chicken feed (grower) in the ratio of 3:1 with adequate attention by changing the water from the culture bowl every alternate day in order to forbid the establishment of any scum on the outer boundary of the water until IV-instar larvae were used for bioassay [28].

2.4. Mosquito-Borne Larvicidal Activity against *Ae. aegypti*

The larvicidal activity of the Nigerian plant extracts was assayed against IV-instar larvae in line with the reference standard (WHO, 2005) [29] and Lame et al., 2015 [30]. The complete test organism was examined at room temperature (26.3 °C) and relative humidity (RH) of (80.4%). To help the plant components dissolve in water, an emulsifier (Tween 80) was used to prepare the stock concentration of each of the plant extracts. A stock solution made up of 1 g of precisely weighed extract and 2 ml of Tween 80 was diluted with 100 ml of tap water. To create the test solutions of 125, 250, 500, and 1000 ppm accessible against the larvicidal activity of *Ae. aegypti*, each stock solution was serially diluted. As a negative control, a solution of 1 ml of Tween 80 and 99 ml of tap water was used. As a positive control, a daktin insecticide (Dichlorvos, 100% EC weight/volume, 2500 ppm) was chosen. Twenty-five (25) early IV instar larvae were placed in a 250 ml beaker along with 100 ml of each test plant extract, and after a 24-hour exposure period, the number of dead larvae for both the test plant solution and the control was recorded. For the adjustment of the observed negative control mortality range of 5-20%, Abbott's formula was suggested. Nevertheless, when bioassay testing revealed >20% negative control mortality, the trials were abandoned and rerun. Larvae that did not respond to light poking with a small needle were deemed dead. (Abbott 1925) [27].

2.5. Phytochemical Screening of Plant Extracts for Larvicidal Efficacy

The phytochemical was subjected to investigation for the possible components causing toxicity in insects, which was carried out in line with the methods of Younoussa et al., 2015 [2015]. These techniques are based on the identification of secondary metabolites such as oils, fats, resins, steroids, saponins, tannins, alkaloids, flavonoids, resins, alkaloids, and flavonoids.

2.6. Data Analysis

ANOVA was used using the Statistical Package for Social Sciences (SPSS 23.0) to examine the data that had been collected. The Student Newman Keuls (SNK) test was used to determine the mean and standard deviation, which were substantially different at $p < 0.05$. In order to compare the larvicidal effects of the test plants on *Ae. aegypti* statistically, probit analysis was used to determine the lethal dosages that result in 90% (LC₉₀) and 50% (LC₅₀) dead rates of larvae within 24 hours post-exposure. Other statistics included 95% lower and upper confidence limits (LCL and UCL), Chi-square, slope, and standard error of the mean in the all-bioassay.

3. Results

3.1. Larvicidal Activity of Different Plant Extracts against *Ae. aegypti*

Twenty plant extracts with concentrations ranging from 125 to 1000 ppm were examined and tested for their ability to kill *Ae. aegypti* larvae. Twenty (20) plants provided the extracts. 15 of the plant extracts had LC₅₀ values that ranged from 412.90 to 17640.41 µg/ml that made them effective against *Ae. aegypti*. After being exposed to adults of *Aedes aegypti* for 24 hours, it was discovered that *Nepeta cataria* stem (412.90 µg/ml) had the highest level of toxicity of the fifteen plant extracts that had larvicidal activity, followed by *Salvia rosmarinus* leaf (473.87 µg/ml), *Salvia rosmarinus* stem (515.632 µg/ml), *Ageratum conyzoides* stem (1612.22 µg/ml). These five plant extracts, *Mentha pulegium* stem, *Mentha piperita* L. (peppermint stem), *Azadirachta indica* (neem) leaf, *Geranium* leaf, and *Azadirachta indica* (neem) stem, did not cause any larvae deaths when used at the same dose (Table 1).

Table 1. Larvicidal potentials of the plant extracts.

Plant Extracts	Conc (ug/ml)	% Mortality (Mean \pm SD)	LC ₅₀ (LCL–UCL) (ppm)	LC ₉₀ (LCL–UCL) (ppm)	Slope \pm SE	χ^2
MUML	125	0 \pm 0 ^a				
	250	3 \pm 1 ^b	1695.51	9643.95	1.70 \pm 0.555	1.41
	500	5 \pm 1 ^c	(941.86-	(2901.56-		
	1000	8 \pm 1 ^d	16527.09)	1953785.82)		
	F-value	45.33				
MISS	125	5 \pm 1 ^a				
	250	10 \pm 1 ^b	515.632	6422.89	1.17 \pm 0.394	0.56
	500	13 \pm 1 ^c	(308.17-1435.39)	(1935.12-		
	1000	15 \pm 1.73 ^c		1738207.52)		
	F-value	37.83				
MISL	125	8 \pm 1 ^a				
	250	10 \pm 2 ^a	473.87	17315.88	0.82 \pm 0.382	0.03
	500	13 \pm 1.73 ^b	(191.64-	(2544.20-		
	1000	15 \pm 1 ^c	18113.64)	5.309E+21)		
	F-value	12.89*				
MNCL	125	0 \pm 0 ^a				
	250	0 \pm 0 ^a	2175.56	4936.21	3.06 \pm 2.476	0.30
	500	0 \pm 0 ^a	-	-		
	1000	3 \pm 1 ^b				
	F-value	27.0				
MNCS	125	0 \pm 0 ^a				
	250	13 \pm 1.73 ^b	412.90	1581.25	2.20 \pm 0.450	9.55
	500	15 \pm 1 ^b	-	-		
	1000	18 \pm 1 ^c				
	F-value	151.2				
MNCR	125	0 \pm 0 ^a				
	250	3 \pm 1 ^b	17640.41	646470.32	0.82 \pm 0.576	2.57
	500	3 \pm 1 ^b	-	-		
	1000	3 \pm 1 ^b				
	F-value	9.0*				
MCGS	125	0 \pm 0 ^a				
	250	0 \pm 0 ^a	1612.22	3555.40	3.73 \pm 1.725	0.84
	500	0 \pm 0 ^a	(1101.48-	(1737.40-		
	1000	6 \pm 1 ^b	870933.150)	3799407100)		
	F-value	108.0				
MUGL	125	0 \pm 0 ^a				
	250	0 \pm 0 ^a	2175.56	4936.21	3.60 \pm 2.476	0.30
	500	0 \pm 0 ^a	-	-		
	1000	3 \pm 1 ^b				
	F-value	27.0				
MELS	125	6 \pm 1.73 ^a				
	250	6 \pm 1 ^a	3463.29	837509.55	0.54 \pm 0.395	0.20
	500	8 \pm 1.73 ^b	-	-		
	1000	10 \pm 2 ^b				
	F-value	4.0*				
MELL	125	3 \pm 1.73 ^a				
	250	6 \pm 1 ^b	1645.17	34761.00	0.967 \pm 0.419	0.22

	500	8 ± 1 ^c	(726.39-	(4239.23-		
	1000	10 ± 1 ^c	5292069.42)	2.481E+15)		
	F-value	17.83*				
MUMS	125	0 ± 0				
	250	0 ± 0	-	-	-	-
	500	0 ± 0				
	1000	0 ± 0				
	F-value	-				
MULL	125	0 ± 0 ^a				
	250	0 ± 0 ^a	2175.56	4936.21	3.60±2.476	0.30
	500	0 ± 0 ^a	-	-		
	1000	3 ± 1.73 ^b				
	F-value	9.0*				
MCSS	125	0 ± 0 ^a				
	250	0 ± 0 ^a	2175.56	4936.21	3.60±2.476	0.30
	500	0 ± 0 ^a	-	-		
	1000	3 ± 1 ^b				
	F-value	27.0				
MOML	125	0 ± 0 ^a				
	250	0 ± 0 ^a	2175.56	4936.21	3.60±2.476	0.30
	500	0 ± 0 ^a	-	-		
	1000	3 ± 1 ^b				
	F-value	27.0				
MOMS	125	0 ± 0				
	250	0 ± 0	-	-	-	-
	500	0 ± 0				
	1000	0 ± 0				
	F-value	-				
MMML	125	0 ± 0 ^a				
	250	0 ± 0 ^a	2175.56	4936.21	3.60±2.476	0.30
	500	0 ± 0 ^a	-	-		
	1000	3 ± 1 ^b				
	F-value	27.0				
MELR	125	0 ± 0 ^a				
	250	3 ± 1 ^b	4438.49	51152.60	1.21±0.571	2.04
	500	3 ± 1 ^b	(1375.81-	(5158.47-3.534E+30		
	1000	5 ± 1 ^c	9.723E+15)			
	F-value	17.0*				
MMNL	125	0 ± 0				
	250	0 ± 0	-	-	-	-
	500	0 ± 0				
	1000	0 ± 0				
	F-value	-				
MMGL	125	0 ± 0				
	250	0 ± 0	-	-	-	-
	500	0 ± 0				
	1000	0 ± 0				
	F-value	-				
MONS	125	0 ± 0				
	250	0 ± 0	-	-	-	-
	500	0 ± 0				
	1000	0 ± 0				

F-value	-
---------	---

Means within a product followed by the same letter do not differ significantly at $p = 0.05$ (Student-Newman-Keuls's test); * $p < 0.001$; LC₅₀ and LC₉₀–Lethal concentrations able to kill 50 and 90% of female adults, respectively; LCL–Lower confidence limit; UCL–Upper confidence limits; Number of replicates–3.

3.2. Phytochemical Screening of Plant Extracts

Table 2 lists the phytochemical components of 20 plant extracts that were found to be present, moderately present, very present, or lacking. The presence of particular plant secretions accounts for the effectiveness of methanol extract on *Ae. aegypti* larvae. Stems from *Nepeta cataria* contained only small quantities of tannins, alkaloids, and steroids, according to a phytochemical examination of the generally hazardous plant extracts. *Salvia rosmarinus* stems exhibited relatively numerous saponins and tannins, as well as present alkaloids, flavonoids, and steroids, in contrast to the leaf's highly concentrated tannins, moderately plentiful alkaloids and steroids, and flavonoids. Plant extracts that contain either flavonoids or resins all showed LC₅₀, which include *Mentha pulegium* leaf (1695.51 ppm), *Salvia rosmarinus* stem (515.632 ppm), *Salvia rosmarinus* leaf (473.87 ppm), *Nepeta cataria* root (17640.41 ppm), *Cymbopogon citratus* leaf (2175.56 ppm), *Mentha piperita* leaf (2175.56 ppm), and *Melissa officinalis* root (4438.49 p) after 24 hours post exposure on the larvae of *Ae. aegypti* mosquitoes.

Table 2. Phytoconstituents of the extracts.

Plant Extracts	Phytochemical					
	Saponins	Tannins	Alkaloids	Flavonoids	Resins	Steroids
MUML	-	+++	++	-	+	+
MISS	++	++	+	+	-	+
MISL	-	+++	++	+	-	++
MNCL	-	++	++	-	-	++
MNCS	-	+++	+++	-	-	++
MNCR	-	+++	++	+	-	+
MCGS	-	++	+	-	-	++
MUGL	-	+	++	-	-	++
MELS	-	++	+	-	-	+
MELL	+++	++	+	-	-	+
MUMS	-	+++	+	-	-	+++
MULL	-	++	++	-	+	+
MCSS	-	+	+	-	-	++
MOML	+++	++	++	-	+	+
MOMS	-	+	+	-	-	+
MMML	+++	++	-	-	-	+
MELR	-	-	-	+	-	-
MMNL	++	++	+	-	-	+
MMGL	+	++	+++	-	-	-
MONS	+++	++	+++	-	-	-

+indicates present, ++ indicates moderately present, ++ indicates highly present, – indicates absent.

4. Discussion

Based on the reality that chemicals present in botanicals are safer to employ without spreading contaminated activities to humans and their environments, secondary metabolites are utilized against pests and numerous insects that transmit human-borne viruses. Numerous studies on natural products have demonstrated their effectiveness against a variety of mosquito and bug species [32–34].

Despite displaying larvicidal activity with the same extraction procedure in multiple studies, many adulticidal effects of plant extracts were not identified against pests including mosquito insects.

Choochote's study found that *Kaempferia galangal* had larvicidal potential but was unable to detect adulticidal efficiency [35]. Additionally, Lee and Chiang confirmed that *Stemona tuberosa* has larvicidal action but no adulticidal potential [36]. Since many studies aim to eliminate the larvae, including adult mosquitoes, in order to decrease the number of dengue vector insect-transmitted diseases, it is possible that some plants with larvicidal action did not even harm adult mosquitoes. In the current investigation, following 24 hours of postexposure, comparative evaluations of the toxicity of 20 different plant extracts against *Ae. aegypti*, a dengue carrier, were conducted. The measured toxicity varies according to the various components found in various plant extracts. The most toxic and efficient was *Nepeta cataria* stem, followed by *Salvia rosmarinus* leaf, *Salvia rosmarinus* stem, *Ageratum conyzoides* stem, *Lantana camara* leaf, *Mentha pulegium* leaf, while six plant including *Nepeta cataria* leaf, *Psidium guajava* leaf, *Cymbopogon citratus* leaf, *Ocimum gratissimum* stem, *Mentha piperita* leaf, *Ageratum houstonianum* leaf showed same LC₅₀, followed by *Lantana camara* stem, *Melissa officinalis* root, *Nepeta cataria* root, and at the same concentration, no larval death was recorded for these five plant extracts, including *Mentha pulegium* stem, *Mentha piperita* L. (peppermint) stem, *Azadirachta indica* (neem) leaf, *Geranium* leaf, and *Azadirachta indica* (neem) stem (Table 1). There were no toxicological differences seen in the control group. By exhibiting ovipositor attractant, repellent, larvicidal, and adulticidal effects on insect growth that have been reported by various activities noted by numerous studies, phytochemicals present in plants can be employed for the benefit of the public [32,37]. In contrast, pesticides with a botanical origin have mostly been utilized against agricultural pests and, to a much lesser extent, against important insect vectors for public health. Further analysis of the plant extracts' phytochemical components led to the identification of at least three or more of these seven bioactive substances, which include saponins, tannins, alkaloids, flavonoids, resins, and steroids (Table 2). All plant extracts with flavonoids and resins have larvicidal effects on *Ae. aegypti* larvae. Alkaloids, glycosides, saponins, tannins, cardiac glycosides, flavonoids, and terpenoids found in plant extracts have been implicated in several insecticidal, antibacterial, antidiabetic, antihyperlipidemic, and antioxidant actions [38,39]. Phenolic, flavonoid, cardiac glycosides, steroids, tannins, alkaloids, and terpenoid compounds are regarded as significant types of phytochemicals because of their enormous health-related value [39,40].

Due to their phytochemical components and general demonstrated eco-friendliness, plant extracts can be used as an alternative source for mosquito biocides and insecticides, according to this study. This could lower the cost of synthetic insecticides and the environmental risks associated with synthetic chemicals. The mechanism of action and active ingredients that suppress adult mosquitoes should be further studied.

5. Conclusions

This study has shown that methanolic plant extracts include a variety of phytochemicals. This study laid a strong platform for further research into the compound responsible for adulticidal activity and supported the use of plants to control the spread of the dengue vector, *Aedes aegypti*.

Author Contributions: Conceptualization, EEA and GTO; methodology, EEA and AMB; software, EEA and AJI; validation, EEA; formal analysis, EEA, AJI and AMB; investigation, EEA, GTO and AMB; resources, EEA and GTO; data curation, EEA, AJI and AMB; writing—original draft preparation, AJI; writing—review and editing, EEA; visualization, EEA; supervision, EEA; project administration, EEA and GTO; funding acquisition, EEA and GTO. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Das, M.K.; Ansari, M.A. Evaluation of repellent action of *Cymbopogon martinii martinii* Stapf var *Sofia* oil against *Anopheles sundiacus* in tribal villages of Car Nicobar Island, Andaman & Nicobar Islands. *India. J Vect Borne Dis* 2003, 40, 101-104.
2. Islam, M.T.; Quispe, C.; Herrera-Bravo, J.; Sarkar, C.; Sharma, R.; Garg, N.; Fredes, L.I.; Martorell, M.;

- Alshehri, M.M.; Sharifi-Rad, J.; Daştan, S.D.; Calina, D.; Alsafi, R.; Alghamdi, S.; Batiha, G.E.; Cruz-Martins, N. Production, Transmission, Pathogenesis, and Control of Dengue Virus: A Literature-Based Undivided Perspective. *Biomed Res Int* 2021, 4224816.
3. World Health Organization. Vector-borne diseases. Available online: <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>. (accessed on 10 March 2022).
 4. Narkhede, C.P.; Patil, C.D.; Suryawanshi, R.K.; Koli, S.H.; Mohite, B.V.; Patil, S.V. Synergistic effect of certain insecticides combined with *Bacillus thuringiensis* on mosquito larvae. *Journal of Entomological and Acarological Research* 2017, 49, 6265.
 5. Braack, L.; GouveiaDeAlmeida, A.P.; Cornel, A.J.; Swanepoel, R.; De Jager, C. Mosquito-borne arboviruses of African origin: Review of key viruses and vectors. *Parasites & Vectors* 2018, 11, 1.
 6. Karunamoorthi, K.; Ilango, K.; Murugan, K. Laboratory evaluation of traditionally used plant-based insect repellent against the malaria vector *Anopheles arabiensis* Patton (Diptera:Culicidae). *Parasitology Research* 2010, 106(5), 1217–1223.
 7. Artzy-Randrup, Y.; Alonso, D.; Pascual, M. Transimssion intensity and drug resistance in malaria population dynamics: implications for climate change. *Plosone* 2010, 5(6).
 8. World Health Organization. Tables of malaria vaccine projects globally: The rainbow tables. 2003. Available online: www.who.int/immunization/research/development/Rainbow_tables/en/ (accessed on 22 December 2022)
 9. Out, A.; Ebenso, B.; Etokidem, A.; Chukwuekezie, O. Dengue fever - an update review and implications for Nigeria, and similar countries. *Afr Health Sci* 2019, 19(2), 2000-2007.
 10. van den, B.H.; Zaim, M.; Yadav, R.S.; Soares, A.; Ameneshewa, B.; Mnzava, A.; Hii, J.; Dash, A.P.; Ejov, M. Global trends in the use of insecticides to control vector-borne diseases. *Environ Health Perspect* 2012, 120(4), 577-82.
 11. Sunaiyana, S.; MonthathiP, K.; Krajana, T.; Kornwika, S.; Unchalee, S.; Michael, J.B; Theeraphap, C. Comparison of Field and Laboratory-Based Tests for Behavioral Response of *Aedes aegypti* (Diptera: Culicidae) to Repellents. *Journal Of Economic Entomology* 2015, 108(6), 2770-2778.
 12. McGaughey, W.H. Insect resistance to the biological insecticide *Bacillus thuringiensis*. *Science* 1985, 229, 193-195.
 13. Drakou, K.; Nikolaou, T.; Vasquez, M.; Petric, D.; Michaelakis, A.; Kapranas, A.; Papatheodoulou, A.; Koliou, M. The Effect of Weather Variables on Mosquito Activity: A Snapshot of the Main Point of Entry of Cyprus. *Int J Environ Res Public Health* 2020, 17(4), 1403.
 14. Warikoo, R.; Kumar, S. Impact of the *Argemone mexicana* Stem Extracts on the Reproductive Fitness and Behavior of Adult Dengue Vector, *Aedes aegypti* L. (Diptera: Culicidae). *International Journal of Insect Science* 2014, 6, 71–78.
 15. Paaijmans, K.P.; Huijben, S. Taking the 'T' out of LLINs: using insecticides in vector control tools other than long-lasting nets to fight malaria. *Malar J* 2020, 19, 73.
 16. da Silva Sá, G.C.; Bezerra, P.V.V.; Alves da, M.F.; Barboza da, L.; Batista, P.; de Melo, M.F.F.; Uchôa, A.F. Arbovirus vectors insects: are botanical insecticides an alternative for its management. *Journal of Pest Science* 2022, 96(1), 1-20.
 17. Karunamoorthi, K.; Sabesan, S.; Jegajeevanram, K.; Vijayalakshmi, J. The role of traditional anti-malarial plants in the battle against global malaria burden. *Vect.-Borne Zoonot Disease* 2013, 13(8), 521–544.
 18. Souto, A.L.; Sylvestre, M.; Tölke, E.D.; Tavares, J.F.; Barbosa-Filho, J.M.; Cebrián-Torrejón, G. Plant-Derived Pesticides as an Alternative to Pest Management and Sustainable Agricultural Production: Prospects, Applications and Challenges. *Molecules* 2021, 26(16), 4835.
 19. Kaliyaperumal, K.; Askual, G.; Hayleeyesus, S.F. Mosquito repellent activity of essential oil of Ethiopian ethnomedicinal plant against Afro-tropical malarial vector *Anopheles arabiensis*. *Journal of King Saud University – Science* 2014, 26, 305–310.
 20. Sasidharan, S.; Chen, Y.; Saravanan, D.; Sundram, K.M.; Yoga, L.L. Extraction, isolation and characterization of bioactive compounds from plants' extracts. *Afr J Tradit Complement Altern Med* 2011, 8(1), 1-10.
 21. Altemimi, A.; Lakhssassi, N.; Baharlouei, A.; Watson, D.G.; Lightfoot, D.A. Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants (Basel)* 2017, 6(4), 42.
 22. Demarque, D.P., Espindola, L.S. Challenges, Advances and Opportunities in Exploring Natural Products to Control Arboviral Disease Vectors. *Front Chem* 2021, 9, 779049.
 23. Ramzi, A.; El Ouali Lalami, A.; Annemer, S.; Ez zoubi, Y.; Assouguem, A.; Almutairi, M.H.; Kamel, M.; Peluso, I.; Ercisli, S.; Farah, A. Synergistic Effect of Bioactive Monoterpenes against the Mosquito, *Culex pipiens* (Diptera: Culicidae). *Molecules* 2022, 27, 4182.
 24. Ayed, R.B.; Moreau, F.; Hlima, H.B.; Rebai, A.; Ercisli, S.; Kadoo, N.; Hanana, M.; Assouguem, A.; Ullah, R.; Ali, E.A. SNP discovery and structural insights into OeFAD2 unravelling high oleic/linoleic ratio in olive

- oil. *Comput. Struct. Biotechnol. J.* 2022, 20, 1229–1243.
25. Onah, G.T.; Ajaegbu, E.E.; Ezeagha, C.C.; Chigozie, V.U.; Bello, A.M.; Ezeagwu, P.C.; Nwigwe, J.O. Larvicidal and synergistic potentials of some plant extracts against *Aedes aegypti*. *Journal of Entomology and Zoology Studies* 2022, 10(2), 177-180.
 26. Ajaegbu, E.E.; Onah, G.T.; Ikuesan, A.J.; Bello, A.M. Larvicidal synergistic efficacy of plant parts of *Lantana camara* against *Aedes aegypti*. *Journal of Entomology and Zoology Studies* 2022, 10(1), 187-192.
 27. Ibe, I.C.; Ajaegbu, E.E.; Younoussa, L.; Danga, S.P.Y.; Ezugwu, C.O. Larvicidal Property of the Extract and Fractions of *Hannoa klaineana* against the Larvae of *Aedes aegypti*. *Current Journal of Applied Science and Technology* 2020, 39(17), 127-132.
 28. Danga, S.P.Y.; Aboubakar, O.B.F.; Ndouwe, H.M.T.; Yonki, B.; Ngadvou, D.; Younoussa, L.; Ajaegbu, E.E.; Esimone, C.O.; Nukenine, E.N. Towards the use of extracts from *Plectranthus glandulosus* (Lamiaceae) and *Callistemon rigidus* (Myrtaceae) leaves to indoor-spray (control) Malaria and other arboviral diseases vector mosquitoes. *Journal of Entomology and Zoology Studies* 2020, 8(5), 2049-2054.
 29. World Health Organization. Guidelines for laboratory and field testing of mosquito larvicides. Geneva. Available online: <https://www.who.int/publications/i/item/WHO-CDS-WHOPES-GCDPP-2005.13> (accessed on 28 January 2023).
 30. Younoussa, L.; Nukenin, N.E.; Danga, Y.S.P.; Ajaegbu, E.E.; Esimone, C.O. Laboratory Evaluations of the Fractions Efficacy of *Annona senegalensis* (Annonaceae) Leaf Extract on Immature Stage Development of Malarial and Filarial Mosquito Vectors. *J Arthropod-Borne Dis* 2015, 9(2), 226–237.
 31. Abbott, W.S. A method for computing the effectiveness of an insecticide. *J Econ Entomol* 1925, 18, 265-267.
 32. Rajasekaran, A.; Duraikannan, G. Larvicidal activity of plant extracts on *Aedes Aegypti* L. *Asian Pacific Journal of Tropical Biomedicine* 2012, S1578-S1582.
 33. Ajaegbu, E.E.; Uzochukwu, I.C.; Danga, S.P.Y.; Okoye, F. Mosquito adulticidal activity of the leaf extracts of *Spondias mombin* L. against *Aedes aegypti* L. and isolation of active principles. *Journal of vector borne diseases* 2016, 53, 17–22.
 34. Radhika, W.; Naim, W.; Sarita K. Larvicidal potential of commercially available pine (*Pinus longifolia*) and cinnamon (*Cinnamomum zeylanicum*) oils against dengue fever mosquito, *Aedes aegypti* L. (Diptera; Culicidae) *Acta Entomologica Sinica* 2011, 54(7), 793-799.
 35. Choochote, W.; Kanjanapothi, D.; Panthong, A.; Taesotikul, T.; Jitpakdi, A.; Chaithong, U. Larvicidal, adulticidal and repellent effects of *Kaempferia galanga*. *Southeast Asian J Trop Med Public Health* 1999, 30(3), 470–6.
 36. Lee, H.L.; Chiang, Y.F. Insecticidal activity of the herbal plant, *Stemona tuberosa* Lour to mosquito larvae. *Trop Biomed* 1994, 11, 87–90.
 37. Eich, E. Solanaceae and Convolvulaceae: Secondary Metabolites, Biosynthesis, Chemotaxonomy, Biological and Economic Significance, 1st ed.; Springer Berlin, Heidelberg, Germany, 2007; pp. 1-9.
 38. Raj, P.D.; Dutt, P.N.; Bahadur, S.D.; Narayan, Y.U.; Prasad, K.D. Phytochemical screening and study of antioxidant, antimicrobial, antidiabetic, anti-inflammatory and analgesic activities of extracts from stem wood of *Pterocarpus marsupium* Roxburgh. *J Intercult Ethnopharmacol* 2017, 6(2).
 39. Agidew, M.G. Phytochemical analysis of some selected traditional medicinal plants in Ethiopia. *Bull Natl Res Cent* 2022, 46, 87.
 40. Roopashree, T.S.; Raman, D.; Rani, R.; Narendra, C. Antibacterial activity of antipsoriatic herbs: *Cassia tora*, *Momordica charantia* and *Calendula officinalis*. *Int J Appl Res Nat Prod* 2008, 1, 20-28.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.