

Communication

Not peer-reviewed version

Heavy Flavor Physics at the sPHENIX Experiment

[Zhaozhong Shi](#) *

Posted Date: 22 January 2024

doi: [10.20944/preprints202401.1526.v1](https://doi.org/10.20944/preprints202401.1526.v1)

Keywords: sPHENIX; heavy-ion collisions; quark-gluon plasma; heavy flavor physics; particle reconstruction; commissioning; data taking; silicon pixel detector; vertexing; streaming readout; cosmic data; b hadrons



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Communication

Heavy Flavor Physics at the sPHENIX Experiment

Zhaozhong Shi ^{+ID} on behalf of the sPHENIX Collaboration

Los Alamos National Laboratory; zhaozhongshi@lanl.gov; Tel.: +1-415-350-3181

+ Current address: Brookhaven National Laboratory.

Abstract: The sPHENIX experiment is a state-of-the-art jet and heavy flavor physics detector which successfully recorded its first Au + Au collision data at 200 GeV at RHIC. sPHENIX will provide heavy flavor physics measurements covering an unexplored kinematic region and unprecedented precision at RHIC to probe the parton energy loss mechanism, parton transport coefficients in quark-gluon plasma, and the hadronization process under various medium conditions. At the center of sPHENIX, the Monolithic-Active-Pixel-Sensor (MAPS) based VerTeX detector (MVTX) is a high-precision silicon pixel detector. The MVTX provides excellent position resolution and the capability of operating in continuous streaming readout mode, allowing precise vertex determination and recording a large data sample, both of which are particularly crucial for heavy flavor physics measurements. In this work, we will show the general performance of heavy-flavor hadron reconstruction. In addition, we will discuss the commissioning experience with sPHENIX. Finally, we will provide the projection of b-hadron and jet observables and discuss the estimated constraints on theoretical models.

Keywords: sPHENIX; heavy-ion collisions; quark-gluon plasma; heavy flavor physics; particle reconstruction; commissioning; data taking; silicon pixel detector; vertexing; streaming readout; cosmic data; b hadrons

1. Introduction

The sPHENIX experiment [1] is the first new experiment at RHIC in over 20 years. According to the 2015 [2] and 2023 [3] United States Nuclear Science Advisory Committee (NSAC) Long Range Plan for Nuclear Science, the sPHENIX is considered as a Department of Energy flagship experiment in heavy-ion physics. The physics goal of sPHENIX is to probe the inner workings of quark-gluon plasma (QGP) by resolving its properties at shorter and shorter length scales and is complementary with experiments at the LHC.

sPHENIX is a state-of-the-art heavy flavor and jet detector at RHIC. It has a 2π angular acceptance over the rapidity range of $|y| < 1$. It consists of tracking and calorimeter systems with excellent capabilities for studying the strongly interacting QCD matter at RHIC. In addition, sPHENIX is equipped with a minimum bias detector (MBD) and zero degree calorimeter (ZDC) in the far forward region and a trigger system for global event characterization. A schematic drawing of the sPHENIX detector is shown in Figure 1 below.

As the inner tracking system, the Monolithic-Active-Pixel-Sensor (MAPS) based VerTeX detector (MVTX) and the intermediate silicon strip tracker (INTT) leverage advanced silicon detector technologies. The outer tracking system is made of a compact Time Projection Chamber (TPC) and the TPC Outer Tracker (TPOT). The sPHENIX tracking system is able to operate in both triggered and continuous streaming readout modes. The streaming readout acts as a triggerless configuration, capable of sending all collisions to the data acquisition system. The INTT provides us with a timing resolution of about 100 ns, which can resolve RHIC bunch crossings. Utilizing the ACTS tracking algorithm with excellent tracking purity and efficiency [4] and thanks to the excellent performance of the tracking detectors, sPHENIX achieves precision momentum resolution and vertexing determination.




Figure 1. Schematic diagram of sPHENIX detector including all its subdetector systems.

The sPHENIX calorimeter system includes the electromagnetic calorimeter (EMCal) and hadronic calorimeter (HCal). Both EMCal and HCal demonstrate excellent energy linearity and resolutions in Fermilab test beam experiments. Based on the hodoscope position dependent correction, the tower averaged energy resolution of the EMCal is $\Delta E/\langle E \rangle = 3.5\% \oplus 13.3\%/\sqrt{E}$ [5]. The energy resolution of the HCal is $\Delta E/E = 11.8\% \oplus 81.8\%/\sqrt{E}$ [6]. Both of them meet the requirements for its physics goals. Moreover, sPHENIX is the first experiment with full 2π azimuthal barrel hadronic calorimeter in the mid-rapidity range ($|\eta| < 1$) at RHIC, enabling full jet reconstruction in heavy-ion collisions.

2. The MVTX Detector

MVTX is a silicon pixel detector with excellent position resolution approaching $5\ \mu\text{m}$ and offers continuous streaming readout option with a strobe¹ length as short as $5\ \mu\text{s}$ [7], both of which are crucial for sPHENIX heavy flavor physics studies. It is adapted from the inner three layers of Inner Tracking System-2 (ITS-2) detector from the ALICE experiment [8]. The sPHENIX MVTX consists of 48 staves² made of nine ALPIDE chips [9] and electronic signals are readout by one front-end readout unit (RU). At the back end, the data will be transmitted to the Front-End Link eXchange (FELIX) board for further processing [10]. 1 FELIX reads out eight RUs. Six FELIX systems are used to service whole MVTX system.

In addition to the detector and readout electronic systems, we employ negative pressure cooling to maintain overall constant temperature of both the silicon sensors and the readout electronics for their normal operation and safety. Finally, slow control and quality control systems are used to operate and monitor the detector status, ensuring high quality data taking.

MVTX staves were hand carried by air from CERN to LBNL. Two half barrels of the MVTX detectors were assembled at LBNL and then shipped to BNL in October 2022. In March 2023, we installed the MVTX cooling and electronic systems and successfully inserted the MVTX detector to the sPHENIX experiment. All 48 staves passed the electrical tests. We were able to readout the entire MVTX and collect fake hit rate and threshold scan data before the RHIC Au ion beam was turned on.

3. First Year Data Taking

sPHENIX anticipates to collect data from 2023 to 2025 [11]. In 2023, the Au + Au collisions at $\sqrt{s_{NN}} = 200\ \text{GeV}$ data taking had to be curtailed due to an unfortunate valve box failure at RHIC on

¹ A readout frame. Here it is set to be $89\ \mu\text{s}$.

² Rectangular flat planes that hold silicon sensors along the beam axis.

August 1st, 2023. Nevertheless, from May 18th to August 1st 2023, sPHENIX has made significant progress in the detector commissioning with beam. All sPHENIX detector subsystems were able to take and record collision data. sPHENIX continued taking another two months of cosmic data after August 1st for alignment and calibration purposes. All sPHENIX detector subsystems were able to take and record data.

Currently, we are carrying out data production and event reconstruction. Offline data analyses are also ongoing. As part of the tracking system, MVTX is able to synchronize with INTT, TPC, and TPOT to record cosmic ray events. A cosmic event display for tracking commissioning is shown in Figure 2.




Figure 2. A cosmic event display of the tracking system taken on 08/23/2023. In this event, the Level 1 physics trigger for cosmic events, which requires the coincident signals between two outer HCal towers, is fired. In addition, the magnetic field is turned off ($B = 0$).

During the run, MVTX demonstrated a low noise level of less than 10^{-6} hot pixels per chip per event after masking hot pixels [12]. The MVTX mostly operated in continuous streaming readout mode with a few runs in triggered mode. For this online display only, to reject the background and obtain pixels fired by cosmic muons, we remove clusters with single pixels. We can see a clear straight line cosmic muon track from the hits of MVTX, INTT, TPOC, and TPOT.

In addition, we analyzed the Au + Au collision data. We correlated the total number of pixels fired between two MVTX layers. In addition, we also correlate clusters of INTT and TPOT. Figure 3 shows the strong correlations from the subsystems.




Figure 3. Correlation between MVTX layer 0 and 1 within one strobe is shown on the left and INTT and TPOT clusters is shown on the right. Strong correlations are observed in MVTX, INTT, and TPOT with beam collisions.

The studies of both Au + Au beam collision and the cosmic events demonstrate the functionality of sPHENIX tracking system and readiness for offline data analysis.

4. Heavy Flavor Physics Program

To complete the RHIC science mission, the sPHENIX physics program consists of jet substructure, open heavy flavor, quarkonia, bulk physics, and cold QCD physics. sPHENIX plans to take Au + Au data at $\sqrt{s_{NN}} = 200$ GeV in 2023, $p^\uparrow + p^\uparrow$ at $\sqrt{s} = 200$ GeV in 2024, and high luminosity Au + Au at $\sqrt{s_{NN}} = 200$ GeV in 2025. With excellent luminosity recorded, and tracking and vertexing capabilities, sPHENIX will be able to perform precise measurements of fully reconstructed open heavy flavor hadrons such as D^0 , B^+ , and Λ_c^+ .

Thanks to their heavy masses, which are much greater than the QCD scale, Λ_{QCD} , and QGP temperature, T_{QGP} , ($m_Q \gg \Lambda_{QCD}$ and $m_Q \gg T_{QGP}$), heavy quarks, such as charm and beauty quarks, are produced in hard scattering processes in the early stage of heavy-ion collisions and have long thermal relaxation times. Their initial production spectra can be calculated by perturbative QCD. Heavy quarks retain their flavor and mass identities as they traverse through the QGP, making them excellent probes to study the transport properties of QGP. A heavy flavor particle widely used to study QGP is the D^0 meson because of its large charm-quark fragmentation fraction and simple hadronic final states in the decay channel of $D^0 \rightarrow K^-\pi^+$. Figure 4 shows fully reconstructed D^0 mesons in the decay channel of $D^0 \rightarrow K^-\pi^+$ with sPHENIX full detector simulations of Au + Au collisions.




Figure 4. Invariant mass distribution of both prompt and non-prompt D^0 in the decay channel of $D^0 \rightarrow K^- \pi^+$ from with HIJING Au + Au event generator with pile up with the sPHENIX detector in GEANT 4 simulations [13]. In the simulations, *KFParticle* package is applied for secondary vertex reconstruction. It should also be noted that this is 25 minutes of full luminosity data.

According to the simulation studies shown in Figure 4, we expect to obtain the statistics that can observe clear D^0 resonance with about 25 minutes of data taking at 15 kHz. Thanks to the large minimum bias $p + p$ and Au + Au datasets, excellent statistics can be achieved for precise and differential D^0 measurements. The projected performance of classic observables such as D^0 nuclear modification factor R_{AA} and elliptic flow v_2 , which characterize the microscopic properties of the QGP such as energy loss and transport coefficients, are shown in Figure 5




Figure 5. The projected performance of nuclear modification factor R_{AA} (left) and elliptic flow v_2 (right) of prompt (black) and non-prompt (blue) D^0 as a function of $D^0 p_T$ with the sPHENIX experiment from simulations [11] are shown above.

With the precision vertex resolution, we can determine prompt and non-prompt D^0 meson using the distance of closest approach (DCA) in a data-driven manner. The studies of R_{AA} and v_2 of prompt D^0 mesons will help us investigate charm-quark thermalization in the QGP and interaction with the medium constituents. From the R_{AA} and v_2 measurements of non-prompt D^0 feed down from b hadrons, we can study flavor dependence of energy loss and constrain beauty-quark transport coefficients in the QGP.

The charm baryon Λ_c^+ has a more complex 3-prong hadronic decay topology: $\Lambda_c^+ \rightarrow pK^-\pi^+$. sPHENIX is also able to fully reconstruct prompt Λ_c^+ . Preliminary studies to tag protons using dE/dx is underway to improve Λ_c^+ reconstruction. Figure 6 shows the projected Λ_c^+/D^0 ratio performance with sPHENIX in p + p and Au + Au collisions




Figure 6. Projected performance Λ_c^+/D^0 as a function of $\Lambda_c^+ p_T$ with full sPHENIX detector simulations in p + p (black) and Au + Au (blue) [11]. The data point of Λ_c^+/D^0 with STAR Au + Au data at 200 GeV (red) is presented [14]. Theoretical model predictions including Tsinghua, Catania, TAMU, and PYTHIA 8 are also overlaid.

The precise and differential measurement of Λ_c^+/D^0 with sPHENIX over a wide range of p_T and event multiplicity or centrality from p + p to Au + Au will pinpoint the fragmentation and recombination mechanisms of charm quark hadronization from vacuum to QGP at RHIC.

In addition to fully reconstructed open heavy flavor hadrons, sPHENIX is capable of studying b-jets substructure and heavy flavor jet physics at RHIC. Figure 7 shows the R_{AA} as a function p_T and jet z_g distributions of b-jet in p + p and Au + Au collisions.




Figure 7. Projected precision of b-jet R_{AA} as a function of p_T (left) and substructure observable $\frac{1}{N_{jet}} \frac{dN}{dz_g}$ as a function jet z_g in p + p and Au + Au with full sPHENIX detector simulations [11].

Equipped with high performance HCal, sPHENIX is expected to deliver the first full b-jet measurements at RHIC. Moreover, sPHENIX will have excellent b-jet reconstruction and tagging capabilities thanks to the calorimeters and MVTX. sPHENIX will be able to perform differential subjet splitting function measurements with good precision at low p_T . The b-jet substructure measurements will test pQCD model calculations in p + p collisions and quantify the medium modification in the unique sPHENIX kinematic region in Au + Au collisions, complementary to LHC jet substructure measurements

Finally, quarkonium spectroscopy stands as one of the flagship measurements of sPHENIX hidden heavy flavor physics. Because of the color screening effect, the binding energy of Y will decrease as the temperature of QGP increases [15]. In experiments, the sequential suppression of Y(1S), Y(2S), and Y(3S) production will be quantified by nuclear modification factor R_{AA} [16]. Hence, Upsilon can be used as a thermometer to measure the temperature of QGP. The excellent performance of the sPHENIX EMCAL and tracking detectors enables outstanding electron identification capabilities to perform precise $Y \rightarrow e^+e^-$ measurements with sufficient invariant mass resolutions to distinguish all three Upsilon states. Figure 8 shows the dielectron invariant mass distribution near the Upsilon resonances in central Au + Au simulations and Y(1S), Y(2S), and Y(3S) R_{AA} as a function of p_T .




Figure 8. Projected performance Y(1S), Y(2S), and Y(3S) invariant mass distribution in 0-10% Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV from dielectron decay channel using sPHENIX simulation scaled up to 24 billion events is shown on the left. The Y(1S), Y(2S), and Y(3S) R_{AA} as a function of p_T of sPHENIX projection (solid circle) as well as the STAR data (open square) [17] are shown on the right.

We can see that sPHENIX will be able to provide very precise and broad p_T coverage (up to about 9 GeV/c) of Y R_{AA} measurements for quarkonia physics studies. In addition, the Y(3S) resonance may potentially be separated from the Y(2S) and observed in Au + Au collisions for the first time by sPHENIX at RHIC, which is complementary to the recent observation and R_{AA} measurements of Y(3S) with the CMS 2018 LHC PbPb data. [18].

5. Summary

We have introduced the sPHENIX experiment, which is the first new experiment at RHIC in over 20 years to study the microscopic properties of QGP and is currently taking data. The sPHENIX physics program consists of jets, open heavy flavor, quarkonia, bulk physics, and cold QCD. In the 2023 Au + Au run, sPHENIX has demonstrated overall detector functionality and readiness for offline data analysis. The correlation studies of tracking and calorimeter systems validate the synchronization among subdetectors. Aside from the 200 GeV Au + Au collision data, sPHENIX have taken large cosmic datasets for detector alignment and calibration studies. In several cosmic ray events, a clear

straight line muon tracks are observed without the magnetic field, showing overall good quality of the sPHENIX data.

We have also reported the projected physics measurements with full sPHENIX detector simulations assuming the luminosity documented in the 2022 sPHENIX Beam Use Proposal [11]. We expect to achieve high statistics, fully reconstructed charm and beauty hadron measurements to study heavy quark diffusion, energy loss, and hadronization in QGP. Moreover, thanks to the excellent calorimeter, tracking, and vertexing performance, sPHENIX will carry out the first inclusive b-jet measurements with high precision at low p_T , complementary to LHC experiments. Finally, Upsilon spectroscopy, one of the major physics topics of the sPHENIX experiment, provides us with precision measurement of QGP temperature. Potential observation and R_{AA} measurements of $\Upsilon(3S)$ for the first time may be accomplished by sPHENIX at RHIC.

Acknowledgments: We would like to express our gratitude to the ISMD 2023 conference organizers for inviting us to present this work and providing useful information for the proceedings submission. We particularly want to thank Professor Máté Csanád for his friendly correspondence and cordial reception. This work is supported by the United States Department of Energy Office of Science and Los Alamos National Laboratory Laboratory Directed Research & Development (LDRD).

References

1. A. Adare (PHENIX Collaboration) et al., "An Upgrade Proposal from the PHENIX Collaboration", arXiv:1501.06197 (2015)
2. A. Aprahamian et al., "Reaching for the horizon: The 2015 long range plan for nuclear science", Nuclear Science Advisory Committee (2015)
3. A. Aprahamian et al., "A New Era of Discovery: The 2023 Long Range Plan for Nuclear Science", Nuclear Science Advisory Committee (2023), <https://nuclearsciencefuture.org/wp-content/uploads/2023/11/NSAC-LRP-2023-v1.3.pdf>
4. J. D. Osborn et al., "Implementation of ACTS into sPHENIX Track Reconstruction", Comput. Softw. Big Sci. 5 (2021) 1, 23
5. C. Aidala et al., "Design and Beam Test Results for the 2-D Projective sPHENIX Electromagnetic Calorimeter Prototype", IEEE Trans. Nucl. Sci. 68 (2021) 2, 173-181
6. C. Aidala et al., "Design and Beam Test Results for the sPHENIX Electromagnetic and Hadronic Calorimeter Prototypes", IEEE Trans. Nucl. Sci. 65 (2018) 12, 2901-2919
7. M. Liu et al., "A Monolithic-Active-Pixel-Sensor-based Vertex Detector (MVTX) for the PHENIX Experiment at RHIC", proposal submitted to DOE Office of Science, Feb 2017 <https://www.phenix.bnl.gov/WWW/publish/barish/publish/mxliu/sPHENIX/MVTX-Project/MVTX-Preproposal-v1.5-02012017.pdf> (Web)
8. F. Reiddt, "Upgrade of the ALICE ITS detector", Nucl. Instrum. Meth. A 1032 (2022) 166632
9. M. Mager, on behalf of ALICE Collaboration, "ALPIDE, the Monolithic Active Pixel Sensor for the ALICE ITS upgrade", Nucl. Instrum. Meth. A 824 (2016) 434-438
10. J. Anderson et al., "FELIX: a High-Throughput Network Approach for Interfacing to Front End Electronics for ATLAS Upgrades", J. Phys. Conf. Ser. 664 (2015) 8, 082050
11. sPHENIX Collaboration, "sPHENIX Beam Use Proposal", sPH-TRG-2022-01, May 2022, https://indico.bnl.gov/event/15845/attachments/40963/68517/sPHENIX_Beam_Use_Proposal_2022.pdf (Web)
12. sPHENIX Collaboration, "Commissioning Status: August 2023", sPHENIX Notes, August 2023, https://indico.bnl.gov/event/20373/attachments/49393/84311/Commissioning_Status_August_2023.pdf (Web)
13. C. Dean (sPHENIX Collaboration), " $D^0 \rightarrow K^-\pi^+$ Modelling and Selection in Min-bias Au+Au Simulations at sPHENIX", sPHENIX-Invenio (2023) <https://sphenix-invenio.sdcc.bnl.gov/records/7c7gv-7qp49>
14. STAR Collaboration, "First measurement of Λ_c^+ baryon production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV", Phys. Rev. Lett. 124 (2020) 17, 172301
15. S. Digal, P. Petreczky, and H. Satz, "Quarkonium feed down and sequential suppression", Phys. Rev. D 64 (2001) 094015
16. CMS Collaboration, "Observation of Sequential Upsilon Suppression in PbPb Collisions", Phys. Rev. Lett. 109 (2012) 222301

17. STAR Collaboration, "Observation of sequential Y suppression in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV with the STAR experiment", Phys. Rev. Lett. 130 (2023) 11, 112301
18. CMS Collaboration, "Observation of the Y(3S) meson and suppression of Y states in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV", CERN-EP-2023-011, submitted to PRL (2023) <https://arxiv.org/abs/2303.17026>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.