

Review

Not peer-reviewed version

BRAF Mutations: Oncogenic Drivers in Solid Tumors

Paola Roa , Nicole Virginia Bremer , Valentina Foglizzo , [Emiliano Cocco](#) *

Posted Date: 29 January 2024

doi: 10.20944/preprints202401.1957.v1

Keywords: BRAF; solid tumors; NGS; targeted therapy; therapy resistance



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

BRAF Mutations: Oncogenic Drivers in Solid Tumors

Paola Roa, Nicole V. Bremer, Valentina Foglizzo and Emiliano Cocco *

University of Miami, Miller School of Medicine, Department of Biochemistry and Molecular Biology, Sylvester Comprehensive Cancer Center, Miami, Florida 33136, USA.

* Correspondence: emiliano.cocco@miami.edu

Simple summary: In this literature review we explored the milestone events from the discovery of *BRAF* mutations to present day clinical intervention strategies. Through this, we delved into the role of the *BRAF* gene in various cancer types such as melanoma, non-small cell lung cancer, colorectal and thyroid cancer. Additionally, we reviewed clinical trials that led to the FDA approval of therapeutic regimens as monotherapy or, more recently, as combinatorial approaches to treat cancer types harboring *BRAF* hotspot mutations.

Abstract: From their discovery in 2002, *BRAF* mutations have been identified as clear drivers of oncogenesis in several cancer types. Currently, their incidence rate is nearly 7% of all solid tumors with *BRAF* V600E constituting approximately 90% of these diagnoses [1]. In melanoma, thyroid cancer, and histiocytic neoplasms, *BRAF* hotspot mutations are found at a rate of about 50%, while in lung and colorectal cancers they range from 3 to 10% of reported cases [2-5]. Though present in other malignancies such as breast and ovarian cancers, they constitute a small portion of diagnoses (<1%). Given their frequency along with advancements in screening technologies, various methods are used for the detection of *BRAF*-mutant cancers. Among these are targeted next-generation sequencing (NGS) on tumor tissue or circulating tumor DNA (ctDNA) and immunohistochemistry (IHC)-based assays. With advancements in detection technologies, several approaches have been taken for the treatment of *BRAF* mutant cancers. In this review we retrace the milestones that led to the clinical development of targeted therapies currently available for these tumors.

Keywords: *BRAF*; solid tumors; NGS; targeted therapy; therapy resistance

1. Introduction

Located on chromosome 7q34, V-Raf Murine Sarcoma Viral Oncogene Homolog B (*BRAF*) codes for a serine-threonine protein kinase (*BRAF*) which belongs to the rapidly accelerated fibrosarcoma (Raf) protein family, which also includes *ARAF* and *CRAF* [6]. These proteins are direct activators of the mitogen-activated protein (MAP) kinase/extracellular signal-regulated kinase (MEK)/extracellular signal-regulated kinase (ERK) signaling pathway as well as effectors of Rat sarcoma (Ras) proteins (**Figure 1**) [7-14]. Importantly, this pathway plays a key role in cell growth and fate in normal as well as in cancerous cells.




Figure 1. MAPK signaling pathway. Schematic depicting physiological activation of BRAF within the MAPK signaling pathway. Position of a wild-type BRAF kinase dimer (either BRAF/BRAF or BRAF/CRAF) within the MAPK signaling pathway. Receptor tyrosine kinases (RTKs) receive signaling from ligands (indicated by blue triangles), which then begin a signaling cascade that passes through RAS, RAF, MEK, ERK, and into the nucleus to impact cell proliferation and survival. In a wild-type cell, these processes can be activated and inactivated through regulation pathways that conserve typical metabolic order, but oncogenic mutations to these kinases can disrupt the cell’s ability to control these signals [15].

Raf proteins contain three conserved regions (CR) with differing functionalities (**Figure 2**): the CR1 domain is divided into two subdomains: a cysteine-rich domain (CRD) involved with Raf kinase domain autoinhibition and needed for Ras protein interaction, and a Ras-binding domain which serves as the interface for Ras proteins [16]. The CR2 domain functions as an inhibitor of Ras protein binding and Raf activation [17] and the CR3 domain possesses serine and threonine-mediated kinase activity [18].




Figure 2. Secondary structure of BRAF. [Top] Locus of BRAF gene on chromosome 7q34. [Middle] Secondary structure of the BRAF protein from amino acid 1 to 766, with conserved regions (CR) 1-3

and functional domains including the RAS-binding domain (RBD), cysteine-rich domain (CRD), phosphate binding loop (PL), catalytic loop (CL), and activation loop (AL). **[Bottom]** Representation of the 18 exons that make up BRAF, with the number of each exon’s first base pair specified next to the gene (only counting the translated sequence, omitting UTR and flanking sequences).




Figure 3. Timeline of key advancements in the understanding of BRAF mutations. Breakthroughs in research, FDA approvals for novel treatment options, and the discovery of the three classes of BRAF mutations [19-29].

Most notably, these mutations are seen at particularly high rates in melanoma, colorectal cancer and papillary thyroid carcinomas (PTC) [26,30-35] (**Figure 4a**). Specifically, BRAF-activating mutations are primarily limited to the kinase domain, encompassing exons 11 to 15 [36] (**Figure 4b**). Among the various *BRAF* mutations, exon 15 p.V600E is by far the most prevalent [26] (**Figure 4b**).




Figure 4. Frequency and types of BRAF alterations across solid tumors. a: Prevalence of BRAF mutations across various cancer histologies (presented as absolute counts). Only published studies on solid tumors with >100 samples, available on cBioPortal [37], were included. The frequency of BRAF mutations varies from around 1% in cancer types such as esophageal, appendiceal, and gallbladder cancers to up to 35% in melanoma. b: Lollipop diagram depicting BRAF mutations

identified in several cancer types. The most prevalent mutation, V600E, is localized in the kinase domain and has been frequently reported in melanoma, thyroid, colorectal and non-small cell lung cancer. BRAF mutations that mapped in the Raf-like Ras binding and C1 domains are also depicted. These mutations are very rare, their oncogenic potential is mainly unclear, and are found in a variety of histologies.

Given over 200 *BRAF* mutant alleles have been identified in human tumors [29], these mutations have been divided into three classes depending on the activity of BRAF protein present [3,29,35,38,39].

Of the three classes of mutations, class I is the most common given it includes exon 15 p.V600 alterations. By inducing elevated levels of kinase activity, these mutations lead to the activation of the MEK/ERK pathways in a manner that is independent of protein dimerization and Ras activation [3,29,35,38,39]. Class II mutations, like class I mutations, are Ras-independent and include gene fusions as well as various point mutations (exon 11 p.G464E/V, exon 11 p. G469A/R/V, exon 15 p. L597Q/V, and exon 15 p.K601E/N/T). However, unlike class I mutations, class II alterations require protein dimerization to induce MEK/ERK pathway activation and they promote both intermediate and high kinase activity [3,29,35,38,39]. In contrast, class III mutations are RAS-dependent and exhibit both increased affinity for Ras-GTP as well as increased binding with CRAF when compared to BRAF wildtype (WT), leading to hyperactivation of the MEK/ERK pathway [29] (Figure 5).




Figure 5. Classes of BRAF mutation/ Representation of the three major classes of oncogenic mutations in BRAF. Class I involves a RAS-independent, kinase-active BRAF monomer with a V600 mutation. Classes II and III are non-V600 mutations, and both act as dimers with either CRAF or a second BRAF. However, Class II is RAS-independent and results in high kinase activity, while Class III is dependent on mutant RAS and results in little to no kinase activity (represented in the figure by arrow thickness) [29,40,41].

2. BRAF Mutations in Cancer

2.1. Melanoma

BRAF mutations are highly prevalent in melanomas, ranging from approximately 40-60% of all reported cases [4]. Of these alterations, 97% occur at exon 15 (codon 600) [1] with up to 90% being just the p.V600E mutant. This mutant arises from a transverse mutation in nucleotide 1799 turning a T to an A (c.1799T>A), ultimately resulting in the valine to glutamic acid substitution (p.V600E) observed [42]. From an epidemiology standpoint, *BRAF*-mutated melanomas often occur in younger

patients, frequently possessing a superficial diffusion or nodular morphology. Unlike WT cases, *BRAF* mutant melanomas are more prone to brain metastasis [30,43] and are often localized in areas with limited sun damage [44].

Initially, tumors harboring the V600E mutation were not treated differently than those with *BRAF* WT. However, given the significantly shorter response to conventional chemotherapy/radiotherapy, a shift in treatment approach was necessary.

Thus came the rise of RAF inhibitors, developed to explore the possibility of targeted therapy in this context. With the onset of targeted therapies, clinical trials such as coBRIM or COLUMBUS revealed the efficacy of specialized *BRAF* inhibitors in cases of *BRAF* exon 15 p.V600 mutated melanomas [45,46]. In addition to V600E, other mutations observed at this codon are p.V600K accounting for about 10% of cases, p.V600R (1%), p.V600M (0.3%), or p.V600D (0.1%) (**Figure 4b**) [42].

Though there is abundant evidence supporting the efficacy of *BRAF* inhibitors against the most common mutants, their effect in rarer cases has yet to be established [47]. In addition, resistance to therapy has begun to emerge in the clinic. To overcome these limitations, recently, combinatorial therapies including RAF and MEK inhibitors as well as immunotherapy have been explored with promising preliminary results. The COMBI-d (NCT01584648) clinical trial was the first *BRAF*-mutant melanoma focused trial to evaluate the efficacy of the combination of dabrafenib plus trametinib against dabrafenib alone. Upon 36-month follow-up, the PFS of the combination group was 22% against 12% for the dabrafenib monotherapy arm [48]. Additionally, in the phase 3 clinical trial CheckMate 067, 6.5 year follow-up analysis revealed that the combination treatment of nivolumab and ipilimumab worked better than nivolumab alone in patients with *BRAF*-mutated melanoma (6-year PFS 38% vs 23% and 6.5-year overall survival (OS) of 57% vs 43%) [49].

2.2. Thyroid

In general, *BRAF* gene mutations occur more frequently in sporadic papillary thyroid carcinomas (PTCs) such as aggressive microcarcinomas and tall cell variant cancers of adult patients [50,51], followed by poorly differentiated carcinomas and PTC-derived anaplastic thyroid carcinomas (ATCs) [52-55]. In other thyroid lesions such as follicular or medullary carcinomas as well as benign neoplasms, *BRAF* mutations are very rare or never seen [52-55]. However, in general, the presence of *BRAF* V600E has been associated from around 18 to 87% of thyroid cancers [54,56].

Through both *in vitro* and *in vivo* work, it was believed that the V600E mutation in particular, associates with an invasive thyroid cancer phenotype and promotes thyroid cancer progression [57,58]. However, a consensus on whether this can be confirmed has not been reached. Nevertheless, the presence of the *BRAF* exon 15p.V600E point mutation does play a critical role in diagnostic and prognostic outlook in patients [59,60]. In addition to associated poorer outcomes and aggressive behavior [51,61-63] when compared to WT, *BRAF* mutations have been linked to higher risk of disease recurrence and persistence [63,64]. Furthermore, other non-*BRAF* factors such as extrathyroidal tumor invasion, older age, lymph node and distant metastases as well as being male have all been tied to poorer prognostic outcome [53,63,65-70].

2.3. Lung Cancer

From the first report of a *BRAF* mutation in non-small cell lung cancer (NSCLC) in 2011 [32], variable mutation frequencies have been reported ranging from 1.5-3.5% to 7-8% [3,71-73]. Among the mutations identified, class I alterations are the most prevalent, though other mutations have been reported [74]. Importantly, nearly all cases of *BRAF*-mutated NSCLC display strong expression of thyroid transcription factor 1 (TTF-1) and feature an adenocarcinoma morphology with a papillary growth pattern [32]. While *BRAF* mutations are primarily associated with a glandular morphology, some cases have been reported of these alterations in small cell carcinoma as well as varying NSCLC subtypes ranging from pulmonary sarcomatoid carcinoma to squamous cell carcinoma [75-77].

Given the variability in hypothesized epidemiological factors affecting the frequency and distribution of *BRAF* mutations in NSCLC, molecular testing through next-generation sequencing (NGS) provides the best prognostic tool for their identification [78]. In some cases, such as with

smokers harboring *BRAF*-mutant lung cancer, the presence of such a mutation is associated with longer median progression free survival (PFS) when compared to non-mutant groups when treated with immune checkpoint inhibitors. In addition to prognostic value, *BRAF* mutations serve as positive predictive markers for the identification of NSCLC patients who could potentially benefit from targeted therapy [79-82]. For instance, in the lung cancer cohort of the phase II, basket, open label AcSé trial, an objective response was observed in 43/96 patients in the *BRAF* V600-mutated group, while no objective response was reached in the *BRAF* non-V600 arm. Thus, in the case of *BRAF* V600 mutant NSCLC, *BRAF* inhibitors such as vemurafenib are an effective clinical option [83].

2.4. Colorectal

Slightly more prevalent than in lung cancer, *BRAF* mutations are found in about 10% of CRC [5,84]. From an epidemiology standpoint, *BRAF* mutations in CRC are more commonly detected in patients 70 or older and in women. As for their morphology, *BRAF* mutant CRCs are typically localized in the proximal colon, displaying a poorly differentiated histology characterized by serrated and mucinous components [85,86]. In contrast to the other cancer types discussed, in CRC, non-p.V600E mutations occur at higher frequencies than *BRAF* p.V600E in younger patients and men, are located on the distal colon, display low-grade histology, and have a longer median OS in response to chemotherapy [87]. Lastly, *BRAF* mutant CRCs often display high microsatellite instability (MSI-H) as well as a high CpG island methylator phenotype (CIMP-H) [88-90]. In prognostic terms, CRC patients possessing *BRAF* exon 15 p.V600E mutations display lower DFS, OS, and cancer-specific survival (CSS) when compared to non-*BRAF* mutated CRC regardless of disease stage or chemotherapeutic intervention [91]. Unlike the other cancer types discussed above, also targeted therapy has shown disappointing results in *BRAF* mutant CRC with monotherapies being mainly inefficacious and combination therapies, including immunotherapy-based regimens, achieving ORR in just about 30% of patients [92].

2.5. Other Cancers

Histiocytic neoplasms such as Langerhans cell histiocytosis (LCH) and Erdheim-Chester disease (ECD) are derived from macrophage/dendritic lineages and known to be enriched for *BRAF* V600 mutations with frequencies up to 50% [2,93]. As such, clinical trials like the VE-BASKET study (NCT01524978), which evaluated the efficacy of vemurafenib in nonmelanoma cancers harboring *BRAF* V600 mutations, revealed that these mutants are highly targetable with an ORR of 61.5% in the larger cohort and an ORR of 54.5% in the ECD cohort [2].

Oncogenic mutations in *BRAF* are also seen at low frequencies in other cancer types (**Figure 4a**). For example, pediatric low-grade gliomas (PLGGs) often harbor the *BRAF* p.V600E point mutation. Importantly, patients with mutated tumors have poorer clinical outcomes than patients presenting with *BRAF* WT disease [94,95]. For example, in a longitudinal study conducted in Ontario in which PLGG patients were treated with a combination of chemo and radiation therapy, 69 of 405 of them were found to harbor the *BRAF* V600E mutation. This cohort exhibited a 10-year PFS of 27% compared to 60.2% in the *BRAF* WT group [95]. In contrast, the presence of the *BRAF* exon 15 p.V600E has no impact on prognosis in patients with glioblastoma even when tumors harbor epithelioid morphology without isocitrate dehydrogenase (*IDH*) alterations [96].

Though frequencies are low, *BRAF* mutations have also been reported in both breast cancer and ovarian cancer [97].

3. Diagnostic Approaches for *BRAF* Mutations

3.1. Detection of mutations

With the advancements in diagnostic technologies, a wide range of methods exist for the detection and diagnosis of *BRAF*-mutant cancers. Among these are sequencing-based and PCR-based methods as well as immunohistochemistry (IHC) [98,99]. Additionally, the onset of liquid biopsy testing technologies has allowed for minimally invasive genetic testing, giving the opportunity to

evaluate clonal evolution as well as resistance mechanisms through the course of treatment [40,100-103]. The shift from genotyping focused on few genes with sequential testing through single gene-assays to next-generation sequencing (NGS) which scans a larger set of genes, has allowed for the detection of previously undiscovered alterations as well as increased the likelihood of detecting rare alterations [74]. As for more studied genes, NGS provides the opportunity to distinguish uncommon genotypes, which are often overlooked in hotspot PCR-based assays [104]. Furthermore, NGS allows for the identification of several classes of alterations such as fusions, amplifications, and mutations [105,106] as well as offers the potential to unveil novel alterations, not yet discovered.

Though DNA-based NGS is usually the main means for genotyping, its variable sensitivity for alternatively spliced transcripts and fusions renders it rather unreliable for analyzing minute changes [107,108]. However, the exploration of RNA-based methods has allowed for the direct assessment of oncogenic RNA transcripts lacking large intronic sequences. This has enabled more sensitive and efficient analyses, leading to the detection of occult kinase fusions which are often overlooked by DNA sequencing [107,108]. In addition to being more efficient, RNA-based analysis allows for higher specificity through confirmation that some fusions produce novel chimeric transcripts while others of uncertain significance detected in DNA are not transcribed into oncogenes [109]. Given RNA sequencing can directly capture aberrant splicing byproducts, it largely outperforms DNA hybrid capture-based target enrichment allowing for the determination of which variants lead to certain exons being skipped [110]. Though RNA-based sequencing offers many advantages when compared to DNA-based assays, RNA is often prone to clinical testing failure given its predisposition to degradation and lability [111,112]. Regarding oncogenic hotspot mutations, a DNA-based technique should be sufficient for a reliable diagnosis. RNA-based assays are instead preferable to detect BRAF fusions which occur at low frequencies in melanoma (3%), glioma (2%) thyroid cancer (1%), pancreatic carcinoma (0.3%), NSCLC (0.2%) and colorectal cancer (0.2%) [113]. Another alternative is IHC utilizing the anti-BRAF V600E (VE1) mouse monoclonal antibody which was generated to specifically recognize the mutated amino acid sequence from amino acids 596 to 606 [114]. Given IHC equipment is both widely available in pathology laboratories and gives much faster results than molecular biology techniques, the VE1 IHC diagnostic method provides a great alternative to conventional BRAF V600E genotyping, particularly for those tumor types in which this mutation is frequently found (e.g., melanoma, histiocytic neoplasms).

While adequate tumor tissue samples are pivotal for successful NGS, they are often not enough for comprehensive testing and collecting them can be invasive for the patient [115]. Through the use of around 3–10 mL of plasma followed by the analysis of circulating tumor DNA (ctDNA), liquid biopsies can provide additional information supplementing tissue-based approaches or replace them altogether in certain instances [116]. While they serve as an alternative approach, liquid biopsies require ultra-deep sequencing (>10,000x depth) and often include fewer genes than tissue-based panels in order to balance sequencing depth and breadth [117]. This, along with the fact that ctDNA in plasma is rather scarce, results in the exclusion of less commonly altered genes as those alterations which are highly recurrent are prioritized.

4. Targeted therapies for BRAF-mutated cancers

Before delving into the currently available FDA-approved therapies for the treatment of BRAF mutant tumors (summarized in **Table 1**), it is important to understand how BRAF inhibitors interact with the BRAF kinase itself. Like most kinases, RAFs are composed of two domains: the N-terminal and C-terminal lobes [118]. These domains are held together by a flexible hinge, and the cleft between them acts as an active site that can bind to substrates such as ATP and kinase inhibitors. Two regulatory elements of BRAF, the α C-helix and the DFG motif, are particularly relevant to the function of its inhibitors [119]. Under typical conditions these elements can switch between inactive-OUT and active-IN conformations, but treatment with RAF inhibitors can cause allosteric structural changes that lock specific conformations into place [28,120]. These conformations can be used to categorize BRAF inhibitors into type I (α C-helix-IN/DFG-IN), type I_{1/2} (α C-helix-OUT/DFG-IN), and type II (α C-helix-IN/DFG-OUT) [120]. BRAF inhibitors can also be categorized by generations, with first-

generation being the least specific treatment and third-generation being the most recent developments.

Table 1. BRAF inhibitors and combination therapies currently FDA-approved to treat BRAF mutated cancers.

BRAF inhibitors			
Drug	Inhibitor type	Approved to treat	Date approved by the FDA
Sorafenib (Nexlavar)	1st-gen BRAFi	Advanced renal cell carcinoma	12/20/2005
Also: Sorafenib tosylate		Unresectable hepatocellular carcinoma	
		Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma (DTC) refractory to radioactive iodine treatment	
Vemurafenib (Zelboraf)	2nd-gen BRAFi	Unresectable or metastatic melanoma with BRAF V600E mutation	8/17/2011
Also: PLX4032		Erdheim-Chester disease with BRAF V600 mutation	
Dabrafenib (Tafinlar)	2nd-gen BRAFi	Unresectable or metastatic melanoma with BRAF V600E mutation	5/29/2013
Also: GSK-2118436			
Encorafenib (Braftovi)	2nd-gen BRAFi	(Not approved for use as single agent)	6/27/2018
Also: LGX818			
Combination Therapies			
Drug	Inhibitor types	Approved to treat	Date approved by the FDA
Dabrafenib + Trametinib (Tafinlar + Mekinist)	Dabrafenib: BRAFi Trametinib: MEK1/2i	Unresectable or metastatic melanoma with BRAF V600E/K	1/9/2014
Also: GSK-2118436 + GSK-1120212		NSCLC with BRAF V600E mutation and involvement of lymph node(s), following complete resection	
		Locally advanced or metastatic anaplastic thyroid cancer with BRAF V600E mutation	
		Unresectable or metastatic solid tumors with BRAF V600E mutation and no satisfactory alternative treatment options	
		Pediatric patients 1 year of age and older with low-grade	

glioma with a BRAF V600E mutation			
Vemurafenib + Cobimetinib (Zelboraf + Cotellic) Also: PLX4032 + GDC-0973	Vemurafenib: BRAFi Cobimetinib: MEK1/2i	Unresectable or metastatic melanoma with BRAF V600E/K	11/10/2015
Encorafenib + Binimetinib (Braftovi + Mektovi) Also: LGX818 + MEK162	Encorafenib: BRAFi Binimetinib: MEK1/2i	Unresectable or metastatic melanoma with BRAF V600E/K Metastatic NSCLC with BRAF V600E	6/27/2018
Encorafenib + Cetuximab (Braftovi + Erbitux) Also: LGX818 + Encorafenib	Encorafenib: BRAFi Cetuximab: monoclonal antibody, EGFR antagonist	Metastatic CRC with a BRAF V600E mutation (after prior therapy)	4/8/2020
Vemurafenib + Cobimetinib + Atezolizumab (Zelboraf + Cotellic + Tecentriq) Also: PLX4032 + GDC-0973 + RG7446	Vemurafenib: BRAFi Cobimetinib: MEK1/2i Atezolizumab: PD-L1 blocking antibody	Unresectable or metastatic melanoma with BRAF V600	7/30/2020

4.1. First generation BRAF inhibitors

First-generation RAF inhibitors were created prior to the discovery of the oncogenicity of BRAF mutations and were initially designed to act as ATP-competitive inhibitors of CRAF [121]. These first-generation drugs stabilize the αC-helix into the active-IN position [122]. Although several iterations of this class of drug have gone through preclinical trials, sorafenib is the only first-generation RAF inhibitor to have gained FDA approval [123]. This particular inhibitor also causes an inactive-OUT conformation of the DFG motif, meaning that it is classified as a type II inhibitor [124]. Sorafenib is not particularly effective against cells that exhibit BRAF V600E mutations; even so, its multikinase inhibition activity makes it effective enough to have been FDA-approved to treat advanced renal cell carcinoma (median months overall survival [MMOS] 17.8 for sorafenib branch vs. 15.1 for placebo, median months progression-free survival [MMPFS] 5.5 vs. 2.8), unresectable hepatocellular carcinoma (MMOS 10.7 vs. 7.9, MMPFS 5.5 vs. 2.8), and metastatic differentiated thyroid carcinoma (MMOS 42.8 vs. 39.4, MMPFS 10.8 vs. 5.8) [19,125,126]. Even with its nonspecific approach to treatment, this pioneer drug helped break ground on a whole new approach to the treatment of BRAF-mutated cancers.

4.2. Second generation BRAF inhibitors

Upon the discovery of BRAF mutations and their roles as oncogenes in 2002 by Davies et al., researchers began to develop more targeted therapies in the form of second-generation RAF inhibitors [125]. These drugs focus on inhibiting BRAF V600E, a mutation that is particularly

prevalent in melanoma, thyroid cancers, and colorectal cancers [26,125]. Specifically, they bind the inactive-OUT position of the α C helix [54]. Second-generation RAF inhibitors can also be effective against other Class I BRAF mutations, which include all V600 variants and make up over 90% of all BRAF mutations [36]. Class I BRAF mutations are highly kinase active, RAS-independent, and signal as monomers [127]. Vemurafenib became the first of these inhibitors to gain FDA approval in 2011, specifically as a monotherapy for the treatment of unresectable or metastatic melanomas with V600E mutations [29]. Currently, it is also used to treat BRAF V600-mutant cases of Erdheim-Chester disease [20]. Dabrafenib, which was FDA-approved as a treatment for unresectable or metastatic melanomas with V600E mutations in 2013, has a similar mechanism of action as vemurafenib and also inhibits the BRAF monomer [20]. These monotherapeutic approaches have been vital in the treatment of patients with Class I BRAF mutations. However, emerging resistance to these drugs—as well as the need for a broader range of treatment options—has led to the advent of several combination therapies [128].

4.3. Combination therapies

A key issue with BRAF inhibitor monotherapy is that it can sometimes lead to increased activation of MEK, which is downstream of BRAF in the MAPK pathway [100]. This can reduce the efficacy of treatments and even cause secondary skin cancers to occur in some patients [129]. The development of combination therapies pairing BRAF and MEK inhibitors is meant to bypass this effect and has led to more targeted treatments for a wider range of histologies. For example, dabrafenib in combination with the MEK1/2 inhibitor trametinib was first approved by the FDA in 2014 for the treatment of unresectable or metastatic melanomas with BRAF V600E and V600K mutations [130]. Currently, this combination is also approved to treat a variety of BRAF V600E-mutant cancers of different histologies including non-small cell lung cancer (NSCLC) with involvement of lymph node(s), locally advanced or metastatic anaplastic thyroid cancer, pediatric low-grade glioma, and any unresectable or metastatic solid tumors with no other satisfactory treatment options. Similarly, the combination of vemurafenib and the MEK1/2 inhibitor cobimetinib has been approved for the treatment of unresectable or metastatic melanomas with BRAF V600E and V600K mutations [21]. Despite not having been approved for monotherapeutic use, the second-generation BRAF inhibitor encorafenib is FDA-approved for the treatment of unresectable or metastatic melanomas with BRAF V600E and V600K mutations [131]. The combination of encorafenib and the anti-EGFR antibody cetuximab, for instance, was FDA-approved in 2020 to treat metastatic colorectal cancer with a BRAF V600E mutation after prior therapy. Combination therapies can also involve three drugs, such as in the case of vemurafenib, cobimetinib, and the PD-L1 blocking antibody atezolizumab; this trio was FDA-approved for treating unresectable or metastatic melanomas with any BRAF V600 mutation [22]. Dual BRAF and MEK inhibition is already known to improve immunity responses against tumors, so the addition of these immunotherapeutic drugs is meant to provide an additional synergistic effect for certain histologies [132]. The use of antibodies in the treatment of BRAF-mutated cancers is reflective of a growing interest in immunotherapy, both within the field of oncology and beyond it.

4.4. Third generation BRAF inhibitors

Despite having promising results in the treatment of Class I BRAF mutations, second-generation RAF inhibitors are not nearly as effective against Class II and III mutations [133]. This is due to a key structural difference in the expression of these mutations; while Class I BRAF functions as a monomer, Class II and III act as homo- or hetero- and heterodimers, respectively [134]. Second-generation BRAF inhibitors such as vemurafenib and dabrafenib bind to the BRAF kinase, stabilizing the α C-helix into its inactive OUT position. This is an effective approach to inhibiting the BRAF monomers associated with Class I BRAF mutations. However, in dimers, this same α C-OUT position causes steric strain between the inhibitor and kinase that leads to a negative allosteric effect in the second protomer [135]. As a result, the monomer-targeting second-generation inhibitors fall flat in the inhibition of dimerized non-V600 mutations. There are currently no FDA-approved BRAF

inhibitors that adequately inhibit Class II and III mutations, but several third-generation BRAF inhibitors are being developed and tested in clinical trials in order to meet this need. Third-generation BRAF inhibitors are characterized by their targeting of both monomers and dimers of the BRAF kinase [133]. One example of this broader approach to treatment is the use of α C-IN RAF inhibitors, such as CEP-32496 and RAF-265 [87,136]. These drugs help to reduce steric hindrance, inhibiting both RAF protomers and reducing the overall activity of dimeric RAF [137]. Unfortunately, this same α C-IN feature also induces RAF priming, which leads to a significant increase in RAF dimerization [135]. α C-OUT inhibitors do not catalyze RAF priming to the same degree, but in cases of heightened activation of the upstream RAS, they can cause RAF priming and dimerization to increase through an effect known as inhibitor-induced paradoxical activation [138]. Future third-generation BRAF inhibitors will have to target both monomeric and dimeric BRAF, as well as avoid paradoxical activation both upstream and downstream of BRAF.

5. Resistance mechanisms and overcoming challenges

5.1. BRAF and the MAPK signaling pathway

BRAF, whether wild-type or mutated, is an inextricable part of the larger MAPK signaling pathway. This means that any attempt at targeting it without disrupting other parts of the pathway can present a challenge. For example, one of the most common resistance mechanisms that tumors develop against BRAF inhibitors is the reactivation of various components of the MAPK pathway [139]. Monotherapeutic administration of BRAF inhibitors often leads to the reactivation of the MEK/ERK effector cascade, which is downstream of BRAF [140]. This is why BRAF/MEK inhibitor combination therapy has become prominent in research efforts and has led to significant improvements in patient response rates, progression-free survival, and median duration of response when compared to monotherapy [141]. As with monotherapy, administration of combination therapies is also subject to the emergence of various mechanisms of resistance. Most of these mechanisms are again MEK/ERK dependent, although MEK/ERK independent variations have also been identified [142].

5.2. MEK/ERK dependent resistance

MEK/ERK dependent forms of BRAF inhibitor resistance can be further subdivided based on whether they are adaptive or acquired. Adaptive resistance occurs without new mutations arising, and its prevalence varies by tumor histology due to the unique variations in receptor expression that these cancers already possess [143]. For example, EGFR is active in both colon cancer and melanoma, but the comparatively higher levels of EGFR expression in colon cancer cause it to develop drug resistance significantly faster than in melanoma. In a similar mechanism involving HER2 and HER3 ligands, V600E thyroid cancers are able to form resistance to vemurafenib more quickly than melanoma can [144,145]. These variations in resistance highlight the importance of taking tumor histology into account when considering treatment options for BRAF-mutated cancers.

In contrast to adaptive resistance, acquired MEK/ERK-dependent resistance arises when changes are made to actual molecular switches of the MAPK pathway [146]. These changes occur as a result of selective pressures caused by treatment with kinase inhibitors. Dependent resistance can arise through alterations directly at the BRAF level, or it can begin upstream or downstream of BRAF such as at RAS, MEK1/2, and ERK 1/2 [147,148]. Similar to adaptive resistance, acquired MEK/ERK-dependent resistance can vary based on tumor histology [149]; this mechanism is more commonly found in skin, colon, and thyroid cancers, though it has also been found to occur in pancreatic, lung, and ovarian tumors [146].

As part of acquired resistance to BRAF inhibitors, emerging alterations in the BRAF gene itself can also occur [150]. For example, amplification of BRAF has been observed in dabrafenib and trametinib combination therapy-resistant melanoma and colon cancer patients, with additional variations in BRAF splicing found in the melanoma patients [148,150-152]. Of the dabrafenib and trametinib resistant melanoma patients, one out of five were found to have a BRAF splicing alteration

associated with a lack of exons 2-10 [148]. A rarer alteration, found in 0.4% of this same subset of dabrafenib and trametinib combination therapy-resistant melanomas, possessed in-frame deletion mutations related to exons 2-8 that are associated with the RAS-binding domain [153]. Deleterious mutations resulting in BRAF activation were also found in 0.6-1% of lung, pancreatic, thyroid, and ovarian cancers [146,154]. These mutations lock the α C-helix into the active-in conformation by shortening the β 3/ α C-helix loop, thereby facilitating the formation of BRAF dimers [146,154]. Lastly, a β 3- α C deletion mutation has been found to diminish the binding abilities of BRAF inhibitors AZ628, dabrafenib, and vemurafenib by increasing the flexibility of the α C-helix [155]. These relatively novel findings show that not all acquired resistance to BRAF inhibitors appears upstream or downstream of BRAF, and that some alterations are capable of directly affecting the structure and function of BRAF itself.

5.3. MEK/ERK independent resistance

Although it is comparatively less common, MEK/ERK independent resistance can also arise due to treatment with BRAF inhibitors [142]. Preclinical research has helped expand our current understanding on this category of resistance, with mechanisms ranging from loss of phosphatase and tensin homolog (PTEN) to changes in metabolic processes [149,156]. However, not much is known about MEK/ERK independent resistance mechanisms in the context of BRAF/MEK inhibitor combination therapy, making this an important area for future research to target [144].

6. Clinical implications and patient outcomes

Understanding the various pathways and molecular mechanisms that affect BRAF is crucial for understanding diseases associated with its mutations. However, it is the clinical trials in which these concepts are put into practice that actually dictate their translational value in patient care. Clinical trials on BRAF-mutated cancers and associated conditions have been the basis on which FDA approvals, new standards of treatment, and novel approaches to research have been made. A table summarizing clinical trials to date concerning BRAF mutations, further subdivided by phase classification is presented in **Supplementary Table 1**.

Another crucial part of patient care is understanding how well treatments can be tolerated, as this factor often dictates whether or not a person can continue medication. Often, adverse effects are dictated by how many off-target effects a drug causes, so more targeted treatments are typically associated with lower toxicity profiles than their more general counterparts [157,158]. BRAF and MEK inhibitors target their respective kinases, but their close association with the larger MAPK signaling pathway means that these treatments can impact several outcomes of cellular signaling, including cell proliferation, differentiation, and even apoptosis [159-161]. These effects can manifest as a variety of different symptoms in varying intensities depending on histology, dosage, mutations, and the specific drug being used [162]. For example, as mentioned previously in this review, the monotherapeutic use of BRAF inhibitors can sometimes lead to re-activation of the MAPK pathway that can lead to paradoxical oncogenesis [163]. This is the reasoning behind the development of combination therapies, but these too come with their own set of symptoms [59]. Clinical trials often report the tolerability of treatments by grading adverse effects. However, due to variations in several aspects of clinical trials such as patient populations and types of reporting, these metrics are not meant to be directly compared across studies [164]. It is important that healthcare providers be able to consider how treatments will affect their patients, which is why many studies use meta-analyses of clinical trials in order to find a reliable way to compare treatments [165]. Below is a table summarizing the results of one such meta-analysis on the grades of adverse effects associated with BRAF mono- and combination therapies (**Table 2**).

Table 2. BRAF inhibitors adverse effects Overall incidence rate of grade 3 to grade 5 (G3-G5), and any grade, adverse effects of BRAFi and BRAFi + MEK1/2i treatments (0-50% in green, 51-75% in orange, 76-100% in red).

Drug(s)	G3-G5	Any grade
Vemurafenib (Zelboraf)	51%	94%
Dabrafenib (Tafinlar)	50%	85%
Encorafenib (Braftovi)	68%	99%
Dabrafenib + Trametinib (Tafinlar + Mekinist)	43%	95%
Vemurafenib + Cobimetinib (Zelboraf + Cotellic)	72%	98%
Encorafenib + Binimetinib (Braftovi + Mektovi)	68%	98%

6. Conclusions and future directions

BRAF hotspot mutations are oncogenic drivers in multiple cancer types. Current approaches include targeted therapies directed against RAF or MEK kinases as well as immunotherapy. While tumors harboring BRAF mutations are highly targetable through these approaches, the opportune therapeutic regimen depends on tumor histology as well as on the class of mutation present. In addition, resistance to therapy is nearly inevitable. Thus, future efforts must be directed towards the development of new strategies to overcome these limitations.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org.

Authors contributions: Conceptualization: P.R., N.V.B., E.C.; writing: P.R., N.V.B., V.F., E.C.; visualization: P.R., N.V.B., E.C.; supervision: E.C.; funding acquisition: E.C. All authors have read and agreed to the submitted version of the manuscript.

Funding: This research was funded by the Woman’s Cancer Association Madelon Ravlin Memorial Grant Award (E.C.), the SCCC Tumor Biology Programmatic Award (E.C.), and the Lung Cancer Research Foundation (LCRF) (E.C.).

Acknowledgments: The authors thank Fabrizio De Andre’, Franco Battiato, Francesco Guccini, Vinicio Capossela, Lucio Dalla, and Francesco De Gregori for the constant support. E.C. also thanks the LCRF for the 2022 William C. Rippe Award.

Conflicts of Interest: E.C. has acted as a consultant for ENTOS Inc., and has received research funding from ERASCA, InnoCare pharma, and Prelude. E.C. is also a committee member of the Clinical Biosafety Service.

References

1. Ihle, M.A.; Fassunke, J.; Konig, K.; Grunewald, I.; Schlaak, M.; Kreuzberg, N.; Tietze, L.; Schildhaus, H.U.; Buttner, R.; Merkelbach-Bruse, S. Comparison of high resolution melting analysis, pyrosequencing, next generation sequencing and immunohistochemistry to conventional Sanger sequencing for the detection of p.V600E and non-p.V600E BRAF mutations. *BMC Cancer* **2014**, *14*, 13, doi:10.1186/1471-2407-14-13.
2. Diamond, E.L.; Subbiah, V.; Lockhart, A.C.; Blay, J.Y.; Puzanov, I.; Chau, I.; Raje, N.S.; Wolf, J.; Erinjeri, J.P.; Torrisi, J.; et al. Vemurafenib for BRAF V600-Mutant Erdheim-Chester Disease and Langerhans Cell Histiocytosis: Analysis of Data From the Histology-Independent, Phase 2, Open-label VE-BASKET Study. *JAMA Oncol* **2018**, *4*, 384-388, doi:10.1001/jamaoncol.2017.5029.
3. Frisone, D.; Friedlaender, A.; Malapelle, U.; Banna, G.; Addeo, A. A BRAF new world. *Crit Rev Oncol Hematol* **2020**, *152*, 103008, doi:10.1016/j.critrevonc.2020.103008.
4. Hayward, N.K.; Wilmott, J.S.; Waddell, N.; Johansson, P.A.; Field, M.A.; Nones, K.; Patch, A.M.; Kakavand, H.; Alexandrov, L.B.; Burke, H.; et al. Whole-genome landscapes of major melanoma subtypes. *Nature* **2017**, *545*, 175-180, doi:10.1038/nature22071.
5. Sanz-Garcia, E.; Argiles, G.; Elez, E.; Tabernero, J. BRAF mutant colorectal cancer: prognosis, treatment, and new perspectives. *Ann Oncol* **2017**, *28*, 2648-2657, doi:10.1093/annonc/mdx401.
6. Matallanas, D.; Birtwistle, M.; Romano, D.; Zebisch, A.; Rauch, J.; von Kriegsheim, A.; Kolch, W. Raf family kinases: old dogs have learned new tricks. *Genes Cancer* **2011**, *2*, 232-260, doi:10.1177/1947601911407323.
7. Dent, P.; Haser, W.; Haystead, T.A.; Vincent, L.A.; Roberts, T.M.; Sturgill, T.W. Activation of mitogen-activated protein kinase kinase by v-Raf in NIH 3T3 cells and in vitro. *Science* **1992**, *257*, 1404-1407, doi:10.1126/science.1326789.
8. Kyriakis, J.M.; App, H.; Zhang, X.F.; Banerjee, P.; Brautigan, D.L.; Rapp, U.R.; Avruch, J. Raf-1 activates MAP kinase-kinase. *Nature* **1992**, *358*, 417-421, doi:10.1038/358417a0.
9. Moodie, S.A.; Willumsen, B.M.; Weber, M.J.; Wolfman, A. Complexes of Ras.GTP with Raf-1 and mitogen-activated protein kinase kinase. *Science* **1993**, *260*, 1658-1661, doi:10.1126/science.8503013.
10. Van Aelst, L.; Barr, M.; Marcus, S.; Polverino, A.; Wigler, M. Complex formation between RAS and RAF and other protein kinases. *Proc Natl Acad Sci U S A* **1993**, *90*, 6213-6217, doi:10.1073/pnas.90.13.6213.
11. Vojtek, A.B.; Hollenberg, S.M.; Cooper, J.A. Mammalian Ras interacts directly with the serine/threonine kinase Raf. *Cell* **1993**, *74*, 205-214, doi:10.1016/0092-8674(93)90307-c.
12. Warne, P.H.; Vician, P.R.; Downward, J. Direct interaction of Ras and the amino-terminal region of Raf-1 in vitro. *Nature* **1993**, *364*, 352-355, doi:10.1038/364352a0.
13. Zhang, W.; Liu, H.T. MAPK signal pathways in the regulation of cell proliferation in mammalian cells. *Cell Res* **2002**, *12*, 9-18, doi:10.1038/sj.cr.7290105.
14. Zhang, X.F.; Settleman, J.; Kyriakis, J.M.; Takeuchi-Suzuki, E.; Elledge, S.J.; Marshall, M.S.; Bruder, J.T.; Rapp, U.R.; Avruch, J. Normal and oncogenic p21ras proteins bind to the amino-terminal regulatory domain of c-Raf-1. *Nature* **1993**, *364*, 308-313, doi:10.1038/364308a0.
15. Poulikakos, P.I.; Sullivan, R.J.; Yaeger, R. Molecular Pathways and Mechanisms of BRAF in Cancer Therapy. *Clin Cancer Res* **2022**, *28*, 4618-4628, doi:10.1158/1078-0432.CCR-21-2138.
16. Tran, N.H.; Wu, X.; Frost, J.A. B-Raf and Raf-1 are regulated by distinct autoregulatory mechanisms. *J Biol Chem* **2005**, *280*, 16244-16253, doi:10.1074/jbc.M501185200.
17. Dhillon, A.S.; Meikle, S.; Yazici, Z.; Eulitz, M.; Kolch, W. Regulation of Raf-1 activation and signalling by dephosphorylation. *EMBO J* **2002**, *21*, 64-71, doi:10.1093/emboj/21.1.64.
18. Chong, H.; Lee, J.; Guan, K.L. Positive and negative regulation of Raf kinase activity and function by phosphorylation. *EMBO J* **2001**, *20*, 3716-3727, doi:10.1093/emboj/20.14.3716.
19. Tecentriq [package insert]. South San Francisco, CA: Genentech, Inc; 2023.
20. Zelboraf [package insert]. Basel, Switzerland: Hoffmann La Roche; 2017. .
21. Mekinist [package insert]. Basel, Switzerland: Novartis Pharmaceuticals; 2023.
22. Braftovi [package insert]. Boulder, CO: Array Biopharma Inc.; 2023.
23. Nexavar [package insert]. Berlin, Germany: Bayer Healthcare Pharmaceuticals LLC; 2023.
24. Tafenlar [package insert]. Basel, Switzerland: Novartis Pharmaceuticals; 2023.
25. Erbitux [package insert]. Bridgewater, New Jersey: ImClone Systems Incorporated; 2021.
26. Davies, H.; Bignell, G.R.; Cox, C.; Stephens, P.; Edkins, S.; Clegg, S.; Teague, J.; Woffendin, H.; Garnett, M.J.; Bottomley, W.; et al. Mutations of the BRAF gene in human cancer. *Nature* **2002**, *417*, 949-954, doi:10.1038/nature00766.

27. Rapp, U.R.; Goldsborough, M.D.; Mark, G.E.; Bonner, T.I.; Groffen, J.; Reynolds, F.H., Jr.; Stephenson, J.R. Structure and biological activity of v-raf, a unique oncogene transduced by a retrovirus. *Proc Natl Acad Sci U S A* **1983**, *80*, 4218-4222, doi:10.1073/pnas.80.14.4218.
28. Yao, H.; Sun, Q.; Zhu, J. Identification and Characterization of Small-Molecule Inhibitors to Selectively Target the DFG-in over the DFG-out Conformation of the B-Raf Kinase V600E Mutant in Colorectal Cancer. *Arch Pharm (Weinheim)* **2016**, *349*, 808-815, doi:10.1002/ardp.201600184.
29. Yao, Z.; Yaeger, R.; Rodrik-Outmezguine, V.S.; Tao, A.; Torres, N.M.; Chang, M.T.; Drosten, M.; Zhao, H.; Cecchi, F.; Hembrough, T.; et al. Tumours with class 3 BRAF mutants are sensitive to the inhibition of activated RAS. *Nature* **2017**, *548*, 234-238, doi:10.1038/nature23291.
30. Cheng, L.; Lopez-Beltran, A.; Massari, F.; MacLennan, G.T.; Montironi, R. Molecular testing for BRAF mutations to inform melanoma treatment decisions: a move toward precision medicine. *Mod Pathol* **2018**, *31*, 24-38, doi:10.1038/modpathol.2017.104.
31. Dhomen, N.; Marais, R. New insight into BRAF mutations in cancer. *Curr Opin Genet Dev* **2007**, *17*, 31-39, doi:10.1016/j.gde.2006.12.005.
32. Marchetti, A.; Felicioni, L.; Malatesta, S.; Grazia Sciarrotta, M.; Guetti, L.; Chella, A.; Viola, P.; Pullara, C.; Mucilli, F.; Buttitta, F. Clinical features and outcome of patients with non-small-cell lung cancer harboring BRAF mutations. *J Clin Oncol* **2011**, *29*, 3574-3579, doi:10.1200/JCO.2011.35.9638.
33. Nikiforov, Y.E.; Nikiforova, M.N. Molecular genetics and diagnosis of thyroid cancer. *Nat Rev Endocrinol* **2011**, *7*, 569-580, doi:10.1038/nrendo.2011.142.
34. Pisapia, P.; Pepe, F.; Malapelle, U.; Troncone, G. BRAF Mutations in Lung Cancer. *Acta Cytol* **2019**, *63*, 247-250, doi:10.1159/000496478.
35. Schirripa, M.; Biason, P.; Lonardi, S.; Pella, N.; Pino, M.S.; Urbano, F.; Antoniotti, C.; Cremolini, C.; Corallo, S.; Pietrantonio, F.; et al. Class 1, 2, and 3 BRAF-Mutated Metastatic Colorectal Cancer: A Detailed Clinical, Pathologic, and Molecular Characterization. *Clin Cancer Res* **2019**, *25*, 3954-3961, doi:10.1158/1078-0432.CCR-19-0311.
36. Singh, A.K.; Sonawane, P.; Kumar, A.; Singh, H.; Naumovich, V.; Pathak, P.; Grishina, M.; Khalilullah, H.; Jaremko, M.; Emwas, A.H.; et al. Challenges and Opportunities in the Crusade of BRAF Inhibitors: From 2002 to 2022. *ACS Omega* **2023**, *8*, 27819-27844, doi:10.1021/acsomega.3c00332.
37. Cheng, D.T.; Mitchell, T.N.; Zehir, A.; Shah, R.H.; Benayed, R.; Syed, A.; Chandramohan, R.; Liu, Z.Y.; Won, H.H.; Scott, S.N.; et al. Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A Hybridization Capture-Based Next-Generation Sequencing Clinical Assay for Solid Tumor Molecular Oncology. *J Mol Diagn* **2015**, *17*, 251-264, doi:10.1016/j.jmoldx.2014.12.006.
38. Bracht, J.W.P.; Karachaliou, N.; Bivona, T.; Lanman, R.B.; Faull, I.; Nagy, R.J.; Drozdowskyj, A.; Berenguer, J.; Fernandez-Bruno, M.; Molina-Vila, M.A.; et al. BRAF Mutations Classes I, II, and III in NSCLC Patients Included in the SLLIP Trial: The Need for a New Pre-Clinical Treatment Rationale. *Cancers (Basel)* **2019**, *11*, doi:10.3390/cancers11091381.
39. Dankner, M.; Rose, A.A.N.; Rajkumar, S.; Siegel, P.M.; Watson, I.R. Classifying BRAF alterations in cancer: new rational therapeutic strategies for actionable mutations. *Oncogene* **2018**, *37*, 3183-3199, doi:10.1038/s41388-018-0171-x.
40. Krebs, M.G.; Malapelle, U.; Andre, F.; Paz-Ares, L.; Schuler, M.; Thomas, D.M.; Vainer, G.; Yoshino, T.; Rolfo, C. Practical Considerations for the Use of Circulating Tumor DNA in the Treatment of Patients With Cancer: A Narrative Review. *JAMA Oncol* **2022**, *8*, 1830-1839, doi:10.1001/jamaoncol.2022.4457.
41. Smiech, M.; Leszczynski, P.; Kono, H.; Wardell, C.; Taniguchi, H. Emerging BRAF Mutations in Cancer Progression and Their Possible Effects on Transcriptional Networks. *Genes (Basel)* **2020**, *11*, doi:10.3390/genes11111342.
42. Bradish, J.R.; Cheng, L. Molecular pathology of malignant melanoma: changing the clinical practice paradigm toward a personalized approach. *Hum Pathol* **2014**, *45*, 1315-1326, doi:10.1016/j.humpath.2014.04.001.
43. Ribas, A.; Flaherty, K.T. BRAF targeted therapy changes the treatment paradigm in melanoma. *Nat Rev Clin Oncol* **2011**, *8*, 426-433, doi:10.1038/nrclinonc.2011.69.
44. Long, G.V.; Menzies, A.M.; Nagrial, A.M.; Haydu, L.E.; Hamilton, A.L.; Mann, G.J.; Hughes, T.M.; Thompson, J.F.; Scolyer, R.A.; Kefford, R.F. Prognostic and clinicopathologic associations of oncogenic BRAF in metastatic melanoma. *J Clin Oncol* **2011**, *29*, 1239-1246, doi:10.1200/JCO.2010.32.4327.

45. Dummer, R.; Ascierto, P.A.; Gogas, H.J.; Arance, A.; Mandala, M.; Liskay, G.; Garbe, C.; Schadendorf, D.; Krajsova, I.; Gutzmer, R.; et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* **2018**, *19*, 603-615, doi:10.1016/S1470-2045(18)30142-6.
46. Larkin, J.; Ascierto, P.A.; Dreno, B.; Atkinson, V.; Liskay, G.; Maio, M.; Mandala, M.; Demidov, L.; Stryakovskiy, D.; Thomas, L.; et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med* **2014**, *371*, 1867-1876, doi:10.1056/NEJMoa1408868.
47. Popescu, A.; Haidar, A.; Anghel, R.M. Treating malignant melanoma when a rare BRAF V600M mutation is present: case report and literature review. *Rom J Intern Med* **2018**, *56*, 122-126, doi:10.1515/rjim-2017-0044.
48. Long, G.V.; Flaherty, K.T.; Stryakovskiy, D.; Gogas, H.; Levchenko, E.; de Braud, F.; Larkin, J.; Garbe, C.; Jouary, T.; Hauschild, A.; et al. Dabrafenib plus trametinib versus dabrafenib monotherapy in patients with metastatic BRAF V600E/K-mutant melanoma: long-term survival and safety analysis of a phase 3 study. *Ann Oncol* **2017**, *28*, 1631-1639, doi:10.1093/annonc/mdx176.
49. Wolchok, J.D.; Chiarion-Sileni, V.; Gonzalez, R.; Grob, J.J.; Rutkowski, P.; Lao, C.D.; Cowey, C.L.; Schadendorf, D.; Wagstaff, J.; Dummer, R.; et al. Long-Term Outcomes With Nivolumab Plus Ipilimumab or Nivolumab Alone Versus Ipilimumab in Patients With Advanced Melanoma. *J Clin Oncol* **2022**, *40*, 127-137, doi:10.1200/JCO.21.02229.
50. Adeniran, A.J.; Zhu, Z.; Gandhi, M.; Steward, D.L.; Fidler, J.P.; Giordano, T.J.; Biddinger, P.W.; Nikiforov, Y.E. Correlation between genetic alterations and microscopic features, clinical manifestations, and prognostic characteristics of thyroid papillary carcinomas. *Am J Surg Pathol* **2006**, *30*, 216-222, doi:10.1097/01.pas.0000176432.73455.1b.
51. Elisei, R.; Ugolini, C.; Viola, D.; Lupi, C.; Biagini, A.; Giannini, R.; Romei, C.; Miccoli, P.; Pinchera, A.; Basolo, F. BRAF(V600E) mutation and outcome of patients with papillary thyroid carcinoma: a 15-year median follow-up study. *J Clin Endocrinol Metab* **2008**, *93*, 3943-3949, doi:10.1210/jc.2008-0607.
52. Cohen, Y.; Rosenbaum, E.; Clark, D.P.; Zeiger, M.A.; Umbricht, C.B.; Tufano, R.P.; Sidransky, D.; Westra, W.H. Mutational analysis of BRAF in fine needle aspiration biopsies of the thyroid: a potential application for the preoperative assessment of thyroid nodules. *Clin Cancer Res* **2004**, *10*, 2761-2765, doi:10.1158/1078-0432.ccr-03-0273.
53. Nikiforova, M.N.; Kimura, E.T.; Gandhi, M.; Biddinger, P.W.; Knauf, J.A.; Basolo, F.; Zhu, Z.; Giannini, R.; Salvatore, G.; Fusco, A.; et al. BRAF mutations in thyroid tumors are restricted to papillary carcinomas and anaplastic or poorly differentiated carcinomas arising from papillary carcinomas. *J Clin Endocrinol Metab* **2003**, *88*, 5399-5404, doi:10.1210/jc.2003-030838.
54. Xing, M. BRAF mutation in thyroid cancer. *Endocr Relat Cancer* **2005**, *12*, 245-262, doi:10.1677/erc.1.0978.
55. Xing, M.; Vasko, V.; Tallini, G.; Larin, A.; Wu, G.; Udelsman, R.; Ringel, M.D.; Ladenson, P.W.; Sidransky, D. BRAF T1796A transversion mutation in various thyroid neoplasms. *J Clin Endocrinol Metab* **2004**, *89*, 1365-1368, doi:10.1210/jc.2003-031488.
56. Trovisco, V.; Soares, P.; Sobrinho-Simoes, M. B-Raf mutations in the etiopathogenesis, diagnosis, and prognosis of thyroid carcinomas. *Hum Pathol* **2006**, *37*, 781-786, doi:10.1016/j.humpath.2006.03.013.
57. Knauf, J.A.; Ma, X.; Smith, E.P.; Zhang, L.; Mitsutake, N.; Liao, X.H.; Refetoff, S.; Nikiforov, Y.E.; Fagin, J.A. Targeted expression of BRAFV600E in thyroid cells of transgenic mice results in papillary thyroid cancers that undergo dedifferentiation. *Cancer Res* **2005**, *65*, 4238-4245, doi:10.1158/0008-5472.CAN-05-0047.
58. Mesa, C., Jr.; Mirza, M.; Mitsutake, N.; Sartor, M.; Medvedovic, M.; Tomlinson, C.; Knauf, J.A.; Weber, G.F.; Fagin, J.A. Conditional activation of RET/PTC3 and BRAFV600E in thyroid cells is associated with gene expression profiles that predict a preferential role of BRAF in extracellular matrix remodeling. *Cancer Res* **2006**, *66*, 6521-6529, doi:10.1158/0008-5472.CAN-06-0739.
59. Bellevicine, C.; Migliatico, I.; Sgariglia, R.; Nacchio, M.; Vigliar, E.; Pisapia, P.; Iaccarino, A.; Bruzzese, D.; Fonderico, F.; Salvatore, D.; et al. Evaluation of BRAF, RAS, RET/PTC, and PAX8/PPARγ alterations in different Bethesda diagnostic categories: A multicentric prospective study on the validity of the 7-gene panel test in 1172 thyroid FNAs deriving from different hospitals in South Italy. *Cancer Cytopathol* **2020**, *128*, 107-118, doi:10.1002/cncy.22217.
60. Bellevicine, C.; Sgariglia, R.; Malapelle, U.; Vigliar, E.; Nacchio, M.; Ciancia, G.; Eszlinger, M.; Paschke, R.; Troncone, G. Young investigator challenge: Can the Ion AmpliSeq Cancer Hotspot Panel v2 be used for next-generation sequencing of thyroid FNA samples? *Cancer Cytopathol* **2016**, *124*, 776-784, doi:10.1002/cncy.21780.

61. Kim, T.H.; Park, Y.J.; Lim, J.A.; Ahn, H.Y.; Lee, E.K.; Lee, Y.J.; Kim, K.W.; Hahn, S.K.; Youn, Y.K.; Kim, K.H.; et al. The association of the BRAF(V600E) mutation with prognostic factors and poor clinical outcome in papillary thyroid cancer: a meta-analysis. *Cancer* **2012**, *118*, 1764-1773, doi:10.1002/cncr.26500.
62. Lee, J.H.; Lee, E.S.; Kim, Y.S. Clinicopathologic significance of BRAF V600E mutation in papillary carcinomas of the thyroid: a meta-analysis. *Cancer* **2007**, *110*, 38-46, doi:10.1002/cncr.22754.
63. Xing, M.; Westra, W.H.; Tufano, R.P.; Cohen, Y.; Rosenbaum, E.; Rhoden, K.J.; Carson, K.A.; Vasko, V.; Larin, A.; Tallini, G.; et al. BRAF mutation predicts a poorer clinical prognosis for papillary thyroid cancer. *J Clin Endocrinol Metab* **2005**, *90*, 6373-6379, doi:10.1210/jc.2005-0987.
64. Kebebew, E.; Weng, J.; Bauer, J.; Ranvier, G.; Clark, O.H.; Duh, Q.Y.; Shibus, D.; Bastian, B.; Griffin, A. The prevalence and prognostic value of BRAF mutation in thyroid cancer. *Ann Surg* **2007**, *246*, 466-470; discussion 470-461, doi:10.1097/SLA.0b013e318148563d.
65. Jo, Y.S.; Li, S.; Song, J.H.; Kwon, K.H.; Lee, J.C.; Rha, S.Y.; Lee, H.J.; Sul, J.Y.; Kweon, G.R.; Ro, H.K.; et al. Influence of the BRAF V600E mutation on expression of vascular endothelial growth factor in papillary thyroid cancer. *J Clin Endocrinol Metab* **2006**, *91*, 3667-3670, doi:10.1210/jc.2005-2836.
66. Kim, J.; Giuliano, A.E.; Turner, R.R.; Gaffney, R.E.; Umetani, N.; Kitago, M.; Elashoff, D.; Hoon, D.S. Lymphatic mapping establishes the role of BRAF gene mutation in papillary thyroid carcinoma. *Ann Surg* **2006**, *244*, 799-804, doi:10.1097/01.sla.0000224751.80858.13.
67. Kim, T.Y.; Kim, W.B.; Rhee, Y.S.; Song, J.Y.; Kim, J.M.; Gong, G.; Lee, S.; Kim, S.Y.; Kim, S.C.; Hong, S.J.; et al. The BRAF mutation is useful for prediction of clinical recurrence in low-risk patients with conventional papillary thyroid carcinoma. *Clin Endocrinol (Oxf)* **2006**, *65*, 364-368, doi:10.1111/j.1365-2265.2006.02605.x.
68. Namba, H.; Nakashima, M.; Hayashi, T.; Hayashida, N.; Maeda, S.; Rogounovitch, T.I.; Ohtsuru, A.; Saenko, V.A.; Kanematsu, T.; Yamashita, S. Clinical implication of hot spot BRAF mutation, V599E, in papillary thyroid cancers. *J Clin Endocrinol Metab* **2003**, *88*, 4393-4397, doi:10.1210/jc.2003-030305.
69. Trovisco, V.; Soares, P.; Preto, A.; de Castro, I.V.; Lima, J.; Castro, P.; Maximo, V.; Botelho, T.; Moreira, S.; Meireles, A.M.; et al. Type and prevalence of BRAF mutations are closely associated with papillary thyroid carcinoma histotype and patients' age but not with tumour aggressiveness. *Virchows Arch* **2005**, *446*, 589-595, doi:10.1007/s00428-005-1236-0.
70. Xu, X.; Quiros, R.M.; Gattuso, P.; Ain, K.B.; Prinz, R.A. High prevalence of BRAF gene mutation in papillary thyroid carcinomas and thyroid tumor cell lines. *Cancer Res* **2003**, *63*, 4561-4567.
71. Cancer Genome Atlas Research, N. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* **2014**, *511*, 543-550, doi:10.1038/nature13385.
72. Imielinski, M.; Berger, A.H.; Hammerman, P.S.; Hernandez, B.; Pugh, T.J.; Hodis, E.; Cho, J.; Suh, J.; Capelletti, M.; Sivachenko, A.; et al. Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* **2012**, *150*, 1107-1120, doi:10.1016/j.cell.2012.08.029.
73. Leonetti, A.; Facchinetti, F.; Rossi, G.; Minari, R.; Conti, A.; Friboulet, L.; Tiseo, M.; Planchard, D. BRAF in non-small cell lung cancer (NSCLC): Pickaxing another brick in the wall. *Cancer Treat Rev* **2018**, *66*, 82-94, doi:10.1016/j.ctrv.2018.04.006.
74. Harada, G.; Yang, S.R.; Cocco, E.; Drilon, A. Rare molecular subtypes of lung cancer. *Nat Rev Clin Oncol* **2023**, *20*, 229-249, doi:10.1038/s41571-023-00733-6.
75. Brustugun, O.T.; Khattak, A.M.; Tromborg, A.K.; Beigi, M.; Beiske, K.; Lund-Iversen, M.; Helland, A. BRAF-mutations in non-small cell lung cancer. *Lung Cancer* **2014**, *84*, 36-38, doi:10.1016/j.lungcan.2014.01.023.
76. George, J.; Lim, J.S.; Jang, S.J.; Cun, Y.; Ozretic, L.; Kong, G.; Leenders, F.; Lu, X.; Fernandez-Cuesta, L.; Bosco, G.; et al. Comprehensive genomic profiles of small cell lung cancer. *Nature* **2015**, *524*, 47-53, doi:10.1038/nature14664.
77. Schrock, A.B.; Li, S.D.; Frampton, G.M.; Suh, J.; Braun, E.; Mehra, R.; Buck, S.C.; Bufill, J.A.; Peled, N.; Karim, N.A.; et al. Pulmonary Sarcomatoid Carcinomas Commonly Harbor Either Potentially Targetable Genomic Alterations or High Tumor Mutational Burden as Observed by Comprehensive Genomic Profiling. *J Thorac Oncol* **2017**, *12*, 932-942, doi:10.1016/j.jtho.2017.03.005.
78. Mosele, F.; Remon, J.; Mateo, J.; Westphalen, C.B.; Barlesi, F.; Lolkema, M.P.; Normanno, N.; Scarpa, A.; Robson, M.; Meric-Bernstam, F.; et al. Recommendations for the use of next-generation sequencing (NGS) for patients with metastatic cancers: a report from the ESMO Precision Medicine Working Group. *Ann Oncol* **2020**, *31*, 1491-1505, doi:10.1016/j.annonc.2020.07.014.

79. Gautschi, O.; Pauli, C.; Strobel, K.; Hirschmann, A.; Printzen, G.; Aebi, S.; Diebold, J. A patient with BRAF V600E lung adenocarcinoma responding to vemurafenib. *J Thorac Oncol* **2012**, *7*, e23-24, doi:10.1097/JTO.0b013e3182629903.
80. Peters, S.; Michielin, O.; Zimmermann, S. Dramatic response induced by vemurafenib in a BRAF V600E-mutated lung adenocarcinoma. *J Clin Oncol* **2013**, *31*, e341-344, doi:10.1200/JCO.2012.47.6143.
81. Robinson, S.D.; O'Shaughnessy, J.A.; Cowey, C.L.; Konduri, K. BRAF V600E-mutated lung adenocarcinoma with metastases to the brain responding to treatment with vemurafenib. *Lung Cancer* **2014**, *85*, 326-330, doi:10.1016/j.lungcan.2014.05.009.
82. Schmid, S.; Siano, M.; Joerger, M.; Rodriguez, R.; Muller, J.; Fruh, M. Response to dabrafenib after progression on vemurafenib in a patient with advanced BRAF V600E-mutant bronchial adenocarcinoma. *Lung Cancer* **2015**, *87*, 85-87, doi:10.1016/j.lungcan.2014.11.008.
83. Mazieres, J.; Cropet, C.; Montane, L.; Barlesi, F.; Souquet, P.J.; Quantin, X.; Dubos-Arvis, C.; Otto, J.; Favier, L.; Avrillon, V.; et al. Vemurafenib in non-small-cell lung cancer patients with BRAF(V600) and BRAF(nonV600) mutations. *Ann Oncol* **2020**, *31*, 289-294, doi:10.1016/j.annonc.2019.10.022.
84. Malapelle, U.; Pisapia, P.; Sgariglia, R.; Vigliar, E.; Biglietto, M.; Carlomagno, C.; Giuffre, G.; Bellevicine, C.; Troncone, G. Less frequently mutated genes in colorectal cancer: evidences from next-generation sequencing of 653 routine cases. *J Clin Pathol* **2016**, *69*, 767-771, doi:10.1136/jclinpath-2015-203403.
85. Clarke, C.N.; Kopetz, E.S. BRAF mutant colorectal cancer as a distinct subset of colorectal cancer: clinical characteristics, clinical behavior, and response to targeted therapies. *J Gastrointest Oncol* **2015**, *6*, 660-667, doi:10.3978/j.issn.2078-6891.2015.077.
86. Gonsalves, W.I.; Mahoney, M.R.; Sargent, D.J.; Nelson, G.D.; Alberts, S.R.; Sinicrope, F.A.; Goldberg, R.M.; Limburg, P.J.; Thibodeau, S.N.; Grothey, A.; et al. Patient and tumor characteristics and BRAF and KRAS mutations in colon cancer, NCCTG/Alliance N0147. *J Natl Cancer Inst* **2014**, *106*, doi:10.1093/jnci/dju106.
87. Jones, J.C.; Renfro, L.A.; Al-Shamsi, H.O.; Schrock, A.B.; Rankin, A.; Zhang, B.Y.; Kasi, P.M.; Voss, J.S.; Leal, A.D.; Sun, J.; et al. (Non-V600) BRAF Mutations Define a Clinically Distinct Molecular Subtype of Metastatic Colorectal Cancer. *J Clin Oncol* **2017**, *35*, 2624-2630, doi:10.1200/JCO.2016.71.4394.
88. Cocco, E.; Benhamida, J.; Middha, S.; Zehir, A.; Mullaney, K.; Shia, J.; Yaeger, R.; Zhang, L.; Wong, D.; Villafania, L.; et al. Colorectal Carcinomas Containing Hypermethylated MLH1 Promoter and Wild-Type BRAF/KRAS Are Enriched for Targetable Kinase Fusions. *Cancer Res* **2019**, *79*, 1047-1053, doi:10.1158/0008-5472.CAN-18-3126.
89. Guinney, J.; Dienstmann, R.; Wang, X.; de Reynies, A.; Schlicker, A.; Soneson, C.; Marisa, L.; Roepman, P.; Nyamundanda, G.; Angelino, P.; et al. The consensus molecular subtypes of colorectal cancer. *Nat Med* **2015**, *21*, 1350-1356, doi:10.1038/nm.3967.
90. Missiaglia, E.; Jacobs, B.; D'Ario, G.; Di Narzo, A.F.; Soneson, C.; Budinska, E.; Popovici, V.; Vecchione, L.; Gerster, S.; Yan, P.; et al. Distal and proximal colon cancers differ in terms of molecular, pathological, and clinical features. *Ann Oncol* **2014**, *25*, 1995-2001, doi:10.1093/annonc/mdu275.
91. Farina-Sarasqueta, A.; van Lijnschoten, G.; Moerland, E.; Creemers, G.J.; Lemmens, V.; Rutten, H.J.T.; van den Brule, A.J.C. The BRAF V600E mutation is an independent prognostic factor for survival in stage II and stage III colon cancer patients. *Ann Oncol* **2010**, *21*, 2396-2402, doi:10.1093/annonc/mdq258.
92. Tabernero, J.; Ros, J.; Elez, E. The Evolving Treatment Landscape in BRAF-V600E-Mutated Metastatic Colorectal Cancer. *Am Soc Clin Oncol Educ Book* **2022**, *42*, 1-10, doi:10.1200/EDBK_349561.
93. Diamond, E.L.; Durham, B.H.; Ulaner, G.A.; Drill, E.; Buthorn, J.; Ki, M.; Bitner, L.; Cho, H.; Young, R.J.; Francis, J.H.; et al. Efficacy of MEK inhibition in patients with histiocytic neoplasms. *Nature* **2019**, *567*, 521-524, doi:10.1038/s41586-019-1012-y.
94. Horbinski, C.; Nikiforova, M.N.; Hagenkord, J.M.; Hamilton, R.L.; Pollack, I.F. Interplay among BRAF, p16, p53, and MIB1 in pediatric low-grade gliomas. *Neuro Oncol* **2012**, *14*, 777-789, doi:10.1093/neuonc/nos077.
95. Lassaletta, A.; Zapotocky, M.; Mistry, M.; Ramaswamy, V.; Honnorat, M.; Krishnatry, R.; Guerreiro Stucklin, A.; Zhukova, N.; Arnoldo, A.; Ryall, S.; et al. Therapeutic and Prognostic Implications of BRAF V600E in Pediatric Low-Grade Gliomas. *J Clin Oncol* **2017**, *35*, 2934-2941, doi:10.1200/JCO.2016.71.8726.
96. Behling, F.; Schittenhelm, J. Oncogenic BRAF Alterations and Their Role in Brain Tumors. *Cancers (Basel)* **2019**, *11*, doi:10.3390/cancers11060794.
97. Pisapia, P.; Pepe, F.; Iaccarino, A.; Sgariglia, R.; Nacchio, M.; Russo, G.; Gagnano, G.; Malapelle, U.; Troncone, G. BRAF: A Two-Faced Janus. *Cells* **2020**, *9*, doi:10.3390/cells9122549.

98. Martins-de-Barros, A.V.; Anjos, R.S.D.; Silva, C.C.G.; Silva, E.; Araujo, F.; Carvalho, M.V. Diagnostic accuracy of immunohistochemistry compared with molecular tests for detection of BRAF V600E mutation in ameloblastomas: Systematic review and meta-analysis. *J Oral Pathol Med* **2022**, *51*, 223-230, doi:10.1111/jop.13278.
99. Vanni, I.; Tanda, E.T.; Spagnolo, F.; Andreotti, V.; Bruno, W.; Ghiorzo, P. The Current State of Molecular Testing in the BRAF-Mutated Melanoma Landscape. *Front Mol Biosci* **2020**, *7*, 113, doi:10.3389/fmolb.2020.00113.
100. Gouda, M.A.; Ong, E.; Huang, H.J.; McPhaul, L.W.; Yoon, S.; Janku, F.; Gianoukakis, A.G. Ultrasensitive detection of BRAF V600E mutations in circulating tumor DNA of patients with metastatic thyroid cancer. *Endocrine* **2022**, *76*, 491-494, doi:10.1007/s12020-022-03004-z.
101. Gouda, M.A.; Polivka, J.; Huang, H.J.; Treskova, I.; Pivovarcikova, K.; Fikrle, T.; Woznica, V.; Dustin, D.J.; Call, S.G.; Meric-Bernstam, F.; et al. Ultrasensitive detection of BRAF mutations in circulating tumor DNA of non-metastatic melanoma. *ESMO Open* **2022**, *7*, 100357, doi:10.1016/j.esmoop.2021.100357.
102. Gouda, M.A.; Subbiah, V. Precision oncology for biliary tract tumors: it's written in blood! *Ann Oncol* **2022**, *33*, 1209-1211, doi:10.1016/j.annonc.2022.09.157.
103. Iyer, P.C.; Cote, G.J.; Hai, T.; Gule-Monroe, M.; Bui-Griffith, J.; Williams, M.D.; Hess, K.; Hofmann, M.C.; Dadu, R.; Zafereo, M.; et al. Circulating BRAF V600E Cell-Free DNA as a Biomarker in the Management of Anaplastic Thyroid Carcinoma. *JCO Precis Oncol* **2018**, *2*, doi:10.1200/PO.18.00173.
104. Passiglia, F.; Malapelle, U.; Normanno, N.; Pinto, C. Optimizing diagnosis and treatment of EGFR exon 20 insertions mutant NSCLC. *Cancer Treat Rev* **2022**, *109*, 102438, doi:10.1016/j.ctrv.2022.102438.
105. Drilon, A.; Cappuzzo, F.; Ou, S.I.; Camidge, D.R. Targeting MET in Lung Cancer: Will Expectations Finally Be MET? *J Thorac Oncol* **2017**, *12*, 15-26, doi:10.1016/j.jtho.2016.10.014.
106. Guo, R.; Luo, J.; Chang, J.; Rekhtman, N.; Arcila, M.; Drilon, A. MET-dependent solid tumours - molecular diagnosis and targeted therapy. *Nat Rev Clin Oncol* **2020**, *17*, 569-587, doi:10.1038/s41571-020-0377-z.
107. Benayed, R.; Offin, M.; Mullaney, K.; Sukhadia, P.; Rios, K.; Desmeules, P.; Ptashkin, R.; Won, H.; Chang, J.; Halpenny, D.; et al. High Yield of RNA Sequencing for Targetable Kinase Fusions in Lung Adenocarcinomas with No Mitogenic Driver Alteration Detected by DNA Sequencing and Low Tumor Mutation Burden. *Clin Cancer Res* **2019**, *25*, 4712-4722, doi:10.1158/1078-0432.CCR-19-0225.
108. Davies, K.D.; Le, A.T.; Sheren, J.; Nijmeh, H.; Gowan, K.; Jones, K.L.; Varella-Garcia, M.; Aisner, D.L.; Doebele, R.C. Comparison of Molecular Testing Modalities for Detection of ROS1 Rearrangements in a Cohort of Positive Patient Samples. *J Thorac Oncol* **2018**, *13*, 1474-1482, doi:10.1016/j.jtho.2018.05.041.
109. Yang, S.R.; Aypar, U.; Rosen, E.Y.; Mata, D.A.; Benayed, R.; Mullaney, K.; Jayakumaran, G.; Zhang, Y.; Frosina, D.; Drilon, A.; et al. A Performance Comparison of Commonly Used Assays to Detect RET Fusions. *Clin Cancer Res* **2021**, *27*, 1316-1328, doi:10.1158/1078-0432.CCR-20-3208.
110. Guo, R.; Offin, M.; Brannon, A.R.; Chang, J.; Chow, A.; Delasos, L.; Girshman, J.; Wilkins, O.; McCarthy, C.G.; Makhnin, A.; et al. MET Exon 14-altered Lung Cancers and MET Inhibitor Resistance. *Clin Cancer Res* **2021**, *27*, 799-806, doi:10.1158/1078-0432.CCR-20-2861.
111. Cohen, D.; Hondelink, L.M.; Solleveld-Westerink, N.; Uljee, S.M.; Ruano, D.; Cleton-Jansen, A.M.; von der Thusen, J.H.; Ramai, S.R.S.; Postmus, P.E.; Graadt van Roggen, J.F.; et al. Optimizing Mutation and Fusion Detection in NSCLC by Sequential DNA and RNA Sequencing. *J Thorac Oncol* **2020**, *15*, 1000-1014, doi:10.1016/j.jtho.2020.01.019.
112. Desmeules, P.; Boudreau, D.K.; Bastien, N.; Boulanger, M.C.; Bosse, Y.; Joubert, P.; Couture, C. Performance of an RNA-Based Next-Generation Sequencing Assay for Combined Detection of Clinically Actionable Fusions and Hotspot Mutations in NSCLC. *JTO Clin Res Rep* **2022**, *3*, 100276, doi:10.1016/j.jtocrr.2022.100276.
113. Ross, J.S.; Wang, K.; Chmielecki, J.; Gay, L.; Johnson, A.; Chudnovsky, J.; Yelensky, R.; Lipson, D.; Ali, S.M.; Elvin, J.A.; et al. The distribution of BRAF gene fusions in solid tumors and response to targeted therapy. *Int J Cancer* **2016**, *138*, 881-890, doi:10.1002/ijc.29825.
114. Capper, D.; Preusser, M.; Habel, A.; Sahm, F.; Ackermann, U.; Schindler, G.; Pusch, S.; Mechttersheimer, G.; Zentgraf, H.; von Deimling, A. Assessment of BRAF V600E mutation status by immunohistochemistry with a mutation-specific monoclonal antibody. *Acta Neuropathol* **2011**, *122*, 11-19, doi:10.1007/s00401-011-0841-z.
115. Al-Kateb, H.; Nguyen, T.T.; Steger-May, K.; Pfeifer, J.D. Identification of major factors associated with failed clinical molecular oncology testing performed by next generation sequencing (NGS). *Mol Oncol* **2015**, *9*, 1737-1743, doi:10.1016/j.molonc.2015.05.004.

116. Leighl, N.B.; Page, R.D.; Raymond, V.M.; Daniel, D.B.; Divers, S.G.; Reckamp, K.L.; Villalona-Calero, M.A.; Dix, D.; Odegaard, J.I.; Lanman, R.B.; et al. Clinical Utility of Comprehensive Cell-free DNA Analysis to Identify Genomic Biomarkers in Patients with Newly Diagnosed Metastatic Non-small Cell Lung Cancer. *Clin Cancer Res* **2019**, *25*, 4691-4700, doi:10.1158/1078-0432.CCR-19-0624.
117. Rose Brannon, A.; Jayakumaran, G.; Diosdado, M.; Patel, J.; Razumova, A.; Hu, Y.; Meng, F.; Haque, M.; Sadowska, J.; Murphy, B.J.; et al. Enhanced specificity of clinical high-sensitivity tumor mutation profiling in cell-free DNA via paired normal sequencing using MSK-ACCESS. *Nat Commun* **2021**, *12*, 3770, doi:10.1038/s41467-021-24109-5.
118. Lavoie, H.; Therrien, M. Regulation of RAF protein kinases in ERK signalling. *Nat Rev Mol Cell Biol* **2015**, *16*, 281-298, doi:10.1038/nrm3979.
119. Kornev, A.P.; Taylor, S.S. Dynamics-Driven Allostery in Protein Kinases. *Trends Biochem Sci* **2015**, *40*, 628-647, doi:10.1016/j.tibs.2015.09.002.
120. Karoulia, Z.; Gavathiotis, E.; Poulikakos, P.I. New perspectives for targeting RAF kinase in human cancer. *Nat Rev Cancer* **2017**, *17*, 676-691, doi:10.1038/nrc.2017.79.
121. Hall-Jackson, C.A.; Evers, P.A.; Cohen, P.; Goedert, M.; Boyle, F.T.; Hewitt, N.; Plant, H.; Hedge, P. Paradoxical activation of Raf by a novel Raf inhibitor. *Chem Biol* **1999**, *6*, 559-568, doi:10.1016/s1074-5521(99)80088-x.
122. Wu, P.; Nielsen, T.E.; Clausen, M.H. FDA-approved small-molecule kinase inhibitors. *Trends Pharmacol Sci* **2015**, *36*, 422-439, doi:10.1016/j.tips.2015.04.005.
123. Adnane, L.; Trail, P.A.; Taylor, I.; Wilhelm, S.M. Sorafenib (BAY 43-9006, Nexavar), a dual-action inhibitor that targets RAF/MEK/ERK pathway in tumor cells and tyrosine kinases VEGFR/PDGFR in tumor vasculature. *Methods Enzymol* **2006**, *407*, 597-612, doi:10.1016/S0076-6879(05)07047-3.
124. Vijayan, R.S.; He, P.; Modi, V.; Duong-Ly, K.C.; Ma, H.; Peterson, J.R.; Dunbrack, R.L., Jr.; Levy, R.M. Conformational analysis of the DFG-out kinase motif and biochemical profiling of structurally validated type II inhibitors. *J Med Chem* **2015**, *58*, 466-479, doi:10.1021/jm501603h.
125. Bayer; Amgen. Study of BAY43-9006 in Patients With Unresectable and/or Metastatic Renal Cell Cancer. **2006**.
126. Tabernero, J.; Garcia-Carbonero, R.; Cassidy, J.; Sobrero, A.; Van Cutsem, E.; Kohne, C.H.; Tejpar, S.; Gladkov, O.; Davidenko, I.; Salazar, R.; et al. Sorafenib in combination with oxaliplatin, leucovorin, and fluorouracil (modified FOLFOX6) as first-line treatment of metastatic colorectal cancer: the RESPECT trial. *Clin Cancer Res* **2013**, *19*, 2541-2550, doi:10.1158/1078-0432.CCR-13-0107.
127. Flaherty, K.T.; Puzanov, I.; Kim, K.B.; Ribas, A.; McArthur, G.A.; Sosman, J.A.; O'Dwyer, P.J.; Lee, R.J.; Grippo, J.F.; Nolop, K.; et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med* **2010**, *363*, 809-819, doi:10.1056/NEJMoa1002011.
128. Hauschild, A.; Grob, J.J.; Demidov, L.V.; Jouary, T.; Gutzmer, R.; Millward, M.; Rutkowski, P.; Blank, C.U.; Miller, W.H., Jr.; Kaempgen, E.; et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. *Lancet* **2012**, *380*, 358-365, doi:10.1016/S0140-6736(12)60868-X.
129. Paraiso, K.H.; Fedorenko, I.V.; Cantini, L.P.; Munko, A.C.; Hall, M.; Sondak, V.K.; Messina, J.L.; Flaherty, K.T.; Smalley, K.S. Recovery of phospho-ERK activity allows melanoma cells to escape from BRAF inhibitor therapy. *Br J Cancer* **2010**, *102*, 1724-1730, doi:10.1038/sj.bjc.6605714.
130. Boussemart, L.; Routier, E.; Mateus, C.; Opletalova, K.; Seville, G.; Kamsu-Kom, N.; Thomas, M.; Vagner, S.; Favre, M.; Tomasic, G.; et al. Prospective study of cutaneous side-effects associated with the BRAF inhibitor vemurafenib: a study of 42 patients. *Ann Oncol* **2013**, *24*, 1691-1697, doi:10.1093/annonc/mdt015.
131. Cotellic [package insert]. South San Francisco, CA: Genentech, Inc.; 2015.
132. Ribas, A.; Lawrence, D.; Atkinson, V.; Agarwal, S.; Miller, W.H., Jr.; Carlino, M.S.; Fisher, R.; Long, G.V.; Hodi, F.S.; Tsoi, J.; et al. Combined BRAF and MEK inhibition with PD-1 blockade immunotherapy in BRAF-mutant melanoma. *Nat Med* **2019**, *25*, 936-940, doi:10.1038/s41591-019-0476-5.
133. Lito, P.; Pratilas, C.A.; Joseph, E.W.; Tadi, M.; Halilovic, E.; Zubrowski, M.; Huang, A.; Wong, W.L.; Callahan, M.K.; Merghoub, T.; et al. Relief of profound feedback inhibition of mitogenic signaling by RAF inhibitors attenuates their activity in BRAFV600E melanomas. *Cancer Cell* **2012**, *22*, 668-682, doi:10.1016/j.ccr.2012.10.009.

134. Lin, Q.; Zhang, H.; Ding, H.; Qian, J.; Lizaso, A.; Lin, J.; Han-Zhang, H.; Xiang, J.; Li, Y.; Zhu, H. The association between BRAF mutation class and clinical features in BRAF-mutant Chinese non-small cell lung cancer patients. *J Transl Med* **2019**, *17*, 298, doi:10.1186/s12967-019-2036-7.
135. Karoulia, Z.; Wu, Y.; Ahmed, T.A.; Xin, Q.; Bollard, J.; Krepler, C.; Wu, X.; Zhang, C.; Bollag, G.; Herlyn, M.; et al. An Integrated Model of RAF Inhibitor Action Predicts Inhibitor Activity against Oncogenic BRAF Signaling. *Cancer Cell* **2016**, *30*, 485-498, doi:10.1016/j.ccell.2016.06.024.
136. Bertazza, L.; Barollo, S.; Radu, C.M.; Cavedon, E.; Simioni, P.; Faggian, D.; Plebani, M.; Pelizzo, M.R.; Rubin, B.; Boscaro, M.; et al. Synergistic antitumour activity of RAF265 and ZSTK474 on human TT medullary thyroid cancer cells. *J Cell Mol Med* **2015**, *19*, 2244-2252, doi:10.1111/jcmm.12612.
137. Peng, S.B.; Henry, J.R.; Kaufman, M.D.; Lu, W.P.; Smith, B.D.; Vogeti, S.; Rutkoski, T.J.; Wise, S.; Chun, L.; Zhang, Y.; et al. Inhibition of RAF Isoforms and Active Dimers by LY3009120 Leads to Anti-tumor Activities in RAS or BRAF Mutant Cancers. *Cancer Cell* **2015**, *28*, 384-398, doi:10.1016/j.ccell.2015.08.002.
138. Jin, T.; Lavoie, H.; Sahmi, M.; David, M.; Hilt, C.; Hammell, A.; Therrien, M. RAF inhibitors promote RAS-RAF interaction by allosterically disrupting RAF autoinhibition. *Nat Commun* **2017**, *8*, 1211, doi:10.1038/s41467-017-01274-0.
139. Pan, J.H.; Zhou, H.; Zhu, S.B.; Huang, J.L.; Zhao, X.X.; Ding, H.; Pan, Y.L. Development of small-molecule therapeutics and strategies for targeting RAF kinase in BRAF-mutant colorectal cancer. *Cancer Manag Res* **2018**, *10*, 2289-2301, doi:10.2147/CMAR.S170105.
140. Nazarian, R.; Shi, H.; Wang, Q.; Kong, X.; Koya, R.C.; Lee, H.; Chen, Z.; Lee, M.K.; Attar, N.; Sazegar, H.; et al. Melanomas acquire resistance to B-RAF(V600E) inhibition by RTK or N-RAS upregulation. *Nature* **2010**, *468*, 973-977, doi:10.1038/nature09626.
141. Poulikakos, P.I.; Zhang, C.; Bollag, G.; Shokat, K.M.; Rosen, N. RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF. *Nature* **2010**, *464*, 427-430, doi:10.1038/nature08902.
142. Flaherty, K.T.; Infante, J.R.; Daud, A.; Gonzalez, R.; Kefford, R.F.; Sosman, J.; Hamid, O.; Schuchter, L.; Cebon, J.; Ibrahim, N.; et al. Combined BRAF and MEK inhibition in melanoma with BRAF V600 mutations. *N Engl J Med* **2012**, *367*, 1694-1703, doi:10.1056/NEJMoa1210093.
143. Shi, H.; Hugo, W.; Kong, X.; Hong, A.; Koya, R.C.; Moriceau, G.; Chodon, T.; Guo, R.; Johnson, D.B.; Dahlman, K.B.; et al. Acquired resistance and clonal evolution in melanoma during BRAF inhibitor therapy. *Cancer Discov* **2014**, *4*, 80-93, doi:10.1158/2159-8290.CD-13-0642.
144. Montero-Conde, C.; Ruiz-Llorente, S.; Dominguez, J.M.; Knauf, J.A.; Viale, A.; Sherman, E.J.; Ryder, M.; Ghossein, R.A.; Rosen, N.; Fagin, J.A. Relief of feedback inhibition of HER3 transcription by RAF and MEK inhibitors attenuates their antitumor effects in BRAF-mutant thyroid carcinomas. *Cancer Discov* **2013**, *3*, 520-533, doi:10.1158/2159-8290.CD-12-0531.
145. Prahallad, A.; Sun, C.; Huang, S.; Di Nicolantonio, F.; Salazar, R.; Zecchin, D.; Beijersbergen, R.L.; Bardelli, A.; Bernards, R. Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. *Nature* **2012**, *483*, 100-103, doi:10.1038/nature10868.
146. Chen, W.; Park, J.I. Tumor Cell Resistance to the Inhibition of BRAF and MEK1/2. *Int J Mol Sci* **2023**, *24*, doi:10.3390/ijms241914837.
147. Heidorn, S.J.; Milagre, C.; Whittaker, S.; Nourry, A.; Niculescu-Duvas, I.; Dhomen, N.; Hussain, J.; Reis-Filho, J.S.; Springer, C.J.; Pritchard, C.; et al. Kinase-dead BRAF and oncogenic RAS cooperate to drive tumor progression through CRAF. *Cell* **2010**, *140*, 209-221, doi:10.1016/j.cell.2009.12.040.
148. Wagle, N.; Van Allen, E.M.; Treacy, D.J.; Frederick, D.T.; Cooper, Z.A.; Taylor-Weiner, A.; Rosenberg, M.; Goetz, E.M.; Sullivan, R.J.; Farlow, D.N.; et al. MAP kinase pathway alterations in BRAF-mutant melanoma patients with acquired resistance to combined RAF/MEK inhibition. *Cancer Discov* **2014**, *4*, 61-68, doi:10.1158/2159-8290.CD-13-0631.
149. Song, Y.; Bi, Z.; Liu, Y.; Qin, F.; Wei, Y.; Wei, X. Targeting RAS-RAF-MEK-ERK signaling pathway in human cancer: Current status in clinical trials. *Genes Dis* **2023**, *10*, 76-88, doi:10.1016/j.gendis.2022.05.006.
150. Long, G.V.; Fung, C.; Menzies, A.M.; Pupo, G.M.; Carlino, M.S.; Hyman, J.; Shahheydari, H.; Tembe, V.; Thompson, J.F.; Saw, R.P.; et al. Increased MAPK reactivation in early resistance to dabrafenib/trametinib combination therapy of BRAF-mutant metastatic melanoma. *Nat Commun* **2014**, *5*, 5694, doi:10.1038/ncomms6694.
151. Ahronian, L.G.; Sennott, E.M.; Van Allen, E.M.; Wagle, N.; Kwak, E.L.; Faris, J.E.; Godfrey, J.T.; Nishimura, K.; Lynch, K.D.; Mermel, C.H.; et al. Clinical Acquired Resistance to RAF Inhibitor Combinations in BRAF-

- Mutant Colorectal Cancer through MAPK Pathway Alterations. *Cancer Discov* **2015**, *5*, 358-367, doi:10.1158/2159-8290.CD-14-1518.
152. Oddo, D.; Sennott, E.M.; Barault, L.; Valtorta, E.; Arena, S.; Cassingena, A.; Filiciotto, G.; Marzolla, G.; Elez, E.; van Geel, R.M.; et al. Molecular Landscape of Acquired Resistance to Targeted Therapy Combinations in BRAF-Mutant Colorectal Cancer. *Cancer Res* **2016**, *76*, 4504-4515, doi:10.1158/0008-5472.CAN-16-0396.
 153. Johnson, D.B.; Childress, M.A.; Chalmers, Z.R.; Frampton, G.M.; Ali, S.M.; Rubinstein, S.M.; Fabrizio, D.; Ross, J.S.; Balasubramanian, S.; Miller, V.A.; et al. BRAF internal deletions and resistance to BRAF/MEK inhibitor therapy. *Pigment Cell Melanoma Res* **2018**, *31*, 432-436, doi:10.1111/pcmr.12674.
 154. Foster, S.A.; Whalen, D.M.; Ozen, A.; Wongchenko, M.J.; Yin, J.; Yen, I.; Schaefer, G.; Mayfield, J.D.; Chmielecki, J.; Stephens, P.J.; et al. Activation Mechanism of Oncogenic Deletion Mutations in BRAF, EGFR, and HER2. *Cancer Cell* **2016**, *29*, 477-493, doi:10.1016/j.ccell.2016.02.010.
 155. Niu, Y.; Zhang, Y.; Yao, X. Resistance mechanism of the oncogenic beta3-alphaC deletion mutation in BRAF kinase to dabrafenib and vemurafenib revealed by molecular dynamics simulations and binding free energy calculations. *Chem Biol Drug Des* **2019**, *93*, 177-187, doi:10.1111/cbdd.13399.
 156. Zuo, Q.; Liu, J.; Huang, L.; Qin, Y.; Hawley, T.; Seo, C.; Merlino, G.; Yu, Y. AXL/AKT axis mediated-resistance to BRAF inhibitor depends on PTEN status in melanoma. *Oncogene* **2018**, *37*, 3275-3289, doi:10.1038/s41388-018-0205-4.
 157. Chabner, B.A.; Roberts, T.G., Jr. Timeline: Chemotherapy and the war on cancer. *Nat Rev Cancer* **2005**, *5*, 65-72, doi:10.1038/nrc1529.
 158. Falzone, L.; Salomone, S.; Libra, M. Evolution of Cancer Pharmacological Treatments at the Turn of the Third Millennium. *Front Pharmacol* **2018**, *9*, 1300, doi:10.3389/fphar.2018.01300.
 159. Kolch, W. Meaningful relationships: the regulation of the Ras/Raf/MEK/ERK pathway by protein interactions. *Biochem J* **2000**, *351 Pt 2*, 289-305.
 160. Leonardi, G.C.; Falzone, L.; Salemi, R.; Zanghi, A.; Spandidos, D.A.; McCubrey, J.A.; Candido, S.; Libra, M. Cutaneous melanoma: From pathogenesis to therapy (Review). *Int J Oncol* **2018**, *52*, 1071-1080, doi:10.3892/ijo.2018.4287.
 161. Morrison, D.K. MAP kinase pathways. *Cold Spring Harb Perspect Biol* **2012**, *4*, doi:10.1101/cshperspect.a011254.
 162. Hoadley, K.A.; Yau, C.; Wolf, D.M.; Cherniack, A.D.; Tamborero, D.; Ng, S.; Leiserson, M.D.M.; Niu, B.; McLellan, M.D.; Uzunangelov, V.; et al. Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell* **2014**, *158*, 929-944, doi:10.1016/j.cell.2014.06.049.
 163. Gibney, G.T.; Messina, J.L.; Fedorenko, I.V.; Sondak, V.K.; Smalley, K.S. Paradoxical oncogenesis--the long-term effects of BRAF inhibition in melanoma. *Nat Rev Clin Oncol* **2013**, *10*, 390-399, doi:10.1038/nrclinonc.2013.83.
 164. NCC MERP, 2008. Statement on Medication Error Rates [WWW Document]. National Coordinating Council for Medication Error Reporting and Prevention. URL <https://www.nccmerp.org/statement-medication-error-rates>.
 165. Garutti, M.; Bergnach, M.; Polesel, J.; Palmero, L.; Pizzichetta, M.A.; Puglisi, F. BRAF and MEK Inhibitors and Their Toxicities: A Meta-Analysis. *Cancers (Basel)* **2022**, *15*, doi:10.3390/cancers15010141.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.