

– Supplementary Information –




Figure S1. Purification of human antibodies against the RBM region of the SARS-CoV-2 spike protein using a recombinant RBM Sepharose 4B affinity column (3 x 1 cm, inner diameter) (a) Evaluation of absorbance at 280 nm of the eluate different fractions, highlighted in green the eluate fractions used to concentrate polyclonal antibodies (b) In-house ELISA showing the specificity of purified antibodies to the multiepitope Dx-SARS-RBD containing RBM fragments.

Table S1: One-dose regime AstraZeneca-Oxford vaccinated serum information.

Sample	Gender	1 st Dose	Collected	Days after vaccination
M1	F	2021-06-18	2021-09-26	100
M3	F	2021-07-20	2021-09-26	70
M4	F	2021-08-07	2021-09-26	50
M5	M	2021-06-21	2021-09-26	97
M7	F	2021-09-09	2021-09-26	17
M9	M	2021-09-02	2021-09-26	24
M10	M	2021-09-15	2021-09-26	11
M11	M	2021-08-11	2021-09-26	46
M12	F	2021-06-26	2021-09-26	92
M13	M	2021-06-27	2021-09-26	91
M14	F	2021-06-28	2021-09-26	90
M17	M	2021-06-29	2021-09-26	89
M20	M	2021-06-30	2021-09-26	88
M21	M	2021-07-01	2021-09-26	87

M23	M	2021-07-28	2021-09-26	60
M24	F	2021-08-03	2021-09-26	54
M25	F	2021-07-04	2021-09-26	84
M27	F	2021-08-18	2021-09-26	39
M28	F	2021-08-20	2021-09-26	37
M29	F	2021-09-09	2021-09-26	17
M30	F	2021-07-08	2021-09-26	80
M31	F	2021-07-16	2021-09-26	72
M34	F	2021-08-17	2021-09-26	40
M37	F	2021-08-20	2021-09-26	37
M39	F	2021-09-10	2021-09-26	16
M40	F	2021-08-30	2021-09-26	27
M41	F	2021-08-13	2021-09-26	44
M43	F	2021-08-22	2021-09-26	35
M46	F	2021-08-24	2021-09-26	33
M47	F	2021-07-17	2021-09-26	71
M48	M	2021-09-12	2021-09-26	14

Table S2: Heterologous booster dose vaccinated serum information.

Sample	Birth data	Gender	Dose regimen	Last dose	Collected	Days after last vaccine
B1	1983-04-08	M	A/A/A/P	2022-07-15	2023-03-08	236
B2	1991-12-03	F	A/A/P/P	2022-06-22	2023-03-08	259
B3	1990-08-22	M	A/A/P/A	2022-10-17	2023-03-08	142
B4	1997-02-15	M	A/A/P/P	2022-11-16	2023-03-08	122
B5	1980-12-11	F	A/A/P/J	2022-06-09	2023-03-08	303
B6	1977-08-03	F	A/A/P	2022-01-10	2023-03-08	401
B7	1992-04-09	F	A/A/P/A	2022-06-20	2023-03-08	261

*A= Astrazeneca viral vector vaccine; P = Pfizer Spike mRNA vaccine; J = Janssen virus-based technology vaccine




Figure S2: Microscale thermophoresis (MST) traces of anti-RBM antibodies binding to different concentrations of S1WT (red) and S2WT (green) peptides by MicroScale Thermophoresis (MST). Relative fluorescence (RF) between the bound and unbound state was determined over a time of 21s with 20s MST-on time for evaluation. The blue bar indicates the Δ RF before the temperature gradient was applied, whereas the red bar shows the Δ RF during the thermophoresis. For interaction experiments, the amount of NT.647-labeled antibodies was kept constant, while the concentration of unlabeled peptides varied from 0.5 μ g/mL–0.12 ng/mL. The assay was performed in PBS containing 0.05% Tween 20 and after a short incubation period, the samples were analyzed in standard glass MST NT.115 capillaries.

IgG Subclasses Vaccinated Serum




Figure S3: IgG subclass neutralizing response against RBM peptides in vaccinated sera. Subclass Immunoglobulin subclasses reactivity of RBM peptides S1WT, S2WT, S8WT, S8BA1, and S8BA5 using a cohort of vaccinated individuals positives for IgG and with the first dose of Oxford-AstraZeneca (n=12) and booster doses. For analysis purposes were done multiple comparisons using Tukey's multiple comparisons test where a $p < 0.05$ was considered a significant difference ($* = p < 0.05$).