

Review

Not peer-reviewed version

Molecular Pathways of Vulnerability Carotid Plaques at Risk for Ischemic Stroke: A Narrative Review

[Giuseppe Miceli](#) , [Maria Grazia Basso](#) , Chiara Pintus , [Andrea Roberta Pennacchio](#) , [Elena Cacciola](#) ,
Mariagiovanna Cuffaro , Martina Profita , [Giuliana Rizzo](#) , [Antonino Tuttolomondo](#) *

Posted Date: 29 February 2024

doi: 10.20944/preprints202402.1730.v1

Keywords: carotid plaque; stroke; vulnerability; inflammation



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Review

Molecular Pathways of Vulnerability Carotid Plaques at Risk for Ischemic Stroke: A Narrative Review

Giuseppe MICELI ^{1,2,*}, Maria Grazia BASSO ^{1,2}, Chiara PINTUS ^{1,2}, Andrea Roberta PENNACCHIO ^{1,2}, Elena COCCIOLA ^{1,2}, Mariagiovanna CUFFARO ^{1,2}, Martina PROFITA ^{1,2}, Giuliana RIZZO ^{1,2} and Antonino TUTTOLOMONDO ^{1,2}

¹ Department of Health promotion, Mother and Child Care, Internal Medicine and Medical specialties (ProMISE) Università degli Studi di Palermo, Piazza delle Cliniche 2, 90127, Palermo, Italy

² Internal Medicine and Stroke Care Ward. University Hospital, Policlinico "P. Giaccone"

* Correspondence: miceli.gpp@gmail.com

Abstract: The concept of vulnerable carotid plaques is pivotal in understanding the pathophysiology of ischemic stroke secondary to large-artery atherosclerosis. At the macroscopic evaluation, vulnerable plaques are characterized by one or more of the following features: microcalcification, neovascularization, lipid-rich necrotic core (LRNC), intraplaque hemorrhage (IPH), thin fibrous caps, plaque surface ulceration, huge dimension determining stenosis, and plaque rupture. Recognizing these macroscopic characteristics is crucial to estimating the risk of cerebrovascular events, also in case of non-significant (less than 50%) stenosis. Inflammatory biomarkers, such as cytokines and adhesion molecules, lipid-related markers like oxidized low-density lipoprotein (LDL), and proteolytic enzymes capable of degrading extracellular matrix components are among the key molecules that will be scrutinized for their associative roles in plaque instability. Through their quantification and evaluation, these biomarkers reveal intricate molecular cross-talking governing plaque inflammation, rupture potential, and thrombogenicity. Current evidence demonstrates that plaque vulnerability phenotypes are multiple and heterogeneous and are associated with as many highly complex molecular pathways that determine the activation of an immune-mediated cascade that culminates in thromboinflammation. This narrative review provides a comprehensive analysis of current knowledge on molecular biomarkers expressed by symptomatic carotid plaques. It explores the association of these biomarkers with the structural and compositional attributes that characterize vulnerable plaques

Keywords: carotid plaque; stroke; vulnerability; inflammation

1. Introduction

Stroke represents one of the leading causes of death and disability worldwide, with carotid artery atherosclerosis being a primary risk factor for its development. The concept of vulnerable carotid plaques is pivotal in understanding the pathophysiology of ischemic stroke secondary to large-artery atherosclerosis (LAAS), one of the most frequent causes according to the trial of ORG 10172 in the acute stroke treatment (TOAST) classification system. Vulnerable plaques are prone to rupture, which can precipitate a cascade leading to thromboembolic episodes and ischemic strokes. Vulnerable plaques are responsible for 0.5-1% of ischemic strokes every year.

Moreover, symptomatic carotid plaques, the ones associated with clinical symptoms such as transient ischemic attacks, particularly underscore the imminent danger of such events. The biological complexity of carotid plaque vulnerability necessitates a multifaceted approach to its analysis. Traditionally, these plaques have been investigated through their anatomical and morphological characteristics. At the macroscopic evaluation, vulnerable plaques are characterized by one or more of the following features: microcalcification, neovascularization, lipid-rich necrotic core (LRNC), intraplaque hemorrhage (IPH), thin fibrous caps, plaque surface ulceration, huge dimension-determining stenosis, and plaque rupture. Recognizing these macroscopic characteristics

is crucial to estimating the risk of cerebrovascular events, also in the case of non-significant (less than 50%) stenosis [5–7].

Delineating these plaques' hallmarks is critical for developing strategies to preempt ischemic stroke and mitigate its hazardous consequences. . However, the advent of new technologies together with artificial intelligence, and the evolution of biochemical insights have shifted the focus towards understanding and identifying molecular patterns that denote plaque instability [9–11]. Inflammatory biomarkers, such as cytokines and adhesion molecules, lipid-related markers like oxidized low-density lipoprotein), and proteolytic enzymes capable of degrading extracellular matrix components are among the key molecules that will be scrutinized for their associative roles in plaque instability. Through their quantification and evaluation, these biomarkers reveal intricate molecular cross-talking governing plaque inflammation, rupture potential, and thrombogenicity . They offer a lens through which the nuances of plaque pathophysiology can be discerned, aiding in the stratification of stroke risk and the personalization of therapeutic interventions. This narrative review provides a comprehensive analysis of current knowledge on molecular biomarkers expressed by symptomatic carotid plaques. It explores the association of these biomarkers with the structural and compositional attributes that characterize vulnerable plaques, and their potential role as predictors of clinical outcomes providing insights into potential future directions in the therapeutic targeting of molecular biomarkers.

2. Vulnerable Plaque in the Proteomic Era

Recent advancements in proteomics have revolutionized our understanding of the molecular intricacies underlying vulnerable carotid plaques, offering unprecedented insights into the pathogenesis of atherosclerosis and its clinical implications . Carotid plaque vulnerability, characterized by the propensity for rupture and subsequent thromboembolic events, represents a critical determinant of stroke risk . Traditional diagnostic modalities, such as imaging techniques and biomarker assays, have provided valuable insights into plaque morphology and composition. However, the identification of specific molecular signatures associated with plaque vulnerability has remained a daunting challenge until the advent of proteomic technologies .

Proteomics, the large-scale study of proteins and their functions, has emerged as a powerful tool for dissecting the complex proteomic landscape of carotid plaques. High-throughput mass spectrometry-based proteomic approaches enable the simultaneous identification and quantification of thousands of proteins within a single sample, providing comprehensive insights into the molecular alterations underlying plaque instability .

Recent proteomic studies have revealed a diverse array of protein alterations associated with vulnerable carotid plaques, encompassing multiple biological processes implicated in plaque pathophysiology. One of the key findings is the dysregulation of proteins involved in extracellular matrix (ECM) remodeling, such as matrix metalloproteinases (MMPs) and their inhibitors, tissue inhibitors of metalloproteinases (TIMPs). MMPs are a family of proteolytic enzymes capable of degrading various components of the ECM, including collagen, elastin, and proteoglycans . Excessive MMP activity has been implicated in plaque destabilization by promoting fibrous cap thinning and collagen degradation, rendering the plaque more prone to rupture. Conversely, the upregulation of TIMPs serves as a counter-regulatory mechanism to inhibit MMP activity and maintain ECM integrity, thus contributing to plaque stabilization .


In addition to ECM remodeling, proteomic studies have identified alterations in proteins involved in inflammation, oxidative stress, lipid metabolism, and thrombosis within vulnerable carotid plaques. Inflammatory mediators, such as cytokines, chemokines, and adhesion molecules, play a crucial role in orchestrating the inflammatory response within the plaque microenvironment, contributing to endothelial dysfunction, leukocyte recruitment, and foam cell formation. Oxidative stress, characterized by an imbalance between reactive oxygen species (ROS) production and antioxidant defense mechanisms, promotes lipid peroxidation, endothelial dysfunction, and vascular inflammation, thereby exacerbating plaque instability. Dysregulation of lipid metabolism-related proteins, such as apolipoproteins and lipid transporters, further contributes to lipid accumulation

and foam cell formation within the plaque. Moreover, alterations in proteins involved in thrombosis, including coagulation factors, platelet receptors, and fibrinolytic enzymes, predispose vulnerable plaques to thrombotic events and subsequent ischemic stroke.

The identification of specific protein biomarkers associated with vulnerable carotid plaques holds immense clinical promise for risk stratification, diagnostic imaging, and targeted therapeutic interventions. Proteomic profiling of plaque specimens obtained via carotid endarterectomy or minimally invasive imaging-guided biopsy techniques allows for the identification of candidate biomarkers indicative of plaque vulnerability. These biomarkers can serve as diagnostic indicators for identifying patients at high risk of stroke and guide personalized treatment strategies aimed at stabilizing vulnerable plaques and preventing adverse cardiovascular events.

3. Pathophysiology of the Vulnerability of Carotid Plaques

The risk stratification for cerebrovascular events is typically based on the identification of symptomatic plaques in patients with carotid atheromas. However, asymptomatic plaques may be responsible for acute cerebrovascular events after their ulceration, rupture, thrombosis, and distal embolization as a consequence of the vulnerability of the lesions. These macroscopic modifications are the result of a complex molecular and endothelial interaction that involves the activation of inflammatory and thrombogenic pathways leading to plaque instability and a high risk of stroke. For this reason, researchers have recently pointed out the importance of discerning the molecular mechanisms accountable for forming vulnerable plaques. Histologically, vulnerable plaques are characterized by a high amount of lipids, macrophages, and T lymphocytes and a lower accumulation of vascular smooth muscle cells (VSMCs) and extracellular matrix compounds [13,14]. The main mechanisms involved in the vulnerability of the carotid plaques are inflammation, degradation of ECM, and lipid metabolism (Figure 1).



3.1. Role of Inflammation in Plaque Vulnerability

Stable and unstable plaques show different cellular compositions and protein expressions on their surface, suggesting the involvement of different inflammatory pathways in determining the instability of atherosclerotic plaques. Several immune cells participate in atherosclerotic disease, such

as macrophages, dendritic cells, monocytes, and T-cell lymphocytes. *Macrophages* are the primary cells involved in the development of atherosclerotic lesions. Different signals from the surrounding microenvironment can induce the differentiation of these cells into different phenotypes responsible for the vulnerability of the plaques. The main macrophage subtypes are M1, M2, M (Hb), Mhem, Mox, and M4. M1 macrophages have a proinflammatory phenotype, secreting some cytokines such as IL-1, IL-6 and tumor necrosis factor (TNF)- α and inducing tissue damage through ROS production. Moreover, M2 macrophages express on their surface some chemokine receptors (CXCL-9, CXCL-10, and CXCL-5), promoting the recruitment of circulating cells into the vascular layer. These mechanisms are responsible for the progression of the atherosclerotic plaques.

On the contrary, M2 macrophages inhibit inflammation. This role is explicated through the production of anti-inflammatory cytokines such as IL-10 and TNF-b and profibrotic proteins such as fibronectin, insulin-like growth factor, and transforming growth factor (TGF)-b, which are responsible for tissue repair.

M (Hb) and Mhem macrophages result from the accumulation of hemoglobin into resident macrophages after IPH. The expression of transporters responsible for cholesterol efflux (LXRa, LXRb, and ABC transporters ABCA1 and ABCG1) prevent the progression of atherosclerosis and stabilize the plaques [20,21].

The expression of antioxidant enzymes in the Mox macrophages, derived from the accumulation of oxidized phospholipids, provides an anti-inflammatory role to these cells, while a proinflammatory phenotype is attributed to M4 macrophages.

Thus, macrophage polarization into the M1 or M2 phenotype determines the vulnerability of the plaque. In the early stages of the atherosclerotic disease, M2 macrophages are the main cells in the lesions. However, with the progression of the disease, M1 cells become predominant, increasing the risk of rupture and cerebral complications [24,25]. Moreover, M1 macrophages are predominantly in the shoulder and the necrotic core of the plaque, while M2 macrophages infiltrate near newly formed blood vessels.

Macrophages can change their polarization depending on the stimuli from the surrounding environment. In his study, Khallou-Laschet demonstrated the induction of polarization from M1 to M2 phenotype from IL-4 and vice versa after lipopolysaccharide and IFN-g induction. The M1/M2 ratio is a mutable condition determining the vulnerability of the atherosclerotic plaques and the risk of acute cerebrovascular events.

Another mechanism through which macrophages induce plaque vulnerability is ferroptosis. Iron accumulation in atherosclerotic lesions is much higher than in healthy vascular walls. The iron overload in macrophages induces ferroptosis, a mechanism of cell death induced by lipid peroxidation leading to mitochondrial dysfunction and ROS production. Foam cells overexpress transferrin receptor 1, a cell surface receptor involved in transferrin-mediated iron uptake and ferritin synthesis, determining a higher iron accumulation into macrophages. Ferroptosis induces macrophage death and increases the dimension of the necrotic core, favoring plaque vulnerability.

Moreover, in the late stages of differentiation, macrophages release the protein YKL-40, which is responsible for different processes such as cell migration, angiogenesis, the regulation of hyaluronic acid synthesis, matrix metalloprotease (MMP)-9 activity, and tissue remodeling. Higher serum levels have recently been related to high plaque vulnerability.

Another molecule produced by macrophages and involved in plaque stability is osteopontin (OPN). OPN is produced by activated macrophages from several stimuli, such as cytokines, ILs, and TGF β . Once released, OPN promotes the recruitment of monocytes and the adhesion to endothelium, induces the expression of MMPs (in particular MMP-9), favoring the degradation of the ECM, and stimulates cytokine expression from macrophages. Moreover, elevated concentrations of OPN have been detected in ulcerated, hemorrhagic, necrotic, and inflamed plaques, proposing that this protein might be used as a biomarker of plaque instability. So, OPN has a crucial role in the progression of the atherosclerotic plaques.

One more signaling involved in the plaque vulnerability is derived from the binding CD40-CD40L. CD40 is expressed in endothelial cells, while CD40L is present in two forms (cell-membrane

bound and soluble) and is released from CD4⁺ T cells, platelets, monocytes, macrophages, B cells, and natural killer cells. After CD40-CD40L binding, endothelial cells express adhesion factors (selectin-E, vascular cell adhesion molecule, and intercellular adhesion molecule-1), recruit proinflammatory cells through the release of chemotactic molecules (IL-8, RANTES-regulated upon activation, normal T-cell expressed and secreted), release proinflammatory cytokines and procoagulant tissue factor (TF). Activated platelets expose on the surface CD40L and release the molecule's soluble form after the binding endothelium. Elevated concentrations of sCD40L have been found in the serum of patients with acute cerebrovascular ischemia compared with controls. In particular, higher levels have been found in the case of LAAS stroke but not in the case of carotid artery stenosis without cerebrovascular disease, underlying the pathogenic role of this pathway in determining the plaque progression and instability.

On the other hand, VSMCs are responsible for plaque stabilization. Inflammation-induced VSMCs activation leads to extracellular matrix and collagen secretion, reinforcing the fibrotic cap. At the same time, the loss of VSMCs is related to the enlargement of the necrotic core and plaque instability. VSMCs produce CTH-H₂S (cystathionine gamma-lyase-hydrogen sulfide), which has a protective role in atherosclerosis and whose levels are decreased in atheromas in comparison with the healthy vascular wall. VSMCs accumulating ox-LDL lack CTH-H₂S and undergo a mechanism of defective autophagy inducing cell death and more vulnerability of atherosclerotic plaques.

Considering the ability of the immune cells to produce proinflammatory molecules, atherosclerotic lesions present different protein expressions on their surface based on the plaque vulnerability. Some of these proteins can be detected in the serum and used as circulating biomarkers for the risk of cerebrovascular events.

Compared with stable plaques, vulnerable lesions show higher expression of CRP, high-sensitivity C-reactive protein (hs-CRP) and TNF- α [39,40] and, in particular, the lesions responsible for acute events express a higher concentration of IL-6, IL-17A, IL-18, IL-21, and IL-23 [41,42] and lower expression of INF- γ in comparison with asymptomatic plaques.

Other proinflammatory proteins detected in higher concentrations in the serum of symptomatic patients compared to asymptomatics are soluble urokinase-type plasminogen activator receptors (suPAR), which correlates with plaque inflammation, and S100A12, a Ca²⁺-binding protein belonging to the calgranulins family whose concentration depends on the INF- γ and IL-1 expression.

Low circulating levels of Fractalkine and increased serum concentration of platelet-derived growth factor B (PDGF-BB) are positively related to plaque calcification and the LNRC and represent circulating biomarkers of plaque instability. Fractalkine and PDGF-BB recruit macrophages on the vascular wall, induce intraplaque neoangiogenesis, and induce the switch of VSMCs into osteogenic phenotype, favoring plaque instability [46,47].

3.2. Extracellular Matrix Degradation

MMPs play a critical role in determining the instability of the plaques because of the capability to degrade the cellular matrix, the vascular wall, and the fibrotic caps, leading to a higher risk of rupture. Different MMP isoforms exist, but not all contribute similarly to atherosclerotic disease.

A higher expression of MMP-9 has been detected on the serum and the surface of unstable plaques. MMP-9 degrades the collagen V, promoting the damage of the fibrotic cap, exposing the subendothelial matrix, and attracting other proinflammatory circulating cells. As a confirmation of the role of MMP-9 in plaque vulnerability, Wang et al. have proved that inhibition of MMP-9 is related to the reduction of plaque instability.

Studies carried out on histopathological slices from carotid endarterectomy in patients with acute cerebrovascular events have shown increased concentrations of MMP-9, with statistically significant differences in case of microembolization or histologic characteristics of plaque instability, suggesting a pathogenetic role of this molecule in determining the vulnerability of the atherosclerotic lesions, with possible novel future therapeutic targets.

Moreover, vulnerable plaques express higher levels of MMP-1, MMP-2, MMP-7, MMP-8, MMP-12, and MMP-14 and lower concentrations of tissue inhibitor of metalloproteinases 3.

A Disintegrin and metalloproteinase with Thrombospondin motifs (ADAMTS)-4 is another molecule involved in many pathways, such as inflammation, angiogenesis, coagulation, and organ development. This protein is released from macrophages, arterial VSMCs, and endothelial cells, and its expression is induced by proinflammatory cytokines such as IL-1, IL-6, TNF- α and leptin [52,53].

ADAMTS4 has been found highly expressed in the shoulder, adjacent lipid core region, macrophage-rich regions, and fibrous caps of carotid plaques, where this protein can degrade versican, which is responsible for the plaque stability and whose levels are decreased in the vulnerable lesions [54,55].

3.3. Lipid Metabolism

The prevalence of lipidic compounds on the fibrotic component is one of the main determinants of the vulnerability of the plaques.

Lectin-like oxidized low-density lipoprotein receptor-1 is a scavenger receptor expressed on the arterial wall that can bind LDLs, inducing endothelial dysfunction and progression of atherosclerosis. The extracellular domain of L-1 is released into soluble form, so increased serum levels are a potential circulating biomarker of plaque instability.

Non-high-density lipoprotein cholesterol particles can activate inflammatory and endothelial cells and can contribute to complicating atheromas. Serum non-high-density lipoprotein cholesterol is considered an independent risk for plaque vulnerability [59,60].

One of the significant features of plaque vulnerability is the intraplaque presence of cholesterol crystals. Lesions containing cholesterol crystals have a proinflammatory phenotype because of the activation of the NOD-like receptor family pyrin domain containing 3 (NLRP3) inflammasome and complement system pathways, leading to the expression of inflammatory cytokines such as IL-1, IL-1 β , and TNF α . The local expression of these proteins induces the accumulation of macrophages and calcifications, responsible for more vulnerability [61,62].

Moreover, high levels of free fatty acids are related to a higher risk of cerebrovascular events due to the accumulation of micelles and fatty acid vesicles driving plaque instability.

4. Plaque Rupture and Atherothrombosis

As the atherosclerotic disease progresses, plaques are at greater risk of developing complications. The main complication of the atherosclerotic vulnerable lesions is rupture and, subsequently, atherothrombosis.

Plaque rupture depends not only on the morphologic characteristics of the lesion but also on extrinsic factors. Thus, local dynamic forces, in particular low shear stress, induce the activation of endothelial cells and leucocytes, upregulating proinflammatory processes, which provokes more vulnerability of the lesions. Moreover, circumference forces acting on the fibrotic cap shoulder, where there is the primary concentration of proinflammatory cells, contribute to determining plaque rupture.

Plaque rupture provokes the lack of endothelial integrity, which is the main determinant of vascular tone, the activation of inflammation, and the diffusion of molecules to the subendothelial layer. The integrity of the endothelium is maintained by the presence of intercellular junctions (occludin, claudin, junctional adhesion molecules, cadherin, and gap junctions) responsible for the interconnection of the endothelial cells, while the integrins link the endothelium to the cellular matrix and subendothelial layers. In integrity, endothelial cells express molecules with antiplatelet, anticoagulant, and fibrinolytic roles.

When the endothelium is damaged, platelets adhere to the vascular wall, inducing the formation of a platelet-rich thrombus through the binding of the platelet glycoprotein (GP) Iba receptor and the von Willebrand factor (vWF), expressed from the damaged endothelium. GPIba receptor also recruits circulating leucocytes binding integrins and P-selectin, perpetuating the inflammatory process.

[67,68]. Two platelet collagen receptors mediate platelet adhesion and activation [GPIa/IIa (integrin $\alpha 2\text{b}1$) and GPVI (immunoglobulin)] favoring the growth of the thrombus .

The exposition of subendothelial TF induces the activation of the coagulation system and the production of thrombin, which cleaves fibrinogen into fibrin, which stabilizes the platelets-rich thrombus . As a confirmation of the crucial role of the TF, it has been shown that inhibition of TF with local administration of TF pathway inhibitors blocks atherothrombosis . Moreover, VSMCs release TF-rich microparticles, increasing the expression of this molecule on the damaged surface .

Thrombin has a direct role in thrombus growth, inducing on platelets the activation of the receptors PAR1 and PAR4 responsible for several intracellular G protein-coupled signaling pathways (Gq, G12/13, Gi and G2) determining platelets degranulation, activation and aggregation .

Other mediators, such as thromboxane A2 and ADP, are involved in perpetuating platelet recruitment and activation, inducing the progression of the clot. Atherothrombosis is maintained from the activation of all these pathways inducing platelet degranulation and the local accumulation of several molecules, such as ATP, serotonin, Ca^{2+} , adhesion proteins (e.g. fibrinogen, fibronectin, vWF, thrombospondin, vitronectin, P-selectin and integrin $\alpha \text{IIb}\beta 3$) and coagulation factors (e.g. FV, FXI, plasminogen activator inhibitor type 1, plasminogen and protein S).

On this concern, inflammation is directly responsible for thrombus growth and different cytokines show a direct prothrombotic role, perpetuating the adhesion of platelets on the vascular wall.

One of the molecules involved in atherothrombosis is CRP: when circulating CRP adheres to the platelets surface, it is cleaved into monomeric form, which has a direct prothrombotic activity, favoring platelets adhesion .

Also, IL-1 induced-inflammation and activation of inflammasome are responsible for atherothrombosis, as demonstrated by the reduction of cardiovascular events using IL-1 antagonist molecules in the CANTOS study .

Recent studies have found a possible role of Neutrophil extracellular traps (NETs), fragments of chromatin surrounded by nuclear proteins released from neutrophils, in the link between inflammation and atherothrombosis .

Apart from known stimuli such as platelet activation, hyperlipidemia, and oxidized phospholipids, NLRP3 inflammasome activation in neutrophils has recently been found to be related to NETosis.

NETs formation is derived from histone citrullination induced by the enzyme peptidylarginine deiminase 4, which is a marker for NET formation . Interestingly, ulcerated and inflamed plaques are associated with a higher neutrophil-to-lymphocyte ratio, a systemic inflammation marker.

Moreover, elevated concentrations of peptidylarginine deiminase 4 have been associated with higher NLR, suggesting the possible role of NETs in the instability of atherosclerotic plaques .

A recent study showed that the deficiency of ABC transporters in the macrophages of knock-out mice was related to the inflammasome activation in macrophages promoting the accumulation of neutrophils and elevated NETs in plaques due to the accumulation of cholesterol into the cells [79–81], favoring the plaque rupture. In addition, IL-1 β intervenes in the crosstalk between macrophages and neutrophils: in fact, the IL-1 β induced-NLRP3 inflammasome activation in the macrophages increases NETosis, but this process is blocked in case of NLRP3 inhibition in neutrophils. So, both NLRP3 inhibition and IL-1 β inhibition block NETosis and reduce the accumulation of neutrophils on the atherosclerotic plaques, thus representing a potential therapeutic strategy for the future .

Plaque rupture represents the final act of a complex molecular mechanism that involves cross-talking between endothelium, immune-mediated response and plaque milieu which can manifest itself with different expressions of vulnerability depending on the patterns involved in this process (Table 1).

Table 1. Principle molecular patterns associated with specific vulnerability characteristics.

FEATURES OF PLAQUE VULNERABILITY	MOLECULAR PATTERNS ASSOCIATED WITH PLAQUE INSTABILITY	REFERENCES
----------------------------------	---	------------

Plaque Rupture And Atherothrombosis	GPIIb, vWF, TF, thrombin, CRP, IL-1, NETs	69, 71, 72, 73,74, 75, 77
Plaque Ulceration	MMP-9, CD40, CD40L, Heat shock protein 27, thrombospondin-1, ApoE, vWF, ADAMTS13	85, 86, 36, 88, 89, 90
Fibrous Cap Thickness	TF, IFN-γ, MMPs, reactive oxygen species, IL-18, growth differentiation factor-15, TNF, PDGF B, Macrophage-produced heparin-binding epidermal growth-factor-like growth factor, Cadherin, prostaglandins, ACE/Ang II, XCR1, CD177	128, 130, 131, 132, 133, 136, 137, 140, 144
Intraplaque Hemorrhage	VEGF, MMPs, bFGF, vWF, VE-cadherin, E-selectin, intercellular adhesion molecule-1, vascular cell adhesion molecule-1, Semaphorin 7A	149, 156, 159, 166, 170, 172, 173, 178, 179
Large Lipidic Core	CD36, ABCA, LP-PLA2, LPA, 7β-OH-cholesterol, C3d, C5b-9 neoantigen, ACAT	186-188, 191, 192, 202, 204, 209, 210
Microcalcification	TNF-α, IL-1β, IL-6, BMP2, IL-8, OPN, OPG, TGF-β	238-240, 244, 245, 246, 249, 257, 261-264, 272
Carotid Stenosis	Lipid hydroperoxide, 8-isoprostane, malondialdehyde, monocyte chemotactic protein-4, amyloid A, hs-CRP, TNF-α, prostaglandin E2, IFN-γ, TSG-6, vWF, CXCL16, VEGF, VEGFR2	4, 276, 277, 278, 281, 284

ABC: ATP binding cassette; ACAT: acylcoenzyme A:cholesterol acyltransferase; ACE/Ang II: angiotensin-converting enzyme/angiotensin II; ADAMTS: A Disintegrin and metalloproteinase with Thrombospondin motifs; ApoE: Apolipoprotein E; bFGF: Basic fibroblast growth factor; BMP2: bone morphogenetic protein 2; CXCL: Chemokine eC-X-C motif chemokine ligand; CRP: C-reactive protein; ECs: endothelial cells; ECM: extracellular matrix; FGF: fibroblast growth factor; HDL: high-density lipoprotein; hs-CRP: high-sensitivity C-reactive protein; IFN-γ: interferon-gamma; IL: Interleukin; IPH: intraplaque hemorrhage; LAAS: large-artery atherosclerosis; LDL: low-density lipoprotein; LNRC: lipid-rich necrotic core; Lp-PLA2: lipoprotein-associated phospholipase A2; LPA: Lysophosphatidic acid; MGP: matrix Gla protein; MES: microembolic signals; MMP: matrix metalloproteinase; NETs: Neutrophil extracellular traps; NLRP3: NOD-like receptor family pyrin domain containing 3; OPN: osteopontin; OPG: osteoprotegerin; OX-LDL: oxidized low-density lipoprotein; GP: platelet glycoprotein; PDGF: platelet-derived growth factor; PMPs: platelets-derived microparticles; RANKL: receptor activator of NF-κB ligand; RBCs: red blood cells; TCD: transcranial Doppler; TF: tissue factor; TGF: transforming growth factor; TOAST: trial of ORG 10172 in the acute stroke treatment; TSG-6: Tumor Necrosis Factor-Stimulated Gene-6; TNF: *Tumor necrosis factor*; VEGF: vascular endothelial growth factor; VSCM: vascular smooth muscle cell; vWF: von Willebrand factor.

5. Plaque Ulceration

Plaque vulnerability is commonly associated with carotid plaque ulceration, an essential indicator of previous plaque rupture and a strong predictor of future cerebrovascular events. Carotid plaque ulceration or surface irregularity is characterized by a fissure or erosion on the surface of a plaque. Using basic and advanced imaging modalities, the surface of a plaque can be defined as smooth, irregular, or ulcerated. Ulceration is reserved for cavities measuring at least 1-2 mm [83].

Ulceration is a significant factor in vulnerability and is linked to the development of neurologic symptoms on the pathological basis of arterial embolism of thrombotic material. Several studies demonstrated that ulcerated carotid plaques are more common in patients with anamnestic stroke or transient ischemic attack, and ulceration is more frequently found in patients with symptoms (ipsilateral or contralateral) and is also associated with the occurrence of new symptoms in asymptomatic patients .

A complex mechanism is responsible for plaque vulnerability characterized by erosion or ulceration. Inflammation accumulation, proteolytic enzyme release by macrophages, and local hemodynamic factors play a role in the pathogenesis of ulceration [85]. It has been shown that the promoters of the ulceration process are intraplaque inflammation and the activity of a series of MMPs, the primary physiological regulators of the extracellular matrix. The study conducted by Loftus et al. [50] about the expression of MMP-9 within this tissue demonstrated a significant increase in MMP-9 localized in the most unstable carotid plaques and the plaques of more symptomatic patients. Interestingly, MMP-9 has been considered an excellent candidate for future pharmacotherapy intended to stabilize carotid plaques and prevent stroke [86].

The complex mechanism of atherosclerotic plaque formation and rupture primarily involves the interaction of lipids and immune-regulated inflammatory pathways. It has been demonstrated that

the CD40–CD40L system could cause plaque instability. Wang et al. showed that CD40L expression on peripheral blood monocytes is associated with carotid atherosclerosis and plaque instability. Furthermore, CD40L expression in circulating monocytes was higher in patients with LAAS, followed by the carotid artery stenosis group [36].

The improvement of proteomics tools has allowed the unraveling of new circulating markers for atherosclerosis that can be targeted for early diagnostic traits, in preventing such life-threatening events, or as targets for new drug therapies [87]. Martin-Ventura et al. showed that heat shock protein 27 secretion into the culture medium was significantly lower in atherosclerotic plaques compared to control arteries, plasma level expression was lower in patients with carotid stenosis concerning healthy controls, and expression of this protein in the unstable human carotid plaque is lower than stable. Rocchiccioli et al. verified a significantly higher concentration of thrombospondin-1 and the vitamin D-binding protein in atherosclerotic subjects. Baart et al. assessed a mouse model for studying plaque ulceration. They showed that the APO E-/- mouse model overexpressed the urokinase-type plasminogen activator, suggesting that the loss of basement-membrane proteins may play a central role in advanced plaque rupture. Finally, some molecular markers such as vWF and ADAMTS13 are known to induce respectively platelet adhesion and aggregation and cleave VWF multimers into smaller sizes. They both have been associated with ischemic stroke but in a recent study VWF antigen level and ADAMTS13 activity seemed not to be related to vulnerable plaque features, such as the volume of IPH and LRNC, and plaque ulceration.

6. Microembolism

Microembolism from carotid plaque is one of the mechanisms responsible for cerebral ischemia. In 1990 Spencer proved that signals registered at the transcranial Doppler (TCD) monitoring in patients before and during endarterectomy procedure were assumed to be microemboli from carotid plaques.

Microembolic signals (MES) can be detected in also asymptomatic patients, even if the highest rate of MES is commonly registered in patients with acute cerebrovascular events or the case of carotid stenosis more than 70%.

On this concern, MES may be considered not only a marker of plaque instability but also an independent risk factor for the incidence and the recurrence of ischemic strokes [93,94]. Even if plaque instability is the primary determinant for microemboli, some specific biomarkers have been associated with a higher risk of distal embolization, supporting the presence of specific mechanisms responsible for this condition.

Commonly, microembolism has been attributed to vulnerable plaque fragmentation; thus, it has been recently proposed that microemboli may derive from platelets aggregation on the ulcerated lesions after rupture.

Recent studies support this hypothesis, showing different protein expressions on platelet surfaces in patients with and without MES. In particular, considering patients with MES on the TCD monitoring, a lower expression of P-selectin on platelets and higher serum concentration of this protein has been detected in the case of symptomatic or asymptomatic stenosis. On the other hand, a lower expression of thrombospondin has been found in patients with symptomatic stenosis and MES. Elevated platelet counts and neutrophil-platelet and lymphocyte-platelet complexes were found in the early and late symptomatic patients compared with asymptomatic ones, in particular if the degree of the stenosis is >70% [98–100] suggesting that platelet activation promotes plaque instability through several mechanisms such as enhanced cell adhesion, smooth muscle cell proliferation and the activation of the coagulation system.

This data confirms that platelet activation plays a crucial role in cerebral microembolism, as demonstrated by the efficacy of antiplatelet drugs in reducing the MES rate [102–104].

Recently, attention was turned to circulating platelets-derived microparticles (PMPs), small vesicles (diameter from 0.05-2 μm) containing phospholipidic compounds from cellular membranes released from platelets. These particles expose antigenic proteins on their surface, which allow specific identification in the serum, distinguishing them from other inflammatory cells. PMPs have

been related to plaque instability and ischemic cerebrovascular events, suggesting a possible role as a marker for cerebrovascular risk. Furthermore, PMP levels were higher in patients with acute ischemic strokes, while no significant differences were found between symptomatic carotid stenosis of more than 60% and healthy controls, suggesting that platelet microparticles may be correlated with microembolism. Supporting this hypothesis, a study carried out by Kandiyil et al. demonstrated that PMPs concentrations were higher in patients with MES on TCD and MRI-documented images of acute ischemic events, underlying the relationship between platelet activation and microembolism and the thrombogenic effect of PMPs in the sites of deposition. Interestingly, antiplatelet medications seem to not influence PMPs levels.

The neutrophil count is notably a marker of atherosclerosis severity and plaque instability. Recent studies have demonstrated that peripheral neutrophilia is associated with a higher risk of microembolism. A higher neutrophil count has been associated with a higher MES rate in patients who underwent carotid stenting. This association was confirmed in patients with recently symptomatic carotid stenosis without revascularization. On this view, neutrophil count can be considered a marker of microembolism, considering the persistent strength of the association even after adjustment for the National Institutes of Health Stroke Scale (NIHSS) and CRP values.

Furthermore, Chemokine eC-X-C motif chemokine ligand (CXCL)-16, a chemokine able to recruit and activate T cells (Th1 and Tc1 cells), has recently been associated with acute cerebrovascular disease. CXCL16 is a protein with several functions: it plays an essential role in inflammation, directing leukocyte migration, inducing the secretion of pro-inflammatory proteins, such as interferon-gamma (IFN- γ) and IL-2, after T cell activation, and acting as a scavenger for oxidized-LDL. It is known that CXCL16 is related to atherosclerotic disease severity: in fact, the surface expression on the carotid plaques and serum concentrations of mRNA are higher in patients with cerebrovascular events compared with asymptomatic plaque. Recent studies have found also a correlation between this protein and microembolization: in fact, higher serum concentrations of CXCL-16 have been detected in patients with MES on the TCD monitoring in LAAS, with a high sensibility and specificity (88.5% and 56.5%, respectively) for a cutoff value of 2.115 ng/ml. This data suggests a potential role of this chemokine in the pathogenesis of cerebral microembolism.

Moreover, higher plasma fibrinogen concentrations and decreased ratio of CD4⁺ CD25^{high} T reg have been found in MES+ compared with MES – patients and carotid atherosclerosis, even if more studies need to be carried out to confirm this data.

Finally, about the link between inflammation and microembolism, Pini et al. have demonstrated that values of Hs-CRP > 5 mg/L and serum amyloid A > 10 mg/L were associated with a higher MES rate in patients with post-carotid artery stenting asymptomatic cerebral ischemia detected by MRI.

7. Fibrous Cap Thickness

The term “vulnerable” plaque is frequently used to refer to a lesion with a fibrous cap less than 65 micrometers thick and infiltrated with macrophages (more than 25 cells per 0.3 mm diameter field). Several studies about plaque rupture suggest that thinner caps atheroma, fibrous caps <65 microns, together with the presence of numerous macrophages within the cap, likely indicate instability and predict plaque rupture [117,118].

As known, the progression of atherosclerotic lesions occurs through three main stages: an initial, asymptomatic, and non-stenotic stage, followed by an advanced phase where thrombi are formed and cause embolism. Infiltrated monocytes are recruited to the lesion area in endothelial cells when modified LDL expresses adhesion molecules and chemokines. When monocytes are differentiated into macrophages, they promote the secretion of connective tissue and the formation of the fibrous cap. If the anti-inflammatory response is not working correctly or is insufficient, atherosclerotic plaques that have been stable for a while may progress and become unstable. The extracellular matrix is hydrolyzed by MMPs and other proteolytic enzymes, causing a loss of thickness of the fibrous cap by necrotic macrophages. LDL and modified LDL could be significant targets for preventing the progression and complications of carotid atherosclerosis.

The fibrous cap serves as a sub-endothelial barrier, separating the vessel lumen from the atherosclerotic necrotic core. Comprised of VSMCs and ECM derived from VSMCs, its primary role is to provide structural support and prevent the exposure of prothrombotic material in the core, thereby averting thrombosis. In response to injury, VSMCs undergo a phenotypic switch to a synthetic state, marked by increased migratory and proliferative activities. Nearby cells initiate the healing process by producing various growth factors, including epidermal growth factor, fibroblast growth factor, insulin-like growth factor, PDGF, TGF- β , and VEGF. In atherosclerosis, stimulated by growth factors from foam cells or endothelial cells (ECs) in the intima, VSMCs from the tunica media migrate to the intima. Macrophage-produced interleukin (IL)-1 enhances PDGF production by VSMCs, leading to autocrine proliferation in the intima. Synthetic VSMCs also increase the production of extracellular matrix components such as interstitial collagen, elastin, and proteoglycans. These proliferating VSMCs, along with ECM production, contribute to the formation of a fibrous cap, crucial for preventing plaque rupture. If mitogen production persists, VSMCs may not revert to the contractile phenotype, facilitating lesion development.

In the fibrous cap, VSMCs and collagen are noted to be present [123.], while the necrotic core of the plaque exhibits minimal collagen but a significant amount of free cholesterol. Transcription factors SP1, SP3, and AP1, interacting with the collagen promoter, play a role in the transcription of type I collagen in these cells. The absence of collagen serves as an indication of VSMC loss, rendering the plaque vulnerable. Reduced metalloproteinase activity may contribute to impaired collagen deposition. Collagen is crucial for providing tensile strength to the fibrous cap, and its absence or minimal presence may result in thinning when VSMCs are depleted from the cap, contributing to a rupture with reduced collagen content. Overproduction of VSMCs can lead to plaque growth dependent on angiogenic factors. Type VIII collagen, typically present in small amounts in normal arteries and produced by macrophages and VSMCs, enhances the migration and growth of VSMCs through the extracellular matrix. Fibrous cap characteristics, including thickness, cellularity, matrix composition, and collagen content, play vital roles in plaque stability. The exact mechanism of plaque rupture involves cap thinning, elevated inflammatory cytokines, matrix-digesting proteases, decreased collagen synthesis, and the presence of injured or apoptotic cells within the necrotic core. Cells contributing to plaque formation are implicated in plaque rupture and subsequent thrombosis. Molecular mediators associated with atherogenesis may alter collagen metabolism, thinning or weakening the fibrous cap. Inflammatory cells secrete cytokines, growth factors, TF, IFN- γ , MMPs, and reactive oxygen species. Accumulation of free cholesterol induces apoptosis of macrophage-derived foam cells, contributing to the formation of the necrotic core. Excess extracellular unesterified cholesterol crystallizes, leading to complicated atheroma and eventual total occlusion of coronary artery branches. Continued inflammatory responses, driven by proinflammatory cytokines, destabilize atherosclerotic plaques, with IFN- γ , IL-18, growth differentiation factor-15, and TNF-like weak inducer of apoptosis promoting destabilization, while TGF- β causes stabilization. Proinflammatory cytokines such as IFN- γ , TNF- α , and IL-1 β stimulate macrophage and smooth muscle cell apoptosis, thinning the fibrous cap. Macrophages infiltrate the thinned fibrous cap, secreting inflammatory cytokines and MMPs, playing a pivotal role in weakening and ultimately rupturing the atherosclerotic plaque. Necrosis of the vulnerable plaque results from a combination of macrophage death and defective phagocytic clearance of apoptotic cells, accelerating or inducing plaque disruption through the release of inflammatory cytokines and matrix proteases. Mechanical stress from the necrotic core on the overlying cap may also contribute to plaque rupture.

The generation of PDGF B chains by macrophages, consistently observed throughout all stages of atherogenesis in both human and experimental models [130,131], likely plays a crucial role in recruiting smooth muscle cells to the intima. Macrophage-produced heparin-binding epidermal growth-factor-like growth factor is also implicated in the migration of smooth muscle cells. Nevertheless, the anticipated outcome involves a mosaic of macrophages and smooth muscle cells. The sorting of cells through cell surface cadherins may contribute to this process, given that smooth muscle cells, expressing N-cadherin, can engage in homotypic binding, forming strong cell-to-cell interactions that exclude macrophages. Notably, N-cadherin is co-expressed with VE-cadherin on

endothelial cells, promoting contact between endothelial cells and smooth muscle cells. The detailed histological characteristics of rupture-prone plaques are extensively addressed in other sections of this volume. In simple terms, plaque rupture tends to occur in cases featuring a large lipid core, a thin cap, and an abundance of macrophages relative to smooth muscle cells. The prevailing perspective suggests that rupture arises from the degradation of an established fibrous cap, a process primarily mediated by macrophages. An alternative interpretation posits that new areas of fatty streak at the peripheries of existing fibrous plaques simply fail to develop, or have not yet developed, a sufficiently robust plaque cap. Various agents, including prostaglandins that elevate cAMP concentrations [199,200], nitric oxide that raises cGMP levels, interferon- γ , and TGF- β , counteract the effects of smooth muscle cell mitogens [136,137]. For instance, the transfer of the VEGF gene leads to a NO-mediated reduction in neointima formation following cuff-induced injury in carotid arteries. The pathways involved are likely diverse, encompassing the downregulation of early events such as MAP kinase activity and subsequent events, including the expression of cyclin D1 and c-myc. Leukocytes are likely the primary source of these inhibitory agents. Studies by Hansson and colleagues have demonstrated that interferon- γ infusion reduces intimal lesions caused by balloon injury, while lesion promotion occurs in lymphocyte-deficient rats, suggesting similar considerations in atherogenesis. If so, the presence of leukocytes in the shoulder regions of advanced plaques may favor the inhibition rather than the proliferation of smooth muscle cells.

In this scenario, the angiotensin-converting enzyme/angiotensin II (ACE/Ang II) system plays a crucial role in forming atherosclerotic lesions. The activation of ACE/Ang II contributes to the formation of a thick fibrous cap. In the early stage of atherosclerotic lesion formation, endothelial cells and macrophages may secrete biologic factors such as ACE/Ang II and induce intimal thickening and the formation of a lipid core. After maturation of the lesion, macrophages may lose the ability to secrete ACE/Ang II, resulting in the thinning of the fibrous cap. A thin fibrous cap and faint expression of ACE/Ang II were observed in the shoulder lesion of unstable plaques. Plaque rupture may be exacerbated by the absence of a correct ACE/Ang II system activity.

A study about the correlation of thin fibrous caps possessing adipophilin-positive macrophages and IPH with high clinical risk for carotid endarterectomy distinguished patients in four groups of risk and demonstrated that plaques in patients in Grade IV (presence of neuro-logical risk) showed more adipophilin-expressing macrophages in the fibrous cap than in those of the other groups. It has been demonstrated that infiltration of adipophilin-positive macrophages in the fibrous cap may be correlated with instability in neurological status. In terms of clinical approaches, this evidence invites us to avoid stenting procedures and perform carotid endarterectomy.

Recently, research studies have investigated the relationship between small non-coding microRNAs' expression and carotid plaques' development and vulnerability. MiRNAs have been shown to play a crucial role in the atherosclerotic process, which suggests that they could be used as potential biomarkers for the diagnosis and prognosis evaluation of cardiovascular events, as demonstrated by multiple recent studies.

VSMC balance can prevent atherosclerosis. In advanced carotid plaques, the expression of miR-145 and miR-210 drives the migration and proliferation of VSMCs, promoting the stability of the fibrous cap and prevention of plaque rupture. The most studied miRs that can reestablish contractile function in VSMCs and potentially be used as therapeutic targets are miR-22 and miR143/145.

In contrast, the predictor of plaque instability seems to be miR-92a, which regulates MMP-9 overexpression, and miR-200c, which was found to be up-regulated in carotid plaques that presented unstable features in imaging studies.

However, studies that have identified specific biomarkers involved in developing a thin fibrous cap are limited, and research is necessary to identify particular microRNAs that could be used in carotid atherosclerosis.

Furthermore, finding circulating biomarkers that can identify patients with thin-cap atherosclerosis lesions at high risk of atherothrombotic stroke remains a challenge.

One important issue that is still unresolved is distinguishing between different lesions by the identification of genes specifically associated with rupture. Faber et al.¹ tried to give a solution

reporting the presence of perilipin exclusively in lesions exhibiting acute rupture. Perilipin was absent in similar vulnerable lesions with necrotic core areas that appeared indistinguishable from those in ruptured lesions expressing the molecule. Consequently, these researchers may have uncovered a potentially significant causative factor for rupture, or alternatively, one among several molecules expressed post-rupture. It is improbable that perilipin serves as the exclusive indicator of rupture. Presenting a single gene among the 35,000 candidates possibly implicated in this process is, at most, an initial stage. While the era of plaque imaging is emerging, the ability to discover individual or even a few new genes, as exemplified in the current study, is approaching its conclusion. Current estimates suggest approximately 35,000 sites of RNA expression in the genome. The technology for representing all these sequences in a hybridization array is already nearly available in the existing generation of microarrays. In theory, one could extract RNA from both a ruptured plaque and a non-ruptured plaque, utilizing a large-scale expression array to meticulously outline all potential differences in expression. Single observations, such as those in the present study, will eventually be considered anecdotal as this technology progresses.

In another recent study, Bazan et al. compared the transcriptomes of asymptomatic and recent ruptured carotid plaques to determine molecular mechanisms active at the time of rupture. The authors found that atherosclerotic plaques that have recently ruptured have shown heightened activity in pro-inflammatory genes responsible for attracting and mobilizing leukocytes to the vessel wall. In the recently ruptured group, there was an increase in the expression of XCR1, a chemokine receptor, and CD177, which facilitates neutrophil activation. Furthermore, the same group exhibited elevated levels of various transcripts linked to B cell function, including numerous immunoglobins, as well as transcripts associated with B cell proliferation (MZB1) and activation (CD79A, SH2D3C, and ZAP70).

The samples of recently ruptured atherosclerotic plaques also displayed alterations in transcript expression, potentially linked to the thinning of the fibrous cap, characterized by reduced VSMC proliferation and migration and increased apoptosis. SDC1 and SIK1, associated with decreased VSMC proliferation and migration, were among the genes showing changes. Additionally, there was an upregulation of transcripts related to the inhibition of cell proliferation (NPDC1, DOK3, and ZBTB17), and a decrease in the expression of a pro-migration transcript (TMSB15B) was observed in the recently ruptured samples. Numerous transcripts encoding Ras GTPase-activating proteins (RASA4, RASA4B, ARHGAP4, HMHA1, and SH3BP1) were identified, promoting the inactivation of Rac1 and cdc42, thereby inhibiting proliferation and migration. Finally, in a fascinating study by Bobryshev et al., a comparison between the relative number of calcified matrix vesicles in fibrous caps of vulnerable (FCT < 100 μ m) and presumably stable atherosclerotic plaques (FCT > 100 μ m) was performed. The quantification of matrix vesicles in stable and vulnerable plaques showed that the relative densities of matrix vesicles were significantly higher in fibrous caps of vulnerable plaques than those in stable plaques. These results suggest that accumulation and calcification of matrix vesicles could be involved in the process of plaque rupture, possibly through the alteration of the texture of connective stroma in plaque fibrous caps.

8. Neo Angiogenesis and Intraplaque Hemorrhage

The regulation of angiogenesis in atherosclerosis involves molecular mechanisms primarily driven by oxygen deficiency. Hypoxia induces monocyte/macrophage survival and the uptake of ox-LDL by macrophages, while also upregulating MMPs expression in various plaque cells, contributing to plaque instability.

Angiogenic sprouting entails the proliferation and migration of ECs into avascular areas. Pathological angiogenesis is a consistent feature of atherosclerotic plaque development, and its progression is linked to a gradient of VEGF triggering EC growth from existing adventitial vasa vasorum. However, the precise origin of plaque neovessels remains incompletely established.

Neoangiogenesis is intimately linked with plaque advancement and likely serves as the primary origin of IPH at locations with microvessel incompetence. Focal aggregations of angiogenic factors derived from T-cells and macrophages contribute to: (1) the branching of vasa vasorum around the

necrotic core; (2) the formation of immature vessels; and (3) the loss of the basement membrane around functional capillaries. This sequence of events initiates the leakage of red blood cells (RBCs) into the plaque and sets off a cycle of inflammation and neovascularization.

Neovessels in vulnerable plaques exhibit immaturity, irregularity, and fragility due to compromised structural integrity. They possess a discontinuous basement membrane and a low number of tight junctions between ECs. Additionally, these premature vessels have limited pericyte coverage and are prone to the leakage of circulating cells, leading to IPH. Similar characteristics are observed in newly formed vessels in tumors, which are heterogeneous, organized chaotically, lack hierarchical branching, and display hyper-permeability.

In advanced atherosclerotic lesions, neovessel leakage becomes the primary entry point for inflammatory cells. RBCs facilitate the extravasation of circulating inflammatory cells by increasing the numbers of rolling and adhering monocytes, potentially enhancing the force and frequency of collisions with the endothelium. Neutrophils and mast cells, associated with neovessels, release granular content rich in serine proteases and MMPs, contributing to elastic fiber and basement membrane degradation. This high proteolytic activity can lead to fibrous cap thinning and plaque erosion.

Moreover, RBC influx and lysis increase the demand for macrophage activity to phagocytose RBC remnants. Combined with impaired efferocytosis in atherosclerotic lesions, hindering the clearance of apoptotic cells by lesional macrophages, this accumulation may explain why macrophages accumulate in the atherosclerotic necrotic core, potentially exacerbating vascular inflammation.

Basic fibroblast growth factor (bFGF) is also involved in promoting intraplaque angiogenesis, and it is stimulated by local hypoxia leading to the formation of neovessels from the vasa vasorum in the adventitia. These newly formed vessels are immature, lacking proper pericyte coverage and tight junctions between endothelial cells, resulting in IPH. This hemorrhage, characterized by the extravasation of RBCs and inflammatory cells, sustains ongoing inflammation. Simultaneously, invading inflammatory cells, especially macrophages, stimulate the synthesis of various angiogenic factors, further promoting plaque angiogenesis. bFGF also facilitates macrophage infiltration into lesions by inducing the expression of chemokines and adhesion molecules. The accumulation of foam cells, derived from macrophages, contributes to lipid storage, atherosclerotic plaque growth, and a pro-inflammatory plaque phenotype. This interplay between intraplaque angiogenesis and inflammation forms a vicious cycle, driving the atherosclerotic plaque toward an unstable phenotype and potential rupture.

bFGF has long been recognized for its role in promoting SMC proliferation and migration, making it a key target in preventing post-interventional intimal hyperplasia. SMCs play a dual role in atherosclerotic lesion stability, contributing to lesion formation through intimal proliferation but also playing a crucial role in forming the fibrous cap that covers the lesion. Following vascular injury or endothelial cell activation, SMCs transition from a contractile to a proliferative phenotype, marked by increased cell proliferation and migration. SMCs in the fibrous cap primarily exhibit a contractile phenotype, and the thickness of the cap is critical for plaque stability, as thinning increases the risk of plaque rupture. Higher expression of bFGF in SMCs is observed in unstable plaques compared to stable plaques, and it is known to stimulate a proliferative phenotype. A recent study hypothesized that blocking bFGF signaling would potentially reduce intraplaque angiogenesis, macrophage infiltration, and SMCs proliferation, resulting in the stabilization of atherosclerotic plaques and demonstrated that K5-mediated bFGF signaling blockade in hypercholesterolemic Apolipoprotein E (ApoE)³*Leiden mice reduces intraplaque angiogenesis, hemorrhage and inflammation.

A primary contributor to micro- and macrohemorrhages within atherosclerotic lesions is an underdeveloped network of blood vessels that emerges within the intima of a plaque. While the majority of these newly formed vessels have endothelial coverage, they seldom feature mural pericytes or smooth muscle cells.

In the context of atherosclerosis, the permeability of intraplaque vasa vasorum, exhibiting leakage, has been characterized through ultrastructural visualization, revealing defects between

endothelial junctions. Alternatively, the identification of perivascular vWf immunoreactivity around vessels has been utilized as an indicator of a less intact or leaky endothelium. The leakiness of these vessels is influenced by factors that induce vessel growth, with VEGF and its related receptors and cofactors serving as the principal regulators of vascular permeability in pathologic angiogenesis. The VEGF signaling pathway involves an interaction between growth factor receptors and specific integrins. The VEGF receptors are likely to play a crucial role in cell-to-cell adhesion and communication by directly interacting with cell-to-cell adhesion molecules.

Observational investigations into the progression of the necrotic core highlight IPH as a crucial factor in the growth and destabilization of atherosclerotic plaques. The swift accumulation of erythrocyte membranes brings about a sudden alteration in plaque composition, characterized by an augmented presence of free cholesterol within the core and an excessive infiltration of macrophages.

Examinations of unstable lesions through pathological analysis have revealed an association between IPH; plaque rupture; and an elevated microvessel density. The precise mechanism by which RBCs leak into the necrotic core remains poorly understood. Our research laboratory has documented widespread perivascular staining of vWF within the vasa vasorum of plaques; along with evidence of erythrocyte membranes within necrotic cores. This observation suggests that microvascular disruption or permeability may contribute to lesion progression by providing cholesterol derived from erythrocytes. Apart from leaky vasa vasorum; the accumulation of erythrocytes can also result from plaque fissuring; a phenomenon described in the coronary vasculature of individuals who succumbed to sudden coronary death.

The count of vasa vasorum was found to be twofold higher in vulnerable plaques and up to fourfold higher in ruptures compared to stable plaques with significant luminal narrowing. Increased T cell numbers are commonly observed at breaks in the medial wall and the base of the necrotic core, as opposed to other plaque regions. It is plausible that T helper cell-driven immune responses, possibly mediated by interferon- γ , may hinder smooth muscle cell proliferation, contributing to medial disruption and the absence of smooth muscle cells in penetrating neovessels.

Recent attention has been directed towards VE-cadherin in carotid atherosclerotic plaque due to its recognized significance in angiogenesis. The VE-cadherin/catenin complex is implicated in the organization and maturation of neovessels and in safeguarding endothelial integrity. Recent evidence, consistent with previous findings by Bobryshev et al, suggests that VE-cadherin is selectively deposited in neovessels surrounded by inflammatory cells, underscoring its central role in this process. Complicated plaques exhibit heightened expression of VE-cadherin, along with analogous expression of beta-catenin, moderate expression of alpha-catenin, and little to no expression of plakoglobin. The reported necessity of plakoglobin for the formation of 'mature' junctions suggests that the absence of plakoglobin in neovessels compromises junction function, increasing vascular permeability.

A significantly elevated expression of VE-cadherin has been demonstrated in complicated plaques, with even higher expression in high-grade stenotic lesions. This finding suggests that VE-cadherin may also play a crucial role in the progression of carotid lesions. Moreover, the strong expression of this molecule in patients with cerebral infarction and clinical symptoms indicates an association between neovascularization and symptomatic carotid disease, as corroborated by previous reports. Consequently, VE-cadherin could be considered a marker of advanced carotid atherosclerotic plaques.

Recently, some researchers employed an innovative translational approach in proteomics to unveil signature molecules associated with the development of carotid atherosclerosis. A captivating study integrating also transcriptomics discovered an abundance of biliverdin reductase B in both plaque tissue and plasma. This association was linked to IPH and processes that were previously correlated with plaque instability.

Although the mechanisms underlying plaque activation have not been completely elucidated, recent data suggest that macrophages and smooth muscle cells release angiogenic factors, causing migration and proliferation of endothelial cells in the intima and expression of VE-cadherin. VE-cadherin is involved in the maturation, expansion, branching, and remodeling of endothelial cells.

into a network of new blood vessels in the plaque. Neovascularization serves as the primary route for migrating inflammatory cells into the intima, involving cell adhesion molecules (E-selectin, intercellular adhesion molecule-1, and vascular cell adhesion molecule-1).

Moreover, a recent study underlined the potential role of in promoting VEGFA/VEGFR2-mediated neovascularization in *ApoE^{-/-}* mice, supporting a pivotal role of Sema7A in the progression of human atherosclerosis and IPH.

Finally, according to recent data, the advancement of IPH is linked to the involvement of endogenous IL-1 β , which inhibits physiological angiogenesis in the atherosclerotic plaque, ultimately resulting in the development of permeable neovessels. Additionally, the induction of MMP-9 expression may play a role in the formation of these leaky neovessels.

9. Lipidic Core

Research on the lipidic core of atherosclerotic plaques is a subject of considerable importance in understanding plaque vulnerability, prognosis, and developing targeted therapies, as its characteristics play a crucial role in determining plaque stability. The presence of a large lipid core occupying more than 40% of the plaque area is considered one of the main criteria for plaque vulnerability [8,180]. Furthermore, previous studies have separated the resulting plaques by stability and measured lipid subcomponents using mass spectrometry showing that the lipid content of unstable plaques was higher than that of stable plaques.

The pathogenesis of this more clinically significant plaque component remains controversial. Lipids, primarily LDL carried by blood, may undergo direct accumulation within the extracellular space or be internalized by macrophages, likely through scavenger receptors following oxidative modification. Subsequently, accumulation may occur indirectly through necrosis of lipid-filled macrophages, also known as foam cells. Nevertheless, findings challenge the notion that the core primarily originates from deceased foam cells in the superficial intima (fatty streaks). Instead, evidence suggests that the core develops predominantly from a gradual buildup of lipids in the extracellular matrix of the deep intima, resulting from intricate interactions between insulating LDL and glycosaminoglycans, collagen, and/or fibrinogen.

The lipid core within a plaque is devoid of supporting collagen, avascular, hypocellular (except at the periphery of the core), and rich in extracellular lipids. By quantitatively analyzing lipid plaque composition, it has been determined that in plaques, cholesteryl ester content is the highest, followed by phospholipids, third-free cholesterol, and triglyceride.

A genome-wide microarray expression study in advanced carotid plaques showed that in transition from being asymptomatic to causing symptoms, there is a concurrent increase in the expression of macrophage adipophilin, a marker indicating the loading of lipids. Moreover, an overexpression of the CD36 and ATP-binding cassette (ABCA) 1 genes in CPs was identified. CD36 and ABCA1 appear to have a crucial role in regulating the intake and release of lipids by macrophages. CD36, known for its role in scavenging oxidized LDL, is a multifunctional membrane glycoprotein considered vital for the formation of foam cells. Its expression is further enhanced by its ligand, creating a potential vicious cycle leading to an excessive intake. However, the detrimental effects of this cycle are mitigated to some extent by ABCA1, a membrane transporter protein belonging to the ABC family. ABCA1 is widely recognized for its atheroprotective properties, as it helps in removing free cholesterol and phospholipids. Additionally, it plays a crucial role in initiating the reverse transport of cholesterol in macrophages, a process essential for maintaining cholesterol homeostasis. Mutations in ABCA1 can lead to a severe deficiency in high-density lipoprotein (HDL), causing cholesterol buildup in tissue macrophages and promoting atherosclerosis. Also, ABCG1 plays a crucial role in the efflux of cellular cholesterol to HDL and its apolipoproteins. Furthermore, both ABCA1 and ABCG1 influence the release of inflammatory signals from cells by adjusting the amount of cholesterol in the cell's outer layer and inside its compartments, reducing the inflammatory stress. Metabolites elevated in metabolic syndrome and diabetes negatively impact ABCA1, reducing cholesterol export from macrophages. Additionally, oxidative modifications of HDL in cardiovascular disease patients hinder apolipoproteins from efficiently removing cellular cholesterol

via the ABCA1 pathway. These findings suggest that a compromised ABCA1 pathway may contribute to increased atherogenesis in common inflammatory and metabolic disorders, making it a crucial target for potential cardiovascular disease therapies .

Other specific biomarkers associated with lipidic core content, such as elevated levels of lipoprotein-associated phospholipase A2 (Lp-PLA2) and lysophosphatic acid, have been studied for their prognostic role in the prediction of cerebrovascular accident. PLA2 is associated with Lp(a). Lp(a) is a type of lipoprotein that promotes the development of arterial plaques and is found only in areas with atherosclerosis. Pathological evidence indicates that Lp-PLA2, associated with Lp(a), might contribute to atherosclerosis; in more advanced plaques, strong Lp-PLA2 is staining in areas rich in lipids and oxidation byproducts [189,190]. As related to lipidic core content, the determination of Lp-PLA2 could be used in conjunction with clinical evaluation to assist in predicting patient risk of cardiovascular disease .

Lysophosphatidic acid (LPA) builds up in the lipid-rich core of carotid atherosclerotic plaques and gets released when the plaque ruptures. This release activates platelets and can lead to the formation of blood clots within the artery. In human atherosclerotic plaques, LPA levels are significantly higher (13-fold) compared to normal arterial tissue, and the highest concentrations are found in the lipid-rich core. The elevated LPA concentration in the plaque's lipid core may be exposed after cap rupture, potentially causing the formation of local blood clots. LPA concentrations were significantly higher and more common in patients with unstable plaque. This means the LPA determination may be a useful biomarker in the clinical identification and prediction of unstable plaque and guiding treatment .

The Multi-Ethnic Study of Atherosclerosis showed that in people with thickened carotid walls, plasma total cholesterol level was strongly associated with lipid core presence by MRI and, as a consequence, high total cholesterol levels may be associated with rupture proneness of atherosclerotic lesions in the general population confirming the pivotal role of the targeted therapy with statins and lipid-lowering drugs.

Furthermore, emerging experimental models are shedding light on the formation of cholesterol-rich vesicles. In the mentioned animal model, rabbits examined 1 to 2 weeks after initiating cholesterol feeding exhibit predominantly vesicular, cholesterol-rich lipid deposits beneath the arterial endothelium . Mitchinson and colleagues have suggested that many vesicles in human atherosclerosis may represent ceroid, a peroxidized and cross-linked complex of lipid and protein. Ceroid can be generated in vitro by macrophages exposed to LDL or polyunsaturated cholesteryl esters. While ceroid is undoubtedly found in the core region of plaques, quantitative assessment is required, as early estimates propose it may only account for 1% to 5% of core lipids, rather than the approximate 25% area fraction occupied by vesicles .

Experiments by Tangirala and coworkers have delved into the mechanisms by which lysosomal hydrolysis of cholesteryl ester contributes to the accumulation of free cholesterol in cultured cells. Rapid loading of macrophages with cholesteryl ester led to a significant accumulation of free cholesterol in lysosomes via hydrolysis, exacerbated by inefficient transfer of cholesterol to extralysosomal locations. Prolonged incubation after loading resulted in the formation of cholesterol crystals within lysosomes, with extracellular crystals appearing in the culture medium at extended time points . VSMCs in culture also took up lipid droplets into lysosomes through phagocytosis, accumulating excess free cholesterol in a compartment identical to or closely linked with the plasma membrane, suggesting inefficient cytoplasmic reesterification . These findings gain additional relevance as the initiation of the cholesterol-rich core in human atherosclerosis is localized to the musculoelastic (deep) intimal sublayer, where the predominant cell type in early lesions is smooth muscle .

The mechanisms through which cellular accumulation of free cholesterol leads to the formation of extracellular cholesterol-rich vesicles in lesions are not fully elucidated. Possible mechanisms include selective or nonselective cell death, detachment of cellular blebs, or extrusion of lysosomal contents. Schmitz, Robenek, and colleagues proposed a model supporting the extrusion hypothesis,

where macrophages loaded with lysosomal lipid were treated with nifedipine in a lipoprotein-free medium, leading to the release of cholesterol-rich lamellar bodies from lysosomes into the medium.

Finally, Chung and colleagues put forth a novel mechanism suggesting that vesicles are formed extracellularly, not from LDL as commonly assumed, but through an interaction of very-LDL, HDL, and lipoprotein lipase. Supporting their proposal, cholesterol-rich vesicles from the human aorta were found to contain apoA-I and several apoC species but only scant apoB.

Brooks et al. first identified oxidized derivatives of cholesterol and cholesteryl esters in human atherosclerosis. Subsequently, oxidized LDL was discovered within human lesions. Myeloperoxidase, an oxidative enzyme produced by monocytes and granulocytes, was detected in both the shoulder regions and the core of fibrous plaques. Intriguingly, the oxidation of LDL can promote aggregation, establishing a connection between these two processes.

In a recent study, Carpenter and colleagues measured oxidized lipids in the core ("necrotic gruel") of advanced human aortic plaques. Hydroperoxy- and hydroxyoctadecadienoic acids, derived from linoleic acid through lipid peroxidation, were present at levels totaling 6% to 7% of the measured linoleic acid in the same material. These percentages fall within the range observed for copper-oxidized LDL. Carpenter et al. also quantified 7 β -OH-cholesterol after saponification and sodium borohydride reduction of core lipids. The resulting concentration of 7 β -OH-cholesterol, representing most oxysterols formed via lipid peroxidation from cholesterol, was only 0.14% of total cholesterol. This concentration is significantly lower than the relative concentration of oxysterols formed in most in vitro models of lipoprotein oxidation. Interestingly, an enzymatically oxidized cholesterol derivative, 26-OH-cholesterol, was approximately fivefold more abundant in core lipids than 7 β -OH-cholesterol. The 7 β -hydroperoxide of cholesterol may exhibit considerably greater bioactivity than the corresponding hydroxy and keto derivatives. Chisolm and colleagues recently identified the hydroperoxide in human plaques. Consequently, substantial evidence of lipid oxidation has been found in the atherosclerotic core, although the formation or retention of oxysterols seems to be less than that observed with the oxidation of lipoproteins in vitro.

Cells bordering and penetrating the atherosclerotic core may not only participate in the deposition or removal of core lipids but may also be influenced by the accumulating lipids and proteins. Complement components are relatively abundant in the core, and complement activation on core lipids may generate both toxic and chemotactic responses. Antigenic markers of complement activation, such as C3d and the terminal C5b-9 neoantigen, have been identified in the atherosclerotic core. Terminal C5b-9 was also found coincident with cholesterol-rich vesicles in the subendothelium of cholesterol-fed rabbits. Cholesterol, oxysterols, and cholesterol-rich vesicles have been shown to activate complement. The cellular effects of complement activation in atherosclerosis require further definition, possibly through transgenic models.

The potential for excessive cholesterol buildup within cell membranes near the core needs evaluation. Under normal circumstances, most cells tightly regulate their plasma membrane cholesterol content between 0.4 and 0.8 mol per mole of phospholipid, depending on the cell type. Phosphatidylcholine, the chief phospholipid of cell membranes, allows ratios of cholesterol to phosphatidylcholine up to 1.0 before phase separation and cholesterol monohydrate crystal formation occur. However, sphingomyelin-rich vesicles in the atherosclerotic core have higher levels of free cholesterol, up to 2.5 mol per mole of total phospholipid. Cholesterol can diffuse at slow but meaningful rates, suggesting that membranes near cholesterol crystals might become saturated with cholesterol at levels much higher than the physiological level for cell membranes.

Moreover, lipoproteins in proximity to cholesterol crystals may become saturated with free cholesterol or undergo phase alteration to become vesicles, potentially impairing reverse cholesterol transport at the tissue level. The impairment of reverse cholesterol transport in atherosclerotic tissue remains a plausible but unproven hypothesis requiring detailed examination.

Considering this environment, cells in atherosclerotic lesions may employ defense mechanisms to prevent excessive free cholesterol accumulation. These mechanisms include unregulated uptake of modified lipoproteins, phagocytosis, passive transfer, and an increase in acylcoenzyme A:cholesterol acyltransferase (ACAT) activity leading to cytoplasmic cholesteryl ester accumulation

. However, the accumulation of cholesteryl ester within cells cannot continue indefinitely, and the pharmacological inhibition of ACAT, while proposed to decrease cholesteryl ester accumulation in atherosclerotic lesions, may have potential adverse effects due to excessive free cholesterol buildup.

A recent adaptation to a high cholesterol content in cells may involve increased synthesis of sphingomyelin. Alteration of sphingomyelin metabolism in vascular cells could have unexpected effects, as sphingomyelin metabolites comprise a signal transduction pathway decreasing the activity of protein kinase C. The observed link between ceramide, a hydrolysis product of sphingomyelin, and apoptosis in the atherosclerotic core suggests a potential connection between these lipids and cell death.

If cellular defenses fail and plasma membrane cholesterol concentrations increase, membrane function derangement is expected. Increases in calcium flux and other alterations have been demonstrated in smooth muscle cells exposed to high concentrations of free cholesterol. Macrophages loaded with cholesterol in the presence of ACAT inhibitors showed evidence of toxicity. The administration of an ACAT inhibitor reduced the content of macrophages in rabbit atherosclerotic lesions. Whether cholesterol-induced toxicity for macrophages might have played a role in this in vivo effect is speculative.

The oxidation of core lipids provides another avenue for cellular effects, with cytotoxicity from oxidized LDL ascribed recently to specific oxidized cholesterol derivatives. These compounds have been identified among atherosclerotic lipids, and cellular responses to lipid oxidation products are well-documented.

Various cell activities have been studied in the context of atherosclerotic lesion development, but not always in specific relation to the core or its lipids. Macrophages are commonly localized at the lateral boundaries and sometimes in the interior of the core. This localization could reflect a contribution by macrophages to lipid deposition, chemoattraction by lipid components or complement fragments for monocyte/macrophages, and perhaps a role for macrophages in lipid removal. Capillarization and calcification of lesions appear closely related to core development, with capillarization being a secondary phenomenon, as the early core does not contain microvessels. Expression of bone-related genes has been found near the core, adjacent to sites of dense calcification. A review of the atherosclerotic core in the future should include new data on these and other cellular responses.

The persistent inflammatory response ultimately destabilizes atherosclerotic plaques through the action of proinflammatory cytokines. Studies have demonstrated that IFN- γ , IL-18, growth differentiation factor-15, and TNF-like weak inducers of apoptosis contribute to plaque destabilization while TGF- β promotes stabilization. IFN- γ , TNF- α , and IL-1 β promote apoptosis of macrophages and foam cells, leading to lipid core enlargement. Moreover, these cytokines induce apoptosis of SMCs, resulting in thinning of the fibrous cap. Proinflammatory cytokines also impede the synthesis of plaque-stabilizing ECM components produced by SMCs. For instance, IFN- γ inhibits collagen synthesis by SMCs.

Macrophages infiltrate the thinned fibrous cap, expressing and secreting numerous inflammatory cytokines and MMPs that digest the stabilizing matrix, playing a pivotal role in weakening and ultimately rupturing the atherosclerotic plaque. It has been reported that necrosis of the vulnerable plaque is attributed to a combination of macrophage death and defective phagocytic clearance of apoptotic cells, accelerating or inducing plaque disruption by releasing inflammatory cytokines and matrix proteases. Finally, the mechanical stress caused by the necrotic core on the overlying cap is also considered an important element that can contribute to plaque rupture.

10. Microcalcification

The process of atherosclerosis, characterized by the accumulation of plaques in arterial walls, is a complex and dynamic phenomenon influenced by various factors, including inflammation and calcification [214,215]. Understanding the interplay between these processes is crucial for unraveling the mechanisms underlying plaque vulnerability and stability.

Recently, there has been a growing interest in understanding the impact of calcification on plaque vulnerability [216,217]. According to the classification proposed by Naghavi et al., plaque microcalcifications are considered a minor criterion for determining plaque vulnerability, corresponding to type Vb lesion as per the American Heart Association's histological classification. [8,218]

The role of calcification in determining plaque vulnerability remains a topic of ongoing debate. Notably, mechanical experiments conducted on human carotid plaques by Mulvihill et al. suggested that calcification within the tissue structure might contribute to an increased vulnerability of the plaque. Conversely, findings by Shaalan et al. indicated that symptomatic plaques tend to be less calcified and more inflamed than asymptomatic plaques, suggesting a potential risk reduction of plaque rupture associated with calcification.

In a study by Wahlgren et al., thirty carotid endarterectomy plaques were classified as noncalcified and calcified, yielding results consistent with the notion that fibrous cap inflammation is more likely to occur in noncalcified plaques compared to calcified ones. This observation implies a protective effect of plaque calcification against rupture.

Computational investigations on microcalcifications have also contributed to the understanding of plaque vulnerability. Kelly-Arnold et al. examined the spatial distribution, clustering, and shape of different microcalcification sizes in fibrous caps, revealing that nearly all fibrous caps contain microcalcifications, but only a small subset has the potential for rupture. Engrenyuk et al. explored stress distribution using multilevel micro-CT-based 3D numerical modeling techniques. Their results indicated that the presence of calcifications (inclusions) increases peak circumferential stress, with elongated microcalcifications potentially further elevating stress levels. In contrast, macrocalcifications in cap shoulders were found to enhance plaque stability.

Microcalcifications, initially smaller than 50 μm , typically emerge from the lipid pool and early necrotic core, with potential formation in the fibrous cap [214,215]. These microcalcifications can progress into larger structures, such as calcified sheets and nodules, ultimately leading to plaque ossification, which is prominently observed in peripheral arteries. Recent evidence suggests that heavily calcified plaques, characterized by sheets of calcification or ossification, exhibit stability, contrasting with plaques featuring small and diffuse calcifications [224,225]. Intravascular imaging techniques, such as intravascular ultrasound and optical coherence tomography, have revealed that vulnerable plaques are often associated with spotty calcifications, whereas larger calcium deposits may indicate stability.

The paradigm of heavily calcified plaques being stable is further supported by the impact of statins on plaque development. Statins, known for reducing adverse cardiovascular events, have been shown to decrease plaque surface while increasing calcification, as demonstrated by IVUS imaging studies [227,228]. This observation has been corroborated by computed tomography and magnetic resonance imaging. Experimental studies in ApoE-deficient mice treated with pravastatin showed coalescence of microcalcifications into larger structures.

The mechanical impact of macrocalcifications on plaque stability appears to be stabilizing rather than harmful, in contrast to the potential danger posed by microcalcifications smaller than 50 μm . Microcalcifications in the fibrous cap, particularly those exceeding 10 μm , have been suggested to exert dangerous mechanical stress, contributing to plaque rupture. Moreover, microcalcifications may not only pose a risk through mechanical stress but also by triggering or amplifying plaque inflammation. Studies using 18F-sodium fluoride PET/computed tomography have associated microcalcifications with macrophage infiltration and apoptosis in carotid plaques, emphasizing their potential role in inflammatory processes. The presence of coronary microcalcifications has been linked to macrophages, further supporting the connection between inflammation and microcalcification. Interestingly, as plaque calcification progresses, macrophage infiltration decreases, and plaques become less symptomatic.

The initiation and evolution of microcalcifications towards larger, potentially more stable structures involve cellular and molecular mechanisms. Inflammation has been implicated as a trigger for microcalcification, supported by clinical studies and animal models. Cell necrosis, especially

within the necrotic core, serves as a potent inducer of pathological calcification. Compromised macrophage phagocytosis of apoptotic bodies, a hallmark of atherosclerosis plaques, may contribute to the formation of early microcalcifications. Additionally, a form of programmed cell necrosis called necroptosis has emerged as a key player in advanced human plaque development, particularly in macrophages.

Inflammatory cytokines, including TNF- α , IL-1 β , and IL-6, stimulate VSMCs to transdifferentiate into chondrocyte-like cells, promoting calcification [238–240]. Understanding these processes is critical, as inflammation appears to precede calcification, and the interaction between inflammation and calcification creates a feedback loop that amplifies both phenomena [241,242].

Repair mechanisms associated with calcification play a role in the terminal phase of atherosclerosis, with large and stable calcification deposits contributing to the resolution of inflammation. Macrophages and VSMCs generate calcification inhibitors, such as matrix Gla protein (MGP), Gla-rich protein, and OPN, to prevent additional crystal growth. Furthermore, VSMCs produce bone morphogenetic protein 2 (BMP2), potentially activating the transition of VSMCs into chondrocytes and creating a cartilage-like environment, ensuring the stability of the calcified tissue.

TNF- α , among other pro-inflammatory cytokines, has the potential to stimulate VSMCs to mineralize independently of their transdifferentiation into mineralizing cells. For example, TNF- α reduces the expression of the inorganic pyrophosphate transporter ANKH and diminishes inorganic pyrophosphate export in VSMCs, a condition linked to calcification, as ANK deficiency in mice leads to soft tissue calcification. Furthermore, TNF- α decreases extracellular inorganic pyrophosphate levels through tissue-nonspecific alkaline phosphatase activation, and it is noteworthy that tissue-nonspecific alkaline phosphatase has roles beyond inorganic pyrophosphate removal in mineralized tissues, including functions in inflammation.

Tissue-nonspecific alkaline phosphatase expression is induced by TNF- α or IL-1 β in human mesenchymal stem cells, leading to increased mineralization, while simultaneously inhibiting the expression of RUNX2 and its transcriptional targets. Additionally, TNF- α may induce calcification in VSMCs by downregulating another crucial calcification inhibitor, MGP. Calcium phosphate crystals smaller than 1 μm activate macrophages, triggering the release of pro-inflammatory cytokines, such as TNF- α , IL-1 β , and IL-8. Macrophages phagocytose these crystals, with the smallest ones being most potent at stimulating TNF- α secretion, suggesting an increase in TNF- α production after internalization. Although the precise mechanisms through which crystals activate TNF- α remain unclear, the activation of IL-1 β secretion is better understood, occurring through the NLRP3 inflammasome.

Macrophages stimulated by crystals may activate VSMCs through the release of inflammatory mediators, and the potential stimulation of VSMCs to transdifferentiate through the release of receptor activator of NF- κB ligand (RANKL) is a question warranting investigation. Inhibition of RANKL-RANK signaling with OPG has been associated with reduced plaque calcification. VSMCs, although not specialized for secreting pro-inflammatory molecules, may release cytokines in response to microcalcifications. VSMCs release IL-1 β in the presence of calcium phosphate crystals, and this release increases as VSMCs progress toward senescence, a phenomenon observed frequently in human plaques. IL-1 β release in VSMCs relies on crystal endocytosis and caspase-1 activity, but not necessarily on NLRP3 activation, as other treatments activating NLRP3-dependent IL-1 β secretion in monocytes/macrophages fail to induce IL-1 β release in VSMCs. Long-term treatment of human VSMCs with calcium phosphate crystals induces crystal endocytosis and cell death. VSMC apoptotic death often leads to secondary necrosis and the release of the alarmin IL-1 α , exacerbating plaque inflammation, while apoptotic bodies from dying VSMCs may serve as a nidus for new calcification, further amplifying inflammation.

Recent findings indicate that monocytes and macrophages can produce γ -carboxylated MGP and Gla-rich protein. Both MGP and Gla-rich protein act as inhibitors of vascular calcification and their production is augmented in the presence of inflammatory molecules or hydroxyapatite crystals. This may contribute to the correlation between the presence of microcalcifications, macrophages, and carboxylated MGP observed in human atherosclerotic plaques. Additionally, VSMCs produce

OPN in response to calcium phosphate crystals . Phosphorylated OPN, an inducible inhibitor of ectopic calcification, binds to crystals and inhibits their growth. OPN stimulates macrophages to phagocytose crystals, similar to its role in stimulating osteoclasts for bone resorption. Notably, OPN deficiency in atherosclerotic mice leads to the development of more calcified plaques .

Many studies have investigated the development of calcification formation and have identified the molecules OPN and osteoprotegerin (OPG) as playing a crucial role in promoting atherosclerosis and modulating vascular mineralization [258–260].

OPG, which is a member of the TNFR family, can inhibit bone resorption and suppress immune responses. In terms of cardiovascular health, OPG plays a crucial role in preventing the development of atherosclerosis. It is synthesized by macrophages and expressed by both endothelial and smooth muscle cells. Additionally, OPG is present in Weibel-Palade bodies and is released into the extracellular environment when the endothelium is activated. OPG safeguards the endothelium from the apoptotic process and binds to TSP-1 to regulate vascular damage and thrombus formation. Furthermore, in the medial layer, OPG prevents the calcification of the ECM without dissolving the calcium that has already been deposited [261–264].

As already underlined, OPN is a proinflammatory cytokine with the ability to bind hydroxyapatite crystals to osteoblasts and is involved in systemic inflammatory and remodeling processes. Fibroblasts and macrophages synthesize this protein. Several studies have highlighted a strong correlation between plasma levels of OPN and atherosclerotic lesions, particularly affecting large vessels. [265,266]

OPG and OPN co-exist in carotid atherosclerotic plaque demonstrating a modulatory role in inflammatory and calcification processes. OPG is strongly expressed in stable, calcified plaques, while OPN is poorly expressed in calcified plaques and plaques without vulnerability signs .

According to research, both OPN and OPG contribute to inflammation in atherosclerotic cardiovascular diseases and interact with vascular calcification. However, more research is required to fully understand the mechanisms involved. The findings of these studies need to be validated further.

The progression from a disorganized, pro-inflammatory tissue containing apatite crystals to the formation of new mineralized tissue may not be exclusive to atherosclerosis plaques but also occurs in bone repair following a fracture. Fracture healing commences with an initial inflammatory phase, crucial for inducing subsequent ossification . The absence of TNF- α , particularly, delays ossification, partly by impeding chondrocyte differentiation . Intriguingly, TNF- α may primarily act through BMP2, as it stimulates BMP2 expression in chondrocytes and endothelial cells , and BMP2 deficiency halts early steps of fracture healing . Consequently, the inflammatory impact of apatite crystals in atherosclerosis plaques might be essential for generating osteochondrocyte-like cells that facilitate the development of extensive mineralized surfaces, contributing to plaque stabilization.

A recent histological study in human carotids proposed that TGF- β expressed by M2 macrophages could link inflammation resolution to osteochondrocyte-like cell differentiation and the growth of macrocalcifications . Thus, M1 macrophages might be implicated in the detrimental cycle connecting microcalcification to inflammation, while M2 macrophages could aid in breaking this cycle and promoting plaque repair through ossification. The spatial distribution of macrophages in atherosclerotic plaques supports this notion, with pro-inflammatory M1 cells located in unstable, rupture-prone areas, while anti-inflammatory M2 macrophages are predominantly found in the vascular adventitia, farther from the lipid core and in more stable plaque regions .

M1 macrophages exhibit increased lipid accumulation, mitochondrial dysfunction, destabilized lysosomes, activated oxidative stress, defective efferocytosis, and elevated release of pro-inflammatory mediators, including IL-6, IL-12, IL-1 β , and TNF- α , as well as reactive oxygen species. In contrast, M2 cells contain small lipid droplets and are thought to foster atherosclerosis regression by supplementing anti-inflammatory factors (IL-10 and TGF- β), facilitating efferocytosis, and promoting tissue repair through collagen formation and angiogenesis . The balance between M1 and M2 macrophages likely plays a pivotal role in transitioning from pro-inflammatory plaques with microcalcifications to inflammation resolution associated with plaque macrocalcification.

Interestingly, a comparable role for M1 and M2 macrophages is also strongly suspected during bone fracture healing .

11. Carotid Stenosis

Traditional methods for identifying atherosclerosis in the carotid artery focus on measuring the narrowing of the artery, identified as luminal stenosis. Severe stenosis is a key factor in defining a vulnerable plaque. Plaques with significant stenosis are at a higher risk of developing thrombosis and sudden blockages due to the influence of shear stress. Additionally, the presence of a stenotic plaque may signal the existence of other non-stenotic or less severely narrowed plaques that could be prone to rupture and thrombosis . Recent research has explored molecular markers associated with the vulnerability of atherosclerotic plaques, particularly those related to severe stenosis. Oxidative stress and inflammatory response are identified as major pathways in the development of atherosclerotic stenosis. In an interesting study , authors demonstrated an elevation of sICAM-1 concentration independently associated with atherosclerosis of the ICA origin and predominantly increased in patients with low-grade lesions and with clinical manifestations of vascular disorders. For instance, studies by Li X and colleagues demonstrated elevated circulating levels of various markers, including total oxidant status, lipid hydroperoxide, 8-isoprostane, malondialdehyde, monocyte chemotactic protein-4, amyloid A, hs-CRP, and TNF- α , in elderly patients with both severe stenosis in the right carotid artery and severe multivessel coronary artery stenosis . Furthermore, other studies found significantly increased circulating levels of malondialdehyde, oxidized-LDL, homocysteine, F2-isoprostanes, TNF- α , hs-CRP, prostaglandin E2, and IFN- γ in patients with combined severe stenosis in both carotid and coronary arteries . In patients with carotid stenosis, levels of TSG-6 (Tumor Necrosis Factor-Stimulated Gene-6) in the serum were higher. TSG-6 expression increased in tissues with moderate and severe stenosis. Patients with symptomatic stenosis had higher serum TSG-6 levels compared to those with asymptomatic stenosis. TSG-6 could potentially be a new, easily measurable biomarker for non-invasive screening of severe and symptomatic carotid stenosis in clinical practice . Moreover, patients with symptomatic carotid stenosis showed significantly higher levels of NETs markers than asymptomatic patients and healthy individuals. Increased levels of neutrophil-platelet aggregates led to the generation of NETs in symptomatic carotid stenosis. NETs contributed to Platelet-derived Microparticle (PCA) formation through TF in patients with carotid stenosis. Inhibiting NETs could be a potential biomarker and therapeutic target for preventing recurring strokes in severe carotid stenosis . Damage to the endothelium is a crucial aspect of atherosclerosis. Researchers hypothesized that increased levels of vWF, thrombomodulin, intercellular adhesion molecule-1, and E-selectin might be related to disease severity in patients with peripheral or carotid atherosclerosis. vWF was identified as the most sensitive marker for peripheral atherosclerosis. However, none of the plasma markers seemed to be useful for indicating the degree of carotid artery stenosis . In individuals with high-grade internal carotid artery (ICA) stenosis, those with recent symptoms showed a higher level of plasma sCD36 compared to those with remote symptoms . Furthermore, serum macrophage CXC-chemokine ligand 16 (CXCL16) is identified as a potential biomarker for carotid-vulnerable plaques.

Serum CXCL16 levels increase with plaque area, lumen stenosis rate, and intima-media thickness. . Patients with stenosis of 70% or more have higher levels of MIP-1 α compared to those with stenosis below 70%. Significantly higher levels of CD14 are observed in patients with hypoechoic (vulnerable) lesions compared to those with hyperechoic (stable) lesions . The platelet-to-lymphocyte ratio has been proposed as a novel indirect marker of inflammation. Yayla C et al. demonstrated that the Platelet-to-lymphocyte ratio significantly increases in parallel with the severity of atherosclerosis . In a compelling investigation , the researchers examined alterations in the concentrations of vascular endothelial growth factor (VEGF) and VEGF Receptor-2 (VEGFR-2) in individuals undergoing carotid endarterectomy. The study encompassed 43 patients with extracranial carotid stenosis exceeding 70%. Exclusion criteria included severe vertebrobasilar stenosis, recent (<1 month) vascular events (such as stroke, coronary infarction, arterial thromboembolism), critical lower extremity ischemia, recent infection, autoimmune disease, or

malignancy. Blood samples were collected before CEA and on the second postoperative day, while a control group comprising thirty healthy blood donors was utilized for comparison. The researchers employed an enzyme-linked immuno-absorbent assay to determine VEGF and VEGFR-2 levels.

The pre-operative levels of VEGF and VEGFR-2 were significantly elevated. Following CEA, there was a noteworthy reduction in both VEGF and VEGFR-2, although they did not return to normal values. Notably, in asymptomatic patients and those with contralateral carotid stenosis exceeding 50%, the decrease in VEGF levels did not reach statistical significance. Conversely, within the same subgroups, a substantial reduction in VEGFR-2 values was observed. The study highlighted a substantial increase in the serum levels of VEGF and VEGFR-2 in patients with severe carotid stenosis. These elevated pre-operative levels significantly decreased after endarterectomy, underscoring the pivotal role of these molecules in the progression of carotid disease. Nuotio et al. , in contrast to earlier studies, found that symptomatic carotid disease was not associated with increased expression of adhesion molecules in the endothelium of advanced carotid plaques or circulation but demonstrated a reduced expression of adhesion molecules in the intima-media of symptomatic carotid plaques. Another research study postulates that environments characterized by severe carotid atherosclerotic disease exacerbate endothelial dysfunction, thereby contributing to an elevated risk of recurrent cerebrovascular events. The researchers utilized nonischemic common carotid arteries from mice, mounted them in tissue baths for isometric contraction force measurements, and exposed them to serum obtained from men with a recent ischemic stroke exhibiting varying degrees of carotid stenosis: low- or moderate-grade stenosis (LMGS; <70%) and high-grade stenosis (HGS; ≥70%). The outcomes revealed that serum from stroke patients induced impairment of acetylcholine relaxations in the carotid arteries of mice, indicative of endothelial dysfunction. This effect was more pronounced when incubated with serum from patients experiencing a recurrent stroke or vascular death within 1 year of follow-up. Stratifying patients based on the degree of stenosis demonstrated that serum from HGS patients induced more prominent endothelial dysfunction in the carotid artery, a phenomenon associated with elevated circulating levels of IL-1 β .

12. Conclusion

The specific mechanisms underlying plaque vulnerability and rupture remain elusive, encompassing various factors such as cap thinning, expansion of the lipid core, presence of microcalcifications, neo-angiogenesis, elevated levels of inflammatory cytokines, and proteases that aid in matrix digestion. All cell types involved in atherosclerotic plaque formation play a role in plaque rupture and subsequent thrombosis. Inflammatory cells within the plaque, which convey molecular signals, are crucial for understanding the pathways leading to vulnerability. For instance, lymphocytes can release factors like CD-40L and macrophage-derived foam cells secrete cytokines, growth factors, tissue factor, IFN- γ , and MMPs, and generate reactive oxygen species. Accumulation of free cholesterol within the plaque strongly induces apoptosis of macrophage-derived foam cells and promotes the enlarging of the necrotic core. Even today, one of the errors in the approach to the study of carotid atherosclerotic pathology remains perhaps considering plaque vulnerability and rupture as a one-dimensional process. Talking about a vulnerable patient rather than a vulnerable plaque helped to broaden the scope of observation and allow an all-around study that also took into consideration the molecular mechanisms underlying the risk associated with potential plaque rupture. Current evidence demonstrates that plaque vulnerability phenotypes are multiple and heterogeneous and are associated with as many highly complex molecular pathways that determine the activation of an immune-mediated cascade that culminates in thromboinflammation. We do not know if there are as many types of vulnerable patients whose exaltation of specific signals and consequent activation of specific pathways induce the manifestation of plaque vulnerability towards one phenotype rather than another. In the future, research will have to focus on the individual molecular triggers underlying the expression of individual plaque vulnerability which may represent diagnostic markers and potential therapeutic targets towards a more personalized medicine.

Abbreviations

ABC: ATP binding cassette; ACAT: acylcoenzyme A:cholesterol acyltransferase; ACE/Ang II: angiotensin-converting enzyme/angiotensin II; ADAMTS: A Disintegrin and metalloproteinase with Thrombospondin motifs; ApoE: Apolipoprotein E; bFGF: Basic fibroblast growth factor; BMP2: bone morphogenetic protein 2; CXCL: Chemokine eC-X-C motif chemokine ligand; CRP: C-reactive protein; ECs: endothelial cells; ECM: extracellular matrix; FGF: fibroblast growth factor; HDL: high-density lipoprotein; hs-CRP: high-sensitivity C-reactive protein; IFN- γ : interferon-gamma; IL: Interleukin; IPH: intraplaque hemorrhage; LAAS: large-artery atherosclerosis; LDL: low-density lipoprotein; LNRC: lipid-rich necrotic core; Lp-PLA2: lipoprotein-associated phospholipase A2; LPA: Lysophosphatidic acid; MGP: matrix Gla protein; MES: microembolic signals; MMP: matrix metalloproteinase; NETs: Neutrophil extracellular traps; NLRP3: NOD-like receptor family pyrin domain containing 3; OPN: osteopontin; OPG: osteoprotegerin; OX-LDL: oxidized low-density lipoprotein; GP: platelet glycoprotein; PDGF: platelet-derived growth factor; PMPs: platelets-derived microparticles; RANKL: receptor activator of NF- κ B ligand; RBCs: red blood cells; TCD: transcranial Doppler; TF: tissue factor; TGF: transforming growth factor; TOAST: trial of ORG 10172 in the acute stroke treatment; TSG-6: Tumor Necrosis Factor-Stimulated Gene-6; TNF: *Tumor necrosis factor*; VEGF: vascular endothelial growth factor; VSMC: vascular smooth muscle cell; vWF: von Willebrand factor.

Author Contributions: Conceptualization, G.M.; methodology, G.M. and M.G.B.; validation, A.T., and G.M.; formal analysis, X.X.; investigation, C.P., M.P., M.C., A.R.P., E.C.; resources, G.R.; data curation, M.G.B.; writing—original draft preparation, C.P., M.G.B., G.M. and A.R.P.; writing—review and editing, E.C., M.P. and M.C.; visualization, A.T.; supervision, G.M. All authors have read and agreed to the published version of the manuscript.”

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Katan, M.; Luft, A. Global burden of stroke. *Seminars in Neurol.* 2018, 38, 208–11, DOI: 10.1055/s-0038-1649503.
2. Virmani, R.; Burke, A.P.; Kolodgie, F.D.; Farb, A. Vulnerable plaque: the pathology of unstable coronary lesions. *J Interv Cardiol.* 2002, 15, 439–46, DOI: 10.1111/j.1540-8183.2002.tb01087.x.
3. Willey, J.Z.; Pasterkamp G. The Role of the Vulnerable Carotid Plaque in Embolic Stroke of Unknown Source. *J. Am. Coll. Cardiol.* 2022, 79, 2200–2202, DOI: 10.1016/j.jacc.2022.04.004
4. Blann, A.D.; Farrel A.; Picton A.; McCollum C.N. Relationship between endothelial cell markers and arterial stenosis in peripheral and carotid artery disease. *Thromb Res.* 2000, 97, 209–216, DOI: 10.1016/s0049-3848(99)00156-5
5. Saba, L.; Saam, T.; Jäger, H.R.; Yuan, C.; Hatsukami, T.S.; Saloner, D.; Wassrman, B.A.; Bonati, L. H.; Wintermark, M. Imaging biomarkers of vulnerable carotid plaques for stroke risk prediction and their potential clinical implications. *Lancet Neurol.* 2019, 18, 559–572, DOI: 10.1016/S1474-4422(19)30035-3
6. Wang, Y.; Qiu, J.; Luo, S.; Xie, X.; Zheng, Y.; Zhang, K.; Ye, Z.; Liu, W.; Gregersen, H.; Wang, G. High shear stress induces atherosclerotic vulnerable plaque formation through angiogenesis. *Regen Biomater.* 2016, 3, 257–267, DOI: 10.1093/rb/rbw021
7. Kashiwazaki, D.; Shiraishi, K.; Yamamoto, S.; Kamo, T.; Uchinom H.; Saito, H.; Akioka, N.; Kuwayama, N.; Noguchi, K.; Kuroda, S. Efficacy of Carotid Endarterectomy for Mild (<50%) Symptomatic Carotid Stenosis with Unstable Plaque. *World Neurosurg.* 2019, 121, e60–e69, DOI: 10.1016/j.wneu.2018.09.013
8. Naghavi, M.; Libby, P.; Falk, E.; Casscells, S. W.; Litovsky, S.; Rumberger, J.; Badimon, J. J.; Stfanadis, C.; Moreno, P.; Pasterkamp, G.; et al. From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: Part I. *Circulation.* 2003, 108, 1664–72, DOI: 10.1161/01.CIR.0000087480.94275.97

9. Miceli, G.; Rizzo, G.; Basso, M. G.; Cocciola, E.; Pennacchio, A. R.; Pintus, C.; Tuttolomondo, A. Artificial Intelligence in Symptomatic Carotid Plaque Detection: A Narrative Review. *Appl. Sci.* 2023,13, 4321, DOI: 10.3390/app13074321
10. Lovett, J.K.; Gallagher, P.J.; Hands, L.J.; Walton, J.; Rothwell, P. M. Histological correlates of carotid plaque surface morphology on lumen contrast imaging. *Circulation.* 2004, 110, 2190-7, DOI: 10.1161/01.CIR.0000144307.82502.32
11. Miceli, G.; Basso, M. G.; Rizzo, G.; Pintus, C.; Cocciola, E.; Pennacchio, A. R.; Tuttolomondo, A. Artificial Intelligence in Acute Ischemic Stroke Subtypes According to Toast Classification: A Comprehensive Narrative Review. *Biomedicines.* 2023, 11(4):1138. DOI: 10.3390/biomedicines11041138
12. Jander, S.; Sitzler, M.; Schumann, R.; Schroeter, M.; Siebler, M.; Steinmetz, H.; Stoll, G. Inflammation in high-grade carotid stenosis. A possible role for macrophages and T cells in plaque de-stabilization. *Stroke.* 1998, 29, 1625–1630, DOI: 10.1161/01.str.29.8.1625
13. Tomas, L.; Edsfieldt, A.; Mollet, I. G.; Matic, L. P.; Prehn, C.; Adamski, J.; Paulsson-Berne, G.; Hedin, U.; Nilsson, J.; Bengtsson, E.; et al. Altered metabolism distinguishes high-risk from stable carotid atherosclerotic plaques. *Eur Heart J.* 2018, 39, 2301–2310, DOI: 10.1093/eurheartj/ehy124
14. Tengryd, C.; Nielsen, S. H.; Cavallera, M.; Bengtsson, E.; Genovese, F.; Karsdal, M.; Dunér, P.; Orholm-Melander, M.; Nilsson, J.; Esfeldt, A.; et al. The proteoglycan mimecan is associated with carotid plaque vulnerability and increased risk of future cardiovascular death. *Atherosclerosis.* 2020, 313, 88–95, DOI: 10.1016/j.atherosclerosis.2020.09.011
15. Jinnouchi, H.; Guo, L.; Sakamoto, A.; Torii, S.; Sato, Y.; Cornelissen, A.; Kuntz, S.; Paek, K. H.; Fernandez, R.; Fuller, D.; et al. Diversity of macrophage phenotypes and responses in atherosclerosis. *Cell Mol Life sciences: CMLS.* 2020, 77(10),1919–32, DOI: 10.1007/s00018-019-03371-3
16. Murray PJ. Macrophage polarization. *Annu Rev Physiol.* 2017, 79, 541–66, DOI: 10.1146/annurev-physiol-022516-034339
17. Shirai, T.; Hilhorst, M.; Harrison, D.G.; Goronzy, J. J.; Wayand, C. M. Macrophages in vascular inflammation—from atherosclerosis to vasculitis. *Autoimmunity.* 2015, 48(3),139–51, DOI: [10.3109/08916934.2015.1027815](https://doi.org/10.3109/08916934.2015.1027815)
18. Arabpour, M.; Saghazadeh, A.; Rezaei, N. Anti-inflammatory and M2 macrophage polarization-promoting effect of mesenchymal stem cell-derived exosomes. *Int Immunopharmac.* 2021, 97, 107823, DOI: 10.1016/j.intimp.2021.107823
19. Jetten, N.; Verbruggen, S.; Gijbels, M. J.; Post, M. J.; De Winther, M. P. J.; Donners, M. M. P. C. Anti-inflammatory M2, but not pro-inflammatory M1 macrophages promote angiogenesis in vivo. *Angiogenesis.* 2014, 17(1), 109–18, DOI: 10.1007/s10456-013-9381-6
20. Finn, A. V.; Nakano, M.; Polavarapu, R.; Karmali, V.; Saeed, O.; Zhao, X. Q.; Yazdani, S.; Otsuka, F.; Davis, T.; Habib, A.; et al. Hemoglobin directs macrophage differentiation and prevents foam cell formation in human atherosclerotic plaques. *J Am Coll Cardiol.* 2012, 59(2), 166–77, DOI: 10.1016/j.jacc.2011.10.852
21. Boyle, J. J.; Johns, M.; Kampfer, T.; Nguyen, T.; Game, L.; Schaer, D. J.; Mason, J. C.; Haskard, D. O. Activating transcription factor 1 directs mhem atheroprotective macrophages through coordinated iron handling and foam cell protection. *Circ Res.* 2012, 110(1), 20–33, DOI: 10.1161/CIRCRESAHA.111.247577
22. Kadl, A.; Meher, A. K.; Sharma, P. R.; Lee, M. Y.; Doran, A. C.; Johnstone, S. R.; Elliott, M. R.; Gruber, F.; Han, J.; Chen, W.; et al. Identification of a novel macrophage phenotype that develops in response to atherogenic phospholipids via Nrf2. *Circ Res.* 2010, 107(6), 737–46, DOI: 10.1161/CIRCRESAHA.109.215715
23. Erbel, C.; Tyka, M.; Helmes, C. M.; Akhavanpoor, M.; Rupp, G.; Domschke, G.; Linden, F.; Wolf, A.; Doesch, A.; Lasitschka, F.; et al. CXCL4-induced plaque macrophages can be specifically identified by co-expression of MMP7+S100A8+ in vitro and in vivo. *Innate Immun.* 2015, 21(3), 255–65, DOI: 10.1177/1753425914526461
24. Zhao, J.; Ling, L.; Zhu, W.; Ying, T.; Yu, T.; Sun, M.; Zhu, X.; Du, Y.; Zhang, L. M1/M2 re-polarization of kaempferol biomimetic NPs in anti-inflammatory therapy of atherosclerosis. *J Control Release.* 2023, 353, 1068–83, DOI: [10.1016/j.jconrel.2022.12.041](https://doi.org/10.1016/j.jconrel.2022.12.041)
25. Bisgaard, L. S.; Mogensen, C. K.; Rosendahl, A.; Cucak, H.; Nielsen, L. B.; Rasmussen, S. E.; Pedersen, T. X. Bone marrow-derived and peritoneal macrophages have different inflammatory response to oxLDL and M1/M2 marker expression - implications for atherosclerosis research. *Sci Rep.* 2016, 6, 35234, DOI:10.1038/srep35234
26. Stöger, J.L.; Gijbels, M. J.; VanderVelden S.; Manca, M.; VanderLoos, C. M.; Biessen, E. A. L.; Daemen, M. J. A. P.; Lutgens, E.; DeWinther, M. P. J. Distribution of macrophage polarization markers in human atherosclerosis. *Atherosclerosis.* 2012, 225(2), 461–8, DOI: [10.1016/j.atherosclerosis.2012.09.013](https://doi.org/10.1016/j.atherosclerosis.2012.09.013)
27. Khallou-Laschet, J.; Varthaman, A.; Fornasa, G.; Compain, C.; Gaston, A. T.; Clement, M.; Dussiot, M.; Levillain, O.; Dubois, S. G.; Nicoletti, A.; et al. Macrophage plasticity in experimental atherosclerosis. *PLoS One.* 2010, 5(1), e8852, DOI: [10.1371/journal.pone.0008852](https://doi.org/10.1371/journal.pone.0008852)

28. Vinchi, F.; Porto, G.; Simmelbauer, A.; Altamura, S.; Passos, S. T.; Garbowski, M.; Silva, A. M. N.; Spaich, S.; Seide, S. E.; Sparla, R.; et al. Atherosclerosis is aggravated by iron overload and ameliorated by dietary and pharmacological iron restriction. *Eur Heart J.* 2020, 41(28), 2681–95, DOI: [10.1093/eurheartj/ehz112](https://doi.org/10.1093/eurheartj/ehz112)
29. Du, J.; Wang, T.; Li, Y.; Zhou, Y.; Wang, X.; Yu, X.; Ren, X.; An, Y.; Wu, Y.; Sun, W.; et al. DHA inhibits proliferation and induces ferroptosis of leukemia cells through autophagy dependent degradation of ferritin. *Free Radic Biol Med.* 2019, 131, 356–69, DOI: [10.1016/j.freeradbiomed.2018.12.011](https://doi.org/10.1016/j.freeradbiomed.2018.12.011)
30. Jiao, Y.; Qin, Y.; Zhang, Z.; Zhang, H.; Liu, H.; Li, C. Early identification of carotid vulnerable plaque in asymptomatic patients. *BMC Cardiovasc Disord.* 2020, 20, 429, DOI: [10.1186/s12872-020-01709-5](https://doi.org/10.1186/s12872-020-01709-5)
31. Butler, W. T. The nature and significance of osteopontin. *Connect Tissue Res.* 1989, 23(2–3), 123–136, DOI: [10.3109/03008208909002412](https://doi.org/10.3109/03008208909002412)
32. Kahles, F.; Findeisen, H. M.; Bruemmer, D. Osteopontin: a novel regulator at the cross roads of inflammation, obesity and diabetes. *Mol Metab.* 2014, 3(4), 384–393, DOI: [10.1016/j.molmet.2014.03.004](https://doi.org/10.1016/j.molmet.2014.03.004)
33. Strobescu-Ciobanu, C.; Giusca, S. E.; Caruntu, I. D.; Amalinei, C.; Rusu, A.; Cojocaru, E.; Popa, R. F.; Lupasacu, C. D. Osteopontin and osteoprotegerin in atherosclerotic plaque – are they significant markers of plaque vulnerability? *Rom J Morphol Embryol.* 2020, 61(3), 793–801, DOI: [10.47162/RJME.61.3.17](https://doi.org/10.47162/RJME.61.3.17)
34. Schönbeck, U.; Libby, P. The CD40/CD154 receptor/ligand dyad. *Cell Mol Life Sci.* 2001, 58, 4–43, DOI: [10.1007/pl00000776](https://doi.org/10.1007/pl00000776)
35. Schönbeck, U.; Libby, P. CD40 signaling and plaque instability. *Circ Res.* 2001, 89, 1092–103, DOI: [10.1161/hh2401.101272](https://doi.org/10.1161/hh2401.101272)
36. Wang, J. H.; Zhang, Y. W.; Zhang, B. Q.; Ding, S.; Wang, Z. K.; Wu, T.; Wang, J. CD40 ligand as a potential biomarker for atherosclerotic instability. *Neurol Res.* 2013, 35(7), 693–700, DOI: [10.1179/1743132813Y.0000000190](https://doi.org/10.1179/1743132813Y.0000000190)
37. Zhang, Y.; Tang, Z. H.; Ren, Z.; Qu, S. L.; Liu, M. H.; Liu, L. S.; Jiang, Z. S. Hydrogen sulfide, the next potent preventive and therapeutic agent in aging and age-associated diseases. *Mol Cell Biol.* 2013, 33(6), 1104–1113, DOI: [10.1128/MCB.01215-12](https://doi.org/10.1128/MCB.01215-12)
38. Ding, Z.; Wang, X.; Schnackenberg, L.; Khaidakov, M.; Liu, S.; Singla, S.; Dai, Y.; Metha, J. L. Regulation of autophagy and apoptosis in response to ox-LDL in vascular smooth muscle cells, and the modulatory effects of the microRNA hsa-let-7 g. *Int J Cardiol.* 2013, 168(2), 1378–1385, DOI: [10.1016/j.ijcard.2012.12.045](https://doi.org/10.1016/j.ijcard.2012.12.045)
39. Garcia, B. A.; Ruiz, C.; Sabin, J. A.; Matas, M. High-sensitivity C-reactive protein in high-grade carotid stenosis: risk marker for unstable carotid plaque. *Journal of Vascular Surgery.* 2003, 38(5), 1018–1024, DOI: [10.1016/s0741-5214\(03\)00709-2](https://doi.org/10.1016/s0741-5214(03)00709-2)
40. Tay, C.; Liu, Y. H.; Hosseini, H.; Kanellakis, P.; Cao, A.; Peter, K.; Tipping, P.; Bobik, A.; Toh, B. H.; Kyaw, T. B-cell-specific depletion of tumour necrosis factor alpha inhibits atherosclerosis development and plaque vulnerability to rupture by reducing cell death and inflammation. *Cardiovasc Res.* 2016, 111, 385–397, DOI: [10.1093/cvr/cvw186](https://doi.org/10.1093/cvr/cvw186)
41. Xiong, X. D.; Xiong, W. D.; Xiong, S. S.; Chen, G. H. Research Progress on the Risk Factors and Outcomes of Human Carotid Atherosclerotic Plaques. *Chin Med J.* 2017, 130, 722–729, DOI: [10.4103/0366-6999.201598](https://doi.org/10.4103/0366-6999.201598)
42. Erbel, C.; Dengler, T. J.; Wangler, S.; Lasitschka, F.; Bea, F.; Wambsganss, N.; Hakimi, M.; Bockler, D.; Katus, H. A.; Gleissner, C. A. Expression of IL-17A in human atherosclerotic lesions is associated with increased inflammation and plaque vulnerability. *Basic Res Cardiol.* 2011, 106, 125–134, DOI: [10.1007/s00395-010-0135-y](https://doi.org/10.1007/s00395-010-0135-y)
43. Edsfeldt, A.; Nitulescu, M.; Grufman, H.; Gronberg, C.; Persson, A.; Nilsson, M.; Persson, M.; Björkbacka, H.; Gonçalves, I. Soluble urokinase plasminogen activator receptor is associated with inflammation in the vulnerable human atherosclerotic plaque. *Stroke.* 2012, 43, 3305–3312, DOI: [10.1161/STROKEAHA.112.664094](https://doi.org/10.1161/STROKEAHA.112.664094)
44. Abbas, A.; Aukrust, P.; Dahl, T. B.; Bjerkeli, V.; Sagen, E. B. L.; Michelsen, A.; Russel, D.; Krohg-Sørensen, K.; Holm, S.; Skjelland, M.; et al. High levels of S100A12 are associated with recent plaque symptomatology in patients with carotid atherosclerosis. *Stroke.* 2012, 43, 1347–1353, DOI: [10.1161/STROKEAHA.111.642256](https://doi.org/10.1161/STROKEAHA.111.642256)
45. Yuan, X.; Shen, G.; Hu, Q.; Xiao, H.; Qin, X. The Circulating Biomarker Fractalkine and Platelet-Derived Growth Factor BB are Correlated with Carotid Plaque Vulnerability Assessed by Computed Tomography Angiography. *Journal of stroke and cerebrovascular diseases.* 2022, 31(5), 1052–3057, DOI: [10.1016/j.jstrokecerebrovasdis.2022.106422](https://doi.org/10.1016/j.jstrokecerebrovasdis.2022.106422)
46. Yang, T.; Guo, L.; Chen, L.; Li, J.; Li, Q.; Pi, Y.; Zhu, J.; Zhang, L. A novel role of FKN/ CX3CR1 in promoting osteogenic transformation of VSMCs and atherosclerotic calcification. *Cell Calcium.* 2020, 91, 102265, DOI: [10.1016/j.ceca.2020.102265](https://doi.org/10.1016/j.ceca.2020.102265)
47. Ledard, N.; Lbioz, A.; Blondeau, B.; Babiak, M.; Moulin, C.; Vallin, B.; Guillas, I.; Mateo, V.; Jumeau, C.; Blirando, K.; et al. Slug, a cancer-related transcription factor, is involved in vascular smooth muscle cell transdifferentiation induced by platelet-derived growth factor-BB during atherosclerosis. *J mA Heart Asoc.* 2020, 9(2), 014276, DOI: [10.1161/JAHA.119.014276](https://doi.org/10.1161/JAHA.119.014276)
48. Müller, A.; Krämer, S. D.; Meletta, R.; Beck, K.; Selinova, S. V.; Rancic, Z.; Kaufmann, P. A.; Vos, B.; Meding, J.; Stellfeld, T.; et al. Gene expression levels of matrix metalloproteinases in human atherosclerotic

- plaques and evaluation of radiolabeled inhibitors as imaging agents for plaque vulnerability. *Nucl Med Biol.* 2014, 41, 562–569, DOI: [10.1016/j.nucmedbio.2014.04.085](https://doi.org/10.1016/j.nucmedbio.2014.04.085)
49. Wang, N.; Zhang, X.; Ma, Z.; Niu, J.; Ma, S.; Wenjie, W.; Chen, J. Combination of tanshinone IIA and astragaloside IV attenuate atherosclerotic plaque vulnerability in ApoE(–/–) mice by activating PI3K/AKT signaling and suppressing TLR4/NF-κB signaling. *Biomed Pharmacother Biomed Pharmacother.* 2020, 123, 109729, DOI: [10.1016/j.biopha.2019.109729](https://doi.org/10.1016/j.biopha.2019.109729)
 50. Loftus, I. M.; Naylor, A. R.; Goodall, S.; Jones, L.; Bell, P. R.; Thompson, M. M. Increased matrix metalloproteinase-9 activity in unstable carotid plaques. A potential role in acute plaque disruption. *Stroke.* 2000, 31(1), 40-7, DOI: [10.1161/01.str.31.1.40](https://doi.org/10.1161/01.str.31.1.40)
 51. Johnson, J. L.; Devel, L.; Czarny, B.; George, S. J.; Jackson, C. L.; Rogakos, V.; Beau, F.; Yiotakis, A.; Newby, A. C.; Dive, V. A selective matrix metalloproteinase-12 inhibitor retards atherosclerotic plaque development in apolipoprotein E-knockout mice. *Arterioscler Thromb Vasc Biol.* 2011, 31, 528–535, DOI: [10.1161/ATVBAHA.110.219147](https://doi.org/10.1161/ATVBAHA.110.219147)
 52. Pelisek, J.; Deutsch, L.; Ansel, A.; Pongratz, J.; Stadlbauer, T.; Gebhard, H.; Matevossian, E.; Eckstein, H. H. Expression of a metalloproteinase family of ADAMTS in human vulnerable carotid lesions. *J Cardiovasc Med.* 2017, 18, 10–18, DOI: [10.2459/JCM.0000000000000254](https://doi.org/10.2459/JCM.0000000000000254)
 53. Ashlin, T. G.; Kwan, A. P.; Ramji, D. P. Regulation of ADAMTS-1, -4 and -5 expression in human macrophages: Differential regulation by key cytokines implicated in atherosclerosis and novel synergism between TL1A and IL-17. *Cytokine.* 2013, 64, 234–242, DOI: [10.2459/JCM.0000000000000254](https://doi.org/10.2459/JCM.0000000000000254)
 54. Dong, H.; Du, T.; Premaratne, S.; Zhao, C. X.; Tian, Q.; Li, Y.; Yan, S.; Zhang, W. W. Relationship between ADAMTS4 and carotid atherosclerotic plaque vulnerability in humans. *J Vasc Surg.* 2018, 67, 1120–1126, DOI: [10.1016/j.jvs.2017.08.075](https://doi.org/10.1016/j.jvs.2017.08.075)
 55. Demircan, K.; Topcu, V.; Takigawa, T.; Akyol, S.; Yonezawa, T.; Ozturk, G.; Ugurcu, V.; Hasgul, R.; Yigitoglu, M. R.; Akyol, O.; et al. ADAMTS4 and ADAMTS5 knockout mice are protected from versican but not aggrecan or brevican proteolysis during spinal cord injury. *Biomed Res Int.* 2014, 2014, 693746, DOI: [10.1155/2014/693746](https://doi.org/10.1155/2014/693746)
 56. Pothineni, N. V. K.; Karathanasis, S. K.; Ding, Z.; Arulandu, A.; Varughese, K. I.; Metha, J. L. LOX-1 in Atherosclerosis and Myocardial Ischemia: Biology, Genetics, and Modulation. *J Am Coll Cardiol.* 2017, 69, 2759–2768, DOI: [10.1016/j.jacc.2017.04.010](https://doi.org/10.1016/j.jacc.2017.04.010)
 57. Jin, P.; Cong, S. LOX-1 and atherosclerotic-related diseases. *Clin Chim Acta Int J Clin Chem.* 2019, 491, 24–29, DOI: [10.1016/j.cca.2019.01.006](https://doi.org/10.1016/j.cca.2019.01.006)
 58. Norata, G. D.; Raselli, S.; Grigore, L.; Garlaschelli, K.; Vianello, D.; Bertocco, S.; Zambon, A.; Catapano, A. L. Small dense LDL and VLDL predict common carotid artery IMT and elicit an inflammatory response in peripheral blood mononuclear and endothelial cells. *Atherosclerosis.* 2009, 206, 556–562, DOI: [10.1016/j.atherosclerosis.2009.03.017](https://doi.org/10.1016/j.atherosclerosis.2009.03.017)
 59. Zambon, A.; Puato, M.; Faggini, E.; Faggini, E.; Grego, F.; Rattazzi, M.; Pauletto, P. Lipoprotein remnants and dense LDL are associated with features of unstable carotid plaque: A flag for non-HDL-C. *Atherosclerosis.* 2013, 230(1), 106–9, DOI: [10.1016/j.atherosclerosis.2013.06.024](https://doi.org/10.1016/j.atherosclerosis.2013.06.024)
 60. Shi, X.; Cai, H.; Wang, F.; Liu, R.; Xu, X.; Li, M.; Han, Y.; Yin, Q.; Ye, R.; Liu, X. Cholesterol Crystals are Associated with Carotid Plaque Vulnerability: An Optical Coherence Tomography Study. *J Stroke Cerebrovasc Dis.* 2020, 29, 104579, DOI: [10.1016/j.jstrokecerebrovasdis.2019.104579](https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.104579)
 61. Janoudi, A.; Shamoun, F. E.; Kalavakunta, J. K.; Abela, G. S. Cholesterol crystal-induced arterial inflammation and destabilization of atherosclerotic plaque. *Eur Heart J.* 2016, 37, 1959–1967, DOI: [10.1093/eurheartj/ehv653](https://doi.org/10.1093/eurheartj/ehv653)
 62. Huang, Y.; Liu, Q.; Xu, J.; Zhu, W.; Jiang, J.; Tang, L.; Chen, M. Contrast-enhanced ultrasound perfusion patterns and serum lipid signatures of vulnerable carotid artery plaque in predicting stroke: A cohort study of carotid stenosis in Chinese patients. *Clin Hemorheol Microcirc.* 2020, 75, 349–359, DOI: [10.3233/CH-190804](https://doi.org/10.3233/CH-190804)
 63. Seneviratne, A.; Hulsmans, M.; Holvoet, P.; Monaco, C. et al. Biomechanical factors and macrophages in plaque stability. *Cardiovasc Res.* 2013, 99, 284–93, DOI: [10.1093/cvr/cvt097](https://doi.org/10.1093/cvr/cvt097)
 64. Lee, R. T.; Kamm, R. D. Vascular mechanics for the cardiologist. *J Am Coll Cardiol.* 1994, 23, 1289–95, DOI: [10.1016/0735-1097\(94\)90369-7](https://doi.org/10.1016/0735-1097(94)90369-7)
 65. Otsuka, F.; Finn, A. V.; Yazdani, S. K.; Nakano, M.; Kolodgie, F. D.; Virmani, R. The importance of the endothelium in atherothrombosis and coronary stenting. *Nat Rev Cardiol.* 2012, 9, 439–53, DOI: [10.1038/nrcardio.2012.64](https://doi.org/10.1038/nrcardio.2012.64)
 66. Freynhofer, M. K.; Bruno, V.; Wojta, J.; Huber, K. The role of platelets in atherothrombotic events. *Curr Pharm Des.* 2012, 18, 5197–214, DOI: [10.2174/138161212803251899](https://doi.org/10.2174/138161212803251899)
 67. Miceli, G.; Basso, M. G.; Rizzo, G.; Pintus, C.; Tuttolomondo, A. The Role of the Coagulation System in Peripheral Arterial Disease: Interactions with the Arterial Wall and Its Vascular Microenvironment and Implications for Rational Therapies. *Int J Mol Sci.* 2022, 23(23), 14914, DOI: [10.3390/ijms232314914](https://doi.org/10.3390/ijms232314914)

68. Penz, S.; Reininger, A. J.; Brandl, R.; Goyal, P.; Rabie, T.; Bernlochner, I.; Rother, E.; Goetz, C.; Engelmann, B.; Smethurst, P. A.; et al. Human atheromatous plaques stimulate thrombus formation by activating platelet glycoprotein VI. *FASEB J.* 2005, 19, 898–909, DOI: [10.1096/fj.04-2748com](https://doi.org/10.1096/fj.04-2748com)
69. Freynhofer, M. K.; Bruno, V.; Wojta, J.; Huber, K. The role of platelets in atherothrombotic events. *Curr Pharm Des.* 2012, 18, 5197–214, DOI: [10.2174/138161212803251899](https://doi.org/10.2174/138161212803251899)
70. Badimon, J. J.; Lettino, M.; Toschi, V.; Fuster, V.; Berrozpe, M.; Chesebro, J. H.; Badimon, L. Local inhibition of tissue factor reduces the thrombogenicity of disrupted human atherosclerotic plaques: effects of tissue factor pathway inhibitor on plaque thrombogenicity under flow conditions. *Circulation.* 1999, 99, 1780–7, DOI: [10.1161/01.cir.99.14.1780](https://doi.org/10.1161/01.cir.99.14.1780)
71. Camino-Lopez, S.; Llorente-Cortes, V.; Sendra, J.; Badimon, L. Tissue factor induction by aggregated LDL depends on LDL receptor-related protein expression (LRP1) and Rho A trans-location in human vascular smooth muscle cells. *Cardiovasc Res.* 2007, 73, 208–16, DOI: [10.1016/j.cardiores.2006.10.017](https://doi.org/10.1016/j.cardiores.2006.10.017)
72. Hemker, H. C.; van Rijn, J. L.; Rosing, J.; van Dieijen, G.; Bevers, E. M.; Zwaal, R. F. Platelet membrane involvement in blood coagulation. *Blood Cells.* 1983, 9, 303–17, PMID: 6661560
73. Grad, E.; Danenberg, H. D. C-reactive protein and atherothrombosis: cause or effect? *Blood Rev.* 2013, 27, 23–9, DOI: [10.1016/j.blre.2012.12.001](https://doi.org/10.1016/j.blre.2012.12.001)
74. Ridker, P. M.; Everett, B. M.; Thuren, T.; MacFadyen, J. G.; Chang, W. H.; Ballantyne, C.; Fonseca, F.; Nicolau, J.; Koenig, W.; Anker, S. D.; et al. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med.* 2017, 377, 1119–1131, DOI: [10.1056/NEJMoa1707914](https://doi.org/10.1056/NEJMoa1707914)
75. Van Avondt, K.; Hartl, D. Mechanisms and disease relevance of neutrophil extracellular trap formation. *Eur J Clin Invest.* 2018, 48(Suppl 2), 12919, DOI: [10.1111/eci.12919](https://doi.org/10.1111/eci.12919)
76. Li, P.; Li, M.; Lindberg, M. R.; Kennett, M. J.; Xiong, N.; Wang, Y. PAD4 is essential for antibacterial innate immunity mediated neutrophil extracellular traps. *J Exp Mde.* 2010, 207, 1853–1862, DOI: [10.1084/jem.20100239](https://doi.org/10.1084/jem.20100239)
77. Shimonaga, K.; Matsushige, T.; Takahashi, H.; Hashimoto, Y.; Yoshiyama, M.; Ono, C.; Sakamoto, S. Peptidylarginine Deiminase 4 as a Possible Biomarker of Plaque Instability in Carotid Artery Stenosis. *Journal of stroke and cerebrovascular diseases: the official journal of National Stroke Association.* 2021, 30(7), 1052–3057, DOI: [10.1016/j.jstrokecerebrovasdis.2021.105816](https://doi.org/10.1016/j.jstrokecerebrovasdis.2021.105816)
78. Kennedy, M. A.; Barrera, G. C.; Nakamura, K.; Baldan, A.; Tarr, P.; Fishbein, M. C.; Frank, J.; Francone, O. L.; Edwards, P. A. ABCG1 Has a critical role in mediating cholesterol efflux to HDL and preventing cellular lipid accumulation. *Cell Metab.* 2005, 1, 121–131, DOI: [10.1016/j.cmet.2005.01.002](https://doi.org/10.1016/j.cmet.2005.01.002)
79. Wang, N.; Lan, D.; Chen, W.; Matsuura, F.; Tall, A. R. ATP-binding cassette transporters G1 and G4 mediate cellular cholesterol efflux to high-density lipoproteins. *Proc Natl Acad Sci.* 2004, 101, 9774–9779, DOI: [10.1073/pnas.0403506101](https://doi.org/10.1073/pnas.0403506101)
80. Wang, N.; Silver, D. L.; Costet, P.; Tall, A. R. Specific binding of ApoA-I, enhanced cholesterol efflux, and altered plasma membrane morphology in cells expressing ABC1. *J Biol Chem.* 2000, 275, 33053–33058, DOI: [10.1074/jbc.M005438200](https://doi.org/10.1074/jbc.M005438200)
81. Yalcinkaya, M.; Fotakis, P.; Liu, W.; Endo-Umeda, K.; Dou, H.; Abramowicz, S.; Xiao, T.; Libby, P.; Wang, N.; Tall, A. R.; et al. Cholesterol accumulation in macrophages drives NETosis in atherosclerotic plaques via IL-1 β secretion. *Cardiovascular Research.* 2023, 119, 969–981, DOI: [10.1093/cvr/cvac189](https://doi.org/10.1093/cvr/cvac189)
82. Yuan, J.; Usman, A.; Das, T.; Patterson, A. J.; Gillard, J. H.; Graves, M. J. Imaging Carotid Atherosclerosis Plaque Ulceration: Comparison of Advanced Imaging Modalities and Recent Developments. *AJNR Am J Neuroradiol.* 2017, 38(4), 664–671, DOI: [10.3174/ajnr.A5026](https://doi.org/10.3174/ajnr.A5026)
83. Rafailidis, V.; Chrysogonidis, I.; Tegos, T.; Kouskouras, K.; Charitanti-Kouridou, A. Imaging of the ulcerated carotid atherosclerotic plaque: a literature review. *Insights Imaging.* 2017, 8(2), 213–225, DOI: [10.1007/s13244-017-0543-8](https://doi.org/10.1007/s13244-017-0543-8)
84. Sayed, S.; Cockerill, G. W.; Torsney, E.; Poston, R.; Thompson, M. M.; Loftus, I. M. Elevated tissue expression of thrombomodulatory factors correlates with acute symptomatic carotid plaque phenotype. *Eur J Vasc Endovasc Surg.* 2009, 38(1), 20–5, DOI: [10.1016/j.ejvs.2009.02.020](https://doi.org/10.1016/j.ejvs.2009.02.020)
85. Fisher, M.; Paganini-Hill, A.; Martin, A.; Cosgrove, M.; Toole, J. F.; Barnett, H. J.; Norris, J. Carotid plaque pathology: thrombosis, ulceration, and stroke pathogenesis. *Stroke.* 2005, 36(2), 253–7, DOI: [10.1161/01.STR.0000152336.71224.21](https://doi.org/10.1161/01.STR.0000152336.71224.21)
86. Nieddu, G.; Formato, M.; Lepedda, A. J. Searching for Atherosclerosis Biomarkers by Proteomics: A Focus on Lesion Pathogenesis and Vulnerability. *Int J Mol Sci.* 2023, 24(20), 15175, DOI: [10.3390/ijms242015175](https://doi.org/10.3390/ijms242015175)
87. Martin-Ventura, J. L.; Duran, M. C.; Blanco-Colio L. M.; Meilhac, O.; Leclercq, A.; Michel, J. B.; Jensen, O. N.; Hernandez-Merida, S.; Tuñón, J.; Vivanco, F.; et al. Identification by a differential proteomic approach of heat shock protein 27 as a potential marker of atherosclerosis. *Circulation.* 2004, 110(15), 2216–9, DOI: [10.1161/01.CIR.0000136814.87170.B1](https://doi.org/10.1161/01.CIR.0000136814.87170.B1)
88. Rocchiccioli, S.; Pelosi, G.; Rosini, S.; Marconi, M.; Viglione, F.; Citti, L.; Ferrari, M.; Trivella, M. G.; Cecchetti, A. Secreted proteins from carotid endarterectomy: an untargeted approach to disclose molecular clues of plaque progression. *J Transl Med.* 2013, 11, 260, DOI: [10.1186/1479-5876-11-260](https://doi.org/10.1186/1479-5876-11-260)

89. van Dijk, A. C.; Donkel, S. J.; Zadi, T.; Sonneveld, M. A. H.; Schreuder, F. H. B. M.; Chohan, M. F.; Koudstaal, P. J.; Leebeek, F. W. G.; Saxena, R.; Hendrikse, J.; et al. Association between fibrinogen and fibrinogen γ' and atherosclerotic plaque morphology and composition in symptomatic carotid artery stenosis: Plaque-At-RISK study. *Thromb Res.* 2019, 177, 130-135, DOI: 10.1016/j.thromres.2019.02.030
90. Spencer, M. P.; Thomas, G. I.; Nicholls, S. C.; Sauvage, L. R. Detection of middle cerebral artery emboli during carotid endarterectomy using transcranial Doppler ultrasonography. *Stroke.* 1990, 21, 415e2, DOI: 10.1161/01.str.21.3.415
91. Puz, P.; Lasek-Bal, A.; Urbanek, T.; Kazibutowska, Z. Assessment of cerebral embolism and vascular reserve parameters in patients with carotid artery stenosis. *Neurol. Neurochir Pol.* 2016, 50, 356-362, DOI: 10.1016/j.pjnns.2016.06.004
92. Jiang, Y.; Jiang, S.; Feng, S.; Sun, D.; Zhuang, A.; Zeng, Q.; Zhang, Y.; Huang, H.; Nie, H.; Zhou, F. Microembolic signal monitoring of TOAST classified cerebral infarction patients. *Mol Med Rep.* 2013, 8, 1135-1142, DOI: 10.3892/mmr.2013.1609
93. Markus, H. S.; Droste, D. W.; Kaps, M.; Larrue, V.; Lees, K. R.; Siebler, M.; Ringelstein, E. B. Dual antiplatelet therapy with clopidogrel and aspirin in symptomatic carotid stenosis evaluated using doppler embolic signal detection: the clopidogrel and aspirin for reduction of emboli in symptomatic carotid stenosis (CARESS) trial. *Circulation.* 2005, 111, 2233-2240, DOI: 10.1161/01.CIR.0000163561.90680.1C
94. Markus, H. S.; Brown, M. M. Differentiation between different pathological cerebral embolic materials using transcranial Doppler in an in vitro model. *Stroke.* 1993, 24, 1-5, DOI: 10.1161/01.str.24.1.1
95. Droste, D. W.; Ritter, M.; Kemeny, V.; Schulte-Altedorneburg, G.; Ringelstein, E. B. Microembolus detections at follow-up in 19 patients with acute stroke: correlation with stroke aetiology and antithrombotic treatment. *Cerebrovasc Dis.* 2000, 10, 272-277, DOI: 10.1159/000016069
96. Ritter, M. A.; Jurk, K.; Schriek, C.; Nabavi, D.; Droste, D. W.; Kherel, B. E.; Ringelstein, E. B. Microembolic signals on transcranial Doppler ultrasound are correlated with platelet activation markers, but not with platelet-leukocyte associates: a study in patients with acute stroke and patients with asymptomatic carotid stenosis. *Neurol Res.* 2009, 31, 11-16, DOI: 10.1179/174313208X331590
97. Kinsella, J. A.; Tobin, W. O.; Tierney, S.; Feeley, T. M.; Egan, B.; Collins, D. R.; Coughlan, T.; O'Neill, D.; Harbison, J.; Madhavan, P.; et al. Increased platelet activation in early symptomatic vs. asymptomatic carotid stenosis and relationship with microembolic status: results from the Platelets and Carotid Stenosis Study. *J Thromb Haemost.* 2013, 11, 1407-1416, DOI: 10.1111/jth.12277
98. Murphy, S. J. X.; Lim, S. T.; Kinsella, J. A.; Murphy, D.; Enright, H. M.; McCabe, D. J. H.; HEIST study group. Increased platelet count and reticulated platelets in recently symptomatic versus asymptomatic carotid artery stenosis and in cerebral microembolic signal-negative patient subgroups: results from the HaEmostasis In carotid STenosis (HEIST) study. *J Neurol.* 2018, 265, 1037-1049, DOI: 10.1007/s00415-018-8797-8
99. Murphy, S. J. X.; Lim, S. T.; Kinsella, J. A.; Tierney, S.; Egan, B.; Feeley, T. M.; Murphy, S. M.; Walsh, R. A.; Collins, D. R.; Coughlan, T.; et al. Increased leucocyte-platelet complex formation in recently symptomatic versus asymptomatic carotid stenosis patients and in micro-emboli negative subgroups. *Thromb Haemost.* 2019, 119, 821-833, DOI: 10.1055/s-0039-1678666
100. Badimon, L.; Padro, T.; Vilahur, G. Atherosclerosis, platelets and thrombosis in acute ischaemic heart disease. *Eur Heart J Acute Cardiovasc Care.* 2012, 1, 60-74, DOI: 10.1177/2048872612441582
101. Wang, X.; Lin, W. H.; Zhao, Y. D.; Chen, X. Y.; Leung, T. W.; Chen, C.; Fu, J.; Markus, H.; Hao, Q.; Wong, K. A. S.; et al. The effectiveness of dual antiplatelet treatment in acute ischemic stroke patients with intracranial arterial stenosis: a subgroup analysis of CLAIR study. *Int J Stroke.* 2013, 8, 663-668, DOI: 10.1111/j.1747-4949.2012.00828.x
102. Goertler, M.; Baeumer, M.; Kross, R.; Blaser, T.; Lutze, G.; Jost, S.; Wallesch, C. W. Rapid decline of cerebral microemboli of arterial origin after intravenous acetylsalicylic acid. *Stroke.* 1999, 30, 66-69, DOI: 10.1161/01.str.30.1.66
103. Libby, P. Inflammation in atherosclerosis. *Nature.* 2002, 420, 868-874, DOI: 10.1038/nature01323
104. Lukasik, M.; Rozalski, M.; Luzak, B.; Michalak, M.; Ambrosius, W.; Watala, C.; Kosubski, W. Enhanced platelet-derived microparticle formation is associated with carotid atherosclerosis in convalescent stroke patients. *Platelets.* 2013, 24(1), 63-70, DOI: 10.3109/09537104.2011.654292
105. Schiro, A.; Wilkinson, F. L.; Weston, R.; Smyth, J. V.; Serracino-Inglott, F.; Alexander, M. Y. Elevated levels of endothelial-derived microparticles, and serum CXCL9 and SCGF- β are associated with unstable asymptomatic carotid plaques. *Sci Rep.* 2015, 5, 16658, DOI: 10.1038/srep16658
106. Kandiyil, N.; MacSweeney, S. T.; Heptinstall, S.; May, J.; Foz, S. C.; Auer, D. P. Circulating Microparticles in Patients with Symptomatic Carotid Disease Are Related to Embolic Plaque Activity and Recent Cerebral Ischaemia. *Cerebrovasc Dis Extra.* 2019, 9, 9-18, DOI: 10.1159/000495942
107. Huang, Z. S.; Jeng, J. S.; Wang, C. H.; Yip, P. K.; Wu, T. H.; Lee, T. K. Correlations between peripheral differential leukocyte counts and carotid atherosclerosis in non-smokers. *Atherosclerosis.* 2001, 158, 431-436, DOI: 10.1016/s0021-9150(01)00445-2

108. Aronow, H. D.; Shishehbor, M.; Davis, D. A.; Katzan, I. L.; Bhatt, D. L.; Abou-Chebl, A.; Derk, K. W.; Whitlow, P. L.; Yadav, H. S. Leukocyte count predicts microembolic Doppler signals during carotid stenting: a link between inflammation and embolization. *Stroke*. 2005, 36, 1910–1914, DOI: [10.1161/01.STR.0000177610.33478.65](https://doi.org/10.1161/01.STR.0000177610.33478.65)
109. Nasr, N.; Ruidavets, J. B.; Arnal, J. F.; Sie, P.; Larrue, V. Association of neutrophil count with microembolization in patients with symptomatic carotid artery stenosis. *Atherosclerosis*. 2009, 207, 519–523, DOI: [10.1016/j.atherosclerosis.2009.05.003](https://doi.org/10.1016/j.atherosclerosis.2009.05.003)
110. Wuttge, D. M.; Zhou, X.; Sheikine, Y.; Wagsater, D.; Stemme, V.; Hedin, U.; Stemme, S.; Hansson, G. K.; Sirsjo, A. CXCL16/SR-PSOX is an interferon-gamma-regulated chemokine and scavenger receptor expressed in atherosclerotic lesions. *Arterioscler Thromb Vasc Biol*. 2004, 24, 750–755, DOI: [10.1161/01.ATV.0000124102.11472.36](https://doi.org/10.1161/01.ATV.0000124102.11472.36)
111. Minami, M.; Kume, N.; Shimaoka, T.; Kataoka, H.; Hayashida, K.; Akiyama, Y.; Nagata, I.; Ando, K.; Nobuyoshi, M.; Hanyuu, M.; et al., Expression of SR-PSOX, a novel cell-surface scavenger receptor for phosphatidylserine and oxidized LDL in human atherosclerotic lesions. *Arterioscler Thromb Vasc Biol*. 2001, 21, 1796–1800, DOI: [10.1161/hq1001.096652](https://doi.org/10.1161/hq1001.096652)
112. Ma, A.; Pan, X.; Xing, X.; Wu, M.; Wang, Y.; Ma, C. Elevation of serum CXCL16 level correlates well with atherosclerotic ischemic stroke. *Arch Med Sci*. 2014, 10, 47–52, DOI: [10.5114/aoms.2013.39200](https://doi.org/10.5114/aoms.2013.39200)
113. Ma, A.; Yang, S.; Wang, Y.; Wang, X.; Pan, X. Increase of serum CXCL16 level correlates well to microembolic signals in acute stroke patients with carotid artery stenosis. *Clin Chim Acta*. 2016, 460, 67–71, DOI: [10.1016/j.cca.2016.06.026](https://doi.org/10.1016/j.cca.2016.06.026)
114. Dong, L.; Ma, A.; Zhao, H.; Li, H.; Song, S.; Pan, X. Correlation between microembolic signal and immune inflammation in acute ischemic stroke. *Int J Cerebrovasc Dis*. 2015, 23, 677–681, ID: wpr-480503
115. Pini, R.; Faggioli, G. L.; Fittipaldi, S.; Pasquinnelli, G.; Tonon, C.; Beltrandi, E.; Mauro, R.; Stella, A. Inflammatory Mediators and Cerebral Embolism in Carotid Stenting: New Markers of Risk. *J ENDOVASC THER*. 2013, 20, 684–694, DOI: [10.1583/13-4354R.1](https://doi.org/10.1583/13-4354R.1)
116. Gerosa, C.; Cerrone, G.; Suri, J. S.; Aimola, V.; Cau, F.; Coni, P.; Piras, M.; Cau, R.; Balestrieri, A.; Scano, A.; et al. The human carotid atherosclerotic plaque: an observational review of histological scoring systems. *Eur Rev Med Pharmacol Sci*. 2023, 27(8), 3784–3792. DOI: [10.26355/eurrev_202304_32179](https://doi.org/10.26355/eurrev_202304_32179)
117. Virmani, R.; Burke, A. P.; Kolodgie, F. D.; Farb, A. Pathology of the thin-cap fibroatheroma: a type of vulnerable plaque. *J Interv Cardiol*. 2003, 16(3), 267–72, DOI: [10.1034/j.1600-0854.2003.8042.x](https://doi.org/10.1034/j.1600-0854.2003.8042.x)
118. Puig, N.; Jiménez-Xarrié, E.; Camps-Renom, P.; Benitez, S. Search for Reliable Circulating Biomarkers to Predict Carotid Plaque Vulnerability. *Int J Mol Sci*. 2020, 21(21), 8236. DOI: [10.3390/ijms21218236](https://doi.org/10.3390/ijms21218236)
119. Jebari-Benslaiman, S.; Galicia-García, U.; Larrea-Sebal, A.; Olaetxea, J. R.; Alloza, I.; Vandenbroeck, K.; Benito-Vicente, A.; Martin, C. Pathophysiology of Atherosclerosis. *Int J Mol Sci*. 2022, 23(6), 3346, DOI: [10.3390/ijms23063346](https://doi.org/10.3390/ijms23063346)
120. Watson, M. G.; Byrne, H. M.; Macaskill, C.; Myerscough, M. R. A two-phase model of early fibrous cap formation in atherosclerosis. *J Theor Biol*. 2018, 456, 123–36, DOI: [10.1016/j.jtbi.2018.08.010](https://doi.org/10.1016/j.jtbi.2018.08.010)
121. Sorokin, V.; Vickneson, K.; Kofidis, T.; Woo, C. C.; Lin, X. Y.; Foo, R.; Sanahan, C. M. Role of vascular smooth muscle cell plasticity and interactions in vessel wall inflammation. *Front Immunol*. 2020, 11, 599415, DOI: [10.3389/fimmu.2020.599415](https://doi.org/10.3389/fimmu.2020.599415)
122. Ulrich, V.; Rotllan, N.; Araldi, E.; Luciano, A.; Skroblin, P.; Abonnenc, M.; Perrotta, P.; Yin, X.; Bauer, A.; Leslie, K. L.; Zhang, P.; et al. Chronic miR-29 antagonism promotes favorable plaque remodeling in atherosclerotic mice. *EMBO Mol Med*. 2016, 8(6), 643–53, DOI: [10.15252/emmm.201506031](https://doi.org/10.15252/emmm.201506031)
123. Rekhter, M. D. Collagen synthesis in atherosclerosis: too much and not enough. *Cardiovasc Res*. 1999, 41(2), 376–84, DOI: [10.1016/s0008-6363\(98\)00321-6](https://doi.org/10.1016/s0008-6363(98)00321-6)
124. Badimon, L.; Vilahur, G. Thrombosis formation on atherosclerotic lesions and plaque rupture. *J Intern Med*. 2014, 276(6), 618–32, DOI: [10.1111/joim.12296](https://doi.org/10.1111/joim.12296)
125. Bentzon, J. F.; Otsuka, F.; Virmani, R.; Falk, E. Mechanisms of plaque formation and rupture. *Circ Res*. 2014, 114(12), 1852–66, DOI: [10.1161/CIRCRESAHA.114.302721](https://doi.org/10.1161/CIRCRESAHA.114.302721)
126. Bennett, M. R.; Sinha, S.; Owens, G. K. Vascular Smooth Muscle Cells in Atherosclerosis. *Circ Res*. 2016, 118(4), 692–702, DOI: [10.1161/CIRCRESAHA.115.306361](https://doi.org/10.1161/CIRCRESAHA.115.306361)
127. Simionescu, M.; Sima, A. V. Morphology of atherosclerotic lesions. *Inflammation and Atherosclerosis*. 2012, 19–37
128. Tabas, I.; Tall, A.; Accili, D. The impact of macrophage insulin resistance on advanced atherosclerotic plaque progression. *Circ Res*. 2010, 106(1), 58–67, DOI: [10.1161/CIRCRESAHA.109.208488](https://doi.org/10.1161/CIRCRESAHA.109.208488)
129. Ross, R.; Masuda, J.; Raines, E. W.; Gown, A. M.; Katsuda, S.; Sasahara, M.; Malden, L. T.; Masuko, H.; Sato, H. Localization of PDGF-B protein in macrophages in all phases of atherogenesis. *Science*. 1990, 248(4958), 1009–12. DOI: [10.1126/science.2343305](https://doi.org/10.1126/science.2343305)
130. Shimokado, K.; Raines, E. W.; Madtes, D. K.; Barrett, T. B.; Benditt, E. P.; Ross, R. A significant part of macrophage-derived growth factor consists of at least two forms of PDGF. *Cell*. 1985, 43(1), 277–86, DOI: [10.1016/0092-8674\(85\)90033-9](https://doi.org/10.1016/0092-8674(85)90033-9)

131. Yap, A. S.; Briehner, W. M.; Gumbiner, B. M. Molecular and functional analysis of cadherin-based adherens junctions. *Annu Rev Cell Dev Biol.* 1997, 13, 119-46, DOI: 10.1146/annurev.cellbio.13.1.119
132. Gilbertson-Beadling, S. K.; Fisher, C. A potential role for N-cadherin in mediating endothelial cell-smooth muscle cell interactions in the rat vasculature. *Lab Invest.* 1993, 69(2), 203-9, PMID: 8350598
133. Mann, J. M.; Davies, M. J. Vulnerable plaque. Relation of characteristics to degree of stenosis in human coronary arteries. *Circulation.* 1996, 94(5), 928-31, DOI: 10.1161/01.cir.94.5.928
134. Libby, P.; Sukhova, G.; Lee, R. T.; Galis, Z. S. Cytokines regulate vascular functions related to stability of the atherosclerotic plaque. *J Cardiovasc Pharmacol.* 1995, 25, 9-12, DOI: 10.1097/00005344-199500252-00003
135. Hansson, G. K.; Hellstrand, M.; Rymo, L.; Rubbia, L.; Gabbiani, G. Interferon gamma inhibits both proliferation and expression of differentiation-specific alpha-smooth muscle actin in arterial smooth muscle cells. *J Exp Med.* 1989, 170(5), 1595-608, DOI: 10.1084/jem.170.5.1595
136. Polyak, K. Negative regulation of cell growth by TGF beta. *Biochim Biophys Acta.* 1996, 242(3), 185-99, DOI: 10.1016/0304-419x(95)00009-5
137. Bennett, M. R.; Evan, G. I.; Newby, A. C. Deregulated expression of the c-myc oncogene abolishes inhibition of proliferation of rat vascular smooth muscle cells by serum reduction, interferon-gamma, heparin, and cyclic nucleotide analogues and induces apoptosis. *Circ Res.* 1994, 74(3), 525-36, DOI: 10.1161/01.res.74.3.525
138. Laitinen, M.; Zachary, I.; Breier, G.; Pakkanen, T.; Häkkinen, T.; Luoma, J.; Abedi, H.; Risau, W.; Soma, M.; Laakso, M.; et al. VEGF gene transfer reduces intimal thickening via increased production of nitric oxide in carotid arteries. *Hum Gene Ther.* 1997, 8(15), 1737-44, DOI: 10.1089/hum.1997.8.15-1737
139. Morioka, M.; Hamada, J.; Hashiguchi, A.; Hasegawa, Y.; Todaka, T.; Yano, S.; Kai, Y.; Miura, M.; Fujioka, S.; Ushio, Y. Contribution of angiotensin-converting enzyme and angiotensin II to ischemic stroke: their role in the formation of stable and unstable carotid atherosclerotic plaques. *Surg Neurol.* 2004, 62(4), 292-301, discussion 301-3, DOI: 10.1016/j.surneu.2004.02.027
140. Hao, H.; Iihara, K.; Ishibashi-Ueda, H.; Saito, F.; Hirota, S. Correlation of thin fibrous cap possessing adipophilin-positive macrophages and intraplaque hemorrhage with high clinical risk for carotid endarterectomy. *J Neurosurg.* 2011, 114(4), 1080-7, DOI: 10.3171/2010.8.JNS10423
141. Badacz, R.; Przewłocki, T.; Legutko, J.; Żmudka, K.; Kablak-Ziembicka, A. microRNAs Associated with Carotid Plaque Development and Vulnerability: The Clinician's Perspective. *Int J Mol Sci.* 2022, 23(24), 15645, DOI: 10.3390/ijms232415645
142. Faber, B. C.; Cleutjens, K. B.; Niessen, R. L.; Aarts, P. L.; Boon, W.; Greenberg, A. S.; Kitslaar, P. J.; TorDOIr, J. H.; Daemen, M. J. Identification of genes potentially involved in rupture of human atherosclerotic plaques. *Circ Res.* 2001, 89(6), 547-54, DOI: 10.1161/hh1801.096340
143. Bazan, H. A.; Brooks, A. J.; Vongbunyong, K.; Tee, C.; Douglas, H. F.; Klingenberg, N. C.; Woods, T. C. A pro-inflammatory and fibrous cap thinning transcriptome profile accompanies carotid plaque rupture leading to stroke. *Sci Rep.* 2022, 12, 13499, DOI:10.1038/s41598-022-17546-9
144. Bobryshev, Y. V.; Killingsworth, M. C.; Lord, R. S. A.; Grabs, A. J. Matrix vesicles in the fibrous cap of atherosclerotic plaque: possible contribution to plaque rupture. *Journal of Cellular and Molecular Medicine.* 2008, 12, 2073-2082, DOI:10.1111/j.1582-4934.2008.00230.x
145. _Sluimer, J. C.; Daemen, M. J. Novel concepts in atherogenesis: angiogenesis and hypoxia in atherosclerosis. *J Pathol.* 2009, 218(1), 7-29, DOI: 10.1002/path.2518
146. Roiniotis, J.; Dinh, H.; Masendycz, P.; Turner, A.; Elsegood, C. L.; Scholz, G. M.; Hamilton, J. A. Hypoxia prolongs monocyte/macrophage survival and enhanced glycolysis is associated with their maturation under aerobic conditions. *J Immunol.* 2009, 182(12), 7974-81, DOI: 10.4049/jimmunol.0804216
147. Nakano, D.; Hayashi, T.; Tazawa, N.; Yamashita, C.; Inamoto, S.; Okuda, N.; Mori, T.; Sohmiya, K.; Kitaura, Y.; Okada, Y.; et al. Chronic hypoxia accelerates the progression of atherosclerosis in apolipoprotein E-knockout mice. *Hypertens Res.* 2005, 28(10), 837-45, DOI: 10.1291/hypres.28.837
148. Hellings, W. E.; Peeters, W.; Moll, F. L.; Piers, S. R.; van Setten, J.; Van der Spek, P. J.; de Vries, J. P.; Seldenrijk, K. A.; De Bruin, P. C.; et al. Composition of carotid atherosclerotic plaque is associated with cardiovascular outcome: a prognostic study. *Circulation.* 2010, 121(17), 1941-50, DOI:10.1161/CIRCULATIONAHA.109.887497
149. van Hinsbergh, V. W.; Eringa, E. C.; Daemen, M. J. Neovascularization of the atherosclerotic plaque: interplay between atherosclerotic lesion, adventitia-derived microvessels and perivascular fat. *Curr Opin Lipidol.* 2015, 26(5), 405-11, DOI: 10.1097/MOL.0000000000000210
150. Xu, J.; Lu, X.; Shi, G. P. Vasa vasorum in atherosclerosis and clinical significance. *Int J Mol Sci.* 2015, 16(5), 11574-608, DOI: 10.3390/ijms160511574
151. Sluimer, J. C.; Kolodgie, F. D.; Bijnens, A. P.; Maxfield, K.; Pacheco, E.; Kutys, B.; Duimel, H.; Frederik, P. M.; van Hinsbergh, V. W.; Virmani, R.; et al. Thin-walled microvessels in human coronary atherosclerotic plaques show incomplete endothelial junctions relevance of compromised structural integrity for intraplaque microvascular leakage. *J Am Coll Cardiol.* 2009, 53(17), 1517-27, DOI: 10.1016/j.jacc.2008.12.056

152. Jeney, V.; Balla, G.; Balla, J. Red blood cell, hemoglobin and heme in the progression of atherosclerosis. *Front Physiol.* 2014, 5, 379, DOI: 10.3389/fphys.2014.00379
153. Nagy, J. A.; Dvorak, A. M.; Dvorak, H. F. Vascular hyperpermeability, angiogenesis, and stroma generation. *Cold Spring Harb Perspect Med.* 2012, 2(2), a006544, DOI: 10.1101/cshperspect.a006544
154. Melder, R. J.; Yuan, J.; Munn, L. L.; Jain, R. K. Erythrocytes enhance lymphocyte rolling and arrest in vivo. *Microvasc Res.* 2000, 59(2), 316-22, DOI: 10.1006/mvre.1999.2223
155. Michel, J. B.; Martin-Ventura, J. L.; Nicoletti, A.; Ho-Tin-Noé, B. Pathology of human plaque vulnerability: mechanisms and consequences of intraplaque haemorrhages. *Atherosclerosis.* 2014, 234(2), 311-9, DOI: 10.1016/j.atherosclerosis.2014.03.020
156. Leclercq, A.; Houard, X.; Philippe, M.; Ollivier, V.; Sebbag, U.; Meilhac, O.; Michel, J. B. Involvement of intraplaque hemorrhage in atherothrombosis evolution via neutrophil protease enrichment. *J Leukoc Biol.* 2007, 82(6), 1420-9, DOI: 10.1189/jlb.1106671
157. Schrijvers, D. M.; De Meyer, G. R.; Kockx, M. M.; Herman, A. G.; Martinet, W. Phagocytosis of apoptotic cells by macrophages is impaired in atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2005, 25(6), 1256-61, DOI: 10.1161/01.ATV.0000166517.18801.a7
158. Mao, Y.; Liu, X.; Song, Y.; Zhai, C.; Zhang, L. VEGF-A/VEGFR-2 and FGF-2/FGFR-1 but not PDGF-BB/PDGFR-beta play important roles in promoting immature and inflammatory intraplaque angiogenesis. *PLoS ONE.* 2018, 13, e0201395, DOI: 10.1371/journal.pone.0201395
159. Parma, L.; Baganha, F.; Quax, P. H. A.; de Vries, M. R. Plaque angiogenesis and intraplaque hemorrhage in atherosclerosis. *Eur. J. Pharmacol.* 2017, 816, 107-115, DOI: 10.1016/j.ejphar.2017.04.028
160. Bobryshev, Y. V.; Ivanova, E. A.; Chistiakov, D. A.; Nikiforov, N. G.; Orekhov, A. N. Macrophages and their role in atherosclerosis: pathophysiology and transcriptome analysis. *Biomed. Res. Int.* 2016, 2016, 9582430, DOI: 10.1155/2016/958243
161. Jackson, C. L.; Reidy, M. A. Basic fibroblast growth factor: its role in the control of smooth muscle cell migration. *Am. J. Pathol.* 1993, 143, 1024-1031, PMC1887069
162. Allahverdian, S.; Chaabane, C.; Boukais, K.; Francis, G. A.; Bochaton-Piallat, M. L. Smooth muscle cell fate and plasticity in atherosclerosis. *Cardiovasc. Res.* 2018, 114, 540-550, DOI: 10.1093/cvr/cvy022
163. Parma, L.; Peters, H. A. B.; Sluiter, T. J.; Simons, K. H.; Lazzari, P.; de Vries, M. R.; Quax, P. H. A. bFGF blockade reduces intraplaque angiogenesis and macrophage infiltration in atherosclerotic vein graft lesions in ApoE3*Leiden mice. *Sci Rep.* 2020, 10(1), 15968, DOI: 10.1038/s41598-020-72992-7
164. Virmani, R.; Narula, J.; Farb, A. When neoangiogenesis ricochets. *Am Heart J.* 1998, 136(6), 937-9, DOI: 10.1016/s0002-8703(98)70144-9
165. Jeziorska, M.; Woolley, D. E. Local neovascularization and cellular composition within vulnerable regions of atherosclerotic plaques of human carotid arteries. *J Pathol.* 1999, 188(2), 189-96, DOI: 10.1002/(SICI)1096-9896(199906)188:2<189::AID-PATH336>3.0.CO;2-N
166. Weis, S. M.; Cheresh, D. A. Pathophysiological consequences of VEGF-induced vascular permeability. *Nature.* 2005, 437(7058), 497-504, DOI: 10.1038/nature03987
167. Davies, M. J.; Thomas, A. C. Plaque fissuring—the cause of acute myocardial infarction, sudden ischaemic death, and crescendo angina. *Br Heart J.* 1985, 53, 363-373, PMC481773
168. Virmani, R.; Kolodgie, F. D.; Burke, A. P.; Finn, A. V.; Gold, H. K.; Tulenko, T. N.; Wrenn, S. P.; Narula, J. Atherosclerotic plaque progression and vulnerability to rupture: angiogenesis as a source of intraplaque hemorrhage. *Arterioscler Thromb Vasc Biol.* 2005, 25(10), 2054-61, DOI: 10.1161/01.ATV.0000178991.71605.18
169. Lampugnani, M. G.; Corada, M.; Caveda, L.; Breviario, F.; Ayalon, O.; Geiger, B.; Dejana, E. The molecular organization of endothelial cell-to-cell junctions: differential association of plakoglobin, beta-catenin, and alpha-catenin with vascular endothelial cadherin (VE-cadherin). *J Cell Biol.* 1995, 129(1), 203-17, DOI: 10.1083/jcb.129.1.203
170. Bobryshev, Y. V.; Cherian, S. M.; Inder, S. J.; Lord, R. S. Neovascular expression of VE-cadherin in human atherosclerotic arteries and its relation to intimal inflammation. *Cardiovasc Res.* 1999, 43(4), 1003-17, DOI: 10.1016/s0008-6363(99)00125-x
171. McCarthy, M. J.; Loftus, I. M.; Thompson, M. M.; Jones, L.; London, N. J.; Bell, P. R.; Naylor, A. R.; Brindle, N. P. Angiogenesis and the atherosclerotic carotid plaque: an association between symptomatology and plaque morphology. *J Vasc Surg.* 1999, 30(2), 261-8, DOI: 10.1016/s0741-5214(99)70136-9
172. Mofidi, R.; Crotty, T. B.; McCarthy, P.; Sheehan, S. J.; Mehigan, D.; Keaveny, T. V. Association between plaque instability, angiogenesis, and symptomatic carotid occlusive disease. *Br J Surg.* 2001, 88(7), 945-50, DOI: 10.1046/j.0007-1323.2001.01823.x
173. Baragetti A, Mattavelli E, Grigore L, Pellegatta F, Magni P, Catapano AL. Targeted Plasma Proteomics to Predict the Development of Carotid Plaques. *Stroke.* 2022 Sep;53(9):e411-e414. doi: 10.1161/STROKEAHA.122.038887. Epub 2022 Jul 1.
174. Matic, L. P.; Iglesias J. M.; Vesterlund, M.; Lengquist, M.; Hong, M. G.; Saieed, S.; Sanchez-Rivera, L.; Berg, M.; Razuvaev, A.; Kronqvist, M.; et al. Novel Multiomics Profiling of Human Carotid Atherosclerotic

- Plaques and Plasma Reveals Biliverdin Reductase B as a Marker of Intraplaque Hemorrhage. *JACC Basic Transl Sci.* 2018, 3(4), 464-480, DOI: 10.1016/j.jacbts.2018.04.001
175. Grosskreutz, C. L.; Anand-Apte, B.; Dupláa, C.; Quinn, T. P.; Terman, B. I.; Zetter, B.; D'Amore, P. A. Vascular endothelial growth factor-induced migration of vascular smooth muscle cells in vitro. *Microvasc Res.* 1999, 58(2), 128-36, DOI: 10.1006/mvre.1999.2171
 176. Carmeliet, P.; Collen, D. Molecular basis of angiogenesis. Role of VEGF and VE-cadherin. *Ann N Y Acad Sci.* 2000, 902, 249-62, discussion 262-4, DOI: 10.1111/j.1749-6632.2000.tb06320.x
 177. O'Brien, K. D.; McDonald, T. O.; Chait, A.; Allen, M. D.; Alpers, C. E. Neovascular expression of E-selectin, intercellular adhesion molecule-1, and vascular cell adhesion molecule-1 in human atherosclerosis and their relation to intimal leukocyte content. *Circulation.* 1996, 93(4), 672-82, DOI: 10.1161/01.cir.93.4.672
 178. Hu, S.; Liu, Y.; You, T.; Zhu, L. Semaphorin 7A Promotes VEGFA/VEGFR2-Mediated Angiogenesis and Intraplaque Neovascularization in *ApoE*^{-/-} Mice. *Front Physiol.* 2018, 9, 1718, DOI: 10.3389/fphys.2018.01718
 179. Suzuki, E.; Imuta, H.; Fujita, D.; Takahashi, M.; Oba, S.; Kiyosue, A.; Nishimatsu, H. Endogenous Interleukin-1 β Is Implicated in Intraplaque Hemorrhage in Apolipoprotein E Gene Null Mice. *Circ J.* 2018, 82(4), 1130-1138, DOI: 10.1253/circj.CJ-17-1023
 180. Naghavi, M.; Libby, P.; Falk, E.; Casscells, S. W.; Litovsky, S.; Rumberger, J.; Badimon, J. J.; Stefanadis, C.; Moreno, P.; Pasterkamp, G.; et al. From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: Part II. *Circulation.* 2003, 108, 1772-8, DOI: 10.1161/01.CIR.0000087481.55887.C9
 181. Witztum, J. L. The oxidation hypothesis of atherosclerosis. *Lancet (London, England).* 1994, 344, 8925:793-5, DOI:10.1016/s0140-6736(94)92346-9
 182. Berliner, J. A.; Navab, M.; Fogelman, A. M.; Frank, J. S.; Demer, L. L.; Edwards, P. A.; Watson, A. D.; Lusis, A. J. Atherosclerosis: basic mechanisms. Oxidation, inflammation, and genetics. *Circulation.* 1995, 91, 9, 2488-96, DOI: 10.1161/01.cir.91.9.2488
 183. Rader, D. J.; Puré, E. Lipoproteins, macrophage function, and atherosclerosis: beyond the foam cell? *Cell Metabolism.* 2005, 1, 223-230, DOI: 10.1016/j.cmet.2005.03.005
 184. Nuotio, K.; Isoviita, P. M.; Saksi, J.; Ijäs, P.; Pitkaniemi, J.; Sonninen, R.; Soinne, L.; Saimanen, E.; Salonen, O.; Kovanen, P. T.; et al. Adipophilin expression is increased in symptomatic carotid atherosclerosis: correlation with red blood cells and cholesterol crystals. *Stroke.* 2007, 38(6), 1791-8, DOI: 10.1161/STROKEAHA.106.478867. Epub 2007 Apr 19. PMID: 17446422
 185. Isoviita, P. M.; Nuotio, K.; Saksi, J.; Turunen, R.; Ijäs, P.; Pitkaniemi, J.; Soinne, L.; Kaste, M.; Kovanen, P. T.; Lindsberg, P. J. An imbalance between CD36 and ABCA1 protein expression favors lipid accumulation in stroke-prone ulcerated carotid plaques. *Stroke.* 2010, 41(2), 389-93, DOI: 10.1161/STROKEAHA.109.567636
 186. Ye, D.; Lammers, B.; Zhao, Y.; Meurs, I.; VanBerker, T. J. C.; VanEck, M. ATP-binding cassette transporters A1 and G1, HDL metabolism, cholesterol efflux, and inflammation: important targets for the treatment of atherosclerosis. *Current drug targets.* 2011, 12.5, 647-60, DOI:10.2174/138945011795378522
 187. Oram, J. F.; Heinecke, J. W. ATP-binding cassette transporter A1: a cell cholesterol exporter that protects against cardiovascular disease. *Physiological reviews.* 2005, 85.4, 1343-72, DOI:10.1152/physrev.00005.2005
 188. Hakkinen, T.; Luoma, J. S.; Hiltunen, M. O.; Macphee, C. H.; Miliner, K. J.; Patel, L.; Rice, S. Q.; Tew, D. G.; Karkola, K.; Yla-Herttuala, S. Lipoprotein-associated phospholipase A2, platelet-activation factor acetylhydrolase, is expressed by macrophages in human and rabbit atherosclerotic lesions. *Arterioscler Thromb Vasc Biol.* 1999, 19, 2909-2917, DOI: 10.1161/01.atv.19.12.2909
 189. Balboa, M. A.; Varela-Nieto, I.; Lucas, K. K.; Dennis, E. A. Expression and function of phospholipase A2 in brain. *FEBS Lett.* 2002, 531, 12-17, DOI: 10.1016/s00145793(02)03481-6
 190. Macphee, C. H.; Suckling, K. E. Lipoprotein-associated phospholipase A2 a target directed at the atherosclerotic plaque. *Expert Opin Ther Targets.* 2002, 6, 309-334, DOI: 10.1517/14728222.6.3.309
 191. Zhou, Z.; Li, X.; Yang, B.; Jiang, D. Relationship between lysophosphatidic acid and matrix metalloproteinase-9 plasma concentrations and carotid atheromatous plaque stability in patients with cerebral infarction. *The Journal of international medical research.* 2014, 42.3, 669-76, DOI:10.1177/0300060514526567
 192. Wasserman, B.; Sharrett, A.; Lai, S.; Gomes, A.; Cushman, M.; Folsom, A.; Bild, D.; Kronmal, R.; Sinha, S.; Bluemke, D. Risk factor associations with the presence of a lipid core in carotid plaque of asymptomatic individuals using high-resolution MRI: the multi-ethnic study of atherosclerosis (MESA). *Stroke.* 2008, 39.2, 329-35, DOI:10.1161/STROKEAHA.107.498634
 193. Guyton, J. R.; Klemp, K. F. Early extracellular and cellular lipid deposits in aorta of cholesterol-fed rabbits. *Am J Pathol.* 1992, 141(4), 925-36, PMC1886647
 194. Mitchinson, M. J.; Ball, R. Y.; Carpenter, K. H.; Enright, J. H.; Brabbs, C. E. Ceroid, macrophages and atherosclerosis. *Biochem Soc Trans.* 1990, 18(6), 1066-9, DOI: 10.1042/bst0181066
 195. Tangirala, R. K.; Mahlberg, F. H.; Glick, J. M.; Jerome, W. G.; Rothblat, G. H. Lysosomal accumulation of unesterified cholesterol in model macrophage foam cells. *J Biol Chem.* 1993, 268(13), 9653-60.

196. Tangirala, R. K.; Jerome, W. G.; Jones, N. L.; Small, D. M.; Johnson, W. J.; Glick, J. M.; Mahlberg, F. H.; Rothblat, G. H. Formation of cholesterol monohydrate crystals in macrophage-derived foam cells. *J Lipid Res.* 1994, 35(1), 93-104.
197. Minor, L. K.; Rothblat, G. H.; Glick, J. M. Triglyceride and cholesteryl ester hydrolysis in a cell culture model of smooth muscle foam cells. *J Lipid Res.* 1989, 30(2), 189-97.
198. Guyton, J. R.; Klemp, K. F. Development of the atherosclerotic core region. Chemical and ultrastructural analysis of microdissected atherosclerotic lesions from human aorta. *Arterioscler Thromb.* 1994, 14(8), 1305-14, DOI: 10.1161/01.atv.14.8.1305
199. Schmitz, G.; Robenek, H.; Beuck, M.; Krause, R.; Schurek, A.; Niemann, R. Ca⁺⁺ antagonists and ACAT inhibitors promote cholesterol efflux from macrophages by different mechanisms. I. Characterization of cellular lipid metabolism. *Arteriosclerosis.* 1988, 8(1), 46-56, DOI: 10.1161/01.atv.8.1.46
200. Chung, B. H.; Tallis, G.; Yalamoori, V.; Anantharamaiah, G. M.; Segrest, J. P. Liposome-like particles isolated from human atherosclerotic plaques are structurally and compositionally similar to surface remnants of triglyceride-rich lipoproteins. *Arterioscler Thromb.* 1994, 14(4), 622-35, DOI: 10.1161/01.atv.14.4.622
201. Carpenter, K. L.; Taylor, S. E.; Ballantine, J. A.; Fussell, B.; Halliwell, B.; Mitchinson, M. J. Lipids and oxidised lipids in human atheroma and normal aorta. *Biochim Biophys Acta.* 1993, 1167(2), 121-30, DOI: 10.1016/0005-2760(93)90151-x
202. Chisolm, G. M.; Ma, G.; Irwin, K. C.; Martin, L. L.; Gunderson, K. G.; Linberg, L. F.; Morel, D. W.; DiCorleto, P. E. 7 beta-hydroperoxycholest-5-en-3 beta-ol, a component of human atherosclerotic lesions, is the primary cytotoxin of oxidized human low-density lipoprotein. *Proc Natl Acad Sci U S A.* 1994, 91(24), 11452-6, DOI: 10.1073/pnas.91.24.11452
203. Hollander, W.; Colombo, M. A.; Kirkpatrick, B.; Paddock, J. Soluble proteins in the human atherosclerotic plaque. With spectral reference to immunoglobulins, C3-complement component, alpha 1-antitrypsin and alpha 2-macroglobulin. *Atherosclerosis.* 1979, 34(4), 391-405, DOI: 10.1016/0021-9150(79)90064-9
204. Seifert, P. S.; Hugo, F.; Hansson, G. K.; Bhakdi, S. Prelesional complement activation in experimental atherosclerosis. Terminal C5b-9 complement deposition coincides with cholesterol accumulation in the aortic intima of hypercholesterolemic rabbits. *Lab Invest.* 1989, 60(6), 747-54, PMID: 2659887
205. Small, D. M. Cellular mechanisms for lipid deposition in atherosclerosis (first of two parts). *N Engl J Med.* 1977, 297(16), 873-7, DOI: 10.1056/NEJM197710202971608
206. Abdulla, Y. H.; Adams, C. W. The action of human high-density lipoprotein on cholesterol crystals. Part 2. Biochemical observations. *Atherosclerosis.* 1978, 31(4), 473-80, DOI: 10.1016/0021-9150(78)90143-0
207. Bottcher, C. J. F.; Vangest, C. M. Changes in the composition of phospholipids and of phospholipid fatty acids associated with atherosclerosis in the human aortic wall. *J Atheroscler Res.* 1961, 1, 36-46, PMID: 6057496
208. Hannun, Y. A.; Bell, R. M. Functions of sphingolipids and sphingolipid breakdown products in cellular regulation. *Science.* 1989, 243(4890), 500-7, DOI: 10.1126/science.2643164
209. Gleason, M. M.; Medow, M. S.; Tulenko, T. N. Excess membrane cholesterol alters calcium movements, cytosolic calcium levels, and membrane fluidity in arterial smooth muscle cells. *Circ Res.* 1991, 69(1), 216-27, DOI: 10.1161/01.res.69.1.216
210. Warner, G. J.; Stoudt, G.; Bamberger, M.; Johnson, W. J.; Rothblat, G. H. Cell toxicity induced by inhibition of acyl coenzyme A:cholesterol acyltransferase and accumulation of unesterified cholesterol. *J Biol Chem.* 1995, 270(11), 5772-8, DOI: 10.1074/jbc.270.11.5772
211. Steinberg, D.; Parthasarathy, S.; Carew TE, Khoo JC, Witztum JL. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med.* 1989, 320(14), 915-24, DOI: 10.1056/NEJM198904063201407
212. Ramji, D. P.; Davies, T. S. Cytokines in atherosclerosis: Key players in all stages of disease and promising therapeutic targets. *Cytokine Growth Factor Rev.* 2015, 26(6), 673-85, DOI: 10.1016/j.cytogfr.2015.04.003. Epub 2015 May 12
213. Mori, H.; Torii, S.; Kutyna, M.; Sakamoto, A.; Finn, A. V.; Virmani, R. Coronary Artery Calcification and its Progression: What Does it Really Mean? *JACC Cardiovasc Imaging.* 2018, 11(1), 127-142, DOI: 10.1016/j.jcmg.2017.10.012
214. Jinnouchi, H.; Sato, Y.; Sakamoto, A.; Cornelissen, A.; Mori, M.; Kawakami, R.; Gadhoke, N. V.; Kolodgie, F. D.; Virmani, R.; Finn, A. V. Calcium deposition within coronary atherosclerotic lesion: Implications for plaque stability. *Atherosclerosis.* 2020, 306, 85-95, DOI: 10.1016/j.atherosclerosis.2020.05.017
215. Vengrenyuk, Y.; Carlier, S.; Xanthos, S.; Cardoso, L.; Ganatos, P.; Virmani, R.; Einav, S.; Gilchrist, L.; Weinbaum, S. A hypothesis for vulnerable plaque rupture due to stress-induced debonding around cellular microcalcifications in thin fibrous caps. *Proc Natl Acad Sci U S A.* 2006, 103(40), 14678-83, DOI: 10.1073/pnas.0606310103
216. Maldonado, N.; Kelly-Arnold, A.; Vengrenyuk, Y.; Laudier, D.; Fallon, J. T.; Virmani, R.; Cardoso, L.; Weinbaum, S. A mechanistic analysis of the role of microcalcifications in atherosclerotic plaque stability:

- potential implications for plaque rupture. *Am J Physiol Heart Circ Physiol.* 2012, 303(5), H619-28, DOI: 10.1152/ajpheart.00036.2012
217. Stary, H. C.; Chandler, A. B.; Dinsmore, R. E. A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis. A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation.* 1995, 92(5), 1355-74, DOI: 10.1161/01.cir.92.5.1355
 218. Mulvihill, J. J.; Cunnane, E. M.; McHugh, S. M.; Kavanagh, E. G.; Walsh, S. R.; Walsh, M. T. Mechanical, biological and structural characterization of in vitro ruptured human carotid plaque tissue. *Acta Biomater.* 2013, 9(11), 9027-35, DOI: 10.1016/j.actbio.2013.07.012
 219. Shaalan, W. E.; Cheng, H.; Gewertz, B.; McKinsey, J. F.; Schwartz, L. B.; Katz, D.; Cao, D.; Desai, T.; Glagov, S.; Bassiouny, H. S. Degree of carotid plaque calcification in relation to symptomatic outcome and plaque inflammation. *J Vasc Surg.* 2004, 40(2), 262-9, DOI: 10.1016/j.jvs.2004.04.025
 220. Wahlgren, C. M.; Zheng, W.; Shaalan, W.; Tang, J.; Bassiouny, H. S. Human carotid plaque calcification and vulnerability: relationship between degree of plaque calcification, fibrous cap inflammatory gene expression and symptomatology. *Cerebrovascular Diseases.* 2009, 27(2), 193-200, DOI: 10.1159/000189204
 221. Kelly-Arnold, A.; Maldonado, N.; Laudier, D.; Aikawa, E.; Cardoso, L.; Weinbaum, S. Revised microcalcification hypothesis for fibrous cap rupture in human coronary arteries. *Proceedings of the National Academy of Sciences of the United States of America.* 2013, 110(26), 10741-10746, DOI: 10.1073/pnas.1308814110
 222. Vengrenyuk, Y.; Cardoso, L.; Weinbaum, S. Micro-CT based analysis of a new paradigm for vulnerable plaque rupture: cellular microcalcifications in fibrous caps. *MCB Molecular and Cellular Biomechanics.* 2008, 5(1), 37-47, PMID: 18524245
 223. Pugliese, G.; Iacobini, C.; Blasetti Fantauzzi, C.; Menini, S. The dark and bright side of atherosclerotic calcification. *Atherosclerosis.* 2015, 238(2), 220-30, DOI: 10.1016/j.atherosclerosis.2014.12.011
 224. Shi, X.; Gao, J.; Lv, Q.; Cai, H.; Wang, F.; Ye, R.; Liu, X. Calcification in Atherosclerotic Plaque Vulnerability: Friend or Foe? *Front Physiol.* 2020, 11, 56, DOI: 10.3389/fphys.2020.00056
 225. Ehara, S.; Kobayashi, Y.; Yoshiyama, M.; Shimada, K.; Shimada, Y.; Fukuda, D.; Nakamura, Y.; Yamashita, H.; Yamagishi, H.; Takeuchi, K.; et al. Spotty calcification typifies the culprit plaque in patients with acute myocardial infarction: an intravascular ultrasound study. *Circulation.* 2004, 110(22), 3424-9, DOI: 10.1161/01.CIR.0000148131.41425.E9
 226. Banach, M.; Serban, C.; Sahebkar, A.; Mikhailidis, D. P.; Ursoniu, S.; Ray, K. K.; Rysz, J.; Toth, P. P.; Muntner, P.; Mosteoru, S.; et al. Lipid and Blood Pressure Meta-analysis Collaboration (LBPMC) Group. Impact of statin therapy on coronary plaque composition: a systematic review and meta-analysis of virtual histology intravascular ultrasound studies. *BMC Med.* 2015, 13, 229, DOI: 10.1186/s12916-015-0459-4
 227. Puri, R.; Nicholls, S. J.; Shao, M.; Kataoka, Y.; Uno, K.; Kapadia, S. R.; Tuzcu, E. M.; Nissen, S. E. Impact of statins on serial coronary calcification during atheroma progression and regression. *J Am Coll Cardiol.* 2015, 65(13), 1273-1282, DOI: 10.1016/j.jacc.2015.01.036
 228. Mujaj, B.; Bos, D.; Selwaness, M.; Leening, M. J. G.; Kavousi, M.; Wentzel, J. J.; van der Lugt, A.; Hofman, A.; Stricker, B. H.; Vernooij, M. W.; et al. Statin use is associated with carotid plaque composition: The Rotterdam Study. *Int J Cardiol.* 2018, 260, 213-218, DOI: 10.1016/j.ijcard.2018.02.111
 229. Xian, J. Z.; Lu, M.; Fong, F.; Qiao, R.; Patel, N. R.; Abeydeera, D.; Iriana, S.; Demer, L. L.; Tintut, Y. Statin Effects on Vascular Calcification: Microarchitectural Changes in Aortic Calcium Deposits in Aged Hyperlipidemic Mice. *Arterioscler Thromb Vasc Biol.* 2021, 41(4), e185-e192, DOI: 10.1161/ATVBAHA.120.315737
 230. Huang, H.; Virmani, R.; Younis, H.; Burke, A. P.; Kamm, R. D.; Lee, R. T. The impact of calcification on the biomechanical stability of atherosclerotic plaques. *Circulation.* 2001, 103(8), 1051-6, DOI: 10.1161/01.cir.103.8.1051
 231. Joshi, N. V.; Vesey, A. T.; Williams, M. C.; Shah, A. S.; Calvert, P. A.; Craighead, F. H.; Yeoh, S. E.; Wallace, W.; Salter, D.; Fletcher, A. M.; et al. 18F-fluoride positron emission tomography for identification of ruptured and high-risk coronary atherosclerotic plaques: a prospective clinical trial. *Lancet.* 2014, 383(9918), 705-13, DOI: 10.1016/S0140-6736(13)61754-7
 232. Reith, S.; Milzi, A.; Dettori, R.; Marx, N.; Burgmaier, M. Predictors for target lesion microcalcifications in patients with stable coronary artery disease: an optical coherence tomography study. *Clin Res Cardiol.* 2018, 107(9), 763-771, DOI: 10.1007/s00392-018-1243-1
 233. Mauriello, A.; Servadei, F.; Zoccai, G. B.; Giacobbi, E.; Anemona, L.; Bonanno, E.; Casella, S. Coronary calcification identifies the vulnerable patient rather than the vulnerable Plaque. *Atherosclerosis.* 2013, 229(1), 124-9, DOI: 10.1016/j.atherosclerosis.2013.03.010
 234. Aikawa, E.; Nahrendorf, M.; Figueiredo, J. L.; Swirski, F. K.; Shtatland, T.; Kohler, R. H.; Jaffer, F. A.; Aikawa, M.; Weissleder, R. Osteogenesis associates with inflammation in early-stage atherosclerosis evaluated by molecular imaging in vivo. *Circulation.* 2007, 116(24), 2841-50, DOI: 10.1161/CIRCULATIONAHA.107.732867

235. Kim, K. M. Apoptosis and calcification. *Scanning Microsc.* 1995, 9(4), 1137-75, discussion 1175-8, PMID: **8819895**
236. Lin, J.; Li, H.; Yang, M.; Ren, J.; Huang, Z.; Han, F.; Huang, J.; Ma, J.; Zhang, D.; Zhang, Z.; et al. A role of RIP3-mediated macrophage necrosis in atherosclerosis development. *Cell Rep.* 2013, 3(1), 200-10, DOI: 10.1016/j.celrep.2012.12.012
237. Lee, H. L.; Woo, K. M.; Ryoo, H. M.; Baek, J. H. Tumor necrosis factor-alpha increases alkaline phosphatase expression in vascular smooth muscle cells via MSX2 induction. *Biochem Biophys Res Commun.* 2010, 391(1), 1087-92, DOI: 10.1016/j.bbrc.2009.12.027
238. Wen, C.; Yang, X.; Yan, Z.; Zhao, M.; Yue, X.; Cheng, X.; Zheng, Z.; Guan, K.; Dou, J.; Xu, T.; et al. Nalp3 inflammasome is activated and required for vascular smooth muscle cell calcification. *Int J Cardiol.* 2013, 168(3), 2242-7, DOI: 10.1016/j.ijcard.2013.01.211
239. Aigner, T.; Neureiter, D.; Câmporean, V.; Soder, S.; Amann, K. Expression of cartilage-specific markers in calcified and non-calcified atherosclerotic lesions. *Atherosclerosis.* 2008, 196(1), 37-41, DOI: 10.1016/j.atherosclerosis.2007.01.020
240. Nadra, I.; Boccaccini, A. R.; Philippidis, P.; Whelan, L. C.; McCarthy, G. M.; Haskard, D. O.; Landis, R. C. Effect of particle size on hydroxyapatite crystal-induced tumor necrosis factor alpha secretion by macrophages. *Atherosclerosis.* 2008, 196(1), 98-105, DOI: 10.1016/j.atherosclerosis.2007.02.005
241. Pazár, B.; Ea, H. K.; Narayan, S.; Kolly, L.; Bagnoud, N.; Chobaz, V.; Roger, T.; Lioté, F.; So, A.; Busso, N. Basic calcium phosphate crystals induce monocyte/macrophage IL-1 β secretion through the NLRP3 inflammasome in vitro. *J Immunol.* 2011, 186(4), 2495-502, DOI: 10.4049/jimmunol.1001284
242. Viegas, C. S. B.; Costa, R. M.; Santos, L.; Videira, P. A.; Silva, Z.; Araújo, N.; Macedo, A. L.; Matos, A. P.; Vermeer, C.; Simes, D. C. Gla-rich protein function as an anti-inflammatory agent in monocytes/macrophages: Implications for calcification-related chronic inflammatory diseases. *PLoS One.* 2018, 13(2), e0192875, DOI: 10.1371/journal.pone.0177829
243. Fakhry, M.; Roszkowska, M.; Briolay, A.; Bougault, C.; Guignandon, A.; Diaz-Hernandez, J. I.; Diaz-Hernandez, M.; Pikula, S.; Buchet, R.; Hamade, E.; et al. TNAP stimulates vascular smooth muscle cell trans-differentiation into chondrocytes through calcium deposition and BMP-2 activation: Possible implication in atherosclerotic plaque stability. *Biochim Biophys Acta Mol Basis Dis.* 2017, 1863(3), 643-653, DOI: 10.1016/j.bbdis.2016.12.003
244. Ho, A. M.; Johnson, M. D.; Kingsley, D. M. Role of the mouse ank gene in control of tissue calcification and arthritis. *Science.* 2000, 289(5477), 265-70, DOI: 10.1126/science.289.5477.265
245. Pettengill, M.; Matute, J. D.; Tresenriter, M.; Hibbert, J.; Burgner, D.; Richmond, P.; Millán, J. L.; Ozonoff, A.; Strunk, T.; Currie, A.; et al. Human alkaline phosphatase dephosphorylates microbial products and is elevated in preterm neonates with a history of late-onset sepsis. *PLoS One.* 2018, 13(5), e0197532, DOI: 10.1371/journal.pone.0175936
246. Ding, J.; Ghali, O.; Lencel, P.; Broux, O.; Chauveau, C.; Devedjian, J. C.; Hardouin, P.; Magne, D. TNF-alpha and IL-1beta inhibit RUNX2 and collagen expression but increase alkaline phosphatase activity and mineralization in human mesenchymal stem cells. *Life Sci.* 2009, 84(15-16), 499-504, DOI: 10.1016/j.lfs.2009.01.013
247. Luo, G.; Ducy, P.; McKee, M. D.; Pinero, G. J.; Loyer, E.; Behringer, R. R.; Karsenty, G. Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature.* 1997, 386(6620), 78-81, DOI: 10.1038/386078a0
248. Nadra, I.; Mason, J. C.; Philippidis, P.; Florey, O.; Smythe, C. D.; McCarthy, G. M.; Landis, R. C.; Haskard, D. O. Proinflammatory activation of macrophages by basic calcium phosphate crystals via protein kinase C and MAP kinase pathways: a vicious cycle of inflammation and arterial calcification? *Circ Res.* 2005, 96(12), 1248-56, DOI: 10.1161/01.RES.0000171451.88616.c2
249. Pazár B, Ea HK, Narayan S, Kolly L, Bagnoud N, Chobaz V, Roger T, Lioté F, So A, Busso N. Basic calcium phosphate crystals induce monocyte/macrophage IL-1 β secretion through the NLRP3 inflammasome in vitro. *J Immunol.* 2011 Feb 15;186(4):2495-502. DOI: 10.4049/jimmunol.1001284
250. Morony, S.; Tintut, Y.; Zhang, Z.; Cattley, R. C.; Van, G.; Dwyer, D.; Stolina, M.; Kostenuik, P. J.; Demer, L. L. Osteoprotegerin inhibits vascular calcification without affecting atherosclerosis in *ldlr*^{-/-} mice. *Circulation.* 2008, 117(3), 411-20, DOI: 10.1161/CIRCULATIONAHA.107.707380
251. Dautova, Y.; Kapustin, A. N.; Pappert, K.; Eppele, M.; Okkenhaug, H.; Cook, S. J.; Shanahan, C. M.; Bootman, M. D.; Proudfoot, D. Calcium phosphate particles stimulate interleukin-1 β release from human vascular smooth muscle cells: A role for spleen tyrosine kinase and exosome release. *J Mol Cell Cardiol.* 2018, 115, 82-93, DOI: 10.1016/j.yjmcc.2017.12.007
252. Ewence, A. E.; Bootman, M.; Roderick, H. L.; Skepper, J. N.; McCarthy, G.; Eppele, M.; Neumann, M.; Shanahan, C. M.; Proudfoot, D. Calcium phosphate crystals induce cell death in human vascular smooth muscle cells: a potential mechanism in atherosclerotic plaque destabilization. *Circ Res.* 2008, 103(5), e28-34, DOI: 10.1161/CIRCRESAHA.108.181305

253. Clarke, M. C.; Talib, S.; Figg, N. L.; Bennett, M. R. Vascular smooth muscle cell apoptosis induces interleukin-1-directed inflammation: effects of hyperlipidemia-mediated inhibition of phagocytosis. *Circ Res.* 2010, 106(2), 363-72, DOI: 10.1161/CIRCRESAHA.109.208389
254. Roijers, R. B.; Debernardi, N.; Cleutjens, J. P.; Schurgers, L. J.; Mutsaers, P. H.; van der Vusse, G. J. Microcalcifications in early intimal lesions of atherosclerotic human coronary arteries. *Am J Pathol.* 2011, 178(6), 2879-87, DOI: 10.1016/j.ajpath.2011.02.004
255. Huang, L. H.; Han, J.; Ouyang, J. M.; Gui, B. S. Shape-dependent adhesion and endocytosis of hydroxyapatite nanoparticles on A7R5 aortic smooth muscle cells. *J Cell Physiol.* 2020, 235(1), 465-479, DOI: 10.1002/jcp.28987
256. Matsui, Y.; Rittling, S. R.; Okamoto, H.; Inobe, M.; Jia, N.; Shimizu, T.; Akino, M.; Sugawara, T.; Morimoto, J.; Kimura, C.; et al. Osteopontin deficiency attenuates atherosclerosis in female apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol.* 2003, 23(6), 1029-34, DOI: 10.1161/01.ATV.0000074878.29805.D0
257. Abedin, M.; Tintut, Y.; Demer, L. L. Vascular calcification: mechanisms and clinical ramifications. *Arterioscler Thromb Vasc Biol.* 2004, 24(7), 1161-1170, DOI: 10.1161/01.ATV.0000133194.94939.42
258. Collin-Osdoby, P. Regulation of vascular calcification by osteoclast regulatory factors RANKL and osteoprotegerin. *Circ Res.* 2004, 95(11), 1046-1057, DOI: 10.1161/01.RES.0000149165.99974.12
259. Kadoglou, N. P. E.; Gerasimidis, T.; Moumtzouoglou, A.; Kapelouzou, A.; Sailer, N.; Fotiadis, G.; Vitta, I.; Katinios, A.; Kougias, P.; Bandios, S.; et al. Intensive lipid-lowering therapy ameliorates novel calcification markers and GSM score in patients with carotid stenosis. *Eur J Vasc Endovasc Surg.* 2008, 35(6), 661-668, DOI: 10.1016/j.ejvs.2007.12.011
260. Reid, P.; Holen, I. Pathophysiological roles of osteoprotegerin (OPG). *Eur J Cell Biol.* 2009, 88(1), 1-17, DOI: 10.1016/j.ejcb.2008.06.004
261. Scatena, M.; Liaw, L.; Giachelli, C. M. Osteopontin: a multifunctional molecule regulating chronic inflammation and vascular disease. *Arterioscler Thromb Vasc Biol.* 2007, 27(11), 2302-2309, DOI: 10.1161/ATVBAHA.107.144824
262. Heymann, M. F.; Herisson, F.; Davaine, J. M.; Charrier, C.; Battaglia, S.; Passuti, N.; Lambert, G.; Gouëffic, Y.; Heymann, D. Role of the OPG/RANK/RANKL triad in calcifications of the atheromatous plaques: comparison between carotid and femoral beds. *Cytokine.* 2012, 58(2), 300-306, DOI: 10.1016/j.cyto.2012.02.004
263. Rochette, L.; Meloux, A.; Rigal, E.; Zeller, M.; Cottin, Y.; Vergely, C. The role of osteoprotegerin and its ligands in vascular function. *Int J Mol Sci.* 2019, 20(3), 705-705, DOI: [10.3390/ijms20030705](https://doi.org/10.3390/ijms20030705)
264. Kwon, H. M.; Hong, B. K.; Kang, T. S.; Kwon, K.; Kim, H. K.; Jang, Y.; Choi, D.; Park, H. Y.; Kang, S. M.; Cho, S. Y.; et al. Expression of osteopontin in calcified coronary atherosclerotic plaques. *J Korean Med Sci.* 2000, 15(5), 485-493, DOI: 10.3346/jkms.2000.15.5.485
265. Momiyama, Y.; Ohmori, R.; Fayad, Z. A.; Kihara, T.; Tanaka, N.; Kato, R.; Taniguchi, H.; Nagata, M.; Nakamura, H.; Ohsuzu, F. Associations between plasma osteopontin levels and the severities of coronary and aortic atherosclerosis. *Atherosclerosis.* 2010, 210(2), 668-670, DOI: 10.1016/j.atherosclerosis.2009.12.024
266. Maruyama, M.; Rhee, C.; Utsunomiya, T.; Zhang, N.; Ueno, M.; Yao, Z.; Goodman, S. B. Modulation of the Inflammatory Response and Bone Healing. *Front Endocrinol (Lausanne).* 2020, 11, 386, DOI: 10.3389/fendo.2020.00386
267. Gerstenfeld, L. C.; Cho, T. J.; Kon, T.; Aizawa, T.; Cruceta, J.; Graves, B. D.; Einhorn, T. A. Impaired intramembranous bone formation during bone repair in the absence of tumor necrosis factor-alpha signaling. *Cells Tissues Organs.* 2001, 169(3), 285-94, DOI: 10.1159/000047893
268. Fukui, N.; Ikeda, Y.; Ohnuki, T.; Hikita, A.; Tanaka, S.; Yamane, S.; Suzuki, R.; Sandell, L. J.; Ochi, T. Pro-inflammatory cytokine tumor necrosis factor-alpha induces bone morphogenetic protein-2 in chondrocytes via mRNA stabilization and transcriptional up-regulation. *J Biol Chem.* 2006, 281(37), 27229-41, DOI: 10.1074/jbc.M603385200
269. Csiszar, A.; Smith, K. E.; Koller, A.; Kaley, G.; Edwards, J. G.; Ungvari, Z. Regulation of bone morphogenetic protein-2 expression in endothelial cells: role of nuclear factor-kappaB activation by tumor necrosis factor-alpha, H2O2, and high intravascular pressure. *Circulation.* 2005, 111(18), 2364-72 DOI: 10.1161/01.CIR.0000164201.40634.1D
270. Tsuji, K.; Bandyopadhyay, A.; Harfe, B. D.; Cox, K.; Kakar, S.; Gerstenfeld, L.; Einhorn, T.; Tabin, C. J.; Rosen, V. BMP2 activity, although dispensable for bone formation, is required for the initiation of fracture healing. *Nat Genet.* 2006, 38(12), 1424-9, DOI: 10.1038/ng1916
271. Montanaro, M.; Scimeca, M.; Anemona, L.; Servadei, F.; Giacobbi, E.; Bonfiglio, R.; Bonanno, E.; Urbano, N.; Ippoliti, A.; Santeusano, G.; et al. The Paradox Effect of Calcification in Carotid Atherosclerosis: Microcalcification is Correlated with Plaque Instability. *Int J Mol Sci.* 2021, 22(1), 395, DOI: 10.3390/ijms22010395
272. Bi, Y.; Chen, J.; Hu, F.; Liu, J.; Li, M.; Zhao, L. M2 Macrophages as a Potential Target for Antiatherosclerosis Treatment. *Neural Plast.* 2019, 2019, 6724903, DOI: 10.1155/2019/6724903

273. Shioi, A.; Ikari, Y. Plaque Calcification During Atherosclerosis Progression and Regression. *J Atheroscler Thromb.* 2018, 25(4), 294-303, DOI: 10.5551/jat.RV17020
274. Okulla, T.; Gass, S.; Böhme, K.; Tiemann, K.; Harbrecht, U.; Klockgether, T.; Pohl, C. Circulating adhesion molecules in patients with internal carotid artery stenosis. *Cerebrovasc Dis.* 2002, 14(1), 9-14, DOI: 10.1159/000063717
275. Li, X.; Guo, D.; Hu, Y.; Chen, Y. Oxidative Stress and Inflammation Are Associated with Coexistent Severe Multivessel Coronary Artery Stenosis and Right Carotid Artery Severe Stenosis in Elderly Patients. *Oxid Med Cell Longev.* 2021, 2021, 2976447, DOI: 10.1155/2021/2976447
276. Li, X.; Guo, D.; Hu, Y.; Zhou, H.; Chen, Y. Potential Biomarkers and Therapeutic Targets: Inflammation and Oxidative Stress in Left Carotid Artery Stenosis with Coronary Artery Disease. *Curr Pharm Des.* 2023, 29(12), 966-979, DOI: 10.2174/1381612829666230417100144
277. Tian, X.; Wang, X.; Shi, Z.; Yu, C.; Li, M.; Chen, L.; Jia, Q.; Liang, G. Tumor necrosis factor-stimulated gene-6-a new serum identification marker to identify severe and symptomatic carotid artery stenosis. *Pathol Res Pract.* 2022, 232, 153838, DOI: 10.1016/j.prp.2022.153838
278. Zhang, S.; Guo, M.; Liu, Q.; Liu, J.; Cui, Y. Neutrophil extracellular traps induce thrombogenicity in severe carotid stenosis. *Immun Inflamm Dis.* 2021, 9(3), 1025-1036, DOI: 10.1002/iid3.466
279. Handberg, A.; Skjelland, M.; Michelsen, A. E.; Sagen, E. L.; Krohg-Sørensen, K.; Russell, D.; Dahl, A.; Ueland, T.; Oie, E.; Aukrust, P.; et al. Soluble CD36 in plasma is increased in patients with symptomatic atherosclerotic carotid plaques and is related to plaque instability. *Stroke.* 2008, 39, 3092-3095, DOI: 10.1161/STROKEAHA.108.517128
280. Jin, G. The relationship between serum CXCL16 level and carotid vulnerable plaque in patients with ischemic stroke. *Eur. Rev. Med. Pharmacol. Sci.* 2017, 21, 3911-3915, PMID: 28975971
281. Profumo, E.; Buttari, B.; Tosti, M. E.; Salvati, B.; Capoano, R.; Riganò, R. Increased circulating levels of MIP-1 α and CD14 are associated with the presence of severe stenosis and hypoechoic plaques in patients with carotid atherosclerosis. *Int J Immunopathol Pharmacol.* 2023, 37, 3946320231160411, DOI: 10.1177/03946320231160411
282. Yayla, Ç.; Açıkgöz, S. K.; Yayla, K. G.; Açıkgöz, E.; Canpolat, U.; Kirbaş, Ö.; Öksüz, F.; Özcan, F.; Akboğa, M. K.; et al. The association between platelet-to-lymphocyte ratio and inflammatory markers with the severity of aortic stenosis. *Biomark Med.* 2016, 10(4), 367-73, DOI: 10.2217/bmm-2015-0016
283. Papalambros, E.; Georgopoulos, S.; Sigala, F.; Vourliotakis, G.; Chrisostomidis, G.; Venetsanou, K.; Hadjoulis, G.; Felekouras, E.; Bastounis, E. Changes in circulating levels of vascular endothelial growth factor and vascular endothelial growth factor receptor-2 after carotid endarterectomy. *Int J Mol Med.* 2004, 14(1), 133-6, DOI: 10.3892/ijmm.14.1.133
284. Nuotio, K.; Lindsberg, P. J.; Carpén, O.; Soinne, L.; Lehtonen-Smeds, E. M.; Saimanen, E.; Lassila, R.; Sairanen, T.; Sarna, S.; Salonen, O.; et al. Adhesion molecule expression in symptomatic and asymptomatic carotid stenosis. *Neurology.* 2003, 60(12), 1890-9, DOI: 10.1212/01.wnl.0000065914.33177.9e
285. Puertas-Umbert, L.; Puig, N.; Camacho, M.; Dantas, A. P.; Marín, R.; Martí-Fàbregas, J.; Jiménez-Xarrié, E.; Benitez, S.; Camps-Renom, P.; Jiménez-Altayó, F. Serum from Stroke Patients with High-Grade Carotid Stenosis Promotes Cyclooxygenase-Dependent Endothelial Dysfunction in Non-ischemic Mice Carotid Arteries. *Transl Stroke Res.* 2024, 15(1), 140-152, DOI: 10.1007/s12975-022-01117-1.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.