

---

# A Secondary Retrospective Analysis of the Predictive Value of Activated Neutrophils (NEUT-RI) in Septic and Non-septic Patients in Intensive Care

---

[Paolo Formenti](#)\*, Letizia Isidori, Stefano Pastori, Vincenzo Roccaforte, Elena Alessandra Mantovani, Massimiliano Iezzi, Alessandro Menozzi, [Rossella Panella](#), Andrea Galimberti, Giovanni Brenna, [Michele Umbrello](#), Angelo Pezzi, Francesco Vetrone, [Giovanni Sabbatini](#), Miriam Gotti

Posted Date: 21 March 2024

doi: 10.20944/preprints202403.1120.v1

Keywords: sepsis; Neutrophil Reactivity (NEUT-RI); C-Reactive Protein; Procalcitonin



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

# A Secondary Retrospective Analysis of the Predictive Value of Neutrophil-Reactive Intensity (NEUT-RI) in Septic and Non-Septic Patients in Intensive Care

Paolo Formenti <sup>1,\*</sup>, Letizia Isidori <sup>1</sup>, Stefano Pastori <sup>2</sup>, Vincenzo Roccaforte <sup>2</sup>, Elena Maria Alessandra Mantovani <sup>1</sup>, Massimiliano Iezzi <sup>3</sup>, Alessandro Menozzi <sup>3</sup>, Rossella Panella <sup>2</sup>, Andrea Galimberti <sup>1</sup>, Giovanni Brenna <sup>1</sup>, Michele Umbrello <sup>4</sup>, Angelo Pezzi <sup>1</sup>, Francesco Vetrone <sup>1</sup>, Giovanni Sabbatini <sup>1</sup> and Miriam Gotti <sup>1</sup>

<sup>1</sup> SC Anestesia, Rianimazione e Terapia Intensiva, ASST Nord Milano, Ospedale Bassini, Cinisello Balsamo, 20097, Italy.

<sup>2</sup> S.C. Analisi Chimico Cliniche e Microbiologiche, ASST Nord Milano, Ospedale Bassini; Cinisello Balsamo, 20097, Italy.

<sup>3</sup> School of Medicine and Surgery, University of Milano-Bicocca, 20126 Milano, Italy.

<sup>4</sup> Department of Intensive Care, New Hospital of Legnano: Ospedale Nuovo di Legnano, 20025, Legnano, Milan, Italy.

\* Correspondence: paolo.formenti@asst-nordmilano.it

**Abstract: Background:** Effective identification and management in the early stages of sepsis are critical for achieving positive outcomes. In this context, Neutrophil-Reactive Intensity (NEUT-RI) emerges as a promising and easily interpretable parameter. This study aimed to assess the predictive value of NEUT-RI in diagnosing sepsis and to evaluate its prognostic significance in distinguishing 28-day mortality outcomes. **Materials:** The study is a secondary retrospective observational analysis. Clinical data upon ICU admission were collected. A control group of critically ill patients without sepsis criteria was included. Patients were divided into subgroups based on renal function for biomarker evaluation, with 28-day outcomes reported for septic and non-septic patients. **Results:** 200 patients were included in the study. A significant difference between the “septic” and “non-septic” groups was detected in the NEUT-RI plasma concentration (53.80 [49.65–59.05] vs. 48.00 [46.00–49.90],  $p < 0.001$ , respectively). NEUT-RI and PCT distinguished between not complicated sepsis and septic shock (PCT 1.71 [0.42 - 12.09] vs 32.59 [8.83 - 100.00],  $< 0.001$  and NEUT-RI 51.50 [47.80 - 56.30] vs 56.20 [52.30 - 61.92],  $p=0.005$ ). NEUT-RI, PCT and CRP values were significantly different in patients with “renal failure”. NEUT-RI and PCT at admission in ICU in septic group were higher in patients who died (58.80 [53.85-73.10] vs 53.05 [48.90-57.22],  $p=0.005$  and 39.56 [17.39-83.72] vs 3.22 [0.59-32.32],  $p=0.002$ , respectively). Both NEUT-RI and PCT showed high negative predictive value and low positive predictive value. **Conclusions:** The inflammatory biomarkers assessed in this study offer valuable support in the early diagnosis of sepsis. NEUT-RI elevation appears particularly promising for early sepsis detection and severity discrimination upon admission.

**Keywords:** sepsis; Neutrophil Reactivity (NEUT-RI); C-reactive protein; procalcitonin

## 1. Introduction

Sepsis represents a severe medical condition marked by significant organ dysfunction and a potentially life-threatening state, arising from an uncontrolled host reaction to infection [1]. Sepsis and septic shock pose significant global health challenges leading to mortality in one third to one sixth of affected individuals [2]. A Sequential Organ Failure Assessment (SOFA) score of  $\geq 2$  indicates a considerable likelihood of mortality and a hospital admission rate surpassing 10% according to

revised sepsis definition [3]. Timely identification and appropriate management during the initial stages of sepsis development are pivotal for favorable outcomes [4]. However, accurate diagnosis is crucial to prevent unwarranted antibiotic use and therefore the outbreak of antibiotic resistance [5]. Currently, blood cultures are considered the gold standard for pathogen isolation and sepsis diagnosis [6]. Routine sepsis markers, such as white blood cell count (WBC) and C-reactive protein (CRP), lack specificity [7]. CRP, commonly used as a biomarker for acute inflammatory states, exhibits increased plasma concentrations parallel to the infection's clinical course, with a decrease indicating resolution [8]. However, its specificity is limited, necessitating the evaluation of alternative markers like procalcitonin (PCT), tumor necrosis factor (TNF), and interleukin 6 (IL-6), although with challenges like cost and processing time [9–13]. Recent research has shifted focus to the role of neutrophil granulocytes in inflammation [14,15], particularly the Neutrophil-Reactive Intensity (NEUT-RI) [16]. NEUT-RI emerges as a promising and easily interpretable parameter, with literature supporting its correlation with increase in immunoglobulins [17]. Notably, NEUT-RI exhibits a significant correlation with infectious status [18], strengthening its potential utility in early sepsis diagnosis, and facilitating prompt initiation of optimal therapeutic interventions. The primary objective of this study was to evaluate the predictive value of NEUT-RI in sepsis diagnosis compared to other commonly used inflammatory parameters based on our previous observation [19]. Additionally, the secondary aim was assessing the prognostic value of NEUT-RI in discriminating 28-day mortality outcomes after intensive care unit (ICU) admission.

## 2. Materials and Methods

A comprehensive retrospective observational analysis of electronic medical records was performed to evaluate the diagnostic efficacy of NEUT-RI. The existing database, including patient information spanning from March 2022 to November 2022, underwent thorough scrutiny and augmentation, with data collection extending until October 2023 [19]. Clinical data of patients upon admission to ICU (clinical diagnosis, proximal and remote medical history for enrollment and exclusion criteria from the study), their respective laboratory values (WBC, PCR, PCT, extended inflammatory parameters, creatinine), blood culture results, SOFA score and SAPS II for severity assessment were collected. NEUT-RI and PCT were also recorded at 48h and 96h from admission. The infection site and the 28-day outcome were added to this existing data. Data were collected retrospectively and stored in a dedicated database for consecutively admitted patients to two ICU in the Milan area (ASST Nord Milano). The protocol for this study was examined and approved by the Ethics Committee of Milano Area 3 (n.114/2023, February 7, 2023). We included consecutively ICU patients admitted between March 2022 and October 2023 with a clinical diagnosis of sepsis, according to the Sepsis-III definition [1]. The diagnosis of sepsis in the study group was based on the clinical diagnosis used in the participating ICUs, encoded in Margherita Prosafe, and centrally validated by the "Istituto Mario Negri" within the PROSAFE project [20]. The control group consisted of the population of critically ill patients admitted to the same ICU during the same period without sepsis diagnostic criteria. Exclusion criteria included age less than 18 years, active neoplasia, chronic myelomonocytic leukemia, chronic myeloid leukemia, chronic corticosteroid therapy (prednisone >10 mg/day or equivalent), immunosuppressive, or immunomodulatory therapy, and congenital immunodeficiency. For each patient, information was gathered concerning age, gender, nosologically code of the clinical record, the originating hospital department, the date of admission to the ICU, the clinical diagnosis upon admission (for the division into "septic" versus "non-septic" study groups), the recent and remote pathological history (for the criteria of enrollment and exclusion from the study), the 28-day outcome, respective laboratory values (WBC, PCR, PCT, extended inflammatory parameters, creatinine), blood culture results, site of infection, and severity level at the time of admission using the SOFA score for the "septic" group. CRP, PCT, and creatinine were measured by using the automated chemical analyzer Beckman Coulter AU 5800 and the immunochemical analyzer Beckman Coulter Unicel DxI800 (Beckman Coulter, Brea; CA, USA) following the manufacturer's recommendations. PCT was measured via chemiluminescence and expressed in ng/mL; CRP was measured through the turbidimetric method and expressed in mg/dL, and creatinine through

immunoenzymatically assay and expressed in mg/dL. The focal point of our study was the extended inflammatory parameter, NEUT-RI. To ascertain complete blood cell counts, a Sysmex XN hematology analyzer (Sysmex, Kobe, Japan) was employed, utilizing cytometry for the enumeration and classification of blood cells. To evaluate the effect of renal function on infection biomarkers, patients were further divided into subgroups "renal insufficiency" and "normal renal function." The "renal insufficiency" subgroup included patients diagnosed with Acute Kidney Injury (AKI) and Chronic Kidney Disease (CKD) according to the Kidney Disease Improving Global Outcomes (KDIGO) classification [21]. "Sepsis" patients were further divided into subgroups "uncomplicated sepsis" and "complicated sepsis" using the definition of septic shock (lactates > 2 mmol/L and the need for vasopressors to maintain MAP > 65 mmHg) [22]. The primary site of infection was recorded, creating seven subgroups for the following infection sites: lungs, abdomen, kidneys and urinary tract, skin and soft tissues, bloodstream, heart valves, CNS. Finally, the 28-day outcome of the two groups of "septic" and "non-septic" patients was reported.


### *Statistical Analysis*

The data were collected in a database built and compiled using Microsoft Excel in compliance with privacy regulations (Law 196/2003 and GDPR EU 679/2016). Statistical analyses of the data were conducted using the RSTUDIO statistical software. Since the Shapiro-Wilk normality test indicated that the NEUT-RI, WBC, PCT, and PCR variables were not normally distributed, a non-parametric statistical analysis was performed. The results were expressed as the median +/- interquartile range [IQR]. The independent samples t-test and non-parametric tests (Wilcoxon Rank Sum Test) were used to investigate significant differences between groups and subgroups. The Chi-squared test with Yates' continuity correction for small samples was employed for the comparison of proportions. The Kruskal-Wallis's rank sum test was conducted for the analysis of variance (for different infection sites). ROC analysis was performed to assess the performance of each biomarker and determine the best cut-off values. DeLong's test for two correlated ROC curves was conducted for comparisons between ROC curves; the sensitivity and specificity tests for correlated ROC curves were conducted for comparison of sensitivity and specificity of the inflammatory parameters. A p-value  $\leq 0.05$  was considered statistically significant. Sensitivity and specificity for the diagnosis of sepsis upon admission to the Intensive Care Unit of the NEUT-RI parameter were compared with those of PCT and PCR, along with their positive and negative predictive values. Between the two subgroups of "septic" patients ("uncomplicated sepsis" versus "complicated sepsis"), the values of each biomarker (NEUT-RI, PCT, and PCR) were compared. The "septic" and "non-septic" groups were further divided into subgroups of "renal insufficiency" and "normal renal function" comparing the values of each biomarker. The trends of NEUT-RI and PCT were analyzed in the first 96h of ICU staying in "septic" group. The "septic" and "non-septic" groups were further divided into subgroups of "alive" versus "deceased" based on the 28-day outcome, and the values of each biomarker were compared. Sensitivity and specificity for predicting the 28-day outcome upon admission to ICU of the NEUT-RI parameter were also compared with those of PCT and PCR, along with their positive and negative predictive values.

### **3. Results**

200 patients were included in the study (Figure 1). The average age was 73 years [59-79], with males representing 50.5% (n=101) and females 49.5% (n = 99). In the "septic" group 89 patients (44.5%) were included vs 111 patients (55.5%) in "non-septic group" (Table 1, Panel A). The admission diagnosis of "septic" patients was Pneumonia or upper respiratory tract infection in 39 patients (42.8%), secondary peritonitis due to cholangitis, intestinal obstruction, or intestinal perforation in 25 patients (27.5%), urinary tract infection or pyelonephritis in 17 patients (18.7%), infection of the skin and soft tissues, including necrotizing fasciitis, gas gangrene, infections of orthopedic prostheses, or submandibular abscesses, in 7 patients (7.7%), heart valve infections in 2 patients (2.2%), and bloodstream infection in 1 patient (1.1%). No patients in our population had CNS infections. Among the "septic" patients, 49 (53.8%) belonged to the "uncomplicated sepsis" subgroup, and 42 (46.2%)

had a diagnosis of septic shock upon admission to ICU (the “complicated sepsis” subgroup) (Table 1, panel B).



**Figure 1.** Study design flow chart.

**Table 1.** Comparisons of patients’ characteristics and inflammatory parameters divided in “septic” and “non-septic” patients (Panel A), and, among septic, in “not complicated sepsis” or “septic shock” patients (Panel B).

	A		P	B		P
	Septic (n=89)	Non septic (n=111)		Not complicated sepsis (n=47)	Septic shock (n=42)	
<b>Age</b>	73.00 [63.00 - 79.00]	73.00 [56.00 - 79.00]	0.454	74.00 [64.00 - 78.00]	73.00 [62.75 - 79.75]	0.808
<b>Male%</b>	48.4	52.3	0.679	59.6	35.7	0.042
<b>Creatinine</b>	1.57 [0.85 - 2.85]	1.00 [0.77 - 1.39]	< 0.001	1.16 [0.77 - 1.64]	2.18 [1.67 - 3.67]	< 0.001
<b>CPR</b>	20.79 [12.54 - 118.91]	6.68 [1.54 - 21.70]	< 0.001	18.79 [7.78 - 149.16]	23.03 [17.97 - 87.09]	0.7244
<b>PCT</b>	8.83 [0.82 - 45.88]	0.48 [0.29 - 1.64]	< 0.001	1.63 [0.40 - 12.09]	32.59 [8.83 - 100.00]	< 0.001
<b>NEUT-RI</b>	53.80 [49.65 - 59.05]	48.00 [46.00 - 49.90]	< 0.001	51.5 [47.80 - 56.30]	56.20 [52.30 - 61.92]	0.0054

### 3.1. Inflammatory Parameters

A significant difference between the “septic” and “non-septic” groups was detected in the NEUT-RI plasma concentration (53.80 [49.65–59.05] vs. 48.00 [46.00–49.90],  $p < 0.001$ , respectively) (Figure 1, panel A). Similar results were detected for PCT and CPR (PCT 8.83 [0.82 - 45.88] vs. 0.48 [0.29 - 1.64],  $p < 0.001$ ; CPR 20.79 [12.54 - 118.91] vs. 6.68 [1.54 - 21.70],  $p < 0.001$ ). The NEUT-RI and PCT were able to discriminate between not complicated sepsis and septic shock (PCT 1.71 [0.42 - 12.09] vs 32.59 [8.83 - 100.00],  $p < 0.001$  and NEUT-RI 51.50 [47.80 - 56.30] vs 56.20 [52.30 - 61.92],  $p = 0.005$ ). There were no differences in NEUT-RI and CPR depending on the site of infection in “septic” group. Regarding the site of infection, PCT was higher in patients with kidney or urinary tract infection

compared to lung infection (88.50 [24.20-113.00] vs 1.64[0.39-12.54],  $p<0.001$ ) (Figure S1 on the Supplementary Material).


### 3.2. Renal Failure and Inflammatory Parameters

Patients affected by renal failure upon admission to ICU were 70 (35%), including 47 AKI and 23 CKD. The inflammatory parameters are described in the Supplementary Materials Table S1. The comparison between patients based upon the presence or the absence of renal insufficiency is described in Table 2. In the “septic” group, 43 patients (48.3%) had renal failure upon admission, specifically 36 with AKI (including 5 with AKI in CKD) and 7 with CKD. In the “non-septic” group, 27 patients (24.3%) had renal failure upon admission, specifically 11 with AKI (including 2 with AKI in CKD) and 16 with CKD. NEUT-RI, PCT and CRP values were significantly different in patients with “renal failure” and those with “normal renal function” in the overall population (Supplementary Materials, Table S1). Within the septic group, no statistically significant difference was found in NEUT-RI and CPR (55.10 [52.15 – 59.05] vs 51.70 [47.82 – 58.65]; 25.53 [17.82 – 148.22] vs 18.80 [7.35 – 103.27], respectively), while PCT resulted significantly different (32.23 [5.86 – 83.72] vs 1.79 [0.39 – 13.04],  $p<0.001$ ). In the “non septic” group, only CRP exhibited a significant difference between patients with or without renal function impairment (13.45 [6.03 - 130.85] vs 3.78 [1.05 - 18.59],  $p=0.003$ ). Moreover, NEUT-RI and PCT were stratified in severity classes according to SAPS II: both inflammatory parameters showed a positive correlation ( $p<0.001$ , linear regression) with SAPS II, but with a great dispersion of data (Figure S2 in the Supplementary Material).

**Table 2.** Comparison of inflammatory parameters among “septic” and “non septic” patients based on renal function.

	Septic			Non Septic		
	Renal Failure (n = 43)	Normal renal function (n = 48)	P	Renal Failure (n = 27)	Normal renal function (n = 82)	P
Age	73.00 [63.00 - 79.00]	72.50 [63.50 - 77.50]	0.443	78.00 [73.00 - 81.00]	68.00 [53.25 - 77.00]	0.004
Male%	37.2	58.3	0.071	51.9	52.4	1
Creatinine	2.81 [1.83 – 4.26]	0.86 [0.74 – 1.28]	$<0.001$	1.60 [1.33 – 2.59]	0.89 [0.68 – 1.19]	$<0.001$
CPR	25.53 [17.82 - 148.22]	18.80 [7.35 - 103.27]	0.164	13.45 [6.03 - 130.85]	3.78 [1.05 - 18.59]	0.003
PCT	32.23 [5.86 - 83.72]	1.79 [0.39 - 13.04]	$<0.001$	1.15 [0.31 - 3.63]	0.47 [0.28 - 1.21]	0.193
NEUT-RI	55.10 [52.15 - 59.05]	51.70 [47.82 - 58.65]	0.101	47.90 [46.40 - 50.05]	48.00 [45.85 - 49.88]	0.886

Both NEUT-RI and PCT showed a statistically significant difference when measured at different timepoint in “septic” group ( $p<0.01$  repeated measures ANOVA for both inflammatory parameters). In particular, both NEUT-RI and PCT were lower at 96h compared to admission time and 48h. Nor NEUT-RI or PCT were lower at 48h compared to admission time (Table S4 on the Supplementary Materials). The performance of each inflammatory parameter for the diagnosis of sepsis and their best cut-off values are described in Supplementary Materials, Table S2, while the comparisons in AUROC are shown in Figure 2. There were not statistically significant differences between AUROC of the three parameters (NEUT-RI vs PCT  $p=0.83$ , NEUT-RI vs CRP  $p=0.29$ , DeLong’s test for two correlated ROC curves). The CPR specificity was statistically different from the NEUT-RI and the PCT specificity ( $p=0.01$  and  $p=0.007$  respectively, specificity test for two correlated ROC curves). Regarding the 28-day outcome, the overall mortality in our population was 17.5%; there was a trend in higher mortality in “septic” compared to “non septic” group (23.1% vs 12.8%,  $p=0.09$ ).



**Figure 2.** Comparison of ROC curves for NEUT-RI (FI), PCT (ng/mL), and PCR (mg/dL) for sepsis; AUC of NEUT-RI higher than those of PCT and PCR. The figure shows the area under the Receiver-operating characteristic (ROC) curve for the distinction of inflammatory parameters for detection of sepsis. The areas under the ROC curves are as follows: NEUT-RI (continue line): 0.79 [95%CI 0.74-0.91]; PCT (dashed line): 0.76 [95%CI 0.77-0.93]; CRP (dots line): 0.73 [95%CI 0.77-0.93];  $p < 0.001$ . NEUT-RI= Neutrophil-Reactive Intensity; PCT= procalcitonin; CRP= C-reactive protein.

### 3.3. 28-Day Outcomes

The comparison of inflammatory parameters for 28-day outcomes between “survivors” and “deceased” in the “septic” and “non-septic” patient group were described in Table 3. We excluded from the analysis 18 patients who were admitted in ICU after cardiac arrest, where poor prognosis was based on neurological data. NEUT-RI and PCT (not CRP) at admission in ICU in septic group were higher in patients who died compared to patients who survived (58.80 [53.85-73.10] vs 53.05 [48.90-57.22],  $p=0.005$  and 39.56 [17.39-83.72] vs 3.22 [0.59-32.32],  $p=0.002$ , respectively). No differences were found in “non-septic” group. A comparison in NEUT-RI and PCT values for discriminating between “alive” and “deceased” at 28 days were shown in Supplementary Materials, Table S3. There were not statistically significant differences between AUROC, sensibility and specificity of the two inflammatory parameters. Both NEUT-RI and PCT showed high negative predictive value (NPV) (93.0 [87.17-96.76] and 95.10 [88.93-98.39], respectively) and low positive predictive value (PPV) (28.30 [16.8-42.34] and 38.09 [23.6-54.3], respectively).

**Table 3.** Comparison of inflammatory parameters for 28-day outcomes between “survivors” and “deceased” in the “septic” and “non-septic” patient group.

	Septic		P	Non-septic		P
	Alive (n=78)	dead (n=11)		Alive (n=95)	dead (n=16)	
PCR	21.52 [11.48-137.89]	19.06 [14.10-36.49]	0.772	7.28 [1.88 - 21.70]	1.06 [0.41 - 80.00]	0.477
PCT	3.22 [0.59 -32.32]	39.56 [17.39-83.72]	0.002	0.46 [0.28 - 1.21]	0.72 [0.57 - 1.10]	0.453
NEUT-RI	53.05 [48.90-57.22]	58.80 [54.45-73.35]	0.005	47.90 [45.80 - 49.82]	45.60 [44.00 - 47.60]	0.184

## 4. Discussion

The main findings of this new retrospective analysis are as follows: 1) NEUT-RI, PCT, and PCR levels were significantly elevated in critically ill patients diagnosed with sepsis compared to those admitted for other causes; 2) Both NEUT-RI and PCT exhibited high accuracy in sepsis diagnosis,

showing an early increase shortly after the onset of systemic inflammation. This could redefine their value as crucial parameters for the early diagnosis of sepsis; 3) NEUT-RI emerged as a more effective biomarker in cases of renal failure; 4) Both NEUT-RI and PCT appeared to predict 28-day mortality upon admission to ICU.

Sepsis, as commonly understood, triggers immune system activation, leading to changes in inflammatory markers [23]. New extended inflammatory parameters like NEUT-RI, which are closely related to the appearance of immature granulocytes and neutrophil activation, are now available with just a complete blood count [16]. Our study corroborates this phenomenon, revealing a significant difference between the “septic” and “non-septic” patient groups in terms of PCT, CRP, and the extended inflammatory parameter NEUT-RI. Through the inclusion of a larger population compared to previous study [19], NEUT-RI has once again demonstrated worthy diagnostic efficacy. The optimal cut-off value is identified as 50.75 FI, surpassing the previous threshold of 51.9 FI. Although sensitivity has decreased slightly to 70.9% from the prior 80.4%, there is a notable improvement in specificity, now standing at 80.7% as opposed to the earlier 76%.

In our investigation, PCT emerged as a valuable biomarker for sepsis diagnosis, maintaining a consistent best cut-off value of 2.17 ng/mL, in line with the previous study [19]. The sensitivity, though slightly reduced to 62.9% from the earlier 69.6%, is counterbalanced by a worthily high specificity of 82.9%, only marginally lower than the previous study’s [19]. CRP, acknowledged as an acute-phase inflammatory marker indicative of the acute phase of sepsis [24], confirmed its diagnostic utility in our study. The results demonstrated a robust sensitivity of 84.6%, surpassing the prior 80.4% [19]. However, this gain in sensitivity is accompanied by a decrease in specificity to 56.2%, compared to the earlier 70.7%. This observation is certainly not new and is consistent with several previously described findings [25]. Regarding positive predictive value, both NEUT-RI and PCT exhibited similar results. However, NEUT-RI show cased a superior negative predictive value of 77.2%, compared to PCT’s 63.7%. This reaffirms the potential diagnostic efficacy of NEUT-RI in minimizing the risk of false negatives and enhancing the likelihood of accurately identifying individuals with sepsis [18,26]. Concerning the discriminatory capacity of various inflammatory biomarkers in assessing the severity of ongoing sepsis, our study, benefiting from an increased sample size, uncovered novel and noteworthy evidence. The extended inflammatory parameter NEUT-RI could help clinicians recognize early-stage sepsis with an uncomplicated course from sepsis that will evolve into septic shock. PCT further demonstrated its utility as a biomarker in this context, revealing a notable distinction between the two examined subgroups “uncomplicated sepsis” and “septic shock.” This insight holds potential for guiding the initial management of septic patients, facilitating more precise and timely therapeutic interventions, and aiding in the prompt identification of complications. We selected renal function as a parameter for sub-analyses due to its known substantial impact on sepsis biomarkers [27]. CRP is acknowledged to have a negative correlation with glomerular filtration rate, yet it remains a reliable predictor of infection in individuals with compromised renal function [28,29]. PCT, being eliminated through renal clearance, may yield elevated values in patients with acute renal failure, potentially leading to increased levels even in the absence of infections [30,31]. Consequently, its sensitivity for bacterial infection diagnosis could be compromised, particularly considering the unknown optimal cut-off value in acute renal failure cases and the unclear relationship between creatinine/urea and PCT values. Our analysis supported the idea that PCT in septic patients is indeed influenced by renal function, challenging its reliability as a predictive biomarker for sepsis. Conversely, as previously highlighted [15,18,19], NEUT-RI appears to be a more reliable biomarker in these scenarios. NEUT-RI exhibited a significant difference between the “renal failure” and “normal renal function” groups, yet this distinction was not corroborated within the individual “septic” and “non-septic” groups. This suggests that NEUT-RI is less susceptible to the influence of renal function, and the initially observed difference may be attributed to the disparate incidence of renal insufficiency between the “septic” and “non-septic” groups, with markedly higher creatinine values in the “septic” group. In our investigations, CRP also emerged as a valuable contributor to sepsis diagnosis, as it did not show a discernible influence by renal function in septic patients. Finally, the analysis of the 28-day outcome revealed that in septic

patients, NEUT-RI exhibited an AUC greater than 0.7, with a sensitivity of 62.5% and specificity of 76.9% at a best cut-off value of 53.6 FI. In comparison, procalcitonin demonstrated an AUC of 0.76, with a sensitivity of 76.2% and specificity of 79.9%, with a best cut-off value of 12.18 ng/mL. No statistically significant differences were found in AUC, sensibility, and specificity. Both NEUT-RI and PCT exhibited a low positive predictive value (28.30 and 38.09, respectively) but a high negative predictive value (93% and 95.1%).

This study has several noteworthy limitations that warrant acknowledgment. Firstly, the comparison of NEUT-RI values with those of PCT and CRP in the early stages of sepsis onset, particularly in the emergency department or ward, was not consistently feasible. This discrepancy arises from the fact that these values were not universally requested at the initial phase but were only analyzed upon admission to ICU. Furthermore, in five septic patients admitted, PCT levels exceeded 250 ng/mL, and in two patients, it surpassed 100 ng/mL. As a result, approximation of these values to 250 ng/mL and 100 ng/mL, respectively, was necessary to facilitate statistical analyses. In exploring the potential correlation between NEUT-RI and PCT with the 28-day mortality of septic patients admitted to the ICU, stratification into severity classes (such as with SAPS II) could be not statistically significant due to the small sample size. This limitation underscores the need for further investigations in the future. Additionally, it would be valuable to correlate the insights derived from NEUT-RI with IL-6 levels for early sepsis diagnosis. As indicated in the literature, IL-6 plays a pivotal role in determining the prognosis of patients in the ICU [32]. Integrating IL-6 data with the previously obtained information on NEUT-RI and PCT has the potential to enhance the specificity and/or sensitivity of the outcome, offering a more comprehensive understanding.

## 5. Conclusions

The inflammatory biomarkers analyzed in this study have proven effective in supporting physicians in the initial phase of sepsis diagnosis. In particular, the increase in NEUT-RI values seems to be a useful tool in the early diagnosis of sepsis, and it has proven effective in discriminating the severity of sepsis at the time of admission, and it would help identify septic patients who may progress to septic shock with a higher likelihood of an unfavorable long term outcome. However, these results should further its integration into other studies to confirm the reliability and generalizability of our results. This section is not mandatory but may be added if there are patents resulting from the work reported in this manuscript.

**Supplementary Materials:** The following supporting information can be downloaded at the website of this paper posted on Preprints.org. Table S1: Comparison of inflammatory parameters between patients with “renal failure” and patients with “normal renal function”; Table S2: NEUT-RI, PCT and CPR values for the detection of septic and non-septic patients; Table S3. NEUT-RI and PCT values for discriminating between “survivors” and “deceased” at 28 days; Table S4. NEUT-RI and PCT at ICU admission time, after 48 and 96 hours in septic patients; Figure S1. NEUT-RI and CPR depending on the site of infection in “septic” group; Figure S2. NEUT-RI and PCT stratified for severity classes based on SAPS II.

**Author Contributions:** L.I., E.A.M., G.S., S.P., V.R., and A.P. Conceptualization, L.I., G.S., E.S.M., S.P. and V.R.; methodology, S.P., V.R., G.S.; software, G.S.; formal analysis, M.G.; investigation, L.I., E.S.M., M.G. and F.V.; data curation, L.I., E.A.M.; writing—original draft preparation, P.F., L.I., G.S., M.U.; writing—review and editing, G.S., P.F., M.U.; supervision, A.P. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted in accordance with the Declaration of Helsinki and approved by the local Ethical Review Board (Ethics Committee of Milano Area 3 n. 114/2023, February 7, 2023).

**Informed Consent Statement:** Written informed consent was waived because of the retrospective nature of the study and data were treated anonymously according to the General Data Protection Regulation-GDPR UE 679/2016.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Singer, M.; Deutschman, C.S.; Seymour, C.W.; Shankar-Hari, M.; Annane, D.; Bauer, M.; Bellomo, R.; Bernard, G.R.; Chiche, J.-D.; Coopersmith, C.M.; et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016, 315, 801–810, doi:10.1001/jama.2016.0287.
2. Hu, B.; Ji, W.; Bo, L.; Bian, J. How to Improve the Care of Septic Patients Following “Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock 2021”? *J Intensive Med* 2023, 3, 144–146, doi:10.1016/j.jointm.2022.08.001.
3. Lambden, S.; Laterre, P.F.; Levy, M.M.; Francois, B. The SOFA Score-Development, Utility and Challenges of Accurate Assessment in Clinical Trials. *Crit Care* 2019, 23, 374, doi:10.1186/s13054-019-2663-7.
4. Evans, L.; Rhodes, A.; Alhazzani, W.; Antonelli, M.; Coopersmith, C.M.; French, C.; Machado, F.R.; McIntyre, L.; Ostermann, M.; Prescott, H.C.; et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock 2021. *Intensive Care Med* 2021, 47, 1181–1247, doi:10.1007/s00134-021-06506-y.
5. Muteeb, G.; Rehman, M.T.; Shahwan, M.; Aatif, M. Origin of Antibiotics and Antibiotic Resistance, and Their Impacts on Drug Development: A Narrative Review. *Pharmaceuticals (Basel)* 2023, 16, 1615, doi:10.3390/ph16111615.
6. Lamy, B.; Dargère, S.; Arendrup, M.C.; Parienti, J.-J.; Tattevin, P. How to Optimize the Use of Blood Cultures for the Diagnosis of Bloodstream Infections? A State-of-the Art. *Front Microbiol* 2016, 7, 697, doi:10.3389/fmicb.2016.00697.
7. Castelli, G.P.; Pognani, C.; Meisner, M.; Stuani, A.; Bellomi, D.; Sgarbi, L. Procalcitonin and C-Reactive Protein during Systemic Inflammatory Response Syndrome, Sepsis and Organ Dysfunction. *Crit Care* 2004, 8, R234-242, doi:10.1186/cc2877.
8. Qu, R.; Hu, L.; Ling, Y.; Hou, Y.; Fang, H.; Zhang, H.; Liang, S.; He, Z.; Fang, M.; Li, J.; et al. C-Reactive Protein Concentration as a Risk Predictor of Mortality in Intensive Care Unit: A Multicenter, Prospective, Observational Study. *BMC Anesthesiol* 2020, 20, 292, doi:10.1186/s12871-020-01207-3.
9. de Werra, I.; Jaccard, C.; Corradin, S.B.; Chioloro, R.; Yersin, B.; Gallati, H.; Assicot, M.; Bohuon, C.; Baumgartner, J.D.; Glauser, M.P.; et al. Cytokines, Nitrite/Nitrate, Soluble Tumor Necrosis Factor Receptors, and Procalcitonin Concentrations: Comparisons in Patients with Septic Shock, Cardiogenic Shock, and Bacterial Pneumonia. *Crit Care Med* 1997, 25, 607–613, doi:10.1097/00003246-199704000-00009.
10. Picod, A.; Morisson, L.; de Roquetaillade, C.; Sadoune, M.; Mebazaa, A.; Gayat, E.; Davison, B.A.; Cotter, G.; Chousterman, B.G. Systemic Inflammation Evaluated by Interleukin-6 or C-Reactive Protein in Critically Ill Patients: Results From the FROG-ICU Study. *Front Immunol* 2022, 13, 868348, doi:10.3389/fimmu.2022.868348.
11. Wiedermann, F.J.; Kaneider, N.; Egger, P.; Tiefenthaler, W.; Wiedermann, C.J.; Lindner, K.H.; Schobersberger, W. Migration of Human Monocytes in Response to Procalcitonin. *Crit Care Med* 2002, 30, 1112–1117, doi:10.1097/00003246-200205000-00025.
12. Wacker, C.; Prkno, A.; Brunkhorst, F.M.; Schlattmann, P. Procalcitonin as a Diagnostic Marker for Sepsis: A Systematic Review and Meta-Analysis. *Lancet Infect Dis* 2013, 13, 426–435, doi:10.1016/S1473-3099(12)70323-7.
13. Brunkhorst, F.M.; Wegscheider, K.; Forycki, Z.F.; Brunkhorst, R. Procalcitonin for Early Diagnosis and Differentiation of SIRS, Sepsis, Severe Sepsis, and Septic Shock. *Intensive Care Med* 2000, 26 Suppl 2, S148-152, doi:10.1007/BF02900728.
14. E, U.; O, B.; U, A. Role of Leucocytes Cell Population Data in the Early Detection of Sepsis. *Journal of clinical pathology* 2018, 71, doi:10.1136/jclinpath-2017-204524.
15. Zhang, W.; Zhang, Z.; Pan, S.; Li, J.; Yang, Y.; Qi, H.; Xie, J.; Qu, J. The Clinical Value of Hematological Neutrophil and Monocyte Parameters in the Diagnosis and Identification of Sepsis. *Ann Transl Med* 2021, 9, 1680, doi:10.21037/atm-21-5639.
16. Cornet, E.; Boubaya, M.; Troussard, X. Contribution of the New XN-1000 Parameters NEUT-RI and NEUT-WY for Managing Patients with Immature Granulocytes. *Int J Lab Hematol* 2015, 37, e123-126, doi:10.1111/ijlh.12372.
17. Park, D.; Chang, J.; Kahng, J.; Park, H.; Jo, I.; Kim, Y.; Han, K. Development of a Novel Flow Cytometry-Based System for White Blood Cell Differential Counts: 10-Color LeukoDiff. *Ann Lab Med* 2019, 39, 141–149, doi:10.3343/alm.2019.39.2.141.
18. Lee, J.; Gu, J.; Seo, J.E.; Kim, J.W.; Kim, H.K. Diagnostic and Prognostic Values of Neutrophil Reactivity Intensity (NEUT-RI) in Pediatric Systemic Inflammatory Response Syndrome and Sepsis. *Ann Clin Lab Sci* 2023, 53, 173–180.
19. Mantovani, E.M.A.; Formenti, P.; Pastori, S.; Roccaforte, V.; Gotti, M.; Panella, R.; Galimberti, A.; Costagliola, R.; Vetrone, F.; Umbrello, M.; et al. The Potential Role of Neutrophil-Reactive Intensity (NEUT-RI) in the Diagnosis of Sepsis in Critically Ill Patients: A Retrospective Cohort Study. *Diagnostics (Basel)* 2023, 13, 1781, doi:10.3390/diagnostics13101781.

20. Finazzi, S.; Paci, G.; Antiga, L.; Brissy, O.; Carrara, G.; Crespi, D.; Csato, G.; Csomos, A.; Duek, O.; Facchinetti, S.; et al. PROSAFE: A European Endeavor to Improve Quality of Critical Care Medicine in Seven Countries. *Minerva Anesthesiol* 2020, 86, 1305–1320, doi:10.23736/S0375-9393.20.14112-9.
21. Levey, A.S. Defining AKD: The Spectrum of AKI, AKD, and CKD. *Nephron* 2022, 146, 302–305, doi:10.1159/000516647.
22. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3) - PubMed Available online: <https://pubmed.ncbi.nlm.nih.gov/26903338/> (accessed on 22 January 2024).
23. Delano, M.J.; Ward, P.A. The Immune System's Role in Sepsis Progression, Resolution and Long-Term Outcome. *Immunol Rev* 2016, 274, 330–353, doi:10.1111/imr.12499.
24. Clyne, B.; Olshaker, J.S. The C-Reactive Protein. *J Emerg Med* 1999, 17, 1019–1025, doi:10.1016/s0736-4679(99)00135-3.
25. Lelubre, C.; Anselin, S.; Zouaoui Boudjeltia, K.; Biston, P.; Piagnerelli, M. Interpretation of C-Reactive Protein Concentrations in Critically Ill Patients. *Biomed Res Int* 2013, 2013, 124021, doi:10.1155/2013/124021.
26. Dinsdale, R.J.; Devi, A.; Hampson, P.; Wearn, C.M.; Bamford, A.L.; Hazeldine, J.; Bishop, J.; Ahmed, S.; Watson, C.; Lord, J.M.; et al. Changes in Novel Haematological Parameters Following Thermal Injury: A Prospective Observational Cohort Study. *Sci Rep* 2017, 7, 3211, doi:10.1038/s41598-017-03222-w.
27. Doi, K. Role of Kidney Injury in Sepsis. *J Intensive Care* 2016, 4, 17, doi:10.1186/s40560-016-0146-3.
28. Han, S.; Kim, M.-J.; Ko, H.-J.; Lee, E.-J.; Kim, H.-R.; Jeon, J.-W.; Ham, Y.-R.; Na, K.-R.; Lee, K.-W.; Lee, S.-I.; et al. Diagnostic and Prognostic Roles of C-Reactive Protein, Procalcitonin, and Presepsin in Acute Kidney Injury Patients Initiating Continuous Renal Replacement Therapy. *Diagnostics (Basel)* 2023, 13, 777, doi:10.3390/diagnostics13040777.
29. Stuvelling, E.M.; Hillege, H.L.; Bakker, S.J.L.; Gans, R.O.B.; De Jong, P.E.; De Zeeuw, D. C-Reactive Protein Is Associated with Renal Function Abnormalities in a Non-Diabetic Population. *Kidney Int* 2003, 63, 654–661, doi:10.1046/j.1523-1755.2003.00762.x.
30. Lu, X.-L.; Xiao, Z.-H.; Yang, M.-Y.; Zhu, Y.-M. Diagnostic Value of Serum Procalcitonin in Patients with Chronic Renal Insufficiency: A Systematic Review and Meta-Analysis. *Nephrol Dial Transplant* 2013, 28, 122–129, doi:10.1093/ndt/gfs339.
31. Steinbach, G.; Bölke, E.; Grünert, A.; Störck, M.; Orth, K. Procalcitonin in Patients with Acute and Chronic Renal Insufficiency. *Wien Klin Wochenschr* 2004, 116, 849–853, doi:10.1007/s00508-004-0279-6.
32. Vivas, M.C.; Villamarin Guerrero, H.F.; Tascon, A.J.; Valderrama-Aguirre, A. Plasma Interleukin-6 Levels Correlate with Survival in Patients with Bacterial Sepsis and Septic Shock. *Interv Med Appl Sci* 2021, 11, 224–230, doi:10.1556/1646.2020.00006.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.