

Review

Not peer-reviewed version

Estimated Cardiorespiratory Fitness and Metabolic Risks

[Robert A. Sloan](#) *

Posted Date: 2 April 2024

doi: 10.20944/preprints202404.0172.v1

Keywords: cardiorespiratory fitness; estimated CRF; metabolic risk factors; noncommunicable diseases; public health; eCRF prediction models; hypertension; hyperglycemia; dyslipidemia; obesity



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Review

Estimated Cardiorespiratory Fitness and Metabolic Risks

Robert A. Sloan

Department of Social and Behavioral Medicine, Kagoshima University Graduate Medical School, Kagoshima 890-8520, Japan; rsloan@m.kufm.kagoshima-u.ac.jp; Tel.: +81-99-275-5751; Fax: +81-99-275-5749

Abstract: This review focuses on the emerging evidence for the association between estimated cardiorespiratory fitness (eCRF) and metabolic risk factors. Given the challenges associated with directly measuring cardiorespiratory fitness (CRF) in large populations, eCRF presents a practical alternative for predicting metabolic health risks. A literature search identified six relevant cohort studies from 2021 to 2024 that investigated the association of eCRF with hypertension, hyperglycemia, dyslipidemia, and obesity. The review findings highlight a consistent inverse relationship between higher eCRF and lower incidence of metabolic risks in line with CRF cohort studies. This review highlights the importance of low eCRF as a primordial indicator for metabolic risks and underscores the need for broader application. Future research directions should include exploring eCRF's predictive accuracy across diverse populations and health outcomes and testing its real-world applicability in healthcare and public health settings.

Keywords: cardiorespiratory fitness; estimated CRF; metabolic risk factors; noncommunicable diseases; public health; eCRF prediction models; hypertension; hyperglycemia; dyslipidemia; obesity

1. Introduction

According to the recent Global Burden of Disease Study, noncommunicable diseases (NCDs) are the primary cause of death, accounting for 74% of all annual mortality [1]. Most of these deaths happen prematurely, before the age of 70. Cardiovascular diseases are the leading cause of mortality related to NCDs, resulting in the loss of 17.9 million lives annually. They are followed by cancer, type 2 diabetes (DM), and kidney disease, predominantly caused by DM [1]. Modifiable health behaviors, such as tobacco use, physical inactivity, and poor diet, significantly contribute to an increase in modifiable metabolic risk factors, including hypertension, hyperglycemia, dyslipidemia, and obesity [2]. The interplay of these risk factors increases the risk of NCD morbidity and mortality. With rising global NCD incidence rates, implementing nuanced approaches targeting metabolic risk factors may help with NCD prevention.

Cardiorespiratory Fitness and Metabolic Health

Vast evidence indicates that low cardiorespiratory fitness is a better prognostic marker of the incidence of morbidity and mortality than inactivity, sedentarism, smoking, overweight, high cholesterol, and high blood pressure [3–7]. Objectively measured cardiorespiratory fitness (CRF) is more comprehensive than traditional risk factors, and according to the American Heart Association, "CRF is directly related to the integrated function of numerous systems, and it is thus considered a reflection of *total body health*" [6]. CRF reflects the ability of an individual's cardiovascular and respiratory systems to supply oxygen to muscles during aerobic activities, serving as an indicator of mitochondrial function and efficiency. It is directly affected by physical activity, smoking, sedentary behavior, body weight, genetics, age, health status, and biological sex [4,8–10]. The gold standard to objectively measure CRF is a laboratory-conducted cardiopulmonary exercise test (CPET) measuring peak oxygen consumption expressed as relative VO_2 peak in mL O_2/kg per minute. A highly

correlated ($r = 0.96$) alternative, maximal graded exercise testing (GXT), is often used in clinical settings for diagnostic purposes [11]. A GXT estimates VO_2 peak and is expressed as a peak metabolic equivalent (MET), where 1 MET is equivalent to a resting value of 3.5 mL O_2/kg per minute. [4,6]. In a large meta-analysis of 102980 healthy adults from baseline, Kodama et al. found that low peak CRF of < 8 METS is a risk factor for all-cause mortality, coronary heart, and vascular disease [12]. They also established age and sex-specific low CRF thresholds per decade (e.g., <9 METS for men and <7 METS for women at age 40, reducing to <8 and <6 METS at age 50, and further to <7 and <5 METS by age 60). In epidemiological investigations, researchers typically categorize low CRF by age and sex into the lowest decile, quartile, or tertile. Notably, America and Japan have established reference standards that specify age and sex-specific categories for low CRF [13,14].

Recent meta-analyses highlight the importance of objectively measuring CRF to independently predict NCDs related to poor metabolic health in healthy adults from baseline. The persistent finding is the independent inverse association between CRF and NCDs amongst covariates. Low CRF is associated with a higher incidence of developing cardiovascular disease relative risk (95%CI) 1.56 (1.39-1.75), higher CRF has an inverse association with lower DM incidence, Hazard Ratio (HR) (95% CI) is 0.62 (0.49-0.77), and chronic kidney disease (CKD) incidence 0.58 (0.46-0.73)[12,15,16]. Longitudinal studies collectively show a robust inverse relationship between low CRF and all-cause mortality relative risk (95% CI) 1.70 (1.51 to 1.92), further emphasizing its significance for a healthier and longer lifespan [12]. Despite strong evidence supporting its inclusion, economic and logistical challenges limit the adoption of CRF for routine use in health care and public health.

Using CRF as a prognostic indicator supports a systems-based approach to primordial and primary prevention of metabolic health risks. An adapted version of a conceptual framework presented by Perumal et al. and Zeiher et al. of the determinants of CRF helps illustrate the interconnectedness of physical, social, behavioral, and biological determinants influencing CRF, metabolic health, and NCDs (Figure 1) [17,18]. The framework is based on social-ecological theory and underscores the need for including individual, community, and public health initiatives to improve CRF and reduce metabolic risk factors that lead to NCDs [19,20].




Figure 1. Conceptual framework of the determinants of CRF (adapted from Zeiher et al. [20]).

Limitations of Measured CRF in Healthcare and Public Health

CPET and GXT are the most precise methods for objectively measuring CRF to predict health outcomes. However, their practical application faces challenges that hinder its widespread use. These challenges include clinical guidelines, high costs, time requirements, and the necessity for specialized staff and equipment. Such obstacles make routine CRF assessment impractical in healthcare and community settings [21]. These limitations are also apparent when conducting epidemiological investigations on metabolic health outcomes. About eleven unique cohorts, such as the Aerobic Center Longitudinal Study (ACLS), are available for longitudinal analyses, containing healthy adults and measuring CRF at baseline [22].

In response to these limitations, there has been a growing emphasis on developing non-exercise estimate equations for CRF (eCRF). These equations use readily available data, such as self-reported physical activity levels, weight, and age, often found in electronic health records or collected through population health surveys. Recent reviews by Ross et al. and Wang et al. of eCRF equations have shown that these models yield moderate ($R^2 = .60$) to high correlations ($R^2 = .80$) with directly measured CRF among generally healthy adults [6,21]. Artero et al. conducted a pioneering study in 2014 on the predictability of eCRF concerning all-cause mortality and heart disease among Caucasian Americans, finding that low eCRF predicts health outcomes as effectively as low CRF [23]. However, most equations were developed using samples of Caucasian populations, potentially limiting their applicability across different ethnicities. The 2019 overview by Wang et al. identified that no eCRF studies had been conducted on metabolic health outcomes [21]. Since then, there has been a gradual rise in cohort studies utilizing eCRF to assess the incidence of metabolic health risks.

Given the recent increase in studies since Wang et al.'s 2019 review, the aim of this review is two-fold. First, synthesize the existing longitudinal research on the association between eCRF and metabolic risk factors in adult populations. Second, identify and discuss gaps in the current literature, highlighting areas for future research and practice.

2. Literature Search

This review was conducted in PubMed, Scopus, and Web of Science. The search focused on cohort studies that utilized non-exercise prediction models to estimate CRF and examined the longitudinal relationships between eCRF and metabolic risk factors, including hypertension, hyperglycemia, dyslipidemia, and obesity. Keywords used in the search encompassed combinations of "estimated cardiorespiratory fitness," "non-exercise prediction models," "metabolic health risks," and specific conditions such as "hypertension," "hyperglycemia," "dyslipidemia," and "obesity." Six studies were identified for review and published from 2021 until 2024. Table 1 provides a summary of the literature search.

Table 1. A summary table of included cohort studies.

First Author, Year of Publication	Mean Follow-Up Years from Baseline ($\pm SD$)	Cohort	Location and Sample Size	Sex	Mean Age ($\pm SD$)	eCRF model	Metabolic Risk Outcomes
Lee et al., 2021	15	Framingham Offspring Study (FOS)	America 2962	M&F	66.2 (8.6)	Jackson	Incidence of SBP ≥ 140 / DBP ≥ 90 mm Hg, Incidence of DM fasting glucose level of 126 mg/dL or higher, nonfasting glucose level of 200 mg/dL or higher, or the use of hypoglycemic medications.
Patel et al., 2022	5	Aerobics Center Longitudinal Study (ACLS)	America 5513	M&F	42.8 (9.0)	Jackson	Incidence of resting SBP ≥ 130 /DBP ≥ 80 mm Hg or self-reported, physician-diagnosed hypertension.

Cabanas-Sánchez et al., 2022	5.7 (4.4)	Taiwan MJ Cohort (TMJC)	Taiwan 200039	M&F	38.5 (12.1)	Jackson	Incidence of SBP \geq 140/ DBP \geq 90 mm Hg, serum total cholesterol \geq 240 mg/dL, and fasting blood glucose \geq 126mg/dL. Atherogenic dyslipidemia was defined as triglycerides \geq 150 mg/dL and HDL-C $<$ 40 mg/dL in men and $<$ 50 mg/dL in women.
Zhao et al., 2022	6.01(Median)	Rural Chinese Cohort Study (RCCS)	China 11825	M&F	51.0 (8.5)	Jackson	Incidence of DM was defined as fasting plasma glucose 7.0 mmol/L or current treatment with anti-diabetes medication or a self-reported history of DM, gestational diabetes mellitus, or diabetes due to other causes.
Sloan et al., 2023	4.87 (4.58)	Aerobics Center Longitudinal Study (ACLS)	America 8602	M&F	43.0 (8.9)	Sloan	Incidence of prediabetes (impaired fasting glucose) or DM as fasting plasma glucose concentrations of 100 to 125 and \geq 126 mg/dL, respectively. Those who self-reported DM or hypoglycemic medication during a follow-up were also classified as having abnormal glucose.
Liu et al., 2024	4 (Median)	China Health and Retirement Longitudinal Study (CHARLS)	China 4862	M&F	58.6 (9.4)	Jackson	Change in resting SBP, DBP, fasting triglycerides, high-density lipoprotein, total cholesterol

3. eCRF and the Incidence of Metabolic Risks

Hypertension

According to the WHO, elevated blood pressure is the primary metabolic risk factor responsible for the highest number of deaths worldwide, accounting for 19% of global mortality [24]. Since 2020, four epidemiological eCRF studies have examined the association of eCRF with the incidence of hypertension [25–28]. Cabanas-Sánchez et al. conducted a large cohort study to examine the long-term relationship (5.7 ± 4.4 years) between eCRF and key metabolic risk factors for adult cardiovascular disease. The study encompassed 200 039 healthy adults (38.5 ± 12.1 years) (50% women) from the Taiwan MJ Cohort (TMJC). The ACLS Jackson eCRF equations were used [9]. The sex-specific Jackson equations include age, body mass index, waist circumference, physical activity index, resting heart rate, and smoking as parameters to calculate eCRF (S1). From baseline, per 1-MET increase in eCRF was inversely associated with hypertension in middle-aged men and women, respectively (Hazard ratio, HR = 0.76, 95% CI, 0.75-0.78 & HR= 0.74, 95% CI, 0.72-0.76)[25]. A sub-analysis also found that minor improvements in eCRF overtime were associated with slightly lower incident rates.

In line with the findings of Cabanas-Sánchez et al., Patel et al. specifically investigated the association between eCRF and hypertension incidence within healthy middle-aged adults (42.8 ± 9.0 years) Caucasian population (N=5513, 20.1% women) from the ACLS cohort [26]. The average follow-up time was five years from baseline. Using the Jackson eCRF equation, the results support the inverse association observed by Cabanas-Sánchez et al. Men in the highest eCRF tertile had an HR = 0.74 (95% CI, 0.68-0.81) compared to those in the lowest tertile. Likewise, the risk reduction for high eCRF was greater for women, HR = 0.64 (95% CI, 0.51-0.81). In addition, a dose-response relationship was found in the cohort. Overall, every 1-MET eCRF increment corresponded to an HR = 0.90 (95% CI, 0.87-0.93) decrease in the incidence of hypertension in the overall cohort [26]. Furthermore, when

each parameter of the eCRF equation was also considered, higher fit, non-smoking, and active individuals had the lowest risk.

Lee et al. investigated the association between eCRF in healthy older adults (61.5 ± 9.2 years) and the incidence of cardiometabolic outcomes, including hypertension, across a 15-year follow-up period. The Framingham Offspring (FOS) cohort of 2,962 Caucasian participants (52.7% women) was used [27]. Unique to the study was the association of midlife eCRF with hypertension incidence, which was analyzed using three different methods. First was a single examination of eCRF during the final follow-up period. Second, eCRF trajectories were determined by examining the initial and final periods. Third, risk was determined based on mean eCRF between examination periods. Low eCRF was defined as the lowest tertile reference or trajectory group. When comparing low eCRF with high single examination eCRF, there was a lower risk of developing hypertension, $HR = 0.63$ (95% CI, 0.46-0.85). Additionally, high eCRF trajectories and high mean eCRF were associated with a lower risk of hypertension $HR = 0.54$ (95% CI, 0.34-0.87) and $HR = 0.48$ (95% CI, 0.34-0.68), respectively [27].

Rather than focusing on incidence, Liu et al. conducted a 4-year investigation utilizing data from the China Health and Retirement Longitudinal Study (CHARLS) to examine eCRF and its impact on change in an array of metabolic risk factors, including blood pressure. The population included 4862 (52.6% female) older Chinese adults aged 58.6 (9.4) [28]. Their results indicate that in the total population, those with higher baseline eCRF tend to have better arterial pressure, characterized by lower SBP and DBP per year. Those with higher baseline eCRF had significantly ($p < 0.0001$) lower SBP (β , 95%CI; -0.39, -0.52--0.25) and lower DBP (β , 95%CI; -0.19, -0.28--0.10) per year [28]. The annual change in eCRF per year was similar for DBP but not SBP.

These four studies provide longitudinal evidence for the independent inverse relationship between eCRF and hypertension and elevated blood pressure. Higher eCRF was consistently associated with lower hypertension incidence across different populations, age groups, and time frames. The TMJC used the lowest quintile to define low eCRF, while the ACLS and FOS cohorts used the lowest tertile. The findings also suggest a dose-response relationship; for every 1-MET increase in eCRF, adults had a 10-25% decrease in risk. While none of the investigations were able to cross-validate eCRF with CRF, the overall findings align with a recent systematic review and meta-analysis on the association of measured CRF and the risk of hypertension, where 1-MET increments in CRF corresponded to an 8% decrease in hypertension in adults [29].

A fundamental similarity across the investigations was that the ACLS Jackson equations were used to determine eCRF. Notably, one of the limitations of the Lee et al. investigation was that self-report physical activity data necessary for the Jackson eCRF calculations were unavailable for some of the examination years. Although the equation was initially validated in a large Caucasian population with mortality as the outcome, the overall findings suggest that low eCRF in generally healthy adults at baseline may serve as a predictor of the onset of hypertension later in life.

Hyperglycemia

The global diabetes prevalence rose from 108 million in 1980 to 537 million in 2021. Projections indicate that this figure will increase to 643 million by 2030 and 783 million by 2045. DM has a substantial impact on global mortality and morbidity, resulting in approximately 6.7 million deaths in 2021, and it increases the likelihood of severe complications such as blindness, kidney failure, heart attacks, stroke, and amputation [30]. The International Diabetes Federation highlights the importance of prevention and early detection in addressing the worldwide spread of DM [31]. Blood tests such as impaired glucose tolerance and fasting glycemia can identify prediabetes and aid in the primary prevention of DM [32]. Recent meta-analyses show that early identification of those with combined low CRF and normal blood glucose may show early signs of insulin resistance and provide prevention opportunities [15,22,33]. From the findings of the meta-analyses, researchers estimated that a 1-MET improvement in CRF leads to clinically significant (5-10%) decreases in the risk of developing DM, impacting public health [15,22,33].

Five cohort studies have explored the relationship between eCRF and hyperglycemia, mainly on the development of DM. As previously described, Lee et al. and Cabanas-Sanchez et al. investigated

eCRF and cardiometabolic risks. Lee et al. derived HRs for the onset of DM using three distinct analyses within the FOS cohort. Their findings indicated a significant inverse relationship between eCRF and DM risk; the highest tertile of eCRF was linked to a reduced risk of developing DM (HR = 0.38, 95% CI, 0.23-0.62), with similar protective effects observed across high eCRF trajectories HR = 0.27 (95% CI, 0.15-0.48) and mean eCRF, HR = 0.31 (95% CI, 0.18-0.54) [27]. Cabanas-Sanchez et al. also found evidence that eCRF can predict the incidence of DM in the TMJC. Two separate analyses investigated the incidence of DM from baseline eCRF, and the other analysis investigated the impact of changes in eCRF over time. Overall, for every 1- MET increase in eCRF, there were corresponding reductions in risk in early middle-aged men and women, respectively. From baseline, men had an HR = 0.67 (95% CI, 0.66-0.69), and women had an HR = 0.64 (95% CI, 0.61-0.66) [25]. When considering changes in eCRF over time, men's HR = 0.75 (95% CI, 0.69-0.81) and women's HR = 0.64 (0.57-0.72).

While the TMJC primarily comprised early middle-aged adults of Chinese ethnicity with a higher socioeconomic status, Zhao et al. conducted a comparable study involving 11,825 late middle-aged (51 \pm 8.5y) adults (52% women) from the Rural Chinese Cohort Study (RCCS) of Chinese ethnicity. The average of a six-year follow-up from baseline was used to determine the association of eCRF with the incidence of DM in men and women. Men in the highest eCRF quartile had an HR = 0.37 (0.22-0.62) compared to the lowest. For every 1-MET increase, there was an HR = 0.69 (0.62-0.78) [34]. Women in the highest eCRF quartile had an HR of 0.56 (0.36-0.88) compared to the lowest. For every 1-MET increase, there was an HR = 0.71 (0.62-0.88). For the total adult population, for every 1-MET increase, there was an HR = 0.89 (0.84-0.95). The four-year CHARLS results supported these inverse associations in the oldest Chinese populations studied. Using the Jackson eCRF as the exposure variable, findings indicate that in the total population, older adults with higher baseline eCRF tend to have significantly ($p < 0.0001$) lower (β , 95%CI: -0.037, -0.05--0.03) fasting blood glucose per year [28]. This association was consistent with changes in eCRF over time and in men and women, respectively.

Though the ACLS Jackson eCRF type equations are advantageous, integrating them with electronic health records might encounter challenges related to accessibility, primarily because the entry of self-reported physical activity levels is not universally standard in healthcare settings [35]. To overcome this potential barrier, Sloan et al. developed nuanced eCRF equations designed primarily for electronic health records without using physical activity status as an equation parameter [36]. The sex-specific equations were initially validated from the original ACLS cohort (N = 42676) and compared to measured CRF for accuracy (S1). The ACLS Sloan equations incorporate universal parameters that can be derived from electronic health records, including resting heart rate, height, weight, blood pressure, and smoking status. To test the predictability of eCRF with a health outcome, Sloan et al. investigated the incidence of abnormal blood glucose (prediabetes/DM) in 8602 healthy adults from baseline (17.8% women) with a mean age of 43.03 (\pm 8.94) using the ACLS cohort with an average of 5 years follow up [37]. Separate analyses were conducted for eCRF and CRF to determine the respective incidence of abnormal blood glucose. A significant inverse relationship was found for both fitness measures. Specifically, for every 1-MET increment, HRs for eCRF and CRF were determined to be 0.96, eCRF (95% CI: 0.93-0.99), and CRF (95% CI: 0.94-0.98), respectively.

Overall, these studies suggest that higher eCRF is independently and inversely associated with the development of hyperglycemia. Evidence from various cohorts, including the ACLS, FOS, TMJC, RCCS, and CHARLS, consistently supports this relationship across different populations and age segments. TMJC and RCCS, both six-year cohort studies focusing on Chinese adults, which employed the ACLS Jackson eCRF, demonstrated similar findings. This protective effect was observed regardless of variations in age, health behaviors, socioeconomic status, and environmental conditions. The types of covariates were generally similar across cohorts. Some of the cohorts did not account for prediabetes at baseline, which is a confounder for the increased risk of DM. Though only the Sloan et al. study cross-validated eCRF with CRF, the overall findings from the eCRF studies are generally aligned with CRF meta-analyses [15,22,33]. Notably, from the collective CRF meta-analyses,

women accounted for only ~16% of the general population. This underrepresentation of women is inherent in CRF cohort studies due to the lack of CPET and GXT testing data [37,38].

Dyslipidemia

Dyslipidemia includes elevated cholesterol, low-density lipid cholesterol, triglycerides, or reduced high-density lipid cholesterol [25]. Heredity and unhealthy lifestyle health behaviors increase the chance of developing dyslipidemia, which increases the risk for cardiovascular disease. Hypercholesterolemia (elevated low density-lipids), a leading form of dyslipidemia, has escalated as a risk factor for death globally, moving from the 15th position in 1990 to the 8th by 2019, indicating a growing burden of cardiovascular disease risk [39]. Atherogenic dyslipidemia, marked by high triglyceride and low high-density lipid-cholesterol levels, is especially common in individuals with DM or metabolic syndrome, exacerbating cardiovascular risks [25].

A dearth of research has been conducted on the association between eCRF and dyslipidemia incidence. From the CHARLS cohort, Liu et al. investigated the annual changes in dyslipidemia with eCRF[28]. Their results indicate that older Chinese adults with higher baseline eCRF tend to have better lipid profiles over time. Those with higher baseline eCRF had a significant ($p < 0.0001$) decrease in triglycerides per year of (β , 95%CI; -0.032 mmol·L, -0.04 to -0.03). This effect was observed in males and females. Significant ($p < 0.0001$) positive changes per year in high-density lipoprotein also occurred (β , 95%CI; 0.005 mmol·L, 0.002 to 0.007). Increases were more significant in males than females.

The findings from the TMJC provided outcomes on hypercholesterolemia and atherogenic dyslipidemia [25]. Separate analyses investigated the incidence of each outcome from baseline eCRF, and the other analyses investigated the impact of changes in eCRF per 1 -MET increase over time. From baseline, men had a reduced HR = 0.95 (95% CI, 0.93-0.96), and women had an HR = 0.98 (95% CI, 0.96-1.01) for hypercholesterolemia. When considering changes over time, the findings were similar. For atherogenic dyslipidemia, from baseline, men had an HR = 0.82 (95% CI, 0.80-0.83), and women had an HR = 0.80 (95% CI, 0.78-0.83). When considering per 1-MET changes in eCRF over time, the findings were again similar to those of the baseline analyses.

Collectively, these two investigations in middle-aged and older adults of Chinese ethnicity provide evidence that higher eCRF predicts dyslipidemia. Again, the ACLS Jackson equation was successfully used. The comparability of investigations regarding measured CRF and dyslipidemia is limited. In an ACLS cohort of healthy men (11418) at baseline, higher CRF was inversely associated with low-density lipid cholesterol and positively associated with HDL. When age was factored in, trajectories revealed that higher CRF in young to middle-aged men delayed abnormal low-density lipid cholesterol by 15 years [40]. Breneman et al. also conducted a study using the ACLS cohort of 9,651 patients (15% female). They found that higher baseline CRF and maintaining fitness (~9 years) were associated with a lower likelihood of atherogenic dyslipidemia [41].

Obesity

The World Health Organization released a 2022 report that obesity rates have doubled since 1990 to 12.2% in men and 15.7% in women, and globally 1 billion people have obesity. Globally, obesity is responsible for ~5 million deaths annually and is defined as having a $BMI \geq 30$ [42]. From a metabolic risk standpoint, a more critical measure of obesity and NCD risk is central fatness, typically measured by waist circumference with differing cut points set according to biological sex and ethnicity [43]. The health consequences of obesity include increased incidence of NCDs and premature mortality. Notably, Areto et al. found that eCRF and CRF were superior predictors of all-cause mortality, CVD mortality, and CVD morbidity compared to BMI or waist circumference in the ACLS cohort [23]. Concomitantly, a recent CHARLS cohort investigation showed clear dose-response relationships with progressively higher eCRF, predicting a lower incidence of CVD, heart disease, and stroke [44].

No studies to date have investigated the longitudinal relationship of eCRF with the onset of obesity, and very few CRF studies have investigated this relationship. Ortega et al. conducted two retrospective cohort analyses and found that low CRF is associated with a significantly increased risk

of abdominal obesity and $\text{BMI} \geq 30$ after two years in Spanish adults [45,46]. The limited literature may be partly due to the limited number of cohorts that have measured CRF to conduct this longitudinal analysis. Therefore, eCRF studies using large electronic health records or population data sets may provide a method for further investigation.

4. Discussion

Studies on eCRF and metabolic health risks are scarce. To date, six cohort investigations have been published, providing evidence for the incidence of hypertension, hyperglycemia, and dyslipidemia. No studies have been conducted on eCRF and the incidence of obesity. This review provides emerging evidence for using eCRF as a prognostic indicator for metabolic health risk. Significant inverse and dose-response associations were repeatedly demonstrated between higher eCRF and lower risk of high blood pressure, blood sugar, and abnormal lipids. These findings are aligned with previous studies using measured CRF. Most CRF cohort studies have been limited to primarily male Caucasian populations [12,15]. However, the increased use of eCRF in population health data sets has begun to expand the evidence on age groups, females, ethnicities, and socioeconomic status.

The limitations identified across some of the eCRF cohort studies in this review include concerns about low sample size, measurement accuracy, confounders, covariates, and generalizability of findings. In 2019 Wang et al. provided a scoping review of more than twenty eCRF equations [21]. At the time, five health eCRF outcome studies focused on mortality as the primary outcome. Since then, the literature has expanded to include metabolic health risk outcomes, as discussed in this review. The ACLS Jackson equation was most commonly used to calculate eCRF in five investigations [9]. The Jackson equation uses self-reported physical activity as one of the equation parameters, initially validated using the ACLS physical activity index [47]. Only the investigation by Patel et al. used the ACLS-validated scale [26]. The FOS, TMJC, RCCS, and CHARLS studies used unvalidated domestically designed questionnaires and adapted the parameter into the equation. This adaption method likely resulted in misclassification of eCRF levels in some participants, thereby reducing the accuracy and reliability of findings. It is also important to point out that the self-reported physical activity status is prone to bias, leading to misclassification.

Other commonly cited issues are the homogeneous populations studied, often with high socioeconomic status or specific ethnic backgrounds, limiting the external validity of the results. The Wang et al. review also recommended choosing equations that share the same ethnicity and age group. While there are validated eCRF equations for people of Chinese ethnicity, the CHARLS, TMJC, and RCCS used the Caucasian-validated Jackson equation with promising findings aligned with CRF meta-analyses. Notably, most of the participants in the meta-analyses are Caucasian males [22,33].

Another caution when applying eCRF equations is using redundant covariates or confounders in multivariate analyses. For example, when BMI is a parameter in an eCRF equation and is used again as a covariate during analysis, it could lead to multicollinearity. Multicollinearity occurs when two or more predictor variables in a regression model are highly correlated, meaning that one can be linearly predicted from the others with a substantial degree of accuracy [48]. This redundancy may inflate the variance of the coefficient estimates and the standard errors, making statistical tests less reliable, the model's predictions less precise, and leading to wider confidence intervals. Potential solutions include conducting variance inflation factor (VIF) analysis or transforming a continuous variable by categorizing the covariate or confounder (e.g., 1= $\text{BMI} < 30$, 2= $\text{BMI} \geq 30$) [49]. More recently, the advancement of causal inference through causal machine learning may offer a solution for more accurately accounting for covariates and confounders. Unlike associative studies that incorporate confounding variables to enhance the accuracy of outcome predictions, causal machine-learning models meticulously seek to isolate and exclude the influence of these variables to assess the impact of the exposure variable directly [50]. Furthermore, machine learning methods may be more beneficial when using large real-world data such as electronic health records.

With the current dearth of literature, there are ample opportunities for further study regarding eCRF, metabolic risks, NCDs, and a broad range of health outcomes. Potential areas for future

research may include focusing on larger multiethnic cohorts and young adults and comparing other eCRF equations for their predictive capability. Also, more evidence across diverse ethnicities and women is needed. One drawback is that a limited number of population health data sets or electronic health records contain CRF or all the parameters (e.g., Jackson) needed to calculate eCRF [36]. Different eCRF equations may need to be applied to access more extensive, heterogeneous cohorts over longer durations. As discussed by Wang et al., eCRF models that do not use self-reported physical activity as a parameter may be applied more broadly (e.g., electronic health records) [21,36].

From a metabolic health outcomes perspective, there are various potential cohort studies to consider. eCRF prediction of prehypertension, prediabetes, and borderline dyslipidemia would be helpful to inform primordial prevention initiatives. Studies focused on the incidence of obesity, metabolic syndrome, and NCDs would add significant value to the growing eCRF prediction literature. Prevalence studies for understanding the fitness level of a particular community, region, or company can help map the magnitude and distribution of low fitness and assist with public health planning. Lastly, conducting experimental intervention studies using change in eCRF would provide more evidence for the tool's validity.

A growing and essential area of eCRF research and primary care is net reclassification improvement (NRI) for risk estimation. NRI is a statistical approach that assesses the extent to which incorporating a new biomarker such as eCRF improves the classification accuracy of individuals into more appropriate risk categories [5]. For example, physicians often use the Framingham Risk Score to make patient clinical decisions. To improve the accuracy of the 10-year CHD risk score, Gander et al. applied the Jackson eCRF [51]. The study showed that adding the eCRF improves the overall accuracy of the Framingham Risk Score in Caucasian men for heart disease risk. Similar findings were also found in a nationally representative sample of Koreans and a southern Chinese population for CVD mortality and morbidity, respectively [52,53]. There are numerous other risk prediction tools (e.g., DM, CKD, dementia) where adding eCRF may add predictive value.

Future Directions

CRF has been stipulated as a vital sign by the American Heart Association, and eCRF has been proposed to be used regularly in primary care settings to identify patients with low fitness and provide brief counseling [6,54]. However, a recent meta-analysis and systematic review concluded that the effectiveness of this individualistic approach might not, *on its own*, improve physical activity, a key determinant of fitness, to sustain beyond 6 to 12 months [55,56]. In agreement with this observation, the International Society for Physical Activity and Health states, "Searching for a single solution to increasing physical activity may have hampered progress in this field by encouraging focus on simple, often short-term, individual-level health outcomes, rather than complex, multiple, upstream, population-level actions and outcomes [57]." Brief counseling may be more effective when meshed with the determinants of eCRF in an individual's environment. This method is in line with the conceptual framework of the determinants of CRF (Figure 1) and stems from social ecological theory [18]. Consequently, research has to be done on using this framework with eCRF.

Given the growing sophistication of technology, eCRF has the potential to be utilized as a *population health vital sign* to help prevent metabolic health risks. Electronic health records can auto-populate eCRF for rapid access and review [37]. Integrating geographic information systems with EHR-derived eCRF data can enhance the early identification and mapping of metabolic risk factors [58]. This integration allows public health officials to visually pinpoint areas of low fitness, referred to as hot spots, and further leverage eCRF parameters to segment and target specific populations, such as unfit middle-aged male smokers. Machine learning and artificial intelligence can further augment this process, enabling sophisticated, actionable analyses to guide targeted interventions, potentially maximizing the impact of individual, community, and public health initiatives [59].

The use of eCRF in healthcare and public health settings aligns with the International Society for Physical Activity and Health's Eight Investments That Work for Physical Activity [57]. Both initiatives focus on accessibility to diverse physical activity, exercise, and sports opportunities, facilitating the implementation of strategies like active travel and urban design. Effective

collaboration between healthcare systems and public health is crucial for navigating the complex social ecology of communities. Such partnerships are instrumental in planning and deploying a systems-based approach to reduce metabolic risks.

5. Conclusions

This review underscores the emerging evidence of eCRF as a primordial indicator of metabolic health risks. The current literature affirms a consistent inverse association between higher eCRF and reduced metabolic risks, highlighting eCRF's predictive ability to be concomitant with CRF. Future research should aim to explore eCRF's predictive accuracy across a broader spectrum of populations and outcomes and evaluate its real-world utility in healthcare and public health settings.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org. Jackson et al., and Sloan et al., equations.

Author Contributions: R.A.S.: Conceptualization, methodology, investigation, writing—original draft.

Funding: Not applicable

Institutional Review Board Statement: This review article does not require ethical approval.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: Not applicable.

Conflicts of Interest: The author declares no conflicts of interest.

References

1. Collaborators, G.B.D.R.F. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* **2020**, *396*, 1223-1249, doi:10.1016/S0140-6736(20)30752-2.
2. Li, Y.; Pan, A.; Wang, D.D.; Liu, X.; Dhana, K.; Franco, O.H.; Kaptoge, S.; Di Angelantonio, E.; Stampfer, M.; Willett, W.C.; et al. Impact of Healthy Lifestyle Factors on Life Expectancies in the US Population. *Circulation* **2018**, *138*, 345-355, doi:10.1161/CIRCULATIONAHA.117.032047.
3. Evenson, K.R.; Goto, M.M.; Furberg, R.D. Systematic review of the validity and reliability of consumer-wearable activity trackers. *Int J Behav Nutr Phys Act* **2015**, *12*, 159, doi:10.1186/s12966-015-0314-1.
4. Ross, R.; Myers, J. Cardiorespiratory Fitness and Its Place in Medicine. *RCM* **2023**, *24*, doi:10.31083/j.rcm2401014.
5. Myers, J.; Ross, R. Implementing Cardiorespiratory Fitness as a Routine Measure in Health Care Settings. *Journal of Clinical Exercise Physiology* **2021**, *10*, 62-69, doi:10.31189/2165-6193-10.2.62.
6. Ross, R.; Blair, S.N.; Arena, R.; Church, T.S.; Despres, J.P.; Franklin, B.A.; Haskell, W.L.; Kaminsky, L.A.; Levine, B.D.; Lavie, C.J.; et al. Importance of Assessing Cardiorespiratory Fitness in Clinical Practice: A Case for Fitness as a Clinical Vital Sign: A Scientific Statement From the American Heart Association. *Circulation* **2016**, *134*, e653-e699, doi:10.1161/CIR.0000000000000461.
7. Despres, J.P. Physical Activity, Sedentary Behaviours, and Cardiovascular Health: When Will Cardiorespiratory Fitness Become a Vital Sign? *Can J Cardiol* **2016**, *32*, 505-513, doi:10.1016/j.cjca.2015.12.006.
8. Lin, X.; Zhang, X.; Guo, J.; Roberts, C.K.; McKenzie, S.; Wu, W.C.; Liu, S.; Song, Y. Effects of Exercise Training on Cardiorespiratory Fitness and Biomarkers of Cardiometabolic Health: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *J Am Heart Assoc* **2015**, *4*, doi:10.1161/JAHA.115.002014.
9. Jackson, A.S.; Sui, X.; O'Connor, D.P.; Church, T.S.; Lee, D.C.; Artero, E.G.; Blair, S.N. Longitudinal cardiorespiratory fitness algorithms for clinical settings. *American journal of preventive medicine* **2012**, *43*, 512-519, doi:10.1016/j.amepre.2012.06.032.
10. Prince, S.A.; Dempsey, P.C.; Reed, J.L.; Rubin, L.; Saunders, T.J.; Ta, J.; Tomkinson, G.R.; Merucci, K.; Lang, J.J. The Effect of Sedentary Behaviour on Cardiorespiratory Fitness: A Systematic Review and Meta-Analysis. *Sports Medicine* **2024**, doi:10.1007/s40279-023-01986-y.

11. Blair SN, K.H.I., Paffenbarger RS Jr, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality: a prospective study of healthy men and women. *JAMA* **1989**, *262*, 2395–2401.
12. Kodama S, S.K., Tanaka S, Maki M, Yachi Y, Asumi M, Sugawara A, Totsuka K, Shimano H, Ohashi Y, Yamada N, Sone H. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*. **2009**, *301*, 2024-2035, doi:10.1001/jama.2009.681.
13. Akiyama, H.; Watanabe, D.; Miyachi, M. Estimated standard values of aerobic capacity according to sex and age in a Japanese population: A scoping review. *PLoS One* **2023**, *18*, e0286936, doi:10.1371/journal.pone.0286936.
14. Kaminsky, L.A.; Arena, R.; Myers, J.; Peterman, J.E.; Bonikowske, A.R.; Harber, M.P.; Medina Inojosa, J.R.; Lavie, C.J.; Squires, R.W. Updated Reference Standards for Cardiorespiratory Fitness Measured with Cardiopulmonary Exercise Testing: Data from the Fitness Registry and the Importance of Exercise National Database (FRIEND). *Mayo Clin Proc* **2022**, *97*, 285-293, doi:10.1016/j.mayocp.2021.08.020.
15. Qiu, S.; Cai, X.; Yang, B.; Du, Z.; Cai, M.; Sun, Z.; Zugel, M.; Michael Steinacker, J.; Schumann, U. Association Between Cardiorespiratory Fitness and Risk of Type 2 Diabetes: A Meta-Analysis. *Obesity (Silver Spring)* **2019**, *27*, 315-324, doi:10.1002/oby.22368.
16. Kunutsor, S.K.; Isiozor, N.M.; Myers, J.; Seidu, S.; Khunti, K.; Laukkanen, J.A. Baseline and usual cardiorespiratory fitness and the risk of chronic kidney disease: A prospective study and meta-analysis of published observational cohort studies. *Geroscience* **2023**, *45*, 1761-1774, doi:10.1007/s11357-023-00727-3.
17. Perumal, N.; Mensink, G.B.M.; Keil, T.; Finger, J.D. Why are some people more fit than others? Correlates and determinants of cardiorespiratory fitness in adults: protocol for a systematic review. *Syst Rev* **2017**, *6*, 102, doi:10.1186/s13643-017-0497-4.
18. Zeiher, J.; Ombrellaro, K.J.; Perumal, N.; Keil, T.; Mensink, G.B.M.; Finger, J.D. Correlates and Determinants of Cardiorespiratory Fitness in Adults: a Systematic Review. *Sports Med Open* **2019**, *5*, 39, doi:10.1186/s40798-019-0211-2.
19. Stokols, D. Translating social ecological theory into guidelines for community health promotion. *Am J Health Promot* **1996**, *10*, 282-298, doi:10.4278/0890-1171-10.4.282.
20. Zeiher, J.; Manz, K.; Kuntz, B.; Perumal, N.; Keil, T.; Mensink, G.B.M.; Finger, J.D. Individual and interpersonal correlates of cardiorespiratory fitness in adults - Findings from the German Health Interview and Examination Survey. *Sci Rep* **2020**, *10*, 445, doi:10.1038/s41598-019-56698-z.
21. Wang, Y., Chen, S., Lavie, C.J. et al. An Overview of Non-exercise Estimated Cardiorespiratory Fitness: Estimation Equations, Cross-Validation and Application. *Journal of Science in Sport and Exercise* **2019**, *1*, 38-53, doi:10.1007/s42978-019-0003-x.
22. Tarp, J.; Stole, A.P.; Blond, K.; Grontved, A. Cardiorespiratory fitness, muscular strength and risk of type 2 diabetes: a systematic review and meta-analysis. *Diabetologia* **2019**, *62*, 1129-1142, doi:10.1007/s00125-019-4867-4.
23. Artero, E.G.; Jackson, A.S.; Sui, X.; Lee, D.C.; O'Connor, D.P.; Lavie, C.J.; Church, T.S.; Blair, S.N. Longitudinal algorithms to estimate cardiorespiratory fitness: associations with nonfatal cardiovascular disease and disease-specific mortality. *J Am Coll Cardiol* **2014**, *63*, 2289-2296, doi:10.1016/j.jacc.2014.03.008.
24. Collaboration, N.C.D.R.F. Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet* **2021**, *398*, 957-980, doi:10.1016/S0140-6736(21)01330-1.
25. Cabanas-Sanchez, V.; Artero, E.G.; Lavie, C.J.; Higueras-Fresnillo, S.; Garcia-Esquinas, E.; Sadarangani, K.P.; Ortola, R.; Rodriguez-Artalejo, F.; Martinez-Gomez, D. Prediction of cardiovascular health by non-exercise estimated cardiorespiratory fitness. *Heart* **2020**, *106*, 1832-1838, doi:10.1136/heartjnl-2020-316871.
26. Patel, P.H.; Gates, M.; Kokkinos, P.; Lavie, C.J.; Zhang, J.; Sui, X. Non-Exercise Estimated Cardiorespiratory Fitness and Incident Hypertension. *Am J Med* **2022**, *135*, 906-914, doi:10.1016/j.amjmed.2022.01.048.
27. Lee, J.; Song, R.J.; Musa Yola, I.; Shrout, T.A.; Mitchell, G.F.; Vasan, R.S.; Xanthakis, V. Association of Estimated Cardiorespiratory Fitness in Midlife With Cardiometabolic Outcomes and Mortality. *JAMA Netw Open* **2021**, *4*, e2131284, doi:10.1001/jamanetworkopen.2021.31284.
28. Liu, Y.; Zhu, J.; Yu, J.; Zhang, X. Cardiorespiratory fitness and metabolic risk in Chinese population: evidence from a prospective cohort study. *BMC Public Health* **2024**, *24*, 522, doi:10.1186/s12889-024-17742-4.

29. Cheng, C.; Zhang, D.; Chen, S.; Duan, G. The association of cardiorespiratory fitness and the risk of hypertension: a systematic review and dose-response meta-analysis. *J Hum Hypertens* **2022**, *36*, 744-752, doi:10.1038/s41371-021-00567-8.

30. Collaborators, G.B.D.D. Global, regional, and national burden of diabetes from 1990 to 2021, with projections of prevalence to 2050: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet* **2023**, *402*, 203-234, doi:10.1016/S0140-6736(23)01301-6.

31. Saeedi, P.; Petersohn, I.; Salpea, P.; Malanda, B.; Karuranga, S.; Unwin, N.; Colagiuri, S.; Guariguata, L.; Motala, A.A.; Ogurtsova, K.; et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9(th) edition. *Diabetes Res Clin Pract* **2019**, *157*, 107843, doi:10.1016/j.diabres.2019.107843.

32. Federation, I.D. *IDF Diabetes Atlas*; 2021.

33. Zaccardi, F.; O'Donovan, G.; Webb, D.R.; Yates, T.; Kurl, S.; Khunti, K.; Davies, M.J.; Laukkanen, J.A. Cardiorespiratory fitness and risk of type 2 diabetes mellitus: A 23-year cohort study and a meta-analysis of prospective studies. *Atherosclerosis* **2015**, *243*, 131-137, doi:10.1016/j.atherosclerosis.2015.09.016.

34. Zhao, Y.; Qie, R.; Han, M.; Huang, S.; Wu, X.; Zhang, Y.; Feng, Y.; Yang, X.; Li, Y.; Wu, Y.; et al. Independent and joint associations of non-exercise cardiorespiratory fitness and obesity with risk of type 2 diabetes mellitus in the Rural Chinese Cohort Study. *Nutr Metab Cardiovasc Dis* **2022**, *32*, 929-936, doi:10.1016/j.numecd.2022.01.005.

35. Lindeman, C.; McCurdy, A.; Lamboglia, C.G.; Wohlers, B.; Pham, A.N.Q.; Sivak, A.; Spence, J.C. The extent to which family physicians record their patients' exercise in medical records: a scoping review. *BMJ Open* **2020**, *10*, e034542, doi:10.1136/bmjopen-2019-034542.

36. Sloan, R.; Visentini-Scarzanella, M.; Sawada, S.; Sui, X.; Myers, J. Estimating Cardiorespiratory Fitness Without Exercise Testing or Physical Activity Status in Healthy Adults: Regression Model Development and Validation. *JMIR Public Health Surveill* **2022**, *8*, e34717, doi:10.2196/34717.

37. Sloan, R.A.; Kim, Y.; Kenyon, J.; Visentini-Scarzanella, M.; Sawada, S.S.; Sui, X.; Lee, I.M.; Myers, J.N.; Lavie, C.J. Association between Estimated Cardiorespiratory Fitness and Abnormal Glucose Risk: A Cohort Study. *J Clin Med* **2023**, *12*, doi:10.3390/jcm12072740.

38. Sui X, H.S., Lee IM, Church TS, Colabianchi N. A prospective study of cardiorespiratory fitness and risk of type 2 diabetes in women. *Diabetes Care* **2008**, *31*, 550-555, doi:10.2337/dc07-.

39. Liu, T.; Zhao, D.; Qi, Y. Global Trends in the Epidemiology and Management of Dyslipidemia. *J Clin Med* **2022**, *11*, doi:10.3390/jcm11216377.

40. Park, Y.M.; Sui, X.; Liu, J.; Zhou, H.; Kokkinos, P.F.; Lavie, C.J.; Hardin, J.W.; Blair, S.N. The effect of cardiorespiratory fitness on age-related lipids and lipoproteins. *J Am Coll Cardiol* **2015**, *65*, 2091-2100, doi:10.1016/j.jacc.2015.03.517.

41. Breneman, C.B.; Polinski, K.; Sarzynski, M.A.; Lavie, C.J.; Kokkinos, P.F.; Ahmed, A.; Sui, X. The Impact of Cardiorespiratory Fitness Levels on the Risk of Developing Atherogenic Dyslipidemia. *Am J Med* **2016**, *129*, 1060-1066, doi:10.1016/j.amjmed.2016.05.017.

42. Collaboration, N.C.D.R.F. Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222 million children, adolescents, and adults. *Lancet* **2024**, *403*, 1027-1050, doi:10.1016/S0140-6736(23)02750-2.

43. Kodama, S.; Horikawa, C.; Fujihara, K.; Heianza, Y.; Hirasawa, R.; Yachi, Y.; Sugawara, A.; Tanaka, S.; Shimano, H.; Iida, K.T.; et al. Comparisons of the strength of associations with future type 2 diabetes risk among anthropometric obesity indicators, including waist-to-height ratio: a meta-analysis. *Am J Epidemiol* **2012**, *176*, 959-969, doi:10.1093/aje/kws172.

44. Liu, Y.; Zhu, J.; Guo, Z.; Yu, J.; Zhang, X.; Ge, H.; Zhu, Y. Estimated cardiorespiratory fitness and incident risk of cardiovascular disease in China. *BMC Public Health* **2023**, *23*, 2338, doi:10.1186/s12889-023-16864-5.

45. Ortega, R.; Grandes, G.; Agullo-Ortuno, M.T.; Gomez-Cantarino, S. Changes in Cardiorespiratory Fitness and Probability of Developing Abdominal Obesity at One and Two Years. *Int J Environ Res Public Health* **2023**, *20*, doi:10.3390/ijerph20064754.

46. Ortega, R.; Grandes, G.; Sanchez, A.; Montoya, I.; Torcal, J.; group, P. Cardiorespiratory fitness and development of abdominal obesity. *Prev Med* **2019**, *118*, 232-237, doi:10.1016/j.ypmed.2018.10.020.

47. Blair, S.N.; Kannel, W.B.; Kohl, H.W.; Goodyear, N.; Wilson, P.W. Surrogate measures of physical activity and physical fitness. Evidence for sedentary traits of resting tachycardia, obesity, and low vital capacity. *Am J Epidemiol* **1989**, *129*, 1145-1156, doi:10.1093/oxfordjournals.aje.a115236.

48. Vatcheva, K.P.; Lee, M.; McCormick, J.B.; Rahbar, M.H. Multicollinearity in Regression Analyses Conducted in Epidemiologic Studies. *Epidemiology (Sunnyvale)* **2016**, *6*, doi:10.4172/2161-1165.1000227.
49. C.F. Dormann, J.E., S. Bacher, C. Buchmann, G. Carl, G. Carré, J.R.G. Marquéz, B. Gruber, B. Lafourcade, P.J. Leitão, T. Münkemüller. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* **2013**, *38*, 27-46.
50. Shi, J.; Norgeot, B. Learning Causal Effects From Observational Data in Healthcare: A Review and Summary. *Front Med (Lausanne)* **2022**, *9*, 864882, doi:10.3389/fmed.2022.864882.
51. Gander, J.C.; Sui, X.; Hebert, J.R.; Lavie, C.J.; Hazlett, L.J.; Cai, B.; Blair, S.N. Addition of estimated cardiorespiratory fitness to the clinical assessment of 10-year coronary heart disease risk in asymptomatic men. *Prev Med Rep* **2017**, *7*, 30-37, doi:10.1016/j.pmedr.2017.05.008.
52. Sun, X.Y.; Ma, R.L.; He, J.; Ding, Y.S.; Rui, D.S.; Li, Y.; Yan, Y.Z.; Mao, Y.D.; Liao, S.Y.; He, X.; et al. Updating Framingham CVD risk score using waist circumference and estimated cardiopulmonary function: a cohort study based on a southern Xinjiang population. *BMC Public Health* **2022**, *22*, 1715, doi:10.1186/s12889-022-14110-y.
53. Lee, I.; Kim, J.; Kang, H. Adding Estimated Cardiorespiratory Fitness to the Framingham Risk Score and Mortality Risk in a Korean Population-Based Cohort Study. *Int J Environ Res Public Health* **2022**, *19*, doi:10.3390/ijerph19010510.
54. Franklin, B.A.; Wedig, I.J.; Sallis, R.E.; Lavie, C.J.; Elmer, S.J. Physical Activity and Cardiorespiratory Fitness as Modulators of Health Outcomes: A Compelling Research-Based Case Presented to the Medical Community. *Mayo Clin Proc* **2023**, *98*, 316-331, doi:10.1016/j.mayocp.2022.09.011.
55. Lamming, L.; Pears, S.; Mason, D.; Morton, K.; Bijkker, M.; Sutton, S.; Hardeman, W.; Team, V.B.I.P. What do we know about brief interventions for physical activity that could be delivered in primary care consultations? A systematic review of reviews. *Prev Med* **2017**, *99*, 152-163, doi:10.1016/j.ypmed.2017.02.017.
56. van der Wardt, V.; di Lorito, C.; Viniol, A. Promoting physical activity in primary care: a systematic review and meta-analysis. *Br J Gen Pract* **2021**, *71*, e399-e405, doi:10.3399/BJGP.2020.0817.
57. Milton, K.; Cavill, N.; Chalkley, A.; Foster, C.; Gomersall, S.; Hagstromer, M.; Kelly, P.; Kolbe-Alexander, T.; Mair, J.; McLaughlin, M.; et al. Eight Investments That Work for Physical Activity. *J Phys Act Health* **2021**, *18*, 625-630, doi:10.1123/jpah.2021-0112.
58. Laranjo, L.; Rodrigues, D.; Pereira, A.M.; Ribeiro, R.T.; Boavida, J.M. Use of Electronic Health Records and Geographic Information Systems in Public Health Surveillance of Type 2 Diabetes: A Feasibility Study. *JMIR Public Health Surveill* **2016**, *2*, e12, doi:10.2196/publichealth.4319.
59. Wiemken, T.L.; Kelley, R.R. Machine Learning in Epidemiology and Health Outcomes Research. *Annu Rev Public Health* **2020**, *41*, 21-36, doi:10.1146/annurev-publhealth-040119-094437.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.