

Article

Not peer-reviewed version

Volatile Characterization of Recovery Minority Varieties from Castilla-La Mancha Region (Spain)

[Cristina Cebrián-Tarancón](#) , [A.Sergio Serrano](#) , [Juan L. Chacón-Vozmediano](#) , [Jesús Martínez-Gascuña](#) , [Gonzalo L. Alonso](#) *

Posted Date: 18 April 2024

doi: 10.20944/preprints202404.1220.v1

Keywords: recovered grapevine cultivars; aroma compounds; aroma potential; water stress



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Volatile Characterization of Recovery Minority Varieties from Castilla-La Mancha Region (Spain)

Cristina Cebrián-Tarancón ¹, A.Sergio Serrano ^{1,2}, Juan L. Chacón-Vozmediano ²,
Jesús Martínez-Gascuña ² and Gonzalo Alonso ^{1,*}

¹ Department of Agricultural Chemistry, School of Agricultural and Forestry Engineering and Biotechnology, University of Castilla-La Mancha, Avda. de España s/n, 02071 Albacete, Spain

² Regional Institute of Agri-Food and Forestry Research and Development of Castilla-La Mancha (IRIAF), Ctra. Toledo-Albacete s/n, 13700 Tomelloso, Spain

* Correspondence: gonzalo.alonso@uclm.es

Abstract: Nowadays, the identification and characterization of grapevine cultivars resilient to climate and water stress while preserving quality traits is crucial for the wine industry. Therefore, the objective of this work was to characterize according to their aromatic potential 9 white and 6 red minority cultivars recently recovered from Castilla-La Mancha region (Spain), subjected at two different water deficit regimes: rainfed, with subsistence irrigation, and irrigated. For this, the varietal aromatic potential index (IPAv) and the detailed aromatic composition were analyzed by HS-SBSE-GC/MS, in extracts of two different ph. For IPAv values, red varieties did not show a clear trend with respect to irrigation. However, in white minority varieties, higher values were obtained under irrigation conditions. Thus, a clear differentiation of the minority varieties in comparison to the references was observed, primarily attributed to the content of esters and acids, in both white and red varieties. A notable contrast was observed at different pH, indicating a greater extractability of certain compounds like linalool, under more acidic conditions. This suggests that some recovered minority cultivars could be promising for cultivation in semi-arid regions with limited water, contributing to the wine sector sustainability in the future.

Keywords: recovered grapevine cultivars; aroma compounds; aroma potential; water stress

1. Introduction

In recent years, viticulture has been suffered significant challenges due to the climate change, which has significantly transformed the climatic conditions traditionally associated with grape production. Rising temperatures, extreme weather events, and changes in rainfall patterns are altering the viticultural landscape, affecting both the quality and quantity of the harvest. Historic wine-growing regions are compelled to reconsider their agricultural practices and grape varieties, aiming to adapt to an ever-evolving environment. Thus, the need for sustainable management, innovation in cultivation techniques, and the identification of grape varieties better adapted to heat and drought become crucial for the preservation and ongoing success of the wine industry in this new climatic scenario.

In this context, the recovery of minority varieties is a topic of great relevance for the sector. As industrialization and the standardization of crops progress, many traditional plant varieties have been marginalized and some of them are even on the verge of extinction. However, these varieties often possess unique characteristics and specific adaptations to their development environment, making them valuable from an agronomic perspective in ensuring resilience against climate change. In this line, the application of advanced genomic sequencing techniques has enabled researchers to identify and analyzed specific genetic markers that have allowed highlighting the rich genetic diversity of recovered varieties, which could make them able to adapt to different weather conditions, resist diseases, and improve the quality of wines [1–5]. Likewise, in recent years, numerous studies

have been conducted to explore the agronomic adaptation of these grape varieties under different water regimes. These investigations have studied the relationship between plants and their environment by assessing stable isotope ratios, primarily focusing on carbon ($\delta^{13}C$), and water potentials, among other factors. The findings from these studies highlight the capability of certain cultivars to effectively confront and adapt to diverse climatic conditions while maintaining yields and general enological parameters [6–9].

Regarding to the chemical composition of grapes, understanding the impact of water stress on the phenolic and aromatic composition, both free and bound, of these recovered minority grapes is of great importance, as these compounds play a crucial role in the quality and organoleptic characteristics of the wine. The impact of irrigation on the phenolic composition of berries in traditional varieties has been thoroughly studied [9–12]. Thus, this effect on minority varieties has recently been considered [13], whose results revealed that some of the traditional varieties, cultivated under moderate and severe water stress conditions, are capable of maintaining productions with an adequate phenolic composition, similar to or even more favourable than other varieties more widespread. In the same line, various authors have observed a strong influence on the accumulation of volatiles compounds in different widely cultivated varieties due to the environmental factors, such as water availability, being positively or negatively affected by water stress depends on the family of compounds under study [14–17]. Nevertheless, the response of volatile composition to water deficit in a great number of recently recovered minority varieties remains unknown. However, this is a critical aspect, considering the potential value of these varieties in adapting to current and future climatic conditions. They can function as valuable resources, contributing to the diversity and sensory complexity of the wines produced.

Therefore, considering all of the aforementioned factors, the aim of this work is to characterize the volatile composition under two different water regimes of 9 widespread and 15 recovered/minority grape varieties, both red and white, traditionally cultivated or recovered on the verge of extinction in Castilla-La Mancha region (Spain).

2. Results and Discussion

This section may be divided by subheadings. It should provide a concise and precise description of the experimental results, their interpretation, as well as the experimental conclusions that can be drawn.

2.1. *Ipav*

Water availability is a crucial factor that influences the development of aromatic compounds in grapes. Therefore, understanding the adaptability of numerous varieties to water stress and its impact on the aromatic potential of grapes is essential. In this context, the IPAv parameter serves as an indicator of the global content of glycosylated aroma precursors in grapes, that includes mostly volatile aglycones such as alcohols, terpenes, phenols, and C13-norisoprenoids [21]. Thus, a higher value of this index indicates that grapes could have a greater amount of glycosylated aroma precursors, which could be liberated during the winemaking process, contributing positively to the aromatic characteristics of wines. The IPAv values of white and red grapes, in both water regimes, are shown in Table 2.

Table 2. Varietal aroma potential index (IPAv) of red and white grapes.

	Rainfed	Irrigation	<i>p</i>
<i>Red varieties</i>			
Bobal	10.83±0.11	12.14±0.06	***
Garnacha Tinta	18.37±0.06	14.50±0.08	***
Merlot	10.90±0.04	10.12±0.30	*
Syrah	7.46±0.02	8.79±0.15	***
Tempranillo	14.06±0.01	9.23±0.02	***

Benedicto	11.73±0.03	16.67±0.02	***
Moravia Agria	7.31±0.08	6.88±0.02	***
Moribel	11.83±0.01	9.49±0.01	***
Tinto Fragoso	8.22±0.04	9.92±0.24	***
Tinto Velasco	13.24±0.06	8.48±0.06	***
Tortozona Tinta	4.38±0.02	5.43±0.06	***
<i>White varieties</i>			
Airén	7.93±0.02	19.53±0.31	***
Chardonnay	5.97±0.03	4.88±0.02	***
Macabeo	9.51±0.17	14.85±0.09	***
Riesling	12.02±0.04	15.71±0.08	***
Albillo Dorado	17.61±0.14	21.76±0.26	***
Azargón	13.69±0.06	14.54±0.11	***
Blanca del Tollo	7.98±0.02	7.66±0.02	***
Jarrosuelto	6.41±0.08	6.54±0.02	*
Maquías	7.69±0.04	6.58±0.00	***
Mizancho	8.34±0.01	9.26±0.07	***
Montonera del Casar	5.76±0.04	7.19±0.04	***
Moscatel Serrano	12.08±0.09	20.51±0.06	***
Pintada	7.36±0.06	11.96±0.09	***

For each variety, differences between water regimes are indicated according to Fisher's LSD test.

The highest values are in bold. * p value < 0.05; ** p value < 0.01; *** p value < 0.001. Bold values indicate the highest value within a variety.

In spite of some papers has suggested that water deficit resulted in an improved of grapes aroma [22], the IPAv values in white varieties was higher under the irrigation regime compared to rainfed for most varieties, showing a positive influence of irrigation on the development of glycosylated aroma precursors in grapes. In reference varieties, the higher increases between the two regimes were found in Airén and Macabeo, with values of 19.53 and 14.85 respectively, so irrigation seems to have a more pronounced impact on the aromatic potential of these grape, which has already been pointed by other authors [14]. Regarding to recovered varieties, Moscatel Serrano showed the highest increase in IPAv value under irrigation regime, 20.51 versus 12.08 in rainfed conditions. This variety, together with Albillo Dorado, showed higher IPAv values than the reference ones, in both water regimes, which suggests that they could be a good growing alternative in these climate change conditions, resulting in wines with a different aromatic profile. By contrast, in Azargón, Jarrosuelto, or Mizancho, differences between the two regimes were less notable, so these cultivars have a moderate response to irrigation in terms of aromatic potential. Only the varieties Chardonnay and Maquías exhibited lower IPAv values under the irrigation regime compared to rainfed, 5.97 and 7.69, respectively.

In the case of red varieties, an opposite behavior of white ones was observed, with IPAv values lower under the irrigation regime for most varieties, being Garnacha Tinta and Tempranillo the reference varieties that showed the highest decrease, suggesting a sensible particularly responsive to the additional water supply, resulting in a decrease of the aromatic potential characteristics. On his part, of the recovered varieties, Tinto Velasco was the minority red variety most influenced by the water regime, reducing its IPAv value from 13.24 to 8.48. On the other hand, Benedicto was the red cultivar that showed a highest increase in IPAv value under irrigation regime, from 11.73 to 16.67, showing higher values than any of the reference varieties, which suggest an important aromatic potential of this variety if it is grown under irrigation conditions.

In summary, the data shown that irrigation tended to increase the aromatic potential of white grape varieties. However, this tendency was not as clear for red varieties, as the magnitude of this effect varied depending on the particular variety. Even so, most of the minority varieties studied have shown IPAv values under rainfed conditions similar to, and in some cases even higher, than those of the reference varieties. Therefore, although it is important to consider the individual characteristics

and responses of each grape variety when applying irrigation strategies to enhance their aromatic qualities, these results allow us to consider these varieties as a good alternative for cultivation in wine-growing regions most affected by these climate change conditions.

2.2. Volatile Composition of Grapes

In order to know the similarity and correlation between the aromatic profile of reference and minority varieties, a heatmap with cluster analysis was carried out for white and red grapes. In this way, the smaller the distance separating two clusters, the greater the similarity between the samples contained within these clusters. So, Figures 1 and 2 show the results obtained for the white and red varieties, respectively. In the case of white grapes, we can distinguish two well-differentiated clusters. Once of the cluster, contains a single group (G1) with the reference varieties Macabeo and Chardonnay, in both water regimes. Instead, the second cluster contains four groups (G2, G3, G4, and G5) with all minority varieties and the rest of the reference varieties. In the same way, in the case of red grapes another two main clusters are also differentiated. In the first we find two groups (G1 and G2), which encompass the traditional varieties Tempranillo, Garnacha Tinta, and Bobal. In the second main cluster, there are four groups (G3, G4, G5, and G6) where all the minority varieties are grouped along with Syrah and Merlot among the traditional ones.

In both white and red grape varieties, it is noteworthy that the primary differentiation between these groups primarily hinges on acids, except for hexanoic acid, and esters, except for ethyl decanoate. The heightened concentration of these compounds may be attributed to the genetic adaptation of these minority grape varieties to the specific region. This genetic adaptation could potentially activate distinct metabolic pathways, resulting in the increased accumulation of these compounds as part of their adaptive response. Hence, this grouping implies that the recovered varieties under examination could yield wines with an aromatic profile distinct from those produced with the traditional varieties of the region.




Figure 1. Heatmaps of white grapes aromatic compounds at pH 3.6.




Figure 2. Heatmaps of red grapes aromatic compounds at pH 3.6.

The identified compounds were grouped into acids, alcohols, esters, norisoprenoids, aldehydes, and terpenes. Acids were represented by octanoic, hexanoic, and decanoic acid. Although the sensorial contribution of these compounds is related with unpleasant odour descriptors such as rancid or cheese [23], some authors [24] suggest that they contribute to wine fresh flavour and they also help to modify the perception of other taste sensations. In the case of white grapes (Figure 1), varieties of the group 1 would be the most abundant, followed by groups 4 and 5. Thus, within these, the most acidic variety was Macabeo under irrigation, where its concentration of octanoic acid stood out; however, in rainfed conditions, the content of this compound decreased significantly. The second most acidic variety was Chardonnay, in both regimes, although in this case, the most abundant compound was decanoic acid, mainly in rainfed conditions. Similarly, if we focus on the recovered varieties, Maquías and Mizancho (G5) were the most acidic, but unlike the traditional ones, the most abundant compound was hexanoic acid, although in lower concentrations than the previous ones, which was positively affected (increased) by irrigation in the Maquías variety and negatively affected (decreased) in Mizancho. In red grapes, the traditional varieties Garnacha, Tempranillo, and Bobal (G1 and G2) showed the highest concentration of these compounds. Although some authors have not observed an effect on the concentration of these compounds by water status in the vineyards [25], this study highlighted the total levels in the Tempranillo variety under irrigation, which exhibited concentrations almost ten times higher than the rest of the varieties. On the other hand, the recovered varieties presented very low concentrations of these compounds.

Regarding to alcohols, in the case of white grapes, varieties of groups G2 and G3 showed the highest concentration of these, with 1-hexanol and 2-hexen-1-ol being the two most abundant compounds. Among all the varieties, those that presented a higher concentration of these compounds were the minority ones, such as Blanca del Tollo and Jarrosuelto, being negatively affected by irrigation. This behavior has already been described previously by other authors, who suggested that low water availability increased the concentration of some C6 compounds [25,26]. However, this trend was not as clear in all varieties of this study, since, among the recovered varieties, Pintada and Azargón showed higher concentrations under irrigation conditions. In the case of red grapes, the varieties in groups 5 and 6 showed the highest levels of these compounds, with the highest levels of 1 hexanol and 2-hexen-1-ol in Tinto Velasco, regardless of the water regime, and the 3 hexen-1-ol and benzyl alcohol content in the varieties Moravia Agria and Tinto Frago in irrigation conditions,

respectively. Thus, of the traditional varieties, the presence of phenylethyl alcohol in the Tempranillo variety under irrigation conditions was of note. In terms of the contribution to the aromatic profile of wines of this group of compounds, C6 alcohols are considered to be responsible for the herbaceous aroma of wines [27], but 2-phenylethyl alcohol is an aromatic alcohol that contributes with rose notes to the wine [28].

Terpenes are one of the most important compounds in wine, which contribute to the varietal characteristics of wine thanks to their flowery and sweet aroma nature [29,30] and, in this work, linalool, geraniol, and geranyl acetone were identified. In the case of white varieties (Figure 1), linalool was only detected in the Riesling and Moscatel Serrano varieties, while the latter two (geraniol and geranyl acetone) were found in the remaining varieties. As expected, the most terpene-rich varieties were Moscatel Serrano and Riesling (G4 and G5), attributed to the presence of linalool, which was not found in the rest of the varieties. The concentration of these compounds in the other recovered varieties was very similar, except in Blanca del Tollo, where only geranyl acetone was identified. Moreover, the highest concentration of terpenes in these varieties was observed in Jarrosuelto under irrigation conditions. In terms of the effect of the water supply, despite some authors suggesting that controlled water stress triggers specific biochemical responses favoring terpene synthesis [14,22], this behavior was only observed in the Albillo Dorado and Chardonnay varieties, with even the Mizancho variety decreasing terpene content under water deficit. Therefore, the fact that these recovered varieties show similar or even higher concentrations of terpenes compared to some traditional varieties (Macabeo or Airén), regardless of the water regime, indicates that they could give rise to wines with distinctive and complex aromatic profiles. On the other hand, concerning red grapes (Figure 2), generally the most terpene-rich varieties were found within groups 4, 5, and 6. In this case, the effect of water availability was observed. Thus, the highest concentration of geraniol was found in Tinto Fragoso under irrigation conditions, followed by Tinto Velasco and Moravia Agria, but in this instance under rainfed conditions. Also, among the recovered varieties, the content of geranyl acetone stood out in the Benedicto variety under irrigation conditions. Finally, regarding linalool, Tinto Velasco showed the highest content among the recovered varieties, although it is noteworthy the content in Garnacha Tinta (G1) under irrigation conditions.

Esters compounds have been positively associated in sensory studies with the overall aroma of the wine, including red and dark fruits, among others. In this work, these group of compounds was the most affected by water regime, showing lower concentrations of these compounds under rainfed conditions, except for the Chardonnay and Moscatel Serrano varieties. These results are consistent with the results obtained by (Talaverano et al., (2018), who have suggested that water restricted treatment negatively affects the fatty acid metabolism, from which these esters are derived. Among the minority varieties, Jarrosuelto, Azargón, and Maquíás showed the highest amount of these compounds. In the case of red grapes, the varieties with the highest content of these compounds were the traditional Bobal and Tempranillo, mainly under irrigation conditions. Among the minority varieties, a higher content of ethyl decanoate was observed in the Tinto Velasco variety, regardless of the water regime.

Norisoprenoids represent key odoriferous compounds in wine most of which are characterized by floral and fruity pleasant notes [32] and were represented by β -ionone and β -damascenone in this work. In white grapes (Figure 1), among the minority varieties studied, Mizancho and Maquíás were the ones that showed a higher concentration of these compounds, in line with the values obtained for the Riesling variety. On his part, in the case of red grapes (Figure 2), the varieties of group 6 (Tortozona Tinta, Moribel, and Tinto Fragoso (irrigation)) had the highest concentration of these compounds, with similar ionone levels in all of them and damascenone standing out in Tortozona Tinta in rainfed conditions.

Finally, aldehydes were represented by nonanal, benzaldehyde, and hexenal. In the case of white grapes, varieties of the group 5 showed the highest concentration of these compounds, with 2-hexenal standing out, mainly due to its high concentration in the variety Maquíás in rainfed conditions. Regarding to red grapes, the highest concentrations of these compounds were observed in groups 5

and 6, indicating that the recovered varieties have a higher amount of these compounds. Thus, the concentration of hexenal in Tortozona Tinta, in both regimes, was the highest of all.

To determine the effect of the three main factors considered in this work (pH, water regime, and grape variety) on the volatile composition of grapes, as well as the groups of compounds that are most affected by these factors, a multivariate analysis (MANOVA) was performed (Table S1). Among the individual factors, pH and grape variety had the most significant effect in alcohols, aldehydes, and norisoprenoids groups of compounds.

Finally, with the purpose of obtaining an overall view of the total content of aroma compounds and not only the easily extractable, a principal component analysis were performed with the aromatic compounds groups of the extracts obtained at bot pH (1 and 3.6), in white and red varieties and with both water regimes (Figures 3 and 4).




Figure 3. PCA with the white grapes aromatic compounds groups at pH 1 and pH 3.6 and two water regimes. **a)** rainfed conditions; **b)** irrigation conditions.




Figure 4. PCA with the red grapes aromatic compounds groups at pH 1 and pH 3.6 and two water regimes. **a)** rainfed conditions; **b)** irrigation conditions.

In relation to Figure 3, in both water regimes, a clear separation was observed in component 1 between the traditional Macabeo and Chardonnay varieties and those recovered, indicating the aromatic disparity between the two varieties. Thus, in Figure 3a, when moving to component 2, it becomes evident that the extracts obtained at pH 1 presented a greater concentration of terpenes and norisoprenoids, highlighting the disparity in the Riesling variety between the two regimes. These suggest a more efficient extraction of aromatic compounds from grapes under acidic conditions,

thereby justifying the higher extractability observed at pH 1 compared to neutral or basic extraction conditions. On the other hand, in Figure 3b, focusing on the negative part of component 2, it is noticeable that the extracts obtained at pH 1 were characterized by an increase in the content of norisoprenoids. These results suggest that, in the recovered white varieties, dry conditions favor the enhancement of aromatic precursors of terpenes and norisoprenoids. This implies that the wines obtained could have a richer aromatic profile under water stress conditions and, therefore, confirms that they could be a good alternative for cultivation in areas with water scarcity.

Regarding red varieties (Figure 4), the effect of pH on the increase in the content of certain groups of compounds was not as clear. Thus, under rainfed conditions (Figure 4a), a significant separation was observed in component 1 between the traditional varieties Garnacha, Bobal, and Tempranillo, and the recovered minority varieties. On the other hand, when shifting to component 2, it can be observed that the Tempranillo variety, a reference in Spanish wines, had a less aromatic profile in terms of terpenes and norisoprenoids compared to other traditional varieties. Finally, under irrigation conditions, a clear separation is observed in the extracts at pH 1 of the Tempranillo and Garnacha Tinta varieties, with these showing a higher content of all the studied compound groups.

3. Materials and Methods

3.1. Grapevines and Water Regime

Grapes were sampled in 2021 year from a multivarietal experimental vineyard located in the Regional Institute of Agri-Food and Forestry Research and Development of Castilla-La Mancha (IRIAF), in Tomelloso, Castilla-La Mancha, Spain (latitude 39°10'14'' N, longitude 3°00'16'' W; altitude 660 m.a.s.l.). The climate of the area is continental semi-arid Mediterranean. The soil of the plot is a Petrocalcic calcixerept (USDA soil classification) with a shallow horizon at 30 cm depth that restricts vine roots. The vineyard was planted in 2008 and grafted onto Fercal rootstock. The vines were planted at 2.8 m between rows and 1.2 m between vines (2,976 vines/ha) and were trained using the VSP trellis system and the bilateral Royat cordon formation. For the study, 9 widespread varieties (5 red and 4 white) and 15 minority/recovered ones (6 red and 9 white) were selected (Table 1).

Table 1. Studied varieties.

White		Red	
Reference	Airén (AI)	Reference	Bobal (BO)
	Chardonnay (CH)		Garnacha Tinta (GT)
	Macabeo (MAC)		Merlot (ME)
	Riesling (RI)		Syrah (SY)
Minority/recovered	Albillo Dorado (AD)	Minority/recovered	Tempranillo (TE)
	Azargón (AZ)		Benedicto (BE)
	Blanca del Tollo (BT)		Moravia Agria (MA)
	Jarrosuelto (JA)		Moribel (MO)
	Maquías (MAQ)		Tinto Fragoso (TF)
	Mizancho (MI)		Tinto Velasco (TV)
	Montonera del Casar (MC)		Tortozona Tinta (TT)
	Moscatel Serrano (MS)		
	Pintada (PI)		

For each variety, 50 vines were studied and subjected to two different water deficit regimes: 20 vines were grown under rainfed conditions with survival irrigation (rainfed treatment, 320 m3/ha supplied), and 30 vines were grown under irrigation (irrigation treatment, 980 m3/ha supplied). To determine the harvest date, regular sampling was conducted until the grapes reached a concentration of total soluble solids between 22.5 and 24.5°Brix for red cultivars, and between 20 and 22°Brix for white cultivars. At that time, a representative quantity of grapes of about 1.5 kg from each water treatment was harvested and frozen for subsequent analysis.

3.2. Grape Extraction Procedure

For the extraction, 200 berries randomly selected were crushed and homogenized at 4000 r.p.m. Then, 50 g of the mixture was weighed in two centrifuge flasks and 50 mL of pH 3.6 aqueous solution (5 g/L tartaric acid adjusted to pH 3.6 with NaOH) was added to one and 50 mL of pH 1 aqueous solution (HCl 0.1 N at pH 1) to the other. The solutions were homogenised and macerated for 24 h at room temperature ($21^{\circ}\text{C} \pm 3$) and after centrifugation at 4000 rpm for 10 min the limpid supernatant was recovered. Two extracts were obtained (pH 1: total extraction potential; and pH 3.6: easily extractable compounds) on which analytical determinations were performed.

3.3. Chemical Analysis

3.3.1. Volatile Compounds

Volatiles were determined according to (Sánchez-Gómez et al., (2016). Briefly, the extraction was carried out by means of Headspace Stir bar Sorptive Extraction (HS-SBSE) with a polydimethylsiloxane twister bar (10 mm length, 0.5 mm film thickness). 22 ml of grapes extracts were placed in a 50 mL vial and a polydimethylsiloxane coated stir bar (twister, 0.5 mm film thickness, 10 mm length; Gerstel, Mülheim, and der Ruhr, Germany) was inserted into the twister-headspace vial and hermetically closed. The vial was stirred during 1 h at 500 rpm with a temperature of 60°C . Later, analysis was performed using an automated thermal desorption unit (TDU, Gerstel, Mülheim and der Ruhr, Germany) mounted on an Agilent 7890A gas chromatograph system (GC) coupled to a quadrupole Agilent 5975C electron ionisation mass spectrometric detector (MS, Agilent Technologies, Palo Alto, CA, USA) equipped with a fused silica capillary column (BP21 stationary phase; 30 m length; 0.25 mm I.D.; and 0.25 μm film thickness) (SGE, Ringwood, Australia). The carrier gas was helium with a constant column pressure of 20.75 psi.

The Stir Bars were thermally desorbed in a stream of helium carrier gas at a flow rate of 75 mL/min with the TDU programmed from -10°C to 295°C (held 5 min) at a rate of $60^{\circ}\text{C}/\text{min}$ at splitless desorption mode. The analytes were focused on a programmed temperature vaporizing injector (PTV) (CIS-4, Gerstel), containing a packed liner (20 mg tenax TA), held at -10°C with cryo cooling prior to injection. After desorption and focusing, the CIS-4 was programmed from -10°C to 260°C (held for 5 min) at $12^{\circ}\text{C}/\text{min}$ to transfer the trapped volatiles onto the analytical column. The GC oven temperature was programmed to 40°C (held for 2 min), raised to 80°C ($5^{\circ}\text{C}/\text{min}$, held for 2 min), raised to 130°C ($10^{\circ}\text{C}/\text{min}$, held for 5 min), raised to 150°C ($5^{\circ}\text{C}/\text{min}$, held for 5 min), and then raised to 230°C ($10^{\circ}\text{C}/\text{min}$, held for 5 min). The MS was operated in scan acquisition (27–300 m/z) with an ionization energy of 70 eV. The temperature of the MS transfer line was maintained at 230°C . MS data acquisition was carried out at positive scan mode, although to avoid matrix interferences, the MS quantification was performed in the single ion-monitoring mode using their characteristic m/z values. Information related to analyzed compounds and m/z values are included in greater detail in Sánchez-Gómez et al. (Sánchez-Gómez et al., (2016) work. Compounds identification was performed using the NIST library and confirmed by comparison with the mass spectra and retention time of their pure standards (Sigma-Aldrich, Steinheim, Germany). As internal standard, 3-methyl-1-pentanol was used. Quantification was based on calibration curves of the respective standards at five different concentrations ($R^2 = 0.95\text{--}0.97$). The analyses of each replicate of grape extracts were made in triplicate.

3.3.2. Varietal Aroma Potential Index

The varietal aroma potential index (IPAv) of grapes was analyzed using a commercially available IPAv kit (Teknokroma, Barcelona, Spain) according to Serrano de la Hoz et al. (Serrano de la Hoz et al., (2014). The method, based on Salinas et al. (Salinas et al., (2012), is an spectrophotometric determination of the glucose released from the glycosylated aroma precursors by acid hydrolysis. Samples were analyzed in triplicate.

3.3.3. Statistical Analysis

Heatmaps were performed using MetaboAnalyst 5.0 Web service. Raw data were normalized by median and auto scaled (mean-centered and divided by standard deviation of each variable) prior to the analysis. Moreover, one-way analysis of variance (ANOVA) at a 95% probability level, according to the Fisher posthoc test, to determine the differences between the IPAv grapes values. Finally, a multivariate analysis of variance (MANOVA) and a principal component analysis (PCA) were performed. These analyses were conducted using the Statgraphics Centurion statistical program (version 18.1.12; StatPoint, Inc., The Plains, VA, USA).

4. Conclusions

The results of this work showed significant variations among cultivars and water management approaches investigated, generally indicating elevated levels of aromatic compounds in the rainfed regime. Some recovered minority varieties such as Maquías, Moscatel Serrano, Tinto Frago, and Tortozona Tinta stood out, which grown under low water availability conditions, showed a similar or even more suitable aromatic composition in grapes than other varieties whose cultivation is more widespread, such as Chardonnay, Merlot, and Syrah. This suggests potential adaptability to forthcoming climate change conditions in these specific scenarios.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Table S1. MANOVA of white and red grapes. Factors: pH, water regime and grape variety; Table S2. Component Weights for Figures 3 and 4.

Author Contributions: Conceptualization, C.C.T, G.L.A and J.L.C.-V.; methodology, C.C.T, G.L.A and A.S.S.; software, C.C.T and A.S.S.; validation, C.C.T and A.S.S.; formal analysis, C.C.T, G.L.A and A.S.S.; investigation, C.C.T and A.S.S.; resources, G.L.A. and J.L.C.-V.; data curation, C.C.T. and A.S.S. ; writing—original draft preparation: C.C.T; writing—review and editing: C.C.T, A.S.S., G.L.A., J.L.C.-V. and J.M.-G.; visualization, C.C.T, A.S.S. and G.L.A. ; supervision, C.C.T and G.L.A.; project administration, J.L.C.-V. and G.L.A.; funding acquisition, J.L.C.-V. and G.L.A. All authors have read and agreed to the published version of the manuscript.

Acknowledgments: We are grateful to the Government of Castilla-La Mancha (Spain) in collaboration with the ERDF for funding the project SBPLY/21/180501/000014 and also to the UCLM for funding the project 2023-GRIN-34180. Moreover, C.C.-T., thanks the Government of Castilla-La Mancha for the postdoctoral contract 2023-POST-21499 funded through the ESF+ programme 2021-2027.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Urrestarazu, J.; Miranda, C.; Santesteban, L.G.; Royo, J.B. Recovery and Identification of Grapevine Varieties Cultivated in Old Vineyards from Navarre (Northeastern Spain). *Sci Hort* **2015**, *191*, 65–73. <https://doi.org/10.1016/J.SCIHORT.2015.04.029>.
- García, J.; Peiró, R.; Martínez-Gil, F.; Soler, J.X.; Jiménez, C.; Yuste, A.; Xirivella, C.; Gisbert, C. Recovering Old Grapevine Varieties. *Vitis - Journal of Grapevine Research* **2020**, *59*, 101–103. <https://doi.org/10.5073/vitis.2020.59.101-103>.
- Mena, A.; Martínez, J.; Fernández-González, M. Recovery, Identification and Relationships by Microsatellite Analysis of Ancient Grapevine Cultivars from Castilla-La Mancha: The Largest Wine Growing Region in the World. *Genet Resour Crop Evol* **2014**, *61*, 625–637. <https://doi.org/10.1007/S10722-013-0064-3/FIGURES/2>.
- Fernández-González, M.; Mena, A.; Izquierdo, P.; Martínez, J. Genetic Characterization of Grapevine (*Vitis Vinifera* L.) Cultivars from Castilla La Mancha (Spain) Using Microsatellite Markers. *Vitis - Journal of Grapevine Research* **2007**, *46*, 126–130.
- Gisbert, C.; Soler, J.X.; Fos, M.; Intrigliolo, D.S.; Yuste, A.; Picó, B.; Torrent, D.; Peiró, R. Characterization of Local Mediterranean Grapevine Varieties for Their Resilience to Semi-Arid Conditions under a Rain-Fed Regime. *Agronomy* **2022**, *12*, 2234. <https://doi.org/10.3390/AGRONOMY12092234/S1>.
- Antolín, M.C.; Salinas, E.; Fernández, A.; Gogorcena, Y.; Pascual, I.; Irigoyen, J.J.; Goicoechea, N. Prospecting the Resilience of Several Spanish Ancient Varieties of Red Grape under Climate Change Scenarios. *Plants* **2022**, *11*, 2929. <https://doi.org/10.3390/PLANTS11212929/S1>.

7. A.S., S.; J., M.-G.; J.L., C.-V. Variability in Water Use Behavior during Drought of Different Grapevine Varieties: Assessment of Their Regulation of Water Status and Stomatal Control. *Agric Water Manag* **2024**, *291*, 108642. <https://doi.org/10.1016/j.agwat.2023.108642>.
8. Serrano, A.S.; Martínez-Gascuña, J.; Alonso, G.L.; Cebrián-Tarancón, C.; Carmona, M.D.; Mena Morales, A.; Chacón-Vozmediano, J.L. Variability in the Agronomic Behavior of 12 White Grapevine Varieties Grown under Severe Water Stress Conditions in the La Mancha Wine Region. *Horticulturae* **2023**, *9*, 243. <https://doi.org/10.3390/horticulturae9020243>.
9. Pérez-Álvarez, E.P.; Intrigliolo Molina, D.S.; Vivaldi, G.A.; García-Esparza, M.J.; Lizama, V.; Álvarez, I. Effects of the Irrigation Regimes on Grapevine Cv. Bobal in a Mediterranean Climate: I. Water Relations, Vine Performance and Grape Composition. *Agric Water Manag* **2021**, *248*, 106772. <https://doi.org/10.1016/j.agwat.2021.106772>.
10. Koundouras, S.; Kanakis, I.; Drossou, E.; Kallithraka, S.; Kotseridis, Y. Effects of Postveraison Water Regime on the Phenolic Composition of Grapes and Wines of Cv. Agiorgitiko (Vitis Vinifera L.). *OENO One* **2013**, *47*, 115–128. <https://doi.org/10.20870/OENO-ONE.2013.47.2.1542>.
11. Chacón-Vozmediano, J.L.; Martínez-Gascuña, J.; García-Romero, E.; Gómez-Alonso, S.; García-Navarro, F.J.; Jiménez-Ballesta, R. Effects of Water Stress on the Phenolic Compounds of ‘Merlot’ Grapes in a Semi-Arid Mediterranean Climate. *Horticulturae* **2021**, Vol. 7, Page 161 **2021**, *7*, 161. <https://doi.org/10.3390/HORTICULTURAE7070161>.
12. Pérez-álvarez, E.P.; Intrigliolo, D.S.; Almajano, M.P.; Rubio-Bretón, P.; Garde-Cerdán, T. Effects of Water Deficit Irrigation on Phenolic Composition and Antioxidant Activity of Monastrell Grapes under Semiarid Conditions. *Antioxidants* **2021**, Vol. 10, Page 1301 **2021**, *10*, 1301. <https://doi.org/10.3390/ANTIOX10081301>.
13. Parra, A.S.S.; Cebrian-Tarancon, C.; Gascuena, J.M.; Vozmediano, J.L.C.; Zapata, M.D.C.; Alonso, G.L. Effect of Two Water Deficit Regimes on Phenolic Composition of 15 Recovered Grapevine Varieties in Castilla-La Mancha Region (Spain): A Comparison with National and International Varieties. *BIO Web Conf* **2023**, *56*, 01005. <https://doi.org/10.1051/bioconf/20235601005>.
14. Buesa, I.; Intrigliolo, D.S.; Castel, J.R.; Vilanova, M. Influence of Water Regime on Grape Aromatic Composition of Muscat of Alexandria in a Semiarid Climate. *Sci Hortic* **2021**, *290*, 110525. <https://doi.org/10.1016/j.scienta.2021.110525>.
15. Palai, G.; Caruso, G.; Gucci, R.; D’Onofrio, C. Deficit Irrigation Differently Affects Aroma Composition in Berries of Vitis Vinifera L. (Cvs Sangiovese and Merlot) Grafted on Two Rootstocks. *Aust J Grape Wine Res* **2022**. <https://doi.org/10.1111/ajgw.12562>.
16. Kovalenko, Y.; Tindjau, R.; Madilao, L.L.; Castellarin, S.D. Regulated Deficit Irrigation Strategies Affect the Terpene Accumulation in Gewürztraminer (Vitis Vinifera L.) Grapes Grown in the Okanagan Valley. *Food Chem* **2021**, *341*, 128172. <https://doi.org/10.1016/j.foodchem.2020.128172>.
17. Fandiño, M.; Vilanova, M.; Rodríguez-Febreiro, M.; Teijeiro, M.T.; Rey, B.J.; Cancela, J.J. Effect of Deficit Irrigation on Yield Components and Chemical Composition of Albariño Grapes Grown in Galicia, NW Spain. *Agriculture (Switzerland)* **2022**, *12*, 1522. <https://doi.org/10.3390/agriculture12101522>.
18. Sánchez-Gómez, R.; Zalacain, A.; Alonso, G.L.; Salinas, M.R. Effect of Toasting on Non-Volatile and Volatile Vine-Shoots Low Molecular Weight Phenolic Compounds. *Food Chem* **2016**, *204*, 499–505. <https://doi.org/10.1016/j.foodchem.2016.02.137>.
19. Serrano de la Hoz, K.; Carmona, M.; Zalacain, A.; Alonso, G.L.; Salinas, M.R. The Varietal Aroma Potential Index (IPAv): A Tool to Evaluate the Quality of Grape and Wines, White and Red. *37th World Congress of Vine and Wine* **2014**.
20. Salinas, M.R.; De La Hoz, K.S.; Zalacain, A.; Lara, J.F.; Garde-Cerdán, T. Analysis of Red Grape Glycosidic Aroma Precursors by Glycosyl Glucose Quantification. *Talanta* **2012**, *89*, 396–400. <https://doi.org/10.1016/J.TALANTA.2011.12.050>.
21. Campayo, A.; Serrano de la Hoz, K.; García-Martínez, M.M.; Sánchez-Martínez, J.F.; Salinas, M.R.; Alonso, G.L. Spraying Ozonated Water on Bobal Grapevines: Effect on Grape Quality. *Food Research International* **2019**, *125*. <https://doi.org/10.1016/j.foodres.2019.108540>.
22. Deluc, L.G.; Quilici, D.R.; Decendit, A.; Grimplet, J.; Wheatley, M.D.; Schlauch, K.A.; Mérillon, J.M.; Cushman, J.C.; Cramer, G.R. Water Deficit Alters Differentially Metabolic Pathways Affecting Important Flavor and Quality Traits in Grape Berries of Cabernet Sauvignon and Chardonnay. *BMC Genomics* **2009**, *10*, 1–33. <https://doi.org/10.1186/1471-2164-10-212/TABLES/2>.
23. Rocha, S.M.; Rodrigues, F.; Coutinho, P.; Delgadillo, I.; Coimbra, M.A. Volatile Composition of Baga Red Wine: Assessment of the Identification of the Would-Be Impact Odourants. *Anal Chim Acta* **2004**, *513*, 257–262. <https://doi.org/10.1016/J.ACA.2003.10.009>.
24. Ribéreau-Gayon, P.; Glories, Y.; Maujean, A.; Dubourdiou D *Handbook of Enology, Chemistry of Wine Stabilization and Treat-Ments.*; Wiley, 2001; Vol. 2;.
25. Talaverano, I.; Valdés, E.; Moreno, D.; Gamero, E.; Mancha, L.; Vilanova, M. The Combined Effect of Water Status and Crop Level on Tempranillo Wine Volatiles. *J Sci Food Agric* **2017**, *97*, 1533–1542. <https://doi.org/10.1002/JSFA.7898>.

26. Wang, J.; Abbey, T.; Kozak, B.; Madilao, L.L.; Tindjau, R.; Del Nin, J.; Castellarin, S.D. Evolution over the Growing Season of Volatile Organic Compounds in Viognier (*Vitis Vinifera* L.) Grapes under Three Irrigation Regimes. *Food Research International* **2019**, *125*, 108512. <https://doi.org/10.1016/J.FOODRES.2019.108512>.
27. Ferreira, V.; López, R.; Cacho, J.F. Quantitative Determination of the Odorants of Young Red Wines from Different Grape Varieties. *J Sci Food Agric* **2000**, *80*, 1659–1667.
28. Guth, H. Quantitation and Sensory Studies of Character Impact Odorants of Different White Wine Varieties. *J Agric Food Chem* **1997**, *45*, 3027–3032. <https://doi.org/10.1021/JF970280A>.
29. Salinas, M.R.; Alonso, G.L.; Pardo, F.; Bayonove, C. Free and Bound Volatiles of Monastrell Wines. *Sci Aliments* **1998**, *18*, 223–231.
30. Parker, M.; Capone, D.L.; Francis, I.L.; Herderich, M.J. Aroma Precursors in Grapes and Wine: Flavor Release during Wine Production and Consumption. **2017**. <https://doi.org/10.1021/acs.jafc.6b05255>.
31. Talaverano, I.; Ubeda, C.; Cáceres-Mella, A.; Valdés, M.E.; Pastenes, C.; Peña-Neira, Á. Water Stress and Ripeness Effects on the Volatile Composition of Cabernet Sauvignon Wines. *J Sci Food Agric* **2018**, *98*, 1140–1152. <https://doi.org/10.1002/jsfa.8565>.
32. Asproudi, A.; Petrozziello, M.; Cavalletto, S.; Guidoni, S. Grape Aroma Precursors in Cv. Nebbiolo as Affected by Vine Microclimate. *Food Chem* **2016**, *211*, 947–956. <https://doi.org/10.1016/J.FOODCHEM.2016.05.070>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.