

Article

Not peer-reviewed version

Synthesis, Structural Characterization and Electrochemical Properties of Copper(II) Complexes with Functionalized Thiosemicarbazones Derived from 5-Acetylbarbituric Acid

[Alfonso Castiñeiras](#) , Nuria Fernández-Hermida , [Isabel García-Santos](#) * , Lourdes Gómez-Rodríguez , [Antonio Frontera](#) , [Juan Niclós-Gutiérrez](#)

Posted Date: 22 April 2024

doi: 10.20944/preprints202404.1405.v1

Keywords: thiosemicarbazone complexes, 5-acetylbarbituric derivatives, crystal structures, cooper(II) complexes, electrochemical analysis



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Synthesis, Structural Characterization and Electrochemical Properties of Copper(II) Complexes with Functionalized Thiosemicarbazones Derived from 5-Acetylbarbituric Acid

Alfonso Castiñeiras ¹, Nuria Fernández-Hermida ¹, Isabel García-Santos ^{1,*}, Lourdes Gómez-Rodríguez ¹, Antonio Frontera ² and Juan Niclós-Gutiérrez ³

¹ Universidad de Santiago de Compostela, Departamento de Química Inorgánica, Facultad de Farmacia, Santiago de Compostela, 15782, Spain; alfonso.castineiras@usc.es

² Universitat de les Illes Balears, Departament de Química, Crta de Valldemossa km 7.5, 07122 Palma de Mallorca (Baleares), Spain; toni.frontera@uib.es

³ Department of Inorganic Chemistry, Faculty of Pharmacy, University of Granada, 18071 Granada, Spain; jniclos@ugr.es

* Correspondence: isabel.garcia@usc.es

Abstract: The reaction between 5-acetylbarbituric acid and 4-dimethylthiosemicarbazide or 4-hexamethyleneiminy thiosemicarbazide produces 5-acetylbarbituric-4-dimethylthiosemicarbazone (H_2AcDM) and 5-acetylbarbituric-4N-hexamethyleneiminy thiosemicarbazone ($H_2AcHexim$). Eight new complexes with different copper (II) salts have been prepared and characterized using elemental analysis, molar conductance, UV-vis, ESI-HRMS, FT-IR, magnetic moment, EPR and cyclic voltammetry. In addition, three-dimensional molecular structures of $[Cu(HAcDM)(H_2O)_2](NO_3) \cdot H_2O$ (**3a**), $[Cu(HAcDM)(H_2O)_2]ClO_4$ (**4**), and $[Cu(HAcHexim)Cl]$ (**6**) were determined by single crystal X-ray crystallography, and was carried out an analysis of their supramolecular structure. The H-bonded assemblies were further studied energetically using DFT calculations, MEP surface and QTAIM analyses. In these complexes, the thiosemicarbazone coordinates to the metal ion in an ONS-tridentate manner, in the O-enolate/S-thione form. The electrochemical behavior of the thiosemicarbazones and their copper(II) complexes has been investigated at room temperature using the cyclic voltammetry technique in DMFA. The Cu(II)/Cu(I) redox system was found to be consistent with the quasi-reversible diffusion-controlled process.

Keywords: thiosemicarbazone complexes; 5-acetylbarbituric derivatives; crystal structures; cooper(II) complexes; electrochemical analysis

1. Introduction

It has been known for more than 30 years that climate change threatens the well-being of Earth, which is why since the Berlin Climate Change Conference in 1995, periodic evaluations have been carried out based on the Kyoto Protocol of 1997 and on the progress made towards the Paris Agreement of 2015 and propose a course of action. As we are already observing, energy and environmental issues are likely to dominate science and, consequently, society, in the years to come. In this regard, a huge research effort has been made to develop green energy devices using organic optoelectronics. Compared to conventional inorganic semiconductors, small organic and metal-organic molecules have a higher degree of versatility and can be synthesized and manufactured using different strategies to deliver low-cost mass production in a wide range of applications, including organic light-emitting diodes, energy storage, hybrid photovoltaic cells and artificial photosynthesis. These properties have triggered the rapid development of the synthesis of new transition metal complexes, generating enormous interest for different energy-related applications [1].

Coordination compounds play a critical role in modern inorganic and bioinorganic chemistry. They are considered the backbone of various chemical industries due to their wide application in different areas such as metal detection, bioimaging, drug delivery, chemosensors, and pharmacological and medicinal chemistry. Substantial advances in new concepts and different bonding models have provided a variety of molecules and their molecular structures, responsible for possible breakthroughs in the chemical industry [2]. The use of complexes in processes in the chemical industry, in the broad sense, enjoys increasing recognition from global corporations engaged in the production of various polymeric materials used in many aspects of our lives [3].

An important aspect of chemical technology is the least possible pollution of the environment, i.e. the application of the principles of green chemistry. Performing reactions in organic solvents is replaced by supercritical liquids or ionic liquids, and the development of the application of biotechnological and organocatalytic methods makes it possible to obtain chemical technologies that respect the environment [3].

Because of their utilitarian properties and interesting structures, coordination compounds have attracted increasing interest. The rapid development of materials science and crystal engineering has greatly encouraged the use of complexes as functional materials, such as photocatalysts, supercapacitors, nonlinear optical materials, and porous materials. Thus, pharmacists, biologists and researchers in medicinal chemistry have also embarked on the search for complexes for biomedical applications [4]. From this perspective and from the point of view of sustainability, the metal complexes used in these applications must be manufactured from sufficiently abundant materials and, in order to enter mass market applications, cost-effective metals with a d^{10} orbital configuration –such as Cu(I)– have to be, and currently are, the center of numerous research studies.

Copper, being one of the cheapest and most abundant elements on Earth, is a good choice for developing inexpensive and efficient coordination complexes for all energy-related applications, from energy conversion to energy storage and lighting. These compounds can successfully replace other unsustainable and toxic transition metals (Co, Pt, Ir, and Ru). There is a great need for further development of copper coordination complexes to increase the performance of each energy-relevant application [1]. In addition, copper complexes constitute an important class of molecules from several points of view: bioinorganic, catalysis, and magnetism. Copper performs various bioinorganic functions, is an essential bioelement, responsible for numerous catalytic processes in living systems, where it is usually present in different nuclearities, and is present in the enzymes of biological systems, either alone or in combination with some other metal ions, to perform its function through a redox reactivity [5].

The design of new ligand systems invariably depends on their reactivity to selected metals to provide metal binding modes. In the absence/existence of coordinated anions or small molecules, the stability in solution and characteristics of the immediately formed fragments are transcendental for their life in solution and their separation as disconnected metal complexes. What's more, these fragments can show self-aggregating reactivity, giving rise to ligand aggregates [6].


Redox-active ligands are easily oxidized or reduced compared to classic spectator ligands. Such ligands are a means of extending the redox reactivity of metal atoms in complexes because they are capable of providing additional redox equivalents. Normally, the number of electrons that a mononuclear metal complex can transfer to a substrate is limited to one or two, but in complexes with active redox ligands, the ligands can provide additional electrons [7].

In selecting specific types of ligands, Schiff bases have been studied extensively because of their fundamental role in the coordination chemistry of transition metals and major groups, inherent in their simple preparation method and structural variety. And, in this sense, innumerable applications have been described where they have been used as chelating ligands in coordination chemistry, as catalysts, as dyes, as initiators in polymerization, and as luminescent compounds. Biologically, they have also been tested as antibacterial, antifungal, antitumor, and antiviral agents including insecticides [8].

One group of Schiff bases that have attracted a great deal of research attention due to their medical application in the treatment of such feared diseases as leprosy and tuberculosis in the 1950s

are thiosemicarbazones (TSCs). Subsequent research on TSCs has shown beneficial biological applications in other areas such as antiviral, antibacterial, anticancer, anti-inflammatory, antimalarial, and, to some extent, HIV control [9,10]. In addition, thiosemicarbazones and their derivatives have attracted a great deal of interest in the fields of chemistry and biology and can also be used for metal analysis, device applications related to telecommunications, optical computing, storage, and information processing [11]. TSCs are synthesized by the condensation reaction of an aldehyde or ketone with a thiosemicarbazide, being found as thiol-thione tautomers, although in some cases, especially when coordinated with a metal center, the thiol form can be deprotonated to form thiolate anions [12]. In general, TSCs coordinate with metals in a bidentate manner through thiocarbonyl sulfur and azamethine nitrogen, mainly in a square-planar geometry. However, they can also be coordinated in a tridentate or tetradeятate manner, among other possibilities, when the aldehyde is functionalized, or it is a bis-thiosemicarbazone. Clearly, the biological activity of the resulting complexes will be affected by the type of metal center and functionalization [13].

In previous studies, we have reported on the synthesis and characterization of some novel thiosemicarbazones derived from 5-acetylbarbiturate acid [14] and its Ni(II), Pd(II) and Pt(II) complexes [15,16]. In the work described here we use the potentially tridentate ligands 5-acetylbarbiturate-4-dimethylthiosemicarbazone (H_2AcbDM) and 5-acetylbarbiturate-4N-hexamethyleneiminimyl thiosemicarbazone ($H_2AcbHexim$) (Scheme 1), in whose abbreviation “ H_2 ” denotes its two potentially dissociable acid protons, i.e., the barbiturate proton (enolic) and the hydrazine proton (thiol). Generally, these thiosemicarbazones are expected to bind to a metal center as dianionic tridentate O,N,S donors (a binding mode we have observed in platinum complexes), as monoanionic tridentate O,N,S donors, after the loss of the enolic barbiturate proton, as thiolate ligands, after dissociation of the hydrazine proton N-H, or as neutral ligands. In this study, we report a series of Cu(II) mononuclear complexes with H_2AcbDM and $H_2AcbHexim$. The eight newly prepared complexes were characterized experimentally, using UV-Vis FT-IR, EPR spectral analysis, cyclic voltammetry, magnetic susceptibility, conductivity measurements, and DFT theoretical studies.



Scheme 1. Molecular structures of 5-acetylbarbituric-4-dimethylthiosemicarbazone (H_2AcbDM) and 5-acetylbarbituric-4N-hexamethyleneiminimyl thiosemicarbazone ($H_2AcbHexim$), with the numbering used for the assignment of 1H and ^{13}C NMR signals.

2. Results and Discussion

2.1. Synthesis and Characterization

The complexes synthesized at ambient temperatures under the magnetic stirring of ethanolic solutions containing thiosemicarbazone ligands and copper salts in a 1:1 molar ratio, gave air-stable solids in good yield. The precipitates formed were then collected by filtration and washed thoroughly

with ethanol. The elementary analysis is concordant with the formulas $[\text{Cu}(\text{HAcBDM})(\text{OAc})]\cdot 3\text{H}_2\text{O}$ (1·3H₂O), $[\text{Cu}(\text{HAcBDM})\text{Cl}]\cdot 1/2\text{EtOH}$ (2·1/2EtOH), $[\text{Cu}(\text{HAcBDM})(\text{NO}_3)]$ (3), $[\text{Cu}(\text{HAcBDM})(\text{H}_2\text{O})_2]\text{ClO}_4$ (4), $[\text{Cu}(\text{HAcBHexim})(\text{OAc})]\cdot 5\text{H}_2\text{O}$ (5·5H₂O), $[\text{Cu}(\text{HAcBHexim})\text{Cl}]\cdot 1/2\text{EtOH}$ (6·1/2EtOH), $[\text{Cu}(\text{HAcBHexim})(\text{NO}_3)]\cdot 1/2\text{H}_2\text{O}$ (7·1/2H₂O) and $[\text{Cu}(\text{HAcBHexim})\text{ClO}_4]$ (8). The complexes showed good solubility in acetonitrile, dimethylformamide (DMF), and dimethyl sulfoxide (DMSO), and partial solubility in acetone and chloroform. However, they are insoluble in ether, methanol, and ethanol. The mass spectra of the complexes (FAB) (Figure S1) in some cases show signals for the molecular ion peak, although the most significant peaks correspond to the fragments $[\text{Cu}(\text{L})]^+$ and $[\text{L}]^+$ at m/z 334 and 272 for **1-4**, and at m/z 388 and 326 for compounds **5-8**, respectively. These fragmentation patterns involve the addition or loss of a hydrogen atom from the fragments.

The molar conductivities of freshly prepared solutions of the complexes in DMF are in the range 10–35 $\Lambda_m/\text{S cm}^2 \text{ mol}^{-1}$, which indicates that the compounds are neutral. However, $[\text{Cu}(\text{HAcBDM})(\text{H}_2\text{O})_2]\text{ClO}_4$ is a 1:1 electrolyte (83 $\Lambda_m/\text{S cm}^2 \text{ mol}^{-1}$). The molar conductivity value for $[\text{Cu}(\text{HAcBDM})(\text{NO}_3)]$ is of 54 $\Lambda_m/\text{S cm}^2 \text{ mol}^{-1}$, and this value may correspond to a mixture of $[\text{Cu}(\text{HAcBDM})(\text{NO}_3)]$ (neutral) and $[\text{Cu}(\text{HAcBDM})(\text{H}_2\text{O})_2](\text{NO}_3)\cdot \text{H}_2\text{O}$ (1:1 electrolyte) present in the solution.[17]

2.2. FT-IR Spectra

Spectra in the mid-infrared region (Figure S2) have bands in the range of 3500–3100 cm^{-1} associated with $\nu(\text{NH})$ and $\nu(\text{OH})$ of coordination water molecules and crystallization solvent molecules. The coordination of copper(II) ligands causes shifts in the bands at low frequencies, due to $\nu(\text{CN}) + \nu(\text{CC})$, in the range of 1625–1520 cm^{-1} in the free thiosemicarbazones,. The band due to $\nu(\text{CO})$, which in ligands appears as a strong band, around 1725 cm^{-1} in complexes, is shifted to 1700–1710 cm^{-1} . In addition, the band corresponding to the C=S stretching modes, which in ligands is observed between 830 and 840 cm^{-1} , appears at lower frequencies in complexes. The existence of the perchlorate group is confirmed by two intense bands [18] around 1120 and 625 cm^{-1} , while the appearance of two bands in the range of 1350–1280 cm^{-1} is due to the presence of nitrate groups [19]. In the spectra of acetate complexes, the characteristic bands, $\nu_a(\text{COO}^-)$ and $\nu_s(\text{COO}^-)$, are observed in the range between 1540 and 1515 cm^{-1} and between 1420 and 1400 cm^{-1} , respectively. The value of Δ [the separation between the two components of $\nu(\text{COO}^-)$, $\Delta = \nu_a(\text{COO}^-) - \nu_s(\text{COO}^-)$], is in the range of 90–130 cm^{-1} , these data being consistent with the mode of coordination of the acetate anion in the complexes [20,21]. This behavior is due to the monodeprotonation of thiosemicarbazones by $\text{N}_{\text{azo}}\text{-H}$, and ligand coordination via ONS. This coordination by the carbonyl oxygen atom, the azometin nitrogen atom, and the thiolate sulfur atom is also supported by the position of the bands $\nu(\text{Cu-O})$ at 550 cm^{-1} , $\nu(\text{Cu-N})$ at 425–450 cm^{-1} , $\nu(\text{Cu-S})$ around 315 cm^{-1} , and $\nu(\text{Cu-Cl})$ at 250 cm^{-1} [22] (Figure S3).

2.3. UV-Visible Spectra

The electronic spectra of copper(II) complexes have been studied qualitatively in solid phase. Figure 1 shows the spectra of compounds **2**, **3**, **6**, and **8**, respectively. The spectra of all complexes show different bands in the UV region. A high-energy band appearing between 28400 and 30400 cm^{-1} was assigned to intraligand charge transfer transitions (ILCT), $\pi \rightarrow \pi^*$, while another observed in some complexes between 21700 and 22100 cm^{-1} , was attributed to ligand-to-metal charge transfer transitions (LMCT), $\text{O} \rightarrow \text{Cu}^{II}$. In the visible region, the spectra show two distinct bands that have been assigned to ${}^2\text{E}_g \rightarrow {}^2\text{T}_{2g}$ transitions. In complexes **2**, **6**, and **8** both bands appear between 15450 and 16500 cm^{-1} and between 14350 and 14425 cm^{-1} and fall in the range of 16650–14250 cm^{-1} , usually typical of square plane complexes. In complexes **1**, **3**, **5**, and **7**, one band appears between 17390 and 17640 cm^{-1} and the other between 14150 and 15650 cm^{-1} , suggesting that these complexes have a distorted square pyramidal coordination geometry. [23]




Figure 1. Electronic absorption spectra of complexes **2**, **3**, **6**, and **8**.

2.4. Magnetic Susceptibilities

The magnetic moments of mononuclear copper(II) complexes are generally observed in the range 1.7 to 2.2 B.M, regardless of their stereochemistry. In most of the complexes studied here, the magnetic moments are in the range of 1.85-2.25 BM, which confirms the presence of mononuclear Cu(II) [24], because those of Cu(I) are expected to be diamagnetic. However, [Cu(HAcBHexim)ClO₄] presents a μ_{eff} value of 1.51 MB, which may be due to the existence of interactions between the Cu(II) centers of two neighboring molecules.

2.5. EPR analysis

The EPR spectra of compounds **1-8** in the polycrystalline state performed at room temperature show different types of geometric species and are similar to others previously reported for Cu^{II} complexes with thiosemicarbazones [25]. The spectra of complexes **3-6** are plotted in Figure 2. Compounds **1** and **5-7** present isotropic spectra containing a single broad signal with values for g in the range of 2.108-2.081. On the other hand, the spectra of compounds **2-4** and **8**, present the characteristics of axial anisotropy with well-defined values of g_{\parallel} and g_{\perp} around 2.189-2.372 and 2.043-2.095, respectively, where the value of $g_{\parallel} > g_{\perp} > 2.0023$, which supports that $d_{x^2-y^2}$ is the fundamental term in a square plane geometry. According to Hathway and Billing, [26] the geometrical parameter G is used to measure the exchange interaction between copper centers, which for axial spectra can be calculated using the equation: $G_{(\text{axial})} = g_{\parallel}2.0023 - g_{\perp}2.0023$, so if the value is between 3 and 5, it indicates that the fundamental state of these complexes is $d_{x^2-y^2}$ [27], and this agrees with the intermediate structure between tetrahedral and planar squared. Four coordination in the complexes is also verified by the relatively low g_{\parallel} values, indicating strong coordination and a significant degree of covalence in the bonds. In addition, a slight increase in g_{\parallel} has been observed when the volume of substituents on thioamide nitrogen increases, as occurs in similar complexes.




Figure 2. The EPR spectra of complexes **3–6**.

2.5. Molecular Structures and Supramolecular Analysis

Table 1 summarizes the relevant crystal data and the refinement of the structures of compounds **3a**, **4**, and **6**. Table 2 shows the coordination bond lengths and angles, and Table S1 shows the parameters of the hydrogen bonding interactions of the three compounds. The asymmetric unit of each of the compounds is represented in Figure 3, respectively. In the asymmetric unit of **3a** (Figure 3a), there are two independent complex molecules (I and II) and the coordination number is five due to the $[\text{HAcbDM}]^-$ ligands, which behave as mono-deprotonated tridentate ONS, and the enolate oxygen atoms, azomethine nitrogen and thiolate sulfur. The asymmetric unit of **4** (Figure 3b) is formed by a single complex molecule that is geometrically similar to either of the two **3a** molecules, but, in this case, with $[\text{HAcbHexim}]^-$ as the ligand.

Table 1. Crystal data and structure refinement for $[\text{Cu}(\text{HAcbDM})(\text{H}_2\text{O})_2](\text{NO}_3)\cdot\text{H}_2\text{O}$ (**3a**), $[\text{Cu}(\text{HAcbDM})(\text{H}_2\text{O})_2]\text{ClO}_4$ (**4**), $[\text{Cu}(\text{HAcbHexim})\text{Cl}]$ (**6**).

Compound	3a	4	6
Empirical formula	$\text{C}_9\text{H}_{18}\text{CuN}_6\text{O}_9\text{S}$	$\text{C}_9\text{H}_{16}\text{ClCuN}_5\text{O}_9\text{S}$	$\text{C}_{13}\text{H}_{18}\text{ClCuN}_5\text{O}_3\text{S}$
Formula weight	449.89	469.32	423.37
Temperature / K	293(2)	293(2)	293(2)
Wavelength / Å	1.54184	1.54184	1.54184
Crystal system	Triclinic	Monoclinic	Orthorhombic
Space group	$P\bar{1}$	$P2_1/n$	$Pbca$
Unit cell dimensions			
a / Å	9.039(4)	9.747	12.8807(7)
b / Å	13.289(2)	17.318	12.3524(5)
c / Å	14.880(2)	9.807	20.9062(8)
α / °	81.526(8)	90	90
β / °	78.678(17)	96.26	90
γ / °	76.91(2)	90	90
Volume / Å ³	1697.2(8)	1645.5	3326.3(3)
Z	4	4	8

Calc. density / Mg/m ³	1.761	1.894	1.691
Absorp. coefc. / mm ⁻¹	3.543	5.129	4.723
<i>F</i> (000)	924	956	1736
Crystal size	0.52 × 0.16 × 0.08	0.24 × 0.16 × 0.16	0.40 × 0.32 × 0.16
θ range / °	3.046 – 71.935	5.108 – 72.910	4.229 – 72.923
Limiting indices / <i>h,k,l</i>	0/11, -15/16, -17/18	-12/12, 0/21, 0/12	0/15, 0/15, 0/25
Refl. collect / unique	7116 / 6674	2893 / 2729	3557 / 3319
	[<i>R</i> _{int} = 0.1085]	[<i>R</i> _{int} = 0.0377]	[<i>R</i> _{int} = 0.0258]
Completeness 67.684 °	100.0	81.4	100.0
Absorp. correct.	Psi-scan	Psi-scan	Psi-scan
Max. /min. transm.	1.000 / 0.340	1.000 / 0.753	1.000 / 0.488
Data / parameters	6674 / 469	2729 / 235	3319 / 227
Goodness-of-fit on <i>F</i> ²	1.028	1.190	1.050
Final <i>R</i> indices	<i>R</i> ₁ = 0.0580, <i>wR</i> ₂ = 0.1577	<i>R</i> ₁ = 0.1069, <i>wR</i> ₂ = 0.2123	<i>R</i> ₁ = 0.0392, <i>wR</i> ₂ = 0.1059
<i>R</i> indices (all data)	<i>R</i> ₁ = 0.1168, <i>wR</i> ₂ = 0.1871	<i>R</i> ₁ = 0.1370, <i>wR</i> ₂ = 0.2237	<i>R</i> ₁ = 0.0524, <i>wR</i> ₂ = 0.1123
Largest dif. peak/hole e.Å ⁻³	0.896/-0.780	0.635/-0.733	0.556/-0.699
CCDC number			

In both compounds, the two remaining coordination positions are occupied by the oxygen atoms of two water molecules, originating a cationic complex, where the electroneutrality of the molecule is compensated by an anion nitrate (**3a**) or perchlorate (**4**). In addition, the **3a** asymmetric unit contains a crystallization water molecule for each complex molecule. In the two complexes, the coordination geometry around Cu(II) can be described as from square pyramidal to square pyramidal slightly distorted, according to the Addison parameter τ_5 of 0.07 (molecule I of **3a**), 0.24 (molecule II of **3a**) and 0.02 (**4**) [$\tau = (\beta - \alpha)/60$] where α and β are the two largest angles L-M-L'; τ is 1 for a perfect trigonal bipyramidal and is zero for a perfect square pyramid.[28] For **3a** and **4**, the three atoms of the ligand and the oxygen atom of a coordination water molecule occupy the four corners of the basal plane of the pyramid, at an average distance of 1.936 for Cu-O or Cu-N and 2.247 Å for Cu-S (Table 4), while the oxygen atom of the other coordination water molecule occupies the axial positions at greater bond distances from 2.441(4) and 2.477(4) Å (**3a**) and 2.609(10) Å (**4**) (Table 2).

Single crystals of [Cu(HAc_bDM)(H₂O)₂](NO₃)·H₂O (**3a**), [Cu(HAc_bDM)(H₂O)₂]ClO₄ (**4**) and [Cu(HAc_bHexim)Cl] (**6**) suitable for a X-ray diffractometry, were obtained from the mother liquor by slow evaporation at room temperature.

Table 2. Selected bond lengths [Å] for [Cu(HAc_bDM)(H₂O)₂](NO₃)·H₂O (**3a**), [Cu(HAc_bDM)(H₂O)₂]ClO₄ (**4**), [Cu(HAc_bHexim)Cl] (**6**).

	3a	4	6
Cu(1)-O(11)	1.912(3)	1.897(8)	1.9114(19)
Cu(1)-N(12)	1.961(4)	1.929(9)	1.959(2)
Cu(1)-O(12) / O(1) / Cl(1)	1.964(3)	1.963(8)	2.2338(7)
Cu(1)-S(1)	2.2487(14)	2.247(3)	2.2311(7)
Cu(1)-O(14) / O(2)	2.441(4)	2.609(10)	
Cu(2)-O(21)	1.913(3)		
Cu(2)-N(22)	1.947(4)		
Cu(2)-O(22)	1.945(4)		
Cu(2)-S(2)	2.2470(14)		
Cu(2)-O(24)	2.477(4)		
O(11)-Cu(1)-N(12)	91.03(14)	91.6(3)	87.90(8)

O(11)-Cu(1)-O(12) /O(1) /Cl(1)	85.98(14)	86.1(3)	91.19(6)
N(12)-Cu(1)-O(12) /O(1) /Cl(1)	173.12(17)	170.1(4)	178.41(7)
O(11)-Cu(1)-S(1)	168.82(12)	171.4(3)	172.67(7)
N(12)-Cu(1)-S(1)	88.39(11)	88.6(3)	87.50(6)
O(12) /O(1) /Cl(1)-Cu(1)-S(1)	93.34(10)	92.3(3)	93.28(3)
O(11)-Cu(1)-O(14) /O(2)	92.73(14)	95.0(3)	
N(12)-Cu(1)-O(14) /O(2)	96.05(15)	104.6(3)	
O(12) /O(1) /Cl(1)-Cu(1)-O(14) /O(2)	90.28(14)	85.2(4)	
S(1)-Cu(1)-O(14) /O(2)	98.43(9)	93.3(2)	
O(21)-Cu(2)-O(22)	85.97(16)		
O(21)-Cu(2)-N(22)	91.38(15)		
O(22)-Cu(2)-N(22)	177.17(19)		
O(21)-Cu(2)-S(2)	162.62(13)		
O(22)-Cu(2)-S(2)	94.18(12)		
N(22)-Cu(2)-S(2)	88.04(12)		
O(21)-Cu(2)-O(24)	95.59(14)		
O(22)-Cu(2)-O(24)	93.39(19)		
N(22)-Cu(2)-O(24)	87.87(15)		
S(2)-Cu(2)-O(24)	101.74(10)		




Figure 3. Perspective view of: (a) $[\text{Cu}(\text{HAcbDM})(\text{H}_2\text{O})_2](\text{NO}_3)\cdot\text{H}_2\text{O}$ (3a), (b) $[\text{Cu}(\text{HAcbDM})(\text{H}_2\text{O})_2]\text{ClO}_4$ (4) and (c) $[\text{Cu}(\text{HAcbHexim})\text{Cl}]$ (6), showing the asymmetrical unit and the atom-numbering scheme. Hydrogen bonds are shown as orange dashed lines.

In the crystal structure of 6 the asymmetric unit contains one molecule of complex (Figure 3c). As in the structures described above, the mono-deprotonated thiosemicarbazone ligand is coordinated to the metal in the expected terdentate manner via the thiolate sulfur S(1), the azomethine nitrogen N(12) and the enolate oxygen O(11), creating five- and six-membered chelate rings with N-Cu-S and O-Cu-N bite angles of 87.50(6) and 87.90(8) $^{\circ}$ respectively. The dihedral angle between the mean planes of the two chelate rings are 14.228(2) $^{\circ}$, which contrasts with the values of 8.46(1), 7.77(1), and 6.8(2) $^{\circ}$ found in the 3a and 4 molecules. The Cl⁻ ion is coordinated to the metal in the trans position with respect to the donor nitrogen atom. The Cu(II) is thus nested in an ONSCl core with circum-metallic bond parameters that deviate slightly from ideal square planar geometry.

This description is in agreement with the value of 0.06 calculated for τ_4 , the index for four-coordinate complexes, which is the sum of angles α and β – the two largest theta angles in the four-coordinate species – subtracted from 360 $^{\circ}$ all divided by 141 $^{\circ}$, where τ_4 is 1.00 in a tetrahedral, and 0.00 in a square planar geometries.[29] This core is nevertheless essentially planar, with a mean plane where the maximum deviation is 0.043(1) \AA in N(12), and from which the metal atom deviates 0.0641(1) \AA and forms a dihedral angle of 13.83(9) $^{\circ}$ with the thiosemicarbazone moiety. The Cu–O, Cu–N and Cu–S distances are all within the range that has been observed previously in 3a and 4 complexes (Table 2). The fact that the Cu–S(1) distance [2.2311(7) \AA] is shorter than the distances Cu–S in 3a and 4 [2.2487(14) and 2.247(3) \AA , respectively], and somewhat shorter than the Cu–Cl(1)

distance [2.2338(7) Å], shows that the trans-influence of the imine nitrogen atom is greater than that of the keto oxygen atom. In the thiosemicarbazone, the C–S(1) and C–O(11) bond lengths [1.714(2) and 1.265(3) Å, respectively] attest to the thiolate character of S(1) and the keto character of O(11), but the differences in bond lengths and bond angles with respect to the other complexes [greater for C–S and minor for C–O(11)], explains lower planarity in the thiosemicarbazone. The dihedral angle between the thiocarbazide moiety and the barbiturate ring is just 22.3(1)°, versus 4.9(3), 13.1(3), and 7.6(3)° in **3a** and **4**.

Unlike that detected in the 5-acetyl barbituric based free thiosemicarbazones, where many of their conformational parameters depend on the substituents present on nitrogen thioamide, in the complexes, these parameters are very similar [14]. When free, 5-acetylbarbituric hydrazine-1-carbothioamide (H_2AcbNH_2) is planar and, in N-mono and disubstituted, the plane of the 2,4,6-pyrimidinetrione ring is rotated between 46 and 57° with respect to that of thiosemicarbazone moiety. However, in the complexes studied here, the plane of the 2,4,6-pyrimidinetrione ring forms angles of 4.8(3)/13(3)° at **3a**, 8.2(6)° at **4**, and 23.4(1)° at **6**. Obviously, this deviation from the values found in the free ligands is imposed by the square-planar coordination geometry of the complexes.

On the other hand, it is known that the hydrazine nitrogen atom in the thiosemicarbazone N12-H13-C18 fragment loses the sp^2 hybridization planarity due to a hysterical impediment with the hydrogen atoms of the substituents on the thioamide nitrogen [14]. In complexes this loss is maintained so that the N13 atom is outside the plane formed by N12-H13-C18 between 0.13(3) and 0.33(2) Å for molecules I and II of **3a**, respectively, 0.28(8) Å in **4** and 0.17(1) Å in **6**.

Finally, the bond lengths and bond angles of the 2,4,6-pyrimidinetrione ring in all three complexes are in the ranges allowed for barbituric acid derivatives [30].

In relation to supramolecular packing, and more specifically to hydrogen bond formation, in the complexes all thiosemicarbazones have two H-acceptor oxygen atoms in common, located in the 2,4,6-pyrimidinetrione ring, and three N-H donors, two in the aforementioned ring and the third that corresponds to the hydrogen atom on the hydrazine nitrogen of the thiosemicarbazone moiety.




Figure 4. Crystal packing of the molecules in **3a**, including supramolecular synthons. Hydrogen bonds are shown as orange dashed lines.

In the crystal packing of **3a**, each cation forms a hydrogen bond with one of the oxygen atoms of a nitrate with N13-H13 as a donor, and another with a crystallization water molecule through the O14-H14A bond of the coordination water molecule that occupies the axial position. In turn, the crystallization molecule and nitrate form a new hydrogen bond O1-H1B···O43, forming a heterosynthon with an $R_3^3(11)$ graph set, where the nitrate ion functions as a bidentate bridge (Figure 3a).

In addition, there are other hydrogen bonds that govern packing. Each HAcB4DM ligand is linked to two other nearest neighbors by means of N-H···O bonds, forming two supramolecular heterosynthons with an $R_2^2(8)$ graph set in which the two HN-C(O) groups of the 2,4,6-pyrimidinetrione ring participate, generating planar zig-zag chains in the direction of the bisector of the angle between the *b* and *c* axes. Also, in the same direction, nitrate ions participate in new hydrogen bonds as triple bridges forming new heterosynthons with graph sets $R_3^2(8)$, $R_4^3(14)$, $R_6^6(20)$ and $R_4^4(21)$ (Figure 4). This creates a robust 3D network in which all the possible hydrogen bond donors and acceptors contained in the molecules participate (Figure S6).

The crystal packing of **4**, based on the formation of hydrogen bonds (Table S1), includes dimer units of cations in inverted positions, one relative to the other, linked by hydrogen bonds between the coordination water molecules (Figure 5a) forming a heterosynthon with an $R_2^2(8)$ graph set where the basal water molecule acts as the donor and the axial as the acceptor. These dimers in turn bind to new inverted dimers via thiosemicarbazone ligands with the formation of the $R_2^2(8)$ graph set heterosynthon between NH-C(O) groups described in compound **3a** packaging (Figure 5b).




Figure 5. (a) A partial packing diagram for **4**, showing dimer units of cations in inverted positions, and (b) ladder-like cation chains parallel to the *c*-axis showing the graph set H-bonded motifs. See Table S1 for symmetry codes.

This association is reinforced by a hydrogen bond between an O2-H of the water molecule in the axial position and the oxygen atom O13 in position 4 of the 2,4,6-pyrimidinetrione ring, forming a new supramolecular heterosynthon $R_3^2(10)$ and giving rise to ladder-like cation chains parallel to the *c*-axis (Figure 5b). These chains are held together in planes parallel to the *ab* plane through perchlorate bridges connected by means of new hydrogen bonds, where the second O-H of each water molecule in the axial position of one chain and the water molecule in the basal position of the neighboring chain participate as donors, and the O16 and O18 atoms of the perchlorate anions as acceptors (Figure 6a). Corrugated sheets similar to the arrangement on a tile roof are then formed, propagating parallel to the *b*-axis (Figure 6b).

The crystal packing of this compound is also aided by chelated ring-chelate···ring-chelate stacking interactions [31] between the five- and six-membered chelate rings that form the thiosemicarbazone ligand in coordination (Figure S7), with inter-centroid distances of 3.715-3.853 Å (Table S2). Also related to these interactions, self-assembled dimers present metal···chelate ring type interactions with a Cu-centroid distance of 3.911 Å (Table S2).




Figure 6. (a) A partial packing diagram for **4**, showing the perchlorate bridges connected by means of hydrogen bonds, and (b) corrugated sheets similar to the arrangement on a tile roof.

In compound **6**, only three intermolecular hydrogen bonds of the N-H \cdots O type define crystal packing (Table S1). An N-H corresponding to the hydrazine nitrogen atom of the thiosemicarbazone moiety (N13) acts as a donor against the carbonyl oxygen atom (C13) at position 4 of the 2,4,6-pyrimidinetrione ring of a neighboring molecule (Figure 7a), forming a zig-zag chain running along the *a*-axis (Figure 7b). In addition, the two hydrogen atoms on the pyrimidine nitrogen atoms are involved in two other hydrogen bonds with the other carbonyl oxygen atom (O15) and with the chloride ligand (Cl1) of two new nearest neighboring molecules (Figure 7c) giving rise to supramolecular heterosynthons with an $R_2^2(10)$ graph set in planar chains running parallel to the *b*-axis (Figure 7d). In this way, the conjunction of both chains gives rise to a dense 3D network (Figure S8).




Figure 7. Detail of crystalline packing in **6** showing: (a) the N-H \cdots O=C hydrogen bond, (b) the zig-zag chain along the "a" axis, (c) pairs of molecules forming supramolecular heterosynthons $R_2^2(10)$, and (d) zig-zag chains of molecules along the "b" axis.

2.6. Cyclic Voltammetry Studies

Systematically, the cyclic voltammograms of complexes **1-4** and **6-8** were carried out in a wide range of potential to show all the peaks. These measurements were performed by delimiting the areas where individual peaks appeared to investigate possible connections between the associated electrochemical reactions (Figure S9).

In the cyclic voltammograms (Figure 8a), first, two systems of peaks, I and II, were observed with cathodic form and absence the anodic form in the majority of the cases. At more negative potentials, one or two cathodic peaks (III and IV) without the corresponding anodic peaks were registered. Finally, potentials more positive than those of system I, is observed anodic peak (V) without the corresponding cathodic form. In complex 3, additional minor peaks were observed as shown in Figure 8b.

All complexes with peak I_a present the typical form of an electrode process of a reagent adsorbed on the electrode. In general, using a more negative vertex potential of 1.1 V it brings about the disappearance of I_a, even with long times of delay, (Figure 8a) indicating that it is potentially at this point that adsorption occurs significantly.

System II is very close to the previous one, so that the parameters of II_c can only be estimated approximately. This proximity is an indication of both peak systems are generated by the same type of electrochemical reaction taking place with molecules of slightly different structures. The fact that II_a does not appear, is indicative that the corresponding oxidation reaction is extremely slow or even non-existent.




Figure 8. Cyclic voltammogram of a 1 mM solution in DMF of (a) compound 2. $E_i=E_f=0.9$; delay = 0; $v = 0.2$ V/s; Everte = -2V; (b) compound 3. $E_i=E_f=0.9$; delay = 0; $v = 0.02$ V/s; Everte = -2.2V; (c) compound 2. $E_i=E_f=0.4$; delay = 10; $v = 0.2$ V/s; Everte = -1.3V; and (d) compound 2. $E_i=E_f=0.4$; delay = 10; $v = 0.2$ V/s; Everte = -0.8V.

The position of both systems indicates the reduction and reoxidation of Cu^{II} according to the scheme [32]:




The characteristics of I_a prove that the difference between the structures of the complexes giving rise to I or II, affects the adsorption of their reduced forms on the electrode surface. The almost planar geometry of the ligand would justify the existence of a certain adsorption capacity of ligand and the fact that it only affects the reduced form of one of the variants of the complex (resulting in peak I_a) indicates the existence of an additional factor that would determine the above-mentioned adsorption [33].


In complexes 1 and 7, the fact that the cyclic voltammograms are carried out from very positive initial potentials (2 V vs ECS) favours the appearance of I_c to the detriment of II_c. This behaviour

indicates that L and L' can represent oxidised and reduced forms of some component of the ligand, V_a being probably the peak resultanting from the electrochemistry oxidation of L'Cu²⁺ to LCu²⁺. On the other hand, the appearance of this peak when a voltammogram is performed in a potential region restricted to the range in which appears (Figure 8c and 8d), proves that the corresponding electrolytic process does not depend on the related to the remaining peaks.

All the processes described can be summarized by the following scheme:



There are different possibilities regarding the nature of L and L', but the most probable is to associate the first form with the existence of conjugated double bonds systems as show in Scheme 2, because this feature would favour the adsorption of these species on the electrode surface, which would justify the appearance of peak I_a. Scheme 2 represents the most probable scheme of the conversion reaction of L' in to L.



Scheme 2. Representation of the system of conjugated double bonds in the thiosemicarbazone anion.

Similar to systems I and II, the peaks are almost certainly related to the same type of electrochemical reactions in which the initial reactive species are slightly different. Such a difference would be the same as that gives rise to the existence of systems I and II, as evidenced by the fact that in which this splitting occurs, it also takes place from peak III in III and IV. It is expected that the electrochemical reaction associated with these peaks leads to the formation of species such as SCN⁻, S₈, SO₄²⁻ or some type of cyclisation, detected in many cases by X-ray diffraction [34,35].

The more important parameters to estimate the reduction potentials of Cu²⁺ are the potentials of peak (E_p) or semipeak (E_{p/2}) of I_c or II_c (Table 3). For reversible process the relationship between potential type follows the expression:

$$|E_p - E_{p/2}| = 2,2 \cdot \frac{R \cdot T}{n \cdot F}$$

If n = 1 and temperature is 20°, |E_p - E_{p/2}| has a value of 0.056 V.

In these conditions the formal potential value of $\text{Cu}^{2+}/\text{Cu}^{1+}$ ($E^{\circ'}$) system can be calculated through the expression:

$$E^{\circ'} = E_p + \frac{R \cdot T}{n \cdot F} \cdot \ln \frac{D_{ox}}{D_{Re}} + 1.109 \cdot \frac{R \cdot T}{n \cdot F}$$

Table 3. Electrochemical data for the indicated complexes.

System		$(E_p)_a^{[a]}$	$(E_p)_a^{[a]}$	$(E_p)_a - (E_p)_c^{[b]}$	$[(E_p)_a + (E_p)_c]/2^{[b]}$	$(E_{p/2})_c^{[a]}$	$(E_{p/2})_c + (E_p)_c^{[b]}$	$E^{\circ'}^{[a]}$
1	II	-0.708	-0.371	0.337	-0.539	-0.518	0.190	-0.679
2	I	-0.234				-0.170	0.064	-0.205
2	II	-0.582				-0.494	0.088	-0.553
3	I	0.057						
3	II	-0.580				-0.482	0.098	-0.551
6	I	-0.250				-0.184	0.066	-0.221
6	II	-0.604				-0.516	0.088	-0.575
7	I		0.074					
7	II	-0.628				-0.502	0.126	-0.599
8	I	0.085						
8	II	-0.604				-0.506	0.098	-0.576

[a] V vs. SCE. [b] V.

2.7. DFT Study

This theoretical study undertakes an energetic analysis of hydrogen-bonded synthons as illustrated in Figures 4 (compound **3a**), 5 (compound **4**), and 7 (compound **6**). Initially, the Molecular Electrostatic Potential (MEP) surface of compound **6** was computed. We have only analyzed the Cu-complex due to its neutrality, which facilitates a more accurate comparison of the hydrogen-bond donor/acceptor capabilities of the CO and NH groups of the barbituric ring and thiosemicarbazide group. The other two compounds, being salts, exhibit MEP values that are positive on the entire surface, due to the overall positive charge of the $[\text{Cu}(\text{HAcDM})(\text{H}_2\text{O})_2]^+$ fragment.

The MEP surface, showcased in Figure 8, identifies the MEP minimum at the chlorido ligand (-45.8 kcal/mol), indicating a negative MEP surface surrounding the Cl and extending to the Cu-coordinated O-atom. The carboxylic O-atoms of the barbituric ring exhibit MEP values of -38.3 kcal/mol and -30.4 kcal/mol, (O13 and O13 respectively, see Figure 3 for numbering scheme) highlighting their strong hydrogen-bond acceptor potential. In contrast, the MEP value at the S-atom is considerably lower (-12.5 kcal/mol). The highest MEP is observed at the NH group of the thiosemicarbazide group (64.4 kcal/mol), establishing it as the stronger hydrogen-bond donor. Additionally, the NH groups of the barbituric ring demonstrate significantly positive MEP values (30.9 and 33.6 kcal/mol for N15-H and N11-H, respectively), supporting the expectation of strong hydrogen bond formation. This MEP analysis corroborates the formation of the diverse hydrogen-bonded synthons detailed previously.




Figure 8. MEP surfaces of compound **1** (a) at the PBE0-D3/def2-TZVP level of theory (density isovalue 0.001 a.u.). The energies are given in kcal/mol.

To evaluate the hydrogen bond (H-bond) energies in the assemblies depicted in Figures 9-11, we employed the Quantum Theory of Atoms in Molecules (QTAIM) approach, calculating the strength of each H-bond using the method proposed by Espinosa et al., which is based on electron density (ρ) at the bond critical point (CP) and uses the formula $E = 0.5 \times V$. This method is particularly apt for the systems under consideration (**3a** and **4** are salts), where electrostatic interactions are predominant, allowing us to estimate H-bond contributions distinctly from ion-pair electrostatic forces.

For compound **3a**, the tetrameric assembly in Figure 9 features a $R_3^2(11)$ and two distinct $R_2^2(8)$. Topological analysis highlighted each H-bond with a bond critical point (BCP, illustrated by fuchsia sphere) and a dashed bond indicating the bond path from the H-atom to the O-atom. The strengths of these interactions are denoted in blue, revealing the $R_3^2(11)$ as the strongest (-16.1 kcal/mol), consistent with its formation of three H-bonds. The $R_2^2(8)$ synthons also showed significant interaction energies (-15.4 and -12.1 kcal/mol), noteworthy for forming only two H-bonds. The energy variation between the synthons correlates with experimental H-bond lengths, longer for the $R_2^2(8)$ synthon linking two symmetrically equivalent $[\text{Cu}(\text{HAcbDM})(\text{H}_2\text{O})_2]^+$ fragments (-12.1 kcal/mol).

Similarly, a trimeric assembly of compound **4** (see Figure 10) has been studied that includes the three synthons crucial for forming ladder-like cation chains along the c-axis (see also Figure 5). The $R_2^2(8)$ synthon associated with the barbituric rings has a formation energy of -8.2 kcal/mol, lower than those in compound **3a**, probably because the carbonyl group also participates in $\text{CO}\cdots\text{H}_2\text{O}$ bonds with the Cu-coordinated water molecule of an adjacent unit within the fused $R_3^2(10)$ synthon. This latter synthon is the strongest one (-18.3 kcal/mol) due to the formation of three H-bonds. The $R_2^2(8)$ synthon involving Cu-coordinated water molecules exhibits a significant formation energy, attributed to the enhanced acidity of the water protons when coordinated to the Cu atom.




Figure 9. QTAIM analysis of the tetrameric assembly of compound **3a** where the bond critical points (BCPs, fuchsia spheres) and bond paths (dashed bonds) of the H-bonding interactions are shown. The energy of the synthons is indicated using a dark blue colored font inside the supramolecular rings.




Figure 10. QTAIM analysis of the trimeric assembly of compound **4** where the bond critical points (BCPs, fuchsia spheres) and bond paths (dashed bonds) of the H-bonding interactions are shown. The energy of the synthons is indicated using a dark blue colored font inside the supramolecular rings.

In the case of compound **6**, we focused on a dimeric assembly extracted from the zig-zag molecular chains along the "b" axis, as depicted in Figure 7. These chains extend through the formation of $R_2^2(10)$ synthons. Our QTAIM analysis on this dimer shown in Figure 11 highlighted that the $R_2^2(10)$ synthon essentially comprises two interconnected $R_2^2(8)$ and $R_1^2(4)$ synthons, due to the NH bond forming a bifurcated $\text{NH}\cdots\text{O},\text{Cl}$ hydrogen bond. The computed formation energies are -5.0 kcal/mol for the $R_2^2(8)$ synthon and -2.3 kcal/mol for the $R_1^2(4)$ synthon, indicating these as the less energetically favorable interactions compared to the other compounds studied in this manuscript.




Figure 11. QTAIM analysis of the dimeric assembly of compound **6** where the bond critical points (BCPs, fuchsia spheres) and bond paths (dashed bonds) of the H-bonding interactions are shown. The energy of the synthons is indicated using a dark blue colored font inside the supramolecular rings.

3. Materials and Methods

3.1. Physical Measurements

All reagents and solvents were purchased from Sigma-Aldrich (Sigma-Aldrich. Inc., Tres Cantos, Madrid, Spain) and used as received, without further purification. Elemental analyses were performed with a Fisons-Carlo Erba 1108 microanalyser (CARLO ERBA Reagents SAS, Chaussée du Vexin, France). ^1H and ^{13}C NMR spectra were obtained as DMSO-d₆ solutions with a Varian Mercury 300 instrument (Varian Medical Systems, Inc, Palo Alto, California, U.S). IR spectra were recorded as KBr disks (4000-400 cm^{-1}) or polyethylene-sandwiched Nujol mulls (500-100 cm^{-1}) with a Bruker IFS-66v spectrophotometer (Bruker Corporation, Billerica, Massachusetts, USA). Mass spectra were obtained in a HP5988A spectrometer for EI and a Micromass AUTOSPEC spectrometer (nitrobenzyl alcohol matrix) for FAB (Agilent Technologies, Inc., Santa Clara, CA 95051, USA). Electronic spectra were carried out on a SHIMADZU UV-3101PC spectrophotometer (Izasa Scientific, Barcelona, Spain) equipped with a reflectance accessory. X-band EPR spectra of the complexes were obtained in 3 mm Pyrex tubes with a Bruker EMX spectrometer (Bruker Corporation, Billerica, Massachusetts, USA) using a conventional Dewar insert at liquid nitrogen temperature. Room temperature magnetic moments were determined on a PPMS (Physical Property Measurement System) magnetometer Quantum Design, model SQUID (Quantum Design, San Diego, CA 92121, USA). The conductivity measurements were carried out with a WTW conductivity meter model LF3 using freshly prepared 10⁻³ solutions of the complexes in DMF. Cyclic voltammograms were obtained on a 273 EG&G Princeton Applied Research electrochemical analyser (Artisan Technology Group, Champaign, IL 61822, USA). In the measurements DMF was used as solvent and tetra(*n*-butyl)ammonium perchlorate (TBAP) as the supporting electrolytes.

3.2. Synthesis of Thiosemicarbazone Ligands

The ligands, 5-acetylbarbituric-4N-dimethylthiosemicarbazone, H₂AcbDM and 5-acetylbarbituric-4N-hexamethyleneiminy thiosemicarbazone, H₂AcbHexim· were synthesized in excellent yields following our reported method [14,16], by condensation reactions between the 5-acetylbarbituric acid [36] and 4-dimethyl-thiosemicarbazine or hexamethyleneiminy thiosemicarbazine.

3.3. Synthesis and Crystallization of Complexes

An ethanolic suspension (20 mL) of thiosemicarbazone was added to a stirred solution of correspondent copper(II) salt in ethanol (15 ml) in molar ratio 1:1. The mixture was stirred at room

temperature for a further 7 days. The resulting suspension was filtered off and the solid was washed with ethanol and dried over calcium chloride.

[Cu(HAc_bDM)(OAc)]·3H₂O (1·3H₂O): Yield: 0.22 g (70.8%), m.p. 280 °C. Elemental analysis: Found: C, 29.5; H, 4.1; N, 15.8; S, 8.0. Calc. for C₁₁H₂₁O₈N₅SCu: C, 29.5; H, 4.7; N, 15.7; S, 7.2 %. IR (ν_{max}/cm⁻¹): 3178, 3053 v(NH); 1717, 1637 v(C=O)+δ(NH); 1591, 1529 v(CN+CC); 1295, 1268 v(CS+CC); 1040 v(NN); 810 v(CS); 425 v(Cu-N); 572, 527 v(Cu-O); 312 v(Cu-S). FAB⁺ MS [m/z, assignment]: 334(3), [ML]⁺; 272(2) [L]⁺. UV-vis (λ_{max}, cm⁻¹): 28409, 25445, 17544, 14184. EPR (X-band, solid sample): g_{av} = 2.10. μ_{eff} = 2.07 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 12.

[Cu(HAc_bDM)Cl]·1/2EtOH (2·1/2EtOH): Yield: 0.20 g (76.7%), m.p. 240 °C. Elemental analysis: Found: C, 30.8; H, 3.5; N, 18.2; S, 7.6. Calc. for C₁₀H₁₅ClO_{3.5}N₅SCu: C, 30.6; H, 3.8; N, 17.8; S, 8.2 %. IR (ν_{max}/cm⁻¹): 3156, 3084 v(NH); 1711, 1642 v(C=O)+δ(NH); 1583 v(CN+CC); 1306, 1241 v(CS+CC); 1060 v(NN); 808 v(CS); 449 v(Cu-N); 553 v(Cu-O); 314 v(Cu-S). FAB⁺ MS [m/z, assignment]: 334(9), [ML]⁺; 272(2) [L]⁺. UV-vis (λ_{max}, cm⁻¹): 29326, 16502, 14420. EPR (X-band, solid sample): g_{II} = 2.27, g_⊥ = 2.06, g_{av} = 2.12, G = 4.12. μ_{eff} = 1.85 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 24.

[Cu(HAc_bDM)(NO₃)] (3): Yield: 0.16 g (58.9%), m.p. 260 °C. Elemental analysis: Found: C, 27.8; H, 3.4; N, 20.6; S, 8.3. Calc. for C₉H₁₂O₆N₆SCu: C, 27.3; H, 3.1; N, 21.2; S, 8.1 %. IR (ν_{max}/cm⁻¹): 3402-3190 v(NH); 1709, 1643 v(C=O)+δ(NH); 1586, 1512, v(CN+CC); 1297, 1242 v(CS+CC); 1038 v(NN); 807 v(CS); 1383, 1297 v(NO₃); 425 v(Cu-N); 552 v(Cu-O); 312 v(Cu-S). FAB⁺ MS [m/z, assignment]: 334(9), [ML]⁺; 272(2) [L]⁺. UV-vis (λ_{max}, cm⁻¹): 29325, 21786, 17544, 15649, 14184. EPR (X-band, solid sample): g_{II} = 2.19, g_⊥ = 2.04, g_{av} = 2.06, G = 4.39. μ_{eff} = 2.14 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 54.

Suitable green crystals for X-ray diffraction were grown at room temperature as *[Cu(HAc_bDM)(H₂O)₂](NO₃)·H₂O (3a)*, after two weeks, from the resulting solution by slow evaporation at room temperature.

[Cu(HAc_bDM)(H₂O)]ClO₄ (4): Yield: 0.15 g (45.6%), m.p. >300 °C. Elemental analysis: Found: C, 23.4; H, 3.5; N, 14.9; S, 6.1. Calc. for C₉H₁₆ClO₉N₅SCu: C, 23.1; H, 3.4; N, 14.9; S, 6.8 %. IR (ν_{max}/cm⁻¹): 3398-3159 v(NH); 1701-1640 v(C=O)+δ(NH); 1579 v(CN+CC); 1308-1240 v(CS+CC); 811 v(CS); 1121-1088, 627 v(ClO₄⁻); 446 v(Cu-N); 553 v(Cu-O). FAB⁺ MS [m/z, assignment]: 334(42), [ML]⁺. EPR (X-band, solid sample): g_{II} = 2.20, g_⊥ = 2.05, g_{av} = 2.06, G = 4.34. μ_{eff} = 2.25 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 83.

Single crystals suitable for X-ray diffraction were obtained as (4), from the resulting solution by slow evaporation at room temperature.

[Cu(HAc_bHexim)(OAc)]·5H₂O (5·5H₂O): Yield: 0.26 g (82.6%), m.p. 280 °C. Elemental analysis: Found: C, 33.6; H, 5.5; N, 13.5; S, 5.1. Calc. for C₁₅H₃₁O₁₀N₅SCu: C, 33.6; H, 5.8; N, 13.0; S, 5.9 %. IR (ν_{max}/cm⁻¹): 3429-3042 v(NH); 1702, 1671 v(C=O)+δ(NH); 1587 v(CN+CC); 1297, 1245 v(CS+CC); 1031 v(NN); 812 v(CS); 433 v(Cu-N); 559, 528 v(Cu-O); 321 v(Cu-S). FAB⁺ MS [m/z, assignment]: 437(2) [ML]⁺; 326(2) [L]⁺. UV-vis (λ_{max}, cm⁻¹): 23337, 17637. EPR (X-band, solid sample): g_{av} = 2.08. μ_{eff} = 2.11 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 11.

[Cu(HAc_bHexim)Cl]·1/2EtOH (6·1/2EtOH): Yield: 0.12 g (47.0%), m.p. >300 °C. Elemental analysis: Found: C, 38.1; H, 4.3; N, 16.2; S, 6.9. Calc. for C₁₄H_{20.5}ClO_{3.5}N₅SCu: C, 37.7; H, 4.7; N, 15.7; S, 7.2 %. IR (ν_{max}/cm⁻¹): 3257 v(NH); 1710, 1653 v(C=O)+δ(NH); 1589, 1560 v(CN+CC); 1309-1241 v(CS+CC); 1032 v(NN); 808 v(CS); 433 v(Cu-N); 556 v(Cu-O); 317 v(Cu-S); 244 v(Cu-Cl). FAB⁺ MS [m/z, assignment]: 388(32), [ML]⁺; 326(2) [L]⁺. UV-vis (λ_{max}, cm⁻¹): 30395, 21954, 16260, 15408, 14368. EPR (X-band, solid sample): g_{av} = 2.08. μ_{eff} = 1.89 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 35.

Single brown crystals suitable for X-ray diffraction were obtained as (6), from the resulting solution by slow evaporation at room temperature.

[Cu(HAc_bHexim)(NO₃)]·1/2H₂O (7·1/2H₂O): Yield: 0.04 g (16.7%), m.p. >300 °C. Elemental analysis: Found: C, 34.2; H, 4.2; N, 18.0; S, 6.3. Calc. for C₁₃H₁₉O_{6.5}N₆SCu: C, 34.0; H, 4.2; N, 18.3; S, 6.9 %. IR (ν_{max}/cm⁻¹): 3345, 3185 v(NH); 1711-1619 v(C=O)+δ(NH); 1590, 1565 v(CN+CC); 1311-1247 v(CS+CC); 1092 v(NN); 799 v(CS); 1384, 1247 v(NO₃⁻); 433 v(Cu-N); 532 v(Cu-O); 322 v(Cu-S). FAB⁺ MS [m/z, assignment]: 388(4), [ML]⁺; 336(2) [L]⁺. UV-vis (λ_{max}, cm⁻¹): 29586, 22075, 17391, 15385. EPR (X-band, solid sample): g_{av} = 2.11. μ_{eff} = 2.01 MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 20.

$[Cu(HAcBHexim)ClO_4]$ (8): Yield: 0.06 g (21.1%), m.p. >300 °C. Elemental analysis: Found: C, 32.3; H, 4.1; N, 14.0; S, 6.2. Calc. for $C_{13}H_{18}ClO_7N_5SCu$: C, 32.0; H, 3.7; N, 14.4; S, 6.6 %. IR (ν_{max}/cm^{-1}): 3467-3037 ν (NH); 1711, 1652 ν (C=O)+ δ (NH); 1590, 1561 ν (CN+ CC); 1309-1245 ν (CS+ CC); 1030 ν (NN); 809 ν (CS); 1121-1094, 677 ν (ClO₄⁻); 441 ν (Cu -N); 555 ν (Cu-O); 316 ν (Cu -S). FAB⁺ MS [*m/z*, assignment]: 388(8), [ML]⁺. UV-vis (λ_{max} , cm⁻¹): 29940, 22727, 15468, 14409. EPR (X-band, solid sample): $g_{\parallel} = 2.37$, $g_{\perp} = 2.09$, $g_{av} = 2.17$. $\mu_{eff} = 1.51$ MB. Conductance (Ohm⁻¹·cm²·mol⁻¹) in DMF: 35.

4.4. Single-Crystal X-ray Diffraction

Diffraction data were obtained using an Enraf Nonius MACH3 automatic diffractometer from crystals mounted on glass fibers. Corrections for Lorentz and polarization effects, as well as absorption, were applied using a multi-scan method [37]. The structures were solved by direct methods [38], which revealed the positions of all non-hydrogen atoms. These were refined on F^2 by a full-matrix least-squares procedure using anisotropic displacement parameters [38]. Hydrogen atoms were located in the difference maps and the positions of O-H and N-H hydrogen atoms were refined (other were included as riders). Isotropic displacement parameters of H atoms were constrained to 1.2/1.5 U_{eq} of the carrier atoms. Molecular graphics were generated with DIAMOND software [39]. Crystal data, experimental procedures, and refinement outcomes are summarized in Table 1.

4.5. Theoretical Methods

DFT calculations were performed using Gaussian-16 [41] software at the PBE0-D3/def2-TZVP level of theory [42-44]. The analysis focused on solid-state interactions utilizing crystallographic coordinates, where only the position of the H-atoms was optimized. Bader's "Atoms in molecules" (QTAIM) theory [45] was employed to investigate hydrogen bonding interactions through the AIMall calculation package [46]. To estimate the H-bond energies we used the methodology proposed by Espinosa *et al.* [40] Molecular electrostatic potential (MEP) surfaces (with an isosurface value of 0.001 atomic units) were generated with Gaussian-16. This methodology and level of theory has been previously used to investigate similar interactions [47,48].

4. Conclusions

In this study, we synthesized eight new copper(II) complexes (**1-8**) with the ligands 5-acetylbarbituric-4-diethylthiosemicarbazone (H_2AcBDM) and 5-acetylbarbituric-4N-hexamethyleneiminy thiosemicarbazone ($H_2AcBHexim$). The complexes were characterized by spectroscopy and the crystal structures of three of them $[Cu(HAcBDM)(H_2O)_2](NO_3)_2 \cdot H_2O$ (**3a**), $[Cu(HAcBDM)(H_2O)_2]ClO_4$ (**4**), and $[Cu(HAcBHexim)Cl]$ (**6**) were determined by single-crystal X-ray diffraction. All complexes are monomers and not electrolytes, except $[Cu(HAcBDM)(H_2O)_2]ClO_4$, which is a 1:1 electrolyte, and the nitrate derivative with H_2AcBDM , which is possibly a mixture of $[Cu(HAcBDM)(NO_3)_2]$ (neutral) and $[Cu(HAcBDM)(H_2O)_2](NO_3)_2 \cdot H_2O$ (1:1 electrolyte). In the complexes, thiosemicarbazone ligands coordinate to copper(II) in a tridentate NOS manner and in a monoanionic tiona-keto mode.

The hydrogen bonds present in the crystal structures of complexes **3a**, **4**, and **6** (N-H···O, O-H···O, C-H···O, and N-H···Cl in their case), as well as stacking interactions π - π in the second, led to the stabilization of the 3D crystal packing characterized by the presence of several supramolecular synthons, among which the graph set $R_2^2(8)$ stands out. DFT calculations, MEP surface, and QTAIM analyses emphasize the relevance of hydrogen bonding interactions and the relative importance of the different H-bonded synthons. We shed light on the energetic characteristics of these interactions and disclosed that the $R_2^2(8)$ synthon where only coordinated water molecules participate is the strongest one.

The EPR spectra of complexes **1**, **5-7** show a single isotropic signal with an average g_{iso} value of 2.094, and complexes **2-4**, **8** have two different g -values (average 2.258 (g_{\parallel}) and 2.063 (g_{\perp}) indicating axial symmetry, and as $g_{\parallel} > g_{\perp}$ the presence of free electrons in the ground state $d_{x^2-y^2}$ of a planar

square geometry is confirmed. The fact that most complexes of the HzAcBDM ligand have an isotropic signal and those of the HzAcBhexim ligand are anisotropic may be related to the volume of the substituent on the thioamide nitrogen atom.

Systematically, cyclic voltammograms of complexes **1-4** and **6-8** were performed over a wide potential range in order to show all peaks. Measurements were then made in the potential range that bounded the zones where the individual peaks appeared to investigate possible connections between the associated electrochemical reactions. Accordingly, the redox activity of the complexes is attributed to the metal center. The cyclic voltammograms of all of them at different sweep velocities illustrate a linear relationship between the anodic peak currents and the square root of the sweeping velocities, confirming a quasi-reversible diffusion-controlled monoelectronic process of the Cu(II)/Cu(I) redox pair.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org. Tables S1 and S2: Intermolecular interactions; Figure S1: Mass spectra; Figures S2 and S3: Infrared spectra; Figure S4: Electronic (Reflectance) spectra; Figure S5: EPR spectra; Figures S6- S8. Unit cells; Figure S9: Cyclic voltammograms.

Author Contributions: Conceptualization, A. C. and I. G.-S.; methodology, all authors; computational calculations, A. F.; crystallography, A. C and I. G.-S.; software, A. C. and A. F.; investigation, I. G.-S., N. F.-H. and L. G.-R.; writing—original draft preparation, all authors, writing—review and editing, all authors; visualization, A. C., N. F.-H. J. N.-G. and I. G.-S. project administration, A. C., J. N.-G. and I. G.-S.; funding acquisition, A. C., I. G.-S and L. G.-R.. All authors have read and agreed to the published version of the manuscript.

Funding: The authors thanks to the financial support received from Consellería de Cultura, Educación, Formación Profesional e Universidades, Xunta de Galicia (Spain). GPC GI-2197 (ED481B-2022-068 2023/19).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Benesperi, I.; Singh, R.; Freitag, M. Copper Coordination Complexes for Enzrgy-Relevant Applications. *Energies* 2020, 13, 2198. <https://doi.org/10.3390/en13092198>
2. Singh, V. K.; Singh, V. K.; Mishra, A.; Varsha; Singh, A. A.; Prasad, G.; Singh, A. K. Recent advancements in coordination compounds and their potential clinical application in the management of diseases: An up-to-date review. *Polyhedron* 2023, 241, 116485. <https://doi.org/10.1016/j.poly.2023.116485>
3. Malinowski, J.; Zych, D.; Jacewicz, D.; Gawdzik, B.; Drzeżdżon, J. Application of Coordination Compounds with Transition Metal Ions in the Chemical Industry—A Review. *Int. J. Mol. Sci.* 2020, 21, 5443; <https://doi.org/10.3390/ijms21155443>
4. Chebout, O.; Bouchene, R.; Bouacida, S.; Boudraa, M.; Mazouz, W.; Merzougui, M.; Ouari, K.; Boudaren, C.; Merazig, H. Two copper(II) coordination complexes based on sulfonamide ligand: Synthesis, structure, thermal analyzes, electrochemical properties and biological activities. *J. Mol. Struct.* 2022, 1247, 131346. <https://doi.org/10.1016/j.molstruc.2021.131346>
5. Borthakur, R.; Kumar, A.; De, A. K.; Lal, R. A. Synthesis, characterization and electrochemical properties of copper(II) complexes derived from succinoyldihydrazine Schiff base ligands. *Arab. J. Chem.* 2019, 12, 2192–2205. <http://dx.doi.org/10.1016/j.arabjc.2014.12.040>
6. Peng, Y.-D.; Huang, F.; Du, M-X; La, Y-T; Yu, J; Hu, X; Dong, W-K. Rare formyl-coordinated homotrinuclear copper(II) salamo-based N₂O₃-donor complex: Experimental and theoretical studies. *Inorg. Chim. Acta* 2024, 560, 121815 <https://doi.org/10.1016/j.ica.2023.121815>
7. Gerhards, L.; Werr, M.; Hübner, O.; Solov'yov, I. A.; Himmel, H.-J. Peculiar Differences between Two Copper Complexes Containing Similar Redox-Active Ligands: Density Functional and Multiconfigurational Calculations. *Inorg. Chem.* 2024, 63, 961–975. <https://doi.org/10.1021/acs.inorgchem.3c02949>
8. Yusuf, T. L.; Oladipo, S. D.; Zamisa, S.; Kumalo, H. M.; Lawal, I. A.; Lawal, M. M.; Mabuba, N. Design of New Schiff-Base Copper(II) Complexes: Synthesis, Crystal Structures, DFT Study, and Binding Potency toward Cytochrome P450 3A4. *ACS Omega* 2021, 6, 13704–13718. <https://doi.org/10.1021/acsomega.1c00906>
9. Misigo, W. O.; Njenga, L. W.; Odhiambo, R. A.; Meyer, M.; Julius, L.; Sibuyi, N.; Lalancette, R. A.; Onani, M. O. New thiosemicarbazones and their palladium(II) complexes: Synthesis, spectroscopic characterization, X-ray structure and anticancer evaluation. *Inorg. Chim. Acta* 2023, 558, 121746. <https://doi.org/10.1016/j.ica.2023.121746>

10. Wang, X.; Zhu, M.; Li, S.; Xu, G.; Zhang, Z.; Yang, F. Novel mono-, bi-, tri- and tetra-nuclear copper complexes that inhibit tumor growth through apoptosis and anti-angiogenesis. *J. Inorg. Biochem.* 2024, 250, 112403. <https://doi.org/10.1016/j.jinorgbio.2023.112403>
11. Normaya, E.; Ahmad, M. N.; Farinab, Y.; Bulat, K. H. K. Synthesis, Characterization and Preliminary Study on Acetylpyrazine N(4)Butylthiosemicarbazone as a Potential CDK2 Inhibitor Combined with DFT Calculations. *J. Braz. Chem. Soc.* 2018, 29, 2197-2206. <http://dx.doi.org/10.21577/0103-5053.20180097>
12. Bermejo, E.; Castiñeiras, A.; García, I.; West, D. X. Spectral and structural studies of mercury(II) complexes of 2-pyridineformamide N(4)-dimethylthiosemicarbazone. *Polyhedron*, 2003, 22, 1147-1154. [https://doi.org/10.1016/S0277-5387\(03\)00100-1](https://doi.org/10.1016/S0277-5387(03)00100-1)
13. Sripathi, M. P.; Berely, S.; Reddy, C. V. R. Computational Studies of 4-Formylpyridinethiosemicarbazone and Structural and Biological Studies of Its Ni(II) and Cu(II) Complexes. *Heteroatom Chem.* 2019, 3507837. <https://doi.org/10.1155/2019/3507837>
14. Castiñeiras, A.; Fernández-Hermida, N.; García-Santos, I.; Gómez-Rodríguez, L.; Gil, D. M.; Frontera, A. Supramolecular, spectroscopic and computational analysis of weak interactions in some thiosemicarbazones derived from 5-acetylbarbituric acid. *J. Mol. Struct.* 2021, 1245, 131031. <https://doi.org/10.1016/j.molstruc.2021.131031>
15. Bermejo, E.; Castiñeiras, A.; García-Santos, I.; Gómez-Rodríguez, L.; P. Sevillano. Pt4S4 Clusters with Functionalized Thiosemicarbazones derived from 5-Acetylbarbituric Acid. *Z. Anorg. Allg. Chem.* 2007, 633, 2255-2261. <https://doi.org/10.1002/zaac.200700151>
16. Castiñeiras, A.; Fernández-Hermida, N.; García-Santos, I.; Gómez-Rodríguez, L. Neutral NiII, PdII and PtII ONS-pincer complexes of 5-acetylbarbituric-4Ndimethylthiosemicarbazone: synthesis, characterization and properties. *Dalton Trans.* 2012, 41, 13486-13495. <https://doi.org/10.1039/C2DT31753B>
17. Geary, W. J. The use of conductivity measurements in organic solvents for the characterisation of coordination compounds. *Coord. Chem. Rev.* 1971, 7, 81-122. [https://doi.org/10.1016/S0010-8545\(00\)80009-0](https://doi.org/10.1016/S0010-8545(00)80009-0)
18. Nakamoto, K. Infrared and Raman Spectra of Inorganic and Coordination Compounds, 6th edn, Wiley, Hoboken, NJ, 2009.
19. Addison, C. C.; Logan, N.; Wallwork, C. S.; Garner, C. D. Q. *Rev., Chem. Soc.* 1971, 25, 289-322. <https://doi.org/10.1039/QR9712500289>
20. Deacon, G. B.; Phillips, R. J. Relationships between the carbon-oxygen stretching frequencies of carboxylato complexes and the type of carboxylate coordination. *Coord. Chem. Rev.*, 1980, 33, 227-250. [https://doi.org/10.1016/S0010-8545\(00\)80455-5](https://doi.org/10.1016/S0010-8545(00)80455-5)
21. Bresolin, L.; Burrow, R. A.; Hörner, M.; Bermejo, E.; Castiñeiras, A. Synthesis and crystal structure of di[(μ -acetato) (2-acetylpyridine 4Nethylthiosemicarbazonoato)zinc(II)]. *Polyhedron*, 1997, 16, 3947-3951. [https://doi.org/10.1016/S0277-5387\(97\)00194-0](https://doi.org/10.1016/S0277-5387(97)00194-0)
22. Weidlein, J.; Müller, U.; Deknicke, K. *Schwingungsspektroskopie*, G. Thieme Verlag, Stuttgart, 2nd edn, 1988
23. West, D. X.; Beraldo, H.; Nassar, A. A.; *Transition Met. Chem.* 1999, 24, 421-424. <https://doi.org/10.1023/A:1006967802158>
24. Chandra, S.; Gupta, L.; Tyagi, V. P. *Co(II), Ni(II), Cu(II) and Zn(II) complexes of Thiosemicarbazones. Synth. React. Inorg. Met.-Org. Chem.* 26(1) (1996) 125-137. <https://doi.org/10.1080/00945719608004251>
25. Lobana, T. S.; Kaushal, M.; Virk, R. K.; Garcia-Santos, I.; Jasinski, J. P. Thiosemicarbazonates of copper: Crystal structures of [(furan-2-acetaldehyde-N-phenylthiosemicarbazonoato)][bis(triphenylphosphine)]copper(I) and [bis(furan-2-formaldehyde-N-phenylthiosemicarbazonoato)]copper(II). *Polyhedron* 2018, 152, 49-54. <https://doi.org/10.1016/j.poly.2018.06.022>
26. Hathway, B.J.; Billing, D.E. The Electronic Properties and Stereochemistry of Mono-Nuclear Complexes of the Copper(II) Ion. *Coord. Chem. Rev.* 1970, 5, 143-207. [https://doi.org/10.1016/S0010-8545\(00\)80135-6](https://doi.org/10.1016/S0010-8545(00)80135-6)
27. García-Tojal, J.; Lezama, L.; Pizarro, J. L.; Insausti, M.; Arriourtia, M. I.; Rojo, T. Spectroscopic and magnetic properties of copper(II) complexes derived from pyridine-2-carbaldehyde thiosemicarbazone. Structures of $[\text{Cu}(\text{NO}_3)(\text{C}_7\text{H}_8\text{N}_4\text{S})(\text{H}_2\text{O})](\text{NO}_3)$ and $[\{\text{Cu}(\text{NCS})(\text{C}_7\text{H}_8\text{N}_4\text{S})\}_2]$. *Polyhedron* 1999, 18, 3703-3711. [https://doi.org/10.1016/S0277-5387\(99\)00310-1](https://doi.org/10.1016/S0277-5387(99)00310-1)
28. Addison, A. W.; Rao, T. N.; Reedijk, J.; van Rijn, J.; Verschoor, C. G. Synthesis, structure, and spectroscopic properties of copper(II) compounds containing nitrogen-sulphur donor ligands; the crystal and molecular structure of aqua[1,7-bis(N-methylbenzimidazol-2'-yl)-2,6-dithiaheptane]copper(II) perchlorate. *J. Chem. Soc., Dalton Trans.* 1984, 1349-1356. <https://doi.org/10.1039/DT9840001349>
29. Yang, L.; Powell, D. R.; Houser, R. P. Structural variation in copper(I) complexes with pyridylmethylamide ligands: structural analysis with a new four-coordinate geometry index, τ_4 . *Dalton Trans.*, 2007, 955-964. <https://doi.org/10.1039/B617136B>
30. Lewis, T. C.; Tocher, . D. A.; Price, S. L. An Experimental and Theoretical Search for Polymorphs of Barbituric Acid: The Challenges of Even Limited Conformational Flexibility. *Cryst. Growth. Des.* 2004, 4, 979-987, <https://doi.org/10.1021/cg034209a>

31. D. P. Malenov, S. D. Zarić, Recognizing New Types of Stacking Interactions by Analyzing Data in the Cambridge Structural Database. *Chemistry* 2023, **5**, 2513–2541. <https://doi.org/10.3390/chemistry5040164>
32. A. S. Kumbhar, S. B. Padhye, D. X. West, A. E. liberta Electrochemical studies of copper(II) 2-acetylpyridine4 N-dialkylthiosemicarbazones. Relation to their spectral, magnetic, and biological properties. *Transition Met. Chem.* 16 (1991) 276-279. <https://doi.org/10.1007/BF01032852>
33. V. Eisner, E. K. Eisner, A. J. En Bard, H. Lund. *Encyclopedia of Electrochemistry of Elements*, Marcel Dekker, New York, Vol. 13 1979
34. Y. P. Kitaev, G.K. Budnikov, I. M. Skrebkova. Tautomerism and geometrical isomerism of nitrogen-containing derivatives of carbonyl compounds. VI.The polarographic investigation of semi- and thiosemicarbazones of some aliphatic and alicyclic ketones *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk* 2 (1962) 244-252. CA 56, 15287e
35. Y. P. Kitaev, G.K. Budnikov, A. E. Arbuzov. Tautomerism and geometric isomerism and nitrogenous derivatives of carbonyl compounds. III. Polarographic study of some semi- and thiosemicarbazones in aqueous alcohol solutions. *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk* 5 (1961) 824-31. CA 55, 440f
36. H. Biltz, H. Wittek, Über alkylierte und acylierte Barbitursäuren. *Chem. Ber.* 1991, **54**, 1035-1058. <https://doi.org/10.1002/cber.19210540519>
37. Sheldrick, G.M. SADABS, Program for Empirical Absorption Correction of Area Detector Data; University of Göttingen: Germany, 2001
38. Sheldrick, G.M. SHELXT – Integrated space-group and crystal structure Determination. *Acta Crystallogr.* 2015, **A71**, 3–8. <https://doi.org/10.1107/S2053273314026370>
39. Putz, H.; Brandenburg, K. Diamond - Crystal and Molecular Structure Visualization Version 4.6.2, Crystal Impact GbR: Bonn, Germany, 2020.
40. Espinosa, E.; Molins E.; Lecomte, C. Hydrogen bond strengths revealed by topological analyses of experimentally observed electron densities. *Chem. Phys. Lett.* **1998**, **285**, 170. [https://doi.org/10.1016/S0009-2614\(98\)00036-0](https://doi.org/10.1016/S0009-2614(98)00036-0)
41. Frisch, M.J.; Trucks, G.W.; Schlegel, H.B.; Scuseria, G.E.; Robb, M.A.; Cheeseman, J.R.; Scalmani, G.; Barone, V.; Petersson, G.A.; Nakatsuji, H.; et al. Gaussian 16; Revision A.01; Gaussian, Inc.: Wallingford, CT, USA, 2016.
42. Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. A Consistent and Accurate ab Initio Parametrization of Density Functional Dispersion Correction (DFT-D) for the 94 Elements H-Pu. *J. Chem. Phys.* **2010**, **132**, 154104–154122. <https://doi.org/10.1063/1.3382344>
43. Weigend, F. Accurate Coulomb-fitting basis sets for H to Rn. *Phys. Chem. Chem. Phys.* **2006**, **8**, 1057–1065. <https://doi.org/10.1039/b515623h>
44. Adamo, C.; Barone, V. Toward reliable density functional methods without adjustable parameters: The PBE0 model. *J. Chem. Phys.* **1999**, **110**, 6158–6169. <https://doi.org/10.1063/1.478522>
45. Bader, R.F.W. A quantum theory of molecular structure and its applications. *Chem. Rev.* **1991**, **91**, 893928. <https://doi.org/10.1021/cr00005a013>
46. Keith, T.A. TK Gristmill Software, version 13.05.06; AIMAll: Overland Park, KS, USA, 2013.
47. Mousavi, H.; García-Rubiño, M.E.; Choquesillo-Lazarte, D.; Castiñeiras, A.; Lezama, L.; Frontera, A.; Niclós-Gutiérrez, J. H(N3)dap (Hdap = 2,6-Diaminopurine) Recognition by Cu₂(EGTA): Structure, Physical Properties, and Density Functional Theory Calculations of [Cu₄(μ-EGTA)₂(μ-H(N3)dap)₂(H₂O)₂]·7H₂O. *Molecules* **2023**, **28**, 6263. <https://doi.org/10.3390/molecules28176263>
48. Belmont-Sánchez, J.C.; Choquesillo-Lazarte, D.; García-Rubiño, M.E.; Matilla-Hernández, A.; Niclós-Gutiérrez, J.; Castiñeiras, A.; Frontera, A. Supramolecular Nature of Multicomponent Crystals Formed from 2,2'-Thiodiacetic Acid with 2,6-Diaminopurine or N9-(2-Hydroxyethyl)adenine. *Int. J. Mol. Sci.* **2023**, **24**, 17381. <https://doi.org/10.3390/ijms242417381>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.