
Distinguishing Axillary Lymphadenopathy after COVID-19 Vaccination from Malignant Lymphadenopathy

[Shintaro Yamanaka](#) , [Keiko Tanaka](#) , [Masao Miyagawa](#) , Teruhito Kido , Shinji Hasebe , Shoichiro Yamamoto , Tomomi Fujii , Kazuto Takeuchi , [Yoshihiro Yakushijin](#) *

Posted Date: 23 April 2024

doi: 10.20944/preprints202404.1454.v1

Keywords: COVID-19; vaccine; axillary-lymphadenopathy; 18F-FDG PET



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Distinguishing Axillary Lymphadenopathy after COVID-19 Vaccination from Malignant Lymphadenopathy

Shintaro Yamanaka ¹, Keiko Tanaka ², Masao Miyagawa ³, Teruhito Kido ³, Shinji Hasebe ¹, Shoichiro Yamamoto ¹, Tomomi Fujii ¹, Kazuto Takeuchi ⁴ and Yoshihiro Yakushijin^{1,*}

¹ Department of Clinical Oncology, Ehime University Graduate School of Medicine, Toon, Japan; aidaisintaro515@gmail.com (Shi.Y.); hasebe.shinji.yw@ehime-u.ac.jp (S.H.); shoichiroy@gmail.com (Sho.Y.); tfujii@m.ehime-u.ac.jp (T.F.)

² Department of Epidemiology and Public Health, Ehime University Graduate School of Medicine, Toon, Japan; tanaka.keiko.jn@ehime-u.ac.jp

³ Department of Radiology, Ehime University Graduate School of Medicine, Toon, Japan; miyagawa848@gmail.com (M.M.); kido.teruhito.mu@ehime-u.ac.jp (T.K.)

⁴ Department of Clinical Laboratory, Ehime Prefectural University of Health Sciences, Tobe, Japan; kazutake0415@gmail.com

* Correspondence: yoshiyak@m.ehime-u.ac.jp; Tel.: +81-89-960-5969

Abstract: Objectives: To study the differences between malignant hypermetabolic axillary lymphadenopathy (MHL) and COVID-19 vaccine-associated axillary hypermetabolic lymphadenopathy (VAHL) using clinical imaging. **Methods:** A total of 1,096 patients underwent Positron Emission Tomography-Computed Tomography (PET-CT) between June 1, 2021 and April 30, 2022, at Ehime University Hospital. In total, 188 patients with axillary lymphadenopathy after the COVID-19 vaccination were evaluated. The patients were classified into three groups such as VAHL (n=27), MHL (n=21), and equivocal hypermetabolic axillary lymphadenopathy (EqHL; n=140). Differences in lymph nodes (LNs) swellings were statistically analyzed using clinical imagings (echography, CT, and ¹⁸F-FDG PET) statistically. **Results:** MHL included a higher female population (90.5%) owing to a higher frequency of breast cancer frequency (80.9%). Axillary LNs of MHL did not show any LN fatty hilums (0%); however, those of VAHL and EqHL did (15.8 and 36%, respectively). After the logistic regression analysis of the patients who had axillary lymphadenopathy without any LN fatty hilums, the minor axis length and ellipticity (minor axis/major axis) in the largest axillary LN, SUVmax, and Tissue-to-Background Ratio (TBR) were useful in distinguishing malignant lymphadenopathies. Receiver-operating characteristic (ROC) analysis indicated that a cut-off value of ≥ 7.3 mm for the axillary LN minor axis (sensitivity: 0.714, specificity: 0.684) and of ≥ 0.671 for ellipticity (0.667 and 0.773, respectively) in the largest LN with the highest SUVmax and TBR were predictive of MHL. **Conclusion:** Axillary lymphadenopathy ≥ 7.3 mm of the minor axis and ≥ 0.671 of ellipticity in LN without fatty hilum suggests malignancy even in patients who have received COVID-19 vaccination. Further examinations such as ¹⁸F-FDG PET are recommended for such patients

Keywords: COVID-19; vaccine; axillary-lymphadenopathy; ¹⁸F-FDG PET

1. Introduction

Since the commencement of the COVID-19 vaccination campaign in Japan in February 2021, over 38 million doses of COVID-19 vaccination have been administered, and over 68% of the Japanese population has already been vaccinated three or more times [1]. Among the various side effects, post-vaccination axillary lymphadenopathy, specifically on the side of the inoculation, have been reported as a clinical problem [2–5]. As with other mRNA vaccines, antigen migration induced by vaccination from the injection site to draining nodes can result in lymph node swelling. Recent clinical observations and reports in patients after COVID-19 vaccination have indicated that the frequency of

axillary lymphadenopathy ranges from 0.3% to 16% post-vaccination and the incidence generally increases with the number of subsequent vaccinations [6]. Detection of lymphadenopathy on various imaging modalities, including ultrasonography (US), CT, MRI, and ^{18}F -FDG PET, has been documented in several case reports and small cohort studies, and the incidence [7–14] and the immunological mechanisms behind its development [15–17] are currently being clarified. For patients with cancer or a history of cancer, distinguishing between reactive and malignant lymphadenopathy is a critical issue in clinical practice. However, the differentiation of the imaging characteristics between COVID-19-associated (reactive) lymphadenopathy and malignancy is not entirely understood. The aim of the present study was to discern the differences between reactive (vaccine-associated hypermetabolic lymphadenopathy [VAHL]) and malignant (malignant hypermetabolic lymphadenopathy [MHL]) axillary lymphadenopathy based on the analysis of clinical imaging data.

2. Materials and Methods

2.1. Criteria for Unilateral Lymphadenopathy and Data Collection

Data of patients aged ≥ 20 years who underwent Positron Emission Tomography-Computed Tomography (PET-CT) between June 1, 2021, and April 30, 2022, were collected from the electronic medical records of Ehime University Hospital. To assist in the clinical interpretation of PET-CT scans, the hospital's Department of Nuclear Medicine started routinely documenting details of COVID-19 vaccination (date of vaccination and injection site) from June 2021. These data were used in the present study. During the study period, a total of 1,096 patients underwent PET-CT, of whom 201 were diagnosed with unilateral lymphadenopathy. The diagnosis of significant axillary lymph node swelling or notable FDG accumulation in the axillary lymph node (ALN) was confirmed by two or more specialists in nuclear medicine. Patients were classified into three ALN groups based on these follow-up data: benign lymph nodes associated with COVID-19 vaccination (vaccine-associated hypermetabolic lymphadenopathy, VAHL group), lymph node metastasis from a malignant tumor (malignant hypermetabolic lymphadenopathy, MHL group), and equivocal hypermetabolic lymphadenopathy (EqHL group). The VAHL group included patients who developed lymphadenopathy within 10 weeks following vaccination and met at least one of the following criteria: lymph nodes assessed as benign according to echography and/or biopsy or documentation of decreased nodal size over time. This criterion was based on a previous study that demonstrated a significant reduction in unilateral axillary lymphadenopathy at least 10 weeks post-vaccination [18]. The MHL group consisted of patients with confirmed vaccination status from medical records and a diagnosis of malignant lymphadenopathy based on the findings of echographic examination or lymph node biopsy. The EqHL group included vaccinated patients whose axillary lymphadenopathy could not be conclusively classified as either MHL or VAHL (Figure 1).




Figure 1. Grouping of the three categories: VAHL, MHL, and EqHL. ^{18}F -FDG PET-CT = ^{18}F -fluorodeoxyglucose positron emission tomography-computed tomography; HL = hypermetabolic

lymphadenopathy; ALN = axillary lymph node; VAHL = vaccine-associated hypermetabolic lymphadenopathy; MHL = malignant hypermetabolic lymphadenopathy; EqHL = equivocal hypermetabolic lymphadenopathy.

Patients' sex, age, date and site of vaccination, primary disease, purpose of the PET-CT imaging, and blood glucose levels at the time of imaging were collected from the electronic records and analyzed.

2.2. Echographic Examination

Axillary US was performed using an Aplio a550 CUS-AA550 (Canon Medical Systems Inc., Ohtawara-shi, Japan) unit with a 12-MHz linear transducer (18L7; PLT-1204BT, Canon Medical Systems Inc.). This procedure was performed by surgical oncologists with expertise in breast surgery. Detailed information of the axillary lymph nodes was obtained on B-mode and color Doppler US images. Suspected lymph node (LN) metastasis was evaluated based on the following criteria: nodal roundness, absence of fatty tissue in the hilum and the node, diffuse or localized cortical thickening >3 mm, and peripheral blood flow signals within the lymph nodes.

2.3. ^{18}F -FDG PET-CT Imaging

^{18}F -FDG PET-CT imaging was performed using an integrated digital PET-CT scanner (Discovery MI; GE Healthcare, Milwaukee, WI, USA) in all patients. Patients fasted for at least 6 hours and had a blood glucose level of 80-120 mg/dL before intravenous administration of ^{18}F -FDG (3.7 MBq/kg). The total examination time for PET-CT was approximately 20 minutes. All PET images were reconstructed using a three-dimensional time-of-flight weighted line-of-response row-action maximum-likelihood algorithm, with attenuation correction using a CT attenuation map. Integrated PET and CT images were reviewed on an Advantage Workstation Server 3.2 Ext. 3.2 (GE Healthcare). The display field of view was 60 × 60 cm, consisting of 256 × 256 matrixes. Voxel size was 2.34 × 2.34 × 2.79 mm.

2.4. PET-CT Imaging Analysis

PET and CT images were evaluated both visually and quantitatively. For each PET-CT dataset, the morphological features and ^{18}F -FDG uptake of hot ALNs were carefully identified, and a volumetric region of interest (VOI) was drawn on each detectable ALN. The following measurements were obtained from the PET-CT images: length of minor and major axes, cortical thickness, ellipticity (length of minor axis divided by length of major axis), maximum standardized uptake value (SUVmax) of the ALN (LN-SUVmax), and mean standardized uptake value (SUVmean) of blood in the thoracic aorta (B-SUVmean) as the background [14]. The ratio of tumor uptake to background (Tissue-to-Background Ratio; TBR) was calculated by dividing the LN-SUVmax by the B-SUVmean (Figure 2).




Figure 2. Schemas of the measurement methods. Cortical thickness, length of the minor axis minus the lymphatic hilum; ellipticity, minor axis length divided by major axis length; LN-SUVmax,

SUVmax of hypermetabolic axillary lymphadenopathy; arterial blood-SUVmean, SUVmean of arterial blood in the thoracic aorta at the level of the axilla (B-SUVmean); TBR, LN-SUVmax divided by B-SUVmean. All length and thickness measurements are in millimeters.

2.5. Statistical Analysis

Normally distributed continuous variables were presented as mean \pm standard deviation (SD). Categorical variables are presented as numbers and percentages. The Student's t-test was used for normally distributed continuous variables. One-way analysis of variance (ANOVA) was used to compare the three groups. The chi-squared test was used for categorical variables. A multivariate model of logistic regression model was used to evaluate the differentiating factors between vaccine-associated and malignant lymphadenopathy. Quantification of the effect size of the differentiating factors was performed using the ratio (OR) with a 95% confidence interval (95%CI). Statistical significance was defined as $p < 0.05$. Receiver-operating characteristic (ROC) curve analysis was performed to determine the cut-off value with the best combination of sensitivity and specificity. Statistical analysis was performed using SAS software package version 9.4 (SAS Institute Inc., Cary, NC, USA).

3. Results

3.1. Patient Demographics

Table 1 presents the patient demographics according to the group. The mean age was 63.9 (± 14.6 SD) years, with no significant difference noted across the three groups ($p = 0.42$). The proportion of females was the highest in the MHL group (90.5%), probably because 80.9% of the patients in this group had breast cancer. No significant differences were observed in age, disease distribution, or timing of imaging test between the VAHL and EqHL groups ($p > 0.05$). More than half of the patients in all three groups underwent PET-CT examination before treatment. The vaccination rate in the MHL group was 76.2%.

Table 1. Patient characteristics of 3 groups with ultrasonography and/or CT scan.

	ALL (n=188)	VAHL (n=27)	MHL (n=21)	EqHL (n=140)	Pv ^a
Male/Female	87/101	13/14	2/19	72/68	<0.01
Age Mean (SD), y	63.9 (14.6)	66.6 (14.4)	65.9 (12.8)	63 (14.8)	0.42
Disease, No. (%)					
Malignancy					
Breast cancer	37 (19.7)	3 (11)	17 (81)	18 (12.9)	<0.01
Lymphoma	27 (14.4)	3 (11)	2 (10)	22 (15.7)	0.66
Lung cancer	25 (13.3)	7 (26)	1 (5)	17 (12.1)	0.07
Other malignancy	77 (41)	12 (44)	1 (5)	65 (46.4)	<0.01
Non-malignancy	23 (12.2)	2 (7)	0 (0)	19 (13.6)	0.15
Timing of imaging test, No. (%)					
Pre-treatment	113 (60.1)	15 (56)	13 (62)	85 (60.7)	0.87
During treatment	29 (15.4)	3 (11)	8 (38)	18 (12.9)	<0.01
Post-treatment	46 (24.4)	9 (33)	0 (0)	37 (26.4)	<0.05
Vaccination, No. (%)	183 (97.3)	27 (100)	16 (76)	140 (100)	<0.01

Categorical variables are reported as frequencies and percentages. Continuous variables are reported as median and interquartile range (IQR).^a P-value of the T-test among the three group.

3.2. Qualitative Imaging Analysis (Ultrasonography and/or CT)

Table 2 shows the prevalence of fatty hilum within the LNs of interest in all the groups. Fatty hilum was detected in 23.4% (44/188) of the LNs in all patients, including 30% (8/27) in the VAHL group, 0% (0/21) in the MHL group, and 25.7% (36/140) in the EqHL group. Chi-square testing

indicated significant differences between the VAHL and MHL groups, and between the MHL and EqHL groups ($p < 0.01$). This finding suggests that the presence or absence of a fatty hilum is an important factor in the differential diagnosis of benign and malignant axillary lymphadenopathies.

Table 2. Frequency and chi-square test on fatty hilum presence of 3 groups with ultrasonography and/or CT scan.

	Fatty Hilum, No. (%)	
ALL (n=188)	44 (23.4)	
VAHL (n=27)	8 (30)	
MHL (n=21)	0 (0)	a
EqHL (n=140)	36 (25.7)	a

} b

^a $P < 0.01$, ^b Not statistically significant.

3.3. Quantitative LN Analysis on CT

In evaluating lymph nodes on CT images, the node with the longest minor axis was targeted. Comparison of the VAHL and MHL groups by univariate analysis revealed a statistically significant difference in minor axis length, which was also identified as an independent predictor of MHL by multivariate analysis (Table 3a). Although univariate analysis revealed significant differences in minor axis length and ellipticity between the MHL and EqHL groups, only minor axis length was an independent predictor in the multivariate analysis (Table 3b). ROC analysis identified an optimal cut-off for minor axis length of 7.3 mm (sensitivity 0.714, specificity 0.684, AUC 0.733) for discriminating VAHL from MHL (Figure 3). These results suggest that axillary lymph node length > 7.3 mm is a potential sign of malignancy.

Table 3. CT imaging assessment: univariate and multivariate logistic regression analyses between VAHL and MHL groups (a), and MHL and EqHL groups (b).

(a)				
Variable	Univariate		Multivariate	
	Pv	OR (95% CI)	Pv	OR (95% CI)
Age (y.o.)	0.93	1.00 (0.96-1.04)	0.77	0.99 (0.95-1.04)
Minor axis (mm)	< 0.01	0.71 (0.51-0.90)	< 0.01	0.72 (0.51-0.92)
Ellipticity	0.18	0.14 (0.006-2.43)	0.73	0.56 (0.019-16.23)
(b)				
Variable	Univariate		Multivariate	
	Pv	OR (95% CI)	Pv	OR (95% CI)
Age (y.o.)	0.59	1.01	0.79	1.01

		(0.98-1.05)		(0.97-1.04)
Minor axis (mm)	<0.01	1.18 (1.05-1.35)	0.048	1.14 (1.00-1.32)
Ellipticity	0.037	12.4 (1.16-154.99)	0.23	5.1 (0.36-76.71)




Figure 3. Receiver-operating characteristic (ROC) analysis of minor axis length between the VAHL and MHL groups. AUC indicates area under the ROC curve.

3.4. Quantitative LN Analysis on PET-CT

For the assessment of lymph nodes on ^{18}F -FDG PET-CT images, the node with the highest SUVmax was selected and analyzed. The TBR was measured in addition to the minor axis length and ellipticity. Comparison of the VAHL and the MHL groups using univariate analysis revealed statistically significant differences in minor axis length, ellipticity, and TBR ($p < 0.01$). In the multivariate analysis adjusted for age and for blood glucose level at the time of PET-CT examination, ellipticity was identified as an independent predictor of MHL ($p < 0.05$) (Table 4a). Comparison of the MHL and EqHL groups by univariate analysis also identified statistically significant differences in minor axis length, ellipticity, and TBR ($p < 0.01$), and TBR showed a significant difference between the MHL and EqHL groups by multivariate analysis ($p < 0.05$) (Table 4b). ROC analysis identified optimal cut-off values of 0.67 (sensitivity, 0.667; specificity, 0.773; AUC, 0.764) for LN ellipticity and 4.42 (sensitivity, 0.714; specificity, 0.727; AUC, 0.74) for the TBR of LNs for distinguishing between the VAHL and MHL groups (Figure 4a,b). These results indicate that in ALN swelling, ellipticity values > 0.67 and TBR values > 4.42 are possible signs of malignancy.

Table 4. ^{18}F -PET-CT imaging assessment: univariate and multivariate logistic regression analyses between VAHL and MHL groups (a), and MHL and EqHL groups (b).

(a)				
Variable	Univariate		Multivariate	
	Pv	OR (95% CI)	Pv	OR (95% CI)
Age (y.o.)	0.91	1.00 (0.96-1.05)		
TBR	<0.01	0.78 (0.61-0.95)	0.051	0.81 (0.62-1.00)
Minor axis (mm)	<0.01	0.67 (0.48-0.86)	0.082	0.79 (0.55-1.03)
Ellipticity	<0.01	0.002 (<0.001-0.091)	0.045	0.009 (<0.001-0.896)

Blood glucose level (mg/dL)	0.93	1.00 (0.96-1.04)		
(b)				
	Univariate		Multivariate	
Variable	Pv	OR (95% CI)	Pv	OR (95% CI)
Age (y.o.)	0.42	1.01 (0.98-1.05)		
TBR	<0.01	1.18 (1.06-1.33)	0.024	1.17 (1.02-1.34)
Minor axis (mm)	<0.01	1.21 (1.06-1.39)	0.63	1.04 (0.88-1.23)
Ellipticity	<0.01	61.39 (4.11->999)	1.015	45.38 (2.08->999)
Blood glucose level (mg/dL)	0.56	0.99 (0.96-1.02)		




Figure 4. Receiver-operating characteristic (ROC) analysis of **(a)** ellipticity and **(b)** TBR between the VAHL and MHL groups. AUC indicates area under the ROC curve.

4. Discussion

The present study evaluated whether unilateral axillary VAHL after the COVID-19 vaccination can be differentiated from MHL in a clinical setting. We identified patients who exhibited unilateral axillary lymphadenopathy, confirmed their diagnoses based on complete clinical follow-up, and performed qualitative and quantitative assessments of the lymph nodes using the imaging techniques of ultrasonography, CT, and PET-CT. We then investigated useful factors for differentiating VAHL from MHL and evaluated whether imaging tests alone could facilitate diagnosis.

We first investigated whether the presence of a fatty hilum in the lymph nodes could serve as a useful finding for distinguishing between VAHL and MHL, utilizing non-invasive or minimally invasive imaging techniques (ultrasonography and noncontrast CT). We found that no patient with a fatty hilum presented with MHL (Table 2), suggesting that the presence of a fatty hilum could be a determinant of benign lymphadenopathy. In metastatic lymphadenopathy, the absence of a fatty hilum indicates the disruption of the normal lymph node structures. Previous studies have also reported the utility of the presence of a fatty hilum in differentiating between benign and malignant lymphadenopathy [19,20].

Second, in cases in which the fatty hilum is not detected in a lymph node, objective and quantitative evaluation using noncontrast CT might be necessary for diagnosis. Our analysis indicates that minor axis length is the next most important factor in reaching the diagnosis. The

present analysis found that if the minor axis length of the largest enlarged lymph node was ≥ 7.3 mm on CT, it was significantly more likely to be MHL than VAHL (Figure 3). We consider that this cut-off value is more clinically useful compared to size classified as pathologically enlarged according to the revised RECIST guideline version 1.1 (≥ 10 mm for lymph nodes) [21].

Third, we propose that PET-CT should be performed prior to invasive biopsy in cases where the differentiation between benign and malignant lymphadenopathy remains unclear after ultrasound or CT. In our quantitative evaluation using 18F-FDG PET-CT, we focused on the lymph node with the highest SUVmax and measured its minor axis length, ellipticity, and TBR. Subsequent analysis revealed ellipticity and TBR as significant differentiating factors for predicting benign or malignant lymph nodes (Table 4) ($p=0.045, 0.051$). Considering reproducibility and simplicity, our measurement of ellipticity was conducted in the horizontal section with the greatest minor axis length of the node. Although SUVmax can vary depending on the imaging equipment and conditions, TBR serves as a semi-quantitative index that minimizes these effects [22]. Accordingly, ellipticity and TBR measurements on PET-CT can serve as useful tools for distinguishing between MHL and VAHL. The present study is the first to highlight the utility of ellipticity and TBR in differentiating lymphadenopathy following vaccination.

In conclusion, a flowchart that enables the differentiation of benign and malignant unilateral axillary lymphadenopathy following COVID-19 vaccination was developed based on the results of the present study (Figure 5). In patients with unilateral axillary lymphadenopathy, enquiries should first be made regarding their vaccination history and timing. Ultrasonography should be performed to assess the presence or absence of a fatty hilum within the lymph nodes. Unilateral lymphadenopathy with fatty hilum can be considered to indicate VAHL. In cases where no fatty hilum is detected, non-contrast CT should be performed to measure the minor axis of the lymph node. If the minor axis length is < 7.3 mm, VAHL can be suspected, and the majority of patients could be followed-up by careful observation. On the contrary, if the minor axis length is ≥ 7.3 mm, MHL should be suspected and PET-CT scan should be conducted in addition to noncontrast CT scan to measure ellipticity and the TBR of the lymph node with the highest SUVmax. If ellipticity is ≥ 0.67 or TBR is ≥ 4.42 , MHL should be suspected, and a biopsy should be performed for diagnosis.




Figure 5. Flowchart for differentiating initial presentation of unilateral axillary lymphadenopathy into VAHL or MHL categories.

A limitation of the present study was that we performed a retrospective analysis. Prospective clinical studies are needed to ascertain the practical utility of our flowchart for differential diagnosis in clinical practice.

Author Contributions: SY and YY designed the study, collected the clinical data, and drafted the manuscript. MM, TK, SH, SY, TF, and KT participated in study design and oversaw the clinical data. TK performed statistical analysis. YY conceived the study, participated in its design and coordination, and helped draft the manuscript. All authors have read the draft, revised it critically, and approved the final manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: This study was approved by the Ethics Committee for Clinical Studies at Ehime University Graduate School of Medicine (IRB #2206003), and was conducted in accordance with the ethical standards of the 1995 Declaration of Helsinki (as revised in Brazil in 2013).

Informed Consent Statement: Informed consent was obtained using the opt-out principle. Information about the study was disclosed on the associated website (https://www.m.ehime-u.ac.jp/school/clinical.oncology/?page_id=1487), which provided the opportunity to decline participation in this study.

Data Availability Statement: The datasets analyzed in this study are available from the corresponding author upon reasonable request.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Prime Minister's Office of Japan <https://www.kantei.go.jp/jp/headline/kansensho/vaccine.html> (access 2023-05-23).
2. Noumi, T.; Watanabe, H.; Ninomiya, K.; et al. COVID-19 vaccine-associated lymphadenopathy mimicking regrowth of axillary lymph node metastasis of lung adenocarcinoma. *Acta Med Okayama*. **2022**;76(5): 593-596. doi: 10.18926/AMO/64041
3. Kado, S.; Kamiya, K.; Iwabuchi, S.; et al. Unilateral lymphadenopathy associated with COVID-19 vaccination. *J Cutan Immunol Allergy*. **2022**;5:100-101. doi: 10.5114/pjr.2022.117560.
4. Takayuki, Y.; Susumu, O.; Isolated unilateral axillary lymphadenopathy diagnosed in a teenager by ultrasonography after BNT162b2 COVID-19 vaccination. *The Kitakanto Medical Journal*. **2022**;72(1), 87-89. doi: 10.2974/kmj.72.87.
5. Edmonds, CE.; Zuckerman, SP.; Conant, EF.; Management of unilateral axillary lymphadenopathy detected on breast MRI in the era of COVID-19 vaccination. *AJR Am J Roentgenol*. **2021**;217(4):831-834. doi: 10.2214/AJR.21.25604.
6. Park, JY.; Lee, JY.; Yi, SY.; Axillary lymphadenopathy on ultrasound after COVID-19 vaccination and its influencing factors: a single-center study. *J Clin Med*. **2022**;11:238. doi: 10.3390/jcm11010238.
7. Ha, SM.; Chu, AJ.; Lee, J.; et al. US evaluation of axillary lymphadenopathy following COVID-19 vaccination: a prospective longitudinal study. *Radiology*. **2022**;305(1): 46-53. doi: 10.1148/radiol.220543.
8. Mukai, K.; Tsunoda, H.; Imai, R.; et al. The location of unilateral axillary lymphadenopathy after COVID-19 vaccination compared with that of metastasis from breast cancer without vaccination. *Jpn J Radiol*. **2023**;41(6): 617-624. doi: 10.1007/s11604-023-01387-1.
9. Shin, M.; Hyun, CY.; Choi, YH.; et al. COVID-19 vaccination-associated lymphadenopathy on FDG PET/CT: distinctive features in adenovirus-vectored vaccine. *Clin Nucl Med*. **2021**;46(10):814-819. doi: 10.1097/RLU.0000000000003800.
10. El-Sayed, MS.; Wechie, GN.; Low, CS.; et al. The incidence and duration of COVID-19 vaccine-related reactive lymphadenopathy on ¹⁸F-FDG PET-CT. *Clin Med (Lond)*. **2021**;21(6):e633-e638. doi: 10.7861/clinmed.2021-0420.
11. Skawran, S.; Gennari, AG.; Dittli, M.; et al. [¹⁸F] FDG uptake of axillary lymph nodes after COVID-19 vaccination in oncological PET/CT: frequency, intensity, and potential clinical impact. *Eur Radiol*. **2022**;32(1):508-516. doi: 10.1007/s00330-021-08122-2.
12. Schroeder, DG.; Jang, S.; Johnson, DR.; et al. Frequency and characteristics of nodal and deltoid FDG and ¹¹C-choline uptake on PET performed after COVID-19 vaccination. *AJR Am J Roentgenol*. **2021**;217(5):1206-1216. doi: 10.2214/AJR.21.25928.
13. Calabria, F.; Bagnato, A.; Guadagnino, G.; et al. COVID-19 vaccine related hypermetabolic lymph nodes on PET/CT: Implications of inflammatory findings in cancer imaging. *Oncol Res*. **2023**;31(2):117-124. doi: 10.32604/or.2023.027705.
14. Ho, TC.; Shen, DH.; Chang, CC.; et al. Immune response related to lymphadenopathy post COVID-19 vaccination. *Vaccines (Basel)*. **2023**;11(3):696. doi: 10.3390/vaccines11030696.

15. Lederer, K.; Castaño, D.; Gómez, A.D.; et al. SARS-CoV-2 mRNA vaccines foster potent antigen-specific germinal center responses associated with neutralizing antibody generation. *Immunity*. **2020**;53(6):1281-1295.e5. doi: 10.1016/j.immuni.2020.11.009.
16. Hayashi, H.; Sun, J.; Yanagida, Y.; et al. Intradermal administration of DNA vaccine targeting Omicron SARS-CoV-2 via pyro-drive jet injector provides the prolonged neutralizing antibody production via germinal center reaction. *Sci Rep*. **2023**;13(1):13033. doi: 10.1038/s41598-023-40172-y.
17. Cohen, D.; Krauthammer, S.H.; Wolf, I.; et al. Hypermetabolic lymphadenopathy following administration of BNT162b2 mRNA Covid-19 vaccine: incidence assessed by [¹⁸F] FDG PET-CT and relevance to study interpretation. *Eur J Nucl Med Mol Imaging*. **2021**;48(6):1854-1863. doi: 10.1007/s00259-021-05314-2.
18. Yamanaka, S.; Miyagawa, M.; Sugawara, Y.; et al. The prognostic significance of whole-body and spleen MTV (metabolic tumor volume) scanning for patients with diffuse large B cell lymphoma. *Int J Clin Oncol*. **2021**;26(1):225-232. doi: 10.1007/s10147-020-01807-6.
19. Rahbar, H.; Partridge, S.C.; Javid, S.H.; et al. Imaging axillary lymph nodes in patients with newly diagnosed breast cancer. *Curr Probl Diagn Radiol*. **2012**;41(5):149-58. doi: 10.1067/j.cpradiol.2011.08.002.
20. Choi, H.Y.; Park, M.; Seo, M.; et al. Preoperative axillary lymph node evaluation in breast cancer: current issues and literature review. *Ultrasound Q*. **2017**;33(1):6-14. doi: 10.1097/RUQ.0000000000000277.
21. Eisenhauer, E.A.; Therasse, P.; Bogaerts, J.; et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer*. **2009**;45(2):228-47. doi: 10.1016/j.ejca.2008.10.026.
22. Gheysens, O.; Jamar, F.; Glaudemans, A.W.J.M.; et al. Semi-quantitative and quantitative [¹⁸F] FDG-PET/CT indices for diagnosing large vessel vasculitis: a critical review. *Diagnostics (Basel)*. **2021**;11(12):2355. doi: 10.3390/diagnostics11122355

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.