

Case Report

Not peer-reviewed version

Deep Venous Thrombosis as an Initial Manifestation of Rectal Adenocarcinoma Metastasis to Skeletal Muscle: Case Report

[Tsung-Jung Tsai](#) and [Yu-Yao Chang](#) *

Posted Date: 29 July 2024

doi: 10.20944/preprints202404.1501.v2

Keywords: Rectal cancer; Metastasis; skeletal muscle metastasis; Deep vein thrombosis



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Case Report

Deep Venous Thrombosis as an Initial Manifestation of Rectal Adenocarcinoma Metastasis to Skeletal Muscle: Case Report

Tsung-Jung Tsai ¹ and Yu-Yao Chang ^{1,2,*}

¹ Division of Colon & Rectal Surgery, Department of Surgery, Changhua Christian Hospital 50006, No 135, Nanxiao Street, Changhua City, Taiwan

² Department of Post-Baccalaureate Medicine, College of Medicine, National Chung Hsing University, Taichung, Taiwan

* Correspondence: 177176@cch.org.tw; Tel: +886-4-7238595; Fax: +886-4-7232942

Abstract: Colorectal cancer is third most common malignancy. For metastasis of rectal cancer, metastasis to skeletal muscle tissue is rare. In this case report, we describe a 51-year-old male patient with rectal cancer with neoadjuvant concurrent radiotherapy and chemotherapy followed by transanal total mesorectal excision with ileostomy. However, he had deep venous thrombosis of left leg after operation and skeletal muscle metastasis was diagnosed via tissue proof. Therefore, less possibility of metastasis increased included brain, duodenum, bone, and nervous system. The prognosis of metastasis to skeletal muscle is poor in contrast to liver or lung metastasis.

Keywords: rectal cancer; metastasis; skeletal muscle metastasis; deep vein thrombosis

Introduction

Colorectal cancer is third most common malignancy and second cancer-related mortality [1]. For metastasis of rectal cancer, lung is the most common, followed by lung, and peritoneum [2]. As times goes, the survival was improved attributed to target therapy and chemotherapy. Therefore, less possibility of metastasis increased included brain, duodenum, bone, and nervous system. Metastasis to skeletal muscle tissue is rare. Herein, we reported a case presenting with deep venous thrombosis but skeletal muscle metastasis to quadriceps medialis and adductor magnus as manifestation of adenocarcinoma of rectum.

Case Report

An 66-year-old female has a no medical history. She was presented with diarrhea, change in defecation habit and tenesmus and visited to outpatient department. On rectal examination, a annular mass was palpable at 5 cm from the anal verge. Biopsy from the mass showed adenocarcinoma. Pelvic magnetic resonance imaging and positron emission tomography were ordered and revealed rectal cancer, cT3N2M0. She was received neoadjuvant concurrent radiotherapy and chemotherapy followed by transanal total mesorectal excision with ileostomy. Pathology revealed pT3N2M0, stage III. She regularly visited outpatient department and ileostomy closure was applied after three months later. However, anal pain was complained. Anal nodule was found and biopsy revealed adenocarcinoma. Local recurrence was impressed and laparoscopic abdominoperineal resection was performed. We suggested infusion chemotherapy but she refused. Therefore, she was started on tablet capecitabine. After five months of operation, she complained left leg swelling. Abdominal computed tomography was ordered and it revealed May-Thurner syndrome with left side deep vein thrombosis. (Figure 1) Enoxaparin was prescribed first and rivaroxaban substituted it. Besides, stent was inserted. However, swelling persisted and lower limb computed tomography was ordered again 3 months later. It revealed May-Thurner syndrome post

stenting, collapsed left proximal superficial femoral vein related to external compression, and diffuse calcification and area of low density in the left thigh muscle.(Figure 2,3) Computed tomography biopsy was applied and pathology revealed adenocarcinoma. On the other side, she had peritoneal carcinomatosis and intestinal obstruction and ileocecal bypass was performed. Target therapy with chemotherapy was given and she passed away one year later.

Discussion

Approximately 80% of recurrences manifest with the initial 3 years after surgery and 95% occurring within 5 years. Primary tumor of skeletal muscle metastasis is most common in gastrointestinal, followed by lung and malignant melanoma. Different tumor tends to certain muscles frequently. For skeletal muscle metastasis, trunk is most common, followed by iliopsoas and gluteal muscle [3].

The presentation of computed tomography has five type. Muscle metastasis type 1 and 2 were most common. Type IV was rare and present with calcification of myositis ossificans, calcific tendinitis, and calcific myonecrosis as our case [3].

The etiology of metastasis included hematogeneous, lymphatic, peritoneal spread or direct extension of the tumor. For muscle, muscle sarcolemma is a physical barrier like blood brain barrier to protect from invasion [4]. It can't be metastasized easily without direction invasion. In this case, we suspected it was related to deep venous thrombosis causing blood stasis. The concept was explained by the seed and soil theory [5].

Patients with metastasis to liver or lung have a better prognosis than those with metastasis to brain, bone, and skeletal muscle. The prognosis of metastasis to skeletal muscle is poor with expected survival ranging from 5 to 12 months [6].

Conclusion

Metastasis to skeletal muscle is rare and deep vein thrombosis may be one of risk factors. When the symptoms persist, keep metastasis in mind.

Ethics approval: No ethical approval is required according to institutional policies.

Consent to participate: All patient-related info was de-identified so patient consent was waived.

Consent for publication: All patient-related info was de-identified so patient consent was waived.

Data availability Statement: Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Conflicts of interest: The authors declare that they have no conflict of interest.

References

1. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA A Cancer J. Clin.* **2018**, *68*, 394–424, <https://doi.org/10.3322/caac.21492>.
2. Pan, H.-D.; Zhao, G.; An, Q.; Xiao, G. Pulmonary metastasis in rectal cancer: a retrospective study of clinicopathological characteristics of 404 patients in Chinese cohort. *BMJ Open* **2018**, *8*, e019614, <https://doi.org/10.1136/bmjopen-2017-019614>.
3. Meyer HJ, Ullrich S, Surov A: CT imaging features of skeletal muscle metastasis: A rare tumour group with different patterns. *J Med Imaging Radiat Oncol* 2020, 64:674-678.
4. Landriscina, M.; Gerardi, A.M.T.; Fersini, A.; Modoni, S.; Stoppino, L.P.; Macarini, L.; Sanguedolce, F.; Bufo, P.; Neri, V. Multiple Skeletal Muscle Metastases from Colon Carcinoma Preceded by Paraneoplastic Dermatomyositis. *Case Rep. Med.* **2013**, *2013*, 1–4, <https://doi.org/10.1155/2013/392609>.
5. Langley, R.R.; Fidler, I.J. The seed and soil hypothesis revisited—The role of tumor-stroma interactions in metastasis to different organs. *Int. J. Cancer* **2011**, *128*, 2527–2535, <https://doi.org/10.1002/ijc.26031>.
6. Singh, S.; Bisht, N.; Joshi, R.; Mishra, P.S.; Mulajker, D.; Gupta, S. Musculoskeletal Metastasis from Primary Rectal Cancer: Series of Two Cases of a Very Rare Occurrence with a Short Literature Review. *J. Gastrointest. Cancer* **2018**, *50*, 991–996, <https://doi.org/10.1007/s12029-018-0165-5>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.