

Article

Not peer-reviewed version

---

# Use of the Drone for Cost-Effective Surveys In Natura 2000 Protected Areas: A Case Study on Monitoring Plant Diversity in Sicily (Italy)

---

[Gianmarco Tavilla](#) <sup>\*</sup>, [Alessandro Crisafulli](#), [Pietro Minissale](#), [Valeria Tomaselli](#), [Maria Adamo](#)

Posted Date: 13 May 2024

doi: [10.20944/preprints202405.0886.v1](https://doi.org/10.20944/preprints202405.0886.v1)

Keywords: endemic species; landscape monitoring; Mediterranean flora; remote sensing; Saxifraga; vascular flora



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

## Article

# Use of the Drone for Cost-Effective Surveys in Natura 2000 Protected Areas: A Case Study on Monitoring Plant Diversity in Sicily (Italy)

Gianmarco Tavilla <sup>1,\*</sup>, Alessandro Crisafulli <sup>2</sup>, Pietro Minissale <sup>3</sup>, Valeria Tomaselli <sup>4</sup>  
and Maria Adamo <sup>1</sup>

<sup>1</sup> National Research Council of Italy, Institute of Atmospheric Pollution Research (CNR-IIA), c/o Interateneo Physics Department, 70125 Bari, Italy

<sup>2</sup> Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina, Italy

<sup>3</sup> Department of Biological, Geological and Environmental Sciences, University of Catania, 95125 Catania, Italy

<sup>4</sup> Department of Biosciences, Biotechnologies and Environment, University of Bari "Aldo Moro", 70125 Bari, Italy

\* Correspondence: gianmarcotavilla@cnr.it

**Abstract:** Unmanned aerial vehicles (UAVs), commonly known as drones, present a cost-effective solution for the swift collection of data from vast and remote areas that are otherwise difficult to access. The Mediterranean Basin, known for being a hot spot for plant biodiversity, hosts several habitats and taxa of significant naturalistic value. However, many of these areas are often inaccessible to botanists, making exploration and research challenging. The aim of this paper is to involve the utilization of drone surveys and open-source software for botanical research. Our primary goal is to show the effectiveness of these tools in the field and demonstrate their practical application in Natura 2000 sites. The protected area chosen for this research is Rocca di Novara, situated in northeastern Sicily. Thanks to our drone investigations, we were able to capture images of the mountain's side that is inaccessible to humans. This allowed us to observe the habitat of some species in detail. One of the most fascinating discoveries was the reappearance of *Saxifraga callosa* subsp. *australis*, which had not been confirmed in this area for over 140 years. Using drones for botanical research can boost field research, making monitoring easier and more cost-effective over time, especially in Natura 2000 sites.

**Keywords:** endemic species; landscape monitoring; Mediterranean flora; remote sensing; *Saxifraga*; vascular flora

## 1. Introduction


The task of observing and charting both natural and semi-natural habitats is crucial in protecting ecological diversity and the environment. These procedures are guided by European standards and regulations [1,2], which ensures that conservation efforts are carried out in a methodical and scientific manner. In the Mediterranean basin, the task of habitat mapping and investigations presents a unique challenge due to the wide range of flora combined with varied environmental conditions [3]. This area is a biodiversity hotspot that hosts a variety of habitats, including forests, grasslands, wetlands, and coastal zones, each with its own distinct ecological characteristics [4]. Accurate monitoring and observation of these habitats are necessary to understand the dynamics of the ecosystem and develop effective conservation management strategies, especially for habitats of European interest [5,6]. In recent times, drone technology has significantly transformed habitat mapping and monitoring, especially when it comes to conserving plant biodiversity [7]. Drones, also known as Unmanned Aerial Vehicles (UAVs), provide an innovative perspective on environmental conservation [8,9]. Equipped with advanced imaging capabilities and sensors, drones can capture high-resolution aerial photographs and generate detailed 3D models of habitats, offering a more comprehensive and precise representation of the terrain and vegetation structure compared to traditional ground-based surveys [10,11]. Moreover, the recent advent of Unmanned Aerial Vehicles (UAVs) has introduced the prospect of low-cost vegetation mapping [12,13]. In the vast and varied landscapes of the Mediterranean region, drones are particularly useful because they can cover large areas quickly and efficiently [14]. Drones can access remote and challenging-to-reach areas with minimal disturbance to the ecosystem, making

them a valuable tool for monitoring sensitive habitats and endangered species [15]. They can also track changes in vegetation over time, providing essential data for assessing the impact of environmental changes and the effectiveness of conservation strategies [16]. The use of UAV photogrammetry has significantly enhanced the efficiency of 3D modeling and vegetation monitoring [17]. Furthermore, drone use contributes to democratizing science. As drone technology becomes more affordable and user-friendly, citizens can participate in conservation efforts by collecting and sharing data about [18,19]. This not only expands the scope of habitat mapping and monitoring but also fosters a greater public understanding and appreciation of biodiversity. In conclusion, drone use in habitat mapping and monitoring represents a significant advancement in environmental conservation. As this technology continues to evolve, it promises to play an increasingly important role in preserving the rich biodiversity of the Mediterranean region and beyond. The study aims to use drones to monitor and evaluate rocky habitats in a Natura 2000 site in Sicily and assess the cost-effectiveness of plant monitoring in inaccessible areas using user-friendly and open-source software.

## 2. Materials and Methods

### 2.1. Study Area

Sicily is an island located in the central Mediterranean. It is the largest island in the Mediterranean, covering an area of 25,832.4 km<sup>2</sup> (including smaller islands), and has a coastline of 1637 km [20]. The Tyrrhenian Sea is situated to the north of the island, the Sicilian Channel to the southwest, and the Ionian Sea to the east. The Strait of Messina separates Sicily from the Italian peninsula to the northeast (Figure 1). The island has a rich and diverse flora, with a total of 3569 accepted taxa [21]. Sicily and its islets are a significant biodiversity hotspot with a consistent history of discovering both native and alien species [22–25]. The island's unique natural environment has garnered considerable interest from botanists due to its rich and diverse ecosystem. Consequently, there has been a continuous flow of new floristic records and descriptions of new species [26–31].



**Figure 1.** Map of Sicily (DEM used as basemap [32]); Rocca di Novara (red star).

The main northern mountain range of the island, with the highest peak being Montagna Grande at 1374 m, is the Peloritani Mountains. In particular, for the goals of this study, the Rocca di Novara mount (also known as Rocca Salvatesta) has been chosen as the study area. This peak is situated within the boundaries of Novara di Sicilia, located in the province of Messina [33]. Geologically, it

is predominantly composed of limestone rocks, which makes it a peculiar feature of the Peloritani chain, known for its metamorphic rock formations [34]. The mountain range of Peloritani, including Rocca di Novara, makes it difficult for researchers to study its vegetation using traditional in-field techniques. This is why many regions within these mountains have not been thoroughly researched in terms of their vegetation and flora characteristics. The presence of limestone outcrops has resulted in a distinct and diverse floristic set, setting it apart from other peaks in the area. This area falls within the SAC "Rocca di Novara" (code ITA030006) due to its relevant naturalistic heritage (Figure 2). Moreover, Rocca di Novara belongs to the Important Plant Area (IPA) named "Monti Peloritani e Rupi di Taormina" [35].




**Figure 2.** Rocca di Novara, Sicily (Italy), photo by G. Tavilla.

## 2.2. Drone Surveys and Data Collection

The field investigation was carried out in May 2022 and June 2022 in order to evaluate the floristic biodiversity of the area. In this research, the drone model used to collect the images was DJI Mavic 3. It uses the WGS 84 Geodetic Reference System (SGR). The 20-megapixel RGB camera on the Mavic 3 captures images with a resolution of 5280×3956 pixels. During the flight, the camera was stabilized using a three-axis gimbal that was incorporated with the UAV's inertial navigation system. A Global Navigation Satellite System (GNSS) receiver provides the UAV's geolocation at the moment of image capture, which is stored in the image's EXIF (Exchangeable Image File Format) header. This unmanned aerial vehicle was chosen due to its cost-effectiveness and integration of a high-resolution camera and gimbal. To guarantee correct stitching, at least 80% of the front and 70% of the sides of each image were overlapped when acquired [36]. During the flight, drone photographs were taken manually using a DJI smart controller. We carried out a total of 10 flights between May and June 2022, and each session had a maximum duration of 30 minutes. These flights led to the gathering of more than 1000 photographs from the study area. The field site analysis involved the utilization of aerial pictures processed through

open-source software. The aerial photos of the cliff were transformed into a high-resolution 3D model using WebODM version 2.4.2 software, a free photogrammetric image processing program developed by OpenDroneMap [37]. This tool is specifically designed to process aerial photos captured by drones and is available for both Windows and Mac OS. WebODM is specifically designed for processing aerial photos captured by drones. For each UAV picture uploaded, WebODM retrieves the EXIF header. The Ground Sampling Distance (GSD) refers to the amount of surface area covered by a single image in flight, and it was the parameter used to evaluate our project on WebODM. The GSD value of 1 cm/px is the one recommended for professional surveys to achieve higher accuracy. Additionally, the following options were selected: auto-boundary: true, mesh-octree-depth: 12, use-3dmesh: true, pc-quality: high, mesh-size: 300000. The entire 3D model processing was entirely automatic (Figure 3).



**Figure 3.** Workflow of this study. Step 1, prepare drones including battery check. Step 2, capturing images using manual flight. Step 3, photo selection and processing. Step 4, plant identification.

### 2.3. Vegetation Sampling and Taxa Identification

The captured images were used to create virtual plots that enabled the evaluation of individual species coverage using a phytosociological method. This method, also known as the Braun-Blanquet approach [38], considers the relationships of plant communities with the environment and the interactions within communities. The phytosociological relevés take into account vegetation plot size, vegetation cover, and species occurrence. The Braun-Blanquet scale assigns each species a coverage value ranging from 1 (1-5% coverage) to 5 (75-100% coverage), while the symbol "+" is used for sporadic species with coverage less than 1%. As concerns the species observed in the pictures, the taxonomic identification was carried out using the "Flora of Italy" [39–42], and the life form follows the Raunkiaer system [43]. The nomenclature of taxa follows the "Portal to the Flora of Italy" [21], while the nomenclature of *Saxifraga callosa* subsp. *australis* (Moric.) Pignatti ex Tavilla & Del Guacchio

is in accordance with Tavilla & Del Guacchio [44]. The habitat type code was used in accordance with Directive 92/43/EEC [45].

### 3. Results and Discussion

The survey conducted in this Natura 2000 site utilized drone technology for comprehensive monitoring. We present the outcomes of our drone-based monitoring campaign in Rocca di Novara. Our study used advanced drone technology to gather high-resolution spatial data and imagery, providing unparalleled insight into the biodiversity of the site. A total area equal to 16,039.92 m<sup>2</sup> has been reconstructed with the Ground Sampling Distance (GSD) of 0.63 cm. Through our aerial surveys, image analysis, and interpretation, we have gained valuable insights into essential ecological parameters (Figure 4).



**Figure 4.** Vegetation changes between (A) May and (B) June (Rocca di Novara, Sicily), photos by G. Tavilla.

In particular, managing the 3D rocky face model was easy with WebODM's user-friendly interface (Figure 5). This interface allowed measurements of the diameter of the identified plants. Conducting population studies on rare species and vegetation sampling are now feasible Table 1. Furthermore, thanks to the phytosociological relevés carried out, it was possible to identify the habitat type in which this vegetation occurs. These rocky communities growing on the limestone side are included in habitat 8210 "Calcareous rocky slopes with chasmophytic vegetation", which is listed in Annex I of Directive 92/43/EEC. This is a vegetation of fissures of limestone cliffs, in the Mediterranean region belonging essentially to the *Saxifragion australis* Biondi & Balelli ex Brullo 1984 (*Asplenietea trichomanis* (Br.-Bl. in Meier & Br.-Bl. 1934) Oberdorfer 1977 class) [46].



**Figure 5.** WebODM 3D model interface.

**Table 1.** Phytosociological relevé from drone image.

Relevé n.	1
Date	28 May 2022
Locality	Rocca di Novara
Plot size (m <sup>2</sup> )	15
Cover (%)	35
Species	
<i>Saxifraga callosa</i> subsp. <i>australis</i>	2
<i>Sedum dasyphyllum</i>	+
<i>Hypochaeris laevigata</i>	1
<i>Edraianthus graminifolius</i> subsp. <i>siculus</i>	+
<i>Athamanta sicula</i>	2
<i>Saxifraga rotundifolia</i>	1
<i>Arabis alpina</i> subsp. <i>caucasica</i>	+
<i>Festuca circummediterranea</i>	1
<i>Aubrieta columnae</i> subsp. <i>sicula</i>	+
<i>Poa bivonae</i>	1
<i>Asplenium ceterach</i>	+

### 3.1. *Taxa of Outstanding Interest*

The Rocca di Novara mountain is home to several plant species of high phytogeographic interest that were initially investigated by Nicotra [47]. Thanks to the use of drones, our study has allowed us to highlight specific taxa of naturalistic interest. The predominant life form found on the cliff is the chamaephyte, which is not surprising. However, hemicryptophytes, such as *Hypochaeris laevigata* (L.) Ces., Pass. & Gibelli and *Saxifraga rotundifolia* L. subsp. *rotundifolia*, also thrived on the side. In addition, some Sicilian endemic species grow on the cliffs of Rocca di Novara. The following paragraphs discuss the most noteworthy taxa.

#### 3.1.1. *Lomelosia crenata* (Cirillo) Greuter & Burdet subsp. *crenata*

It is a chamaephyte cespitose found in Albania, Algeria, Greece, Italy, Serbia, Sicily, Spain, and Tunisia [48]. According to Giardina et al. [49]. This taxon is rare across the rest of Sicily, but it is

frequent on the rocky peaks of Northern Sicily, such as Rocca di Novara (Figure 6). Therefore this species has an important phytogeographical value in Sicily.



**Figure 6.** *Lomelosia crenata* (Cirillo) Greuter & Burdet subsp. *crenata* (Rocca di Novara, Sicily), photograph by G. Tavilla.

### 3.1.2. *Saxifraga callosa* subsp. *australis* (Moric.) Pignatti ex Tavilla & Del Guacchio

This chamaephyte grows in shady, mountain environments on the limestone of central and southern Apennines and Western Sicily [44]. The only population known for the eastern part of Sicily is in Rocca di Novara [49]. In the only floristic work ever made for Rocca di Novara by Nicotra, *S. callosa* subsp. *australis* was reported [47]. According to the author, he did not see the plant himself, but it was first noted by G. Seguenza. For over 140 years, there has been no confirmation of this species' presence. Thanks to our drone monitoring, we were able to photograph the plant on an inaccessible side of the rocky cliff, not accessible by humans (Figure 7). This is another excellent example of how drones can be useful in botanical research, especially in cliff ecosystems [50], by avoiding hazards for botanists in rupicolous habitats and unnecessary spending (Figure 8). It should be noted that this is the only plant reported in this paper that can only be identified and reported through drone surveys.




**Figure 7.** *Saxifraga callosa* subsp. *australis* (Moric.) Pignatti ex Tavilla & Del Guacchio (Rocca di Novara, Sicily), photograph by G. Tavilla.



**Figure 8.** Inaccessible side of Rocca di Novara, photograph by G. Tavilla.

### 3.1.3. *Aubrieta columnae* Guss. subsp. *sicula* (Strobl) M.A.Koch, D.A.German & R.Karl

This chamaephyte is endemic to Sicily and is classified as an Endangered species according to IUCN criteria [51,52]. It grows on limestone cliffs of the Madonie massif (north-west Sicily), and only one population in north-east Sicily, i.e., Rocca di Novara (Figure 9). Nicotra [47] reported for the first time this taxa in Peloritani Mountains. Based on our recent field investigations, the majority of individuals bloomed in May. However, in the past, blooming peaks were seen in late June. This observation suggests that individual fertility may be at risk in the future since the Mediterranean mountain regions are most vulnerable to climate change [53].



**Figure 9.** *Aubrieta columnae* Guss. subsp. *sicula* (Strobl) M.A.Koch, D.A.German & R.Karl (Rocca di Novara, Sicily), photograph by G. Tavilla.

### 3.1.4. *Odontites bocconeи* (Guss.) Walp. subsp. *bocconeи*

It is an endemic Sicilian chamaephyte (Figure 10) with a rare distribution on the Island [49]. Currently, it is classified at the national level as Least Concern [54]. Furthermore, it should be noted that *O. bocconeи* is not currently reported in the section "Other important species of flora and fauna" of the Rocca di Novara SAC data form [55].



**Figure 10.** *Odontites bocconeii* (Guss.) Walp. subsp. *bocconeii* (Rocca di Novara, Sicily), photograph by G. Tavilla.

#### 4. Conclusions

This research has enabled us to gain insights into various aspects of biodiversity, habitat structure, and ecosystem health within the Natura 2000 site. Drones have distinct advantages, including rapid and cost-effective data acquisition over expansive and often inaccessible terrain. Our workflow shows an easily applicable way to monitor and assess floristic diversity in protected areas. The use of this tool ensures the safety of botanists by preventing them from exploring inaccessible and dangerous habitats. Furthermore, the aerial view provided by drones has enabled us to comprehensively assess landscape features and ecological processes at various spatial scales, facilitating a more nuanced understanding of ecosystem functioning.

**Author Contributions:** Conceptualization, G.T.; methodology, G.T.; software, G.T.; formal analysis, G.T.; investigation, G.T, A.C., P.M.; data curation, G.T.; writing—original draft preparation, G.T.; writing—review and editing, G.T., A.C., P.M., V.T., M.A.; funding acquisition, P.M. All authors have read and agreed to the published version of the manuscript.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data that support the findings of this study are available from the corresponding author.

**Acknowledgments:** The technical equipment was financed and supported by the project INTERREG V-A ITALIA-MALTA 2014-2020 Axis III – Objective 3.1 FAST - Fight Alien Species Transborder CUP: E99C20000160005

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Directive, H.; others. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Official Journal of the European Union* **1992**, *206*, 50.
2. Evans, D. The Habitats of the European Union Habitats Directive. *Biology and Environment: Proceedings of the Royal Irish Academy* **2006**, *106B*, 167–173.
3. Médail, F.; Baumel, A. Using phylogeography to define conservation priorities: The case of narrow endemic plants in the Mediterranean Basin hotspot. *Biological Conservation* **2018**, *224*, 258–266.
4. Medail, F.; Quezel, P. Biodiversity hotspots in the Mediterranean Basin: setting global conservation priorities. *Conservation biology* **1999**, *13*, 1510–1513.
5. Bunce, R.; Metzger, M.; Jongman, R.; Brandt, J.; De Blust, G.; Elena-Rossello, R.; Groom, G.B.; Halada, L.; Hofer, G.; Howard, D.; others. A standardized procedure for surveillance and monitoring European habitats and provision of spatial data. *Landscape ecology* **2008**, *23*, 11–25.
6. Tavilla, G.; Angiolini, C.; Bagella, S.; Bonini, F.; Cambria, S.; Caria, M.C.; Esposito, A.; Fanfarillo, E.; Ferri, V.; Fiaschi, T.; Gianguzzi, L.; del Galdo, G.G.; Ilardi, V.; Mei, G.; Minissale, P.; Rivieccio, G.; Sciandrello, S.; Stinca, A.; Bazan, G. New national and regional Annex I Habitat records: from #37 to #44. *Plant Sociology* **2022**, *59*, 49–66. doi:10.3897/pls2022591/05.
7. Pimm, S.L.; Alibhai, S.; Bergl, R.; Dehgan, A.; Giri, C.; Jewell, Z.; Joppa, L.; Kays, R.; Loarie, S. Emerging technologies to conserve biodiversity. *Trends in ecology & evolution* **2015**, *30*, 685–696.
8. Jiménez López, J.; Mulero-Pázmány, M. Drones for Conservation in Protected Areas: Present and Future. *Drones* **2019**, *3*. doi:10.3390/drones3010010.
9. Koh, L.P.; Wich, S.A. Dawn of drone ecology: low-cost autonomous aerial vehicles for conservation. *Tropical conservation science* **2012**, *5*, 121–132.
10. Wich, S.A.; Koh, L.P. *Conservation Drones: Mapping and Monitoring Biodiversity*; Oxford University Press, 2018. doi:10.1093/oso/9780198787617.001.0001.
11. Strumia, S.; Buonanno, M.; Aronne, G.; Santo, A.; Santangelo, A. Monitoring of Plant Species and Communities on Coastal Cliffs: Is the Use of Unmanned Aerial Vehicles Suitable? *Diversity* **2020**, *12*. doi:10.3390/d12040149.
12. Bryson, M.; Reid, A.; Hung, C.; Ramos, F.T.; Sukkarieh, S., Cost-Effective Mapping Using Unmanned Aerial Vehicles in Ecology Monitoring Applications. In *Experimental Robotics: The 12th International Symposium on Experimental Robotics*; Khatib, O.; Kumar, V.; Sukhatme, G., Eds.; Springer Berlin Heidelberg: Berlin, Heidelberg, 2014; pp. 509–523. doi:10.1007/978-3-642-28572-1\_35.
13. Green, D.R.; Hagon, J.J.; Gómez, C.; Gregory, B.J. Chapter 21 - Using Low-Cost UAVs for Environmental Monitoring, Mapping, and Modelling: Examples From the Coastal Zone. In *Coastal Management*; Krishnamurthy, R.; Jonathan, M.; Srinivasulu, S.; Glaeser, B., Eds.; Academic Press, 2019; pp. 465–501. doi:https://doi.org/10.1016/B978-0-12-810473-6.00022-4.
14. Dunford, R.; Michel, K.; Gagnage, M.; Piégay, H.; Trémelo, M.L. Potential and constraints of Unmanned Aerial Vehicle technology for the characterization of Mediterranean riparian forest. *International Journal of Remote Sensing* **2009**, *30*, 4915–4935.
15. Eischeid, I.; Soirinen, E.M.; Assmann, J.J.; Ims, R.A.; Madsen, J.; Pedersen, Pirotti, F.; Yoccoz, N.G.; Ravolainen, V.T. Disturbance Mapping in Arctic Tundra Improved by a Planning Workflow for Drone Studies: Advancing Tools for Future Ecosystem Monitoring. *Remote Sensing* **2021**, *13*. doi:10.3390/rs13214466.
16. Bersaglio, B.; Enns, C.; Goldman, M.; Lunstrum, L.; Millner, N. Grounding drones in political ecology: understanding the complexities and power relations of drone use in conservation. *Global Social Challenges Journal* **2023**, *2*, 47 – 67. doi:10.1332/HNEK4485.
17. Li, M.; Yan, E.; Zhou, H.; Zhu, J.; Jiang, J.; Mo, D. A novel method for cliff vegetation estimation based on the unmanned aerial vehicle 3D modeling. *Frontiers in Plant Science* **2022**, *13*. doi:10.3389/fpls.2022.1006795.
18. Pucino, N.; Kennedy, D.M.; Carvalho, R.C.; Allan, B.; Ierodiaconou, D. Citizen science for monitoring seasonal-scale beach erosion and behaviour with aerial drones. *Scientific reports* **2021**, *11*, 3935.
19. Theuerkauf, E.J.; Bunting, E.L.; Mack, E.A.; Rabins, L.A. Initial insights into the development and implementation of a citizen-science drone-based coastal change monitoring program in the Great Lakes region. *Journal of Great Lakes Research* **2022**, *48*, 606–613. doi:https://doi.org/10.1016/j.jglr.2022.01.011.

20. Treccani.it. [https://www.treccani.it/enciclopedia/sicilia\\_res-51b7c2ab-973b-11e5-8844-00271042e8d9/](https://www.treccani.it/enciclopedia/sicilia_res-51b7c2ab-973b-11e5-8844-00271042e8d9/), accessed on 14.04.2024.

21. Portal to the Flora of Italy. <http://dryades.units.it/floritaly>, accessed on 14.04.2024.

22. Minissale, P.; Cambria, S.; Montoleone, E.; Tavilla, G.; Giusso del Galdo, G.; Sciandrello, S.; Badalamenti, E.; Mantia, T.L. The alien vascular flora of the Pantelleria Island National Park (Sicily Channel, Italy): new insights into the distribution of some potentially invasive species. *BioInvasions Records* **2023**, *12*, 861 – 885. doi:10.3391/bir.2023.12.4.01.

23. Sciandrello, S.; Cambria, S.; Giusso del Galdo, G.; Guarino, R.; Minissale, P.; Pasta, S.; Tavilla, G.; Cristaudo, A. Floristic and Vegetation Changes on a Small Mediterranean Island over the Last Century. *Plants* **2021**, *10*. doi:10.3390/plants10040680.

24. Brullo, S.; Minissale, P.; Spampinato, G. Considerazioni fitogeografiche sulla flora della Sicilia. *Ecologia mediterranea* **1995**, *21*, 99–117.

25. Cambria, S.; Brullo, C.; Tavilla, G.; Sciandrello, S.; Minissale, P.; Giusso Del Galdo, G.; Brullo, S. *Ferula sommieriana* (Apiaceae), a new species from Pelagie Islands (Sicily). *Phytotaxa* **2021**, *525*, 89–108.

26. Cambria, S.; Azzaro, D.; Caldarella, O.; Aleo, M.; Bazan, G.; Guarino, R.; Torre, G.; Cristaudo, A.E.; Ilardi, V.; La Rosa, A.; Laface, V.L.A.; Luchino, F.; Mascia, F.; Minissale, P.; Sciandrello, S.; Tosetto, L.; Tavilla, G. New Data on Native and Alien Vascular Flora of Sicily (Italy): New Findings and Updates. *Plants* **2023**, *12*. doi:10.3390/plants12091743.

27. Brullo, S.; Brullo, C.; Cambria, S.; Tavilla, G.; Giusso del Galdo, G.; Bogdanović, S. Taxonomical and chorological remarks on the Mediterranean *Poa maroccana* (Poaceae) and the first record in Italy from the Sicilian flora. *Acta Botanica Croatica* **2021**, *80*, 63 – 73. doi:10.37427/botcro-2021-011.

28. Brullo, S.; Cambria, S.; Crisafulli, A.; Tavilla, G.; Sciandrello, S. Taxonomic remarks on the *Centaurea aeolica* (Asteraceae) species complex. *Phytotaxa* **2021**, *483*, 9–24. doi:<https://doi.org/10.11646/phytotaxa.483.1.2>.

29. Brullo, S.; Brullo, C.; Cambria, S.; Minissale, P.; Sciandrello, S.; Siracusa, G.; Tavilla, G.; Tomaselli, V.; Giusso Del Galdo, G. Taxonomical remarks on *Solenopsis laurentia* (Campanulaceae) in Italy. *Phytotaxa* **2023**, *584*, 59–88. doi:<https://doi.org/10.11646/phytotaxa.584.2.1>.

30. Cambria, S.; Giusso del Galdo, G.; Minissale, P.; Sciandrello, S.; Tavilla, G. *Lablab purpureus* (Fabaceae), a new alien species for Sicily. *Flora Mediterranea* **2022**, *32*, 73 – 78. doi:10.7320/FIMedit32.073.

31. Musarella, C.M.; Laface, V.L.A.; Angiolini, C.; Bacchetta, G.; Bajona, E.; Banfi, E.; Barone, G.; Biscotti, N.; Bonsanto, D.; Calvia, G.; Cambria, S.; Capuano, A.; Caruso, G.; Crisafulli, A.; Del Guacchio, E.; Di Gristina, E.; Domina, G.; Fanfarillo, E.; Fascetti, S.; Fiaschi, T.; Galasso, G.; Mascia, F.; Mazzacuva, G.; Mei, G.; Minissale, P.; Motti, R.; Perrino, E.V.; Picone, R.M.; Pinzani, L.; Podda, L.; Potenza, G.; Rosati, L.; Stinca, A.; Tavilla, G.; Villano, C.; Wagensommer, R.P.; Spampinato, G. New Alien Plant Taxa for Italy and Europe: An Update. *Plants* **2024**, *13*. doi:10.3390/plants13050620.

32. Tarquini, S.; Nannipieri, L. The 10m-resolution TINITALY DEM as a trans-disciplinary basis for the analysis of the Italian territory: Current trends and new perspectives. *Geomorphology* **2017**, *281*, 108–115. doi:<https://doi.org/10.1016/j.geomorph.2016.12.022>.

33. Ruggieri, R. North-Eastern Sicily: Karst Area of the Nebrodi Mts. and the Peloritani Mts. In *Karst of Sicily: A Journey Inside and Outside the Island's Mountains*; Springer, 2022; pp. 403–436.

34. Cirrincione, R.; Fazio, E.; Ortolano, G.; Pezzino, A.; Punturo, R. Fault-related rocks: deciphering the structural–metamorphic evolution of an accretionary wedge in a collisional belt, NE Sicily. *International Geology Review* **2012**, *54*, 940–956.

35. Blasi, C.; Marignani, M.; Copiz, R.; Fipaldini, M.; Bonacquisti, S.; Del Vico, E.; Rosati, L.; Zavattero, L. Important Plant Areas in Italy: From data to mapping. *Biological Conservation* **2011**, *144*, 220–226. doi:<https://doi.org/10.1016/j.biocon.2010.08.019>.

36. Takahashi, N.; Wakutsu, R.; Kato, T.; Wakaizumi, T.; Ooishi, T.; Matsuoka, R. Experiment on UAV photogrammetry and terrestrial laser scanning for ICT-integrated construction. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences* **2017**, *XLII-2/W6*, 371–377. doi:10.5194/isprs-archives-XLII-2-W6-371-2017.

37. WebODM Drone Software. <https://www.opendronemap.org/webodm/>, accessed on 14.04.2024.

38. Braun-Blanquet, J. *Pflanzensoziologie: grundzüge der vegetationskunde*; Springer-Verlag, 2013.

39. Pignatti, S. *Volume 1: Flora d'Italia & Flora Digitale*. In *Flora d'Italia: In 4 Volumi*, 2 ed.; Edagricole-Edizioni Agricole di New Business Media srl: Milano, Italy, 2017.

40. Pignatti, S. *Volume 2: Flora d'Italia & Flora Digitale*. In *Flora d'Italia: In 4 Volumi*, 2 ed.; Edagricole-Edizioni Agricole di New Business Media srl: Milano, Italy, 2017.
41. Pignatti, S. *Volume 3: Flora d'Italia & Flora Digitale*. In *Flora d'Italia: In 4 Volumi*, 2 ed.; Edagricole-Edizioni Agricole di New Business Media srl: Milano, Italy, 2018.
42. Pignatti, S.; Guarino, R.L.R.M. *Volume 4: Flora d'Italia & Flora Digitale*. In *Flora d'Italia: In 4 Volumi*, 2 ed.; Edagricole-Edizioni Agricole di New Business Media srl: Milano, Italy, 2019.
43. Raunkiaer, C. *The Life Forms of Plants and Statistical Plant Geography Being the Collected Papers of C. Raunkiaer*; Clarendon Press, 1934.
44. Tavilla, G.; Del Guacchio, E. An updated taxonomic treatment for *Saxifraga callosa* sensu amplo (Saxifragaceae). *Phytotaxa* **2023**, *622*, 165–171. doi:10.11646/phytotaxa.622.2.6.
45. The European Environment Agency (EEA). <https://eunis.eea.europa.eu/habitats-code.jsp>, accessed on 14.04.2024.
46. Biondi, E.; Blasi, C.; Burrascano, S.; Casavecchia, S.; Copiz, R.; Del Vico, E.; Galderzi, D.; Gigante, D.; Lasen, C.; Spampinato, G.; et al. Manuale Italiano di Interpretazione Degli Habitat Della Direttiva 92/43/CEE. Società Botanica Italiana. Ministero dell'Ambiente e Della Tutela del Territorio e del Mare, D.P.N. 2009. <http://vnr.unipg.it/habitat>, accessed on 14.04.2024.
47. Nicotra, L. Notizie intorno alla vegetazione del Salvatesta. *Nuovo Giornale Botanico Italiano* **1880**, *12*, 366–370.
48. Domina, G. Dipsacaceae. *Euro+ Med Plantbase—The Information Resource for Euro-Mediterranean Plant Diversity*. Available online: <http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp> **2017**.
49. Giardina, G.; Raimondo, F.M.; Spadaro, V. A catalogue of plants growing in Sicily. *Bocconeia* **2007**, *20*, 5–582.
50. Zhou, H.; Zhu, J.; Li, J.; Xu, Y.; Li, Q.; Yan, E.; Zhao, S.; Xiong, Y.; Mo, D. Opening a new era of investigating unreachable cliff flora using smart UAVs. *Remote Sensing in Ecology and Conservation* **2021**, *7*, 638–648. doi:<https://doi.org/10.1002/rse2.214>.
51. Fenu, G.; Cambria, S.; Giacò, A.; Khabibullaev, B.S.; Shomurodov, K.F.; Peruzzi, L.; Porrovecchio, M.; Tavilla, G.; Orsenigo, S. Global and Regional IUCN Red List Assessments: 16. *Italian Botanist* **2023**, *16*, 121–133. doi:10.3897/italianbotanist.16.115947.
52. IUCN Red List of Threatened Species. Version 2022-2. <https://www.iucnredlist.org>, accessed on 14.04.2024.
53. Kazakis, G.; Ghosn, D.; Remoundou, I.; Nyktas, P.; Talias, M.A.; Vogiatzakis, I.N. Altitudinal Vascular Plant Richness and Climate Change in the Alpine Zone of the Lefka Ori, Crete. *Diversity* **2021**, *13*. doi:10.3390/d13010022.
54. Rossi, G.; Orsenigo, S.; Gargano, D.; Montagnani, C.; Peruzzi, L.; Fenu, G.; Abeli, T.; Alessandrini, A.; Astuti, G.; Bacchetta, G.; others. Lista Rossa della Flora Italiana. 2 Endemiti e altre specie minacciate. **2020**.
55. Biodiversity Europa. <https://biodiversity.europa.eu/sites/natura2000/ITA030006>, accessed on 14.04.2024.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.