
Bowman-Birk Inhibitor Mutants of Soybean Generated by CRISPR-Cas9 Reveal Drastic Reductions in Trypsin and Chymotrypsin Inhibitor Activities

Won-Seok Kim , [Jason D. Gillman](#) , Sunhyung Kim , Junqi Liu , Madhusudhana R. Janga , Robert M. Stupar , [Hari B. Krishnan](#) *

Posted Date: 16 May 2024

doi: 10.20944/preprints202405.1076.v1

Keywords: soybean; Glycine max; Antinutritional factors; Bowman-Birk inhibitor; CRISPR-Cas9



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

Bowman-Birk Inhibitor Mutants of Soybean Generated by CRISPR-Cas9 Reveal Drastic Reductions in Trypsin and Chymotrypsin Inhibitor Activities

Won-Seok Kim ¹, Jason D. Gillman ², Sunhyung Kim ³, Junqi Liu ⁴, Madhusudhana R. Janga ⁵, Robert M. Stupar ⁴ and Hari B. Krishnan ^{2,*}

¹ Division of Plant Science and Technology, University of Missouri, Columbia, MO 65211, USA; wonseokk@missouri.edu

² Plant Genetics Research Unit, US Department of Agriculture-Agricultural Research Service, Columbia, MO 65211, USA; Jason.Gillman@usda.gov

³ Division of Plant Science and Technology, University of Missouri, Columbia, MO 65211, USA; jeongs@missouri.edu

⁴ Department of Agronomy and Plant Genetics, University of Minnesota, St. Paul, MN 55108, USA; liuqx162@umn.edu; stup0004@umn.edu

⁵ Institute of Genomics for Crop Abiotic Stress Tolerance, Texas Tech University, Lubbock, TX 79409, USA; mjanga@ttu.edu

* Correspondence: hari.b.krishnan@usda.gov; Tel.: 573-882-8151

Abstract: Despite the high quality of soybean protein, raw soybeans and soybean meal cannot be directly included into animal feed mixtures due to the presence of Kunitz (KTI) and Bowman Birk protease inhibitors (BBI) which reduces animal productivity. Heat treatment can substantially inactivate trypsin and chymotrypsin inhibitors (BBI) but such treatment is energy intensive, adds expense, and negatively impact the quality of seed proteins. As an alternative approach we have employed CRISPR/Cas9 gene editing to create mutations in *BBI* genes to drastically lower the protease inhibitor content in soybean seed. Agrobacterium-mediated transformation was used to generate several stable transgenic soybean events. These independent CRISPR/Cas9 events were examined in comparison to wild type plants using Sanger sequencing, proteomic analysis, trypsin/chymotrypsin inhibitor activity assays and qRT-PCR. Collectively, our results demonstrate the creation of an allelic series of loss of function mutations affecting the major *BBI* gene in soybean. Mutations in two of the highly expressed seed specific *BBI* genes lead to substantial reductions in both trypsin and chymotrypsin inhibitor activities.

Keywords: soybean 1; *Glycine max* (L. (Merr)) 2; Antinutritional factors 3; Bowman-Birk inhibitor 4; CRISPR-Cas9 enabled mutagenesis 5

1. Introduction

Soybean *Glycine max* (L.) Merr) is grown worldwide as a rotation partner for grain crops, particularly maize, due to an ability to fix atmospheric nitrogen, high seed yield potential, relatively few major biotic pathogens and relatively low input costs [1,2]. Soybean seeds contain far more protein than other legumes (on average 40%) and soybean meal is often the first choice for swine and poultry feed mixtures due to a balanced (albeit suboptimal) amino acid profile. Soybean meal, produced after extraction of seed oil, has a relatively low cost and is of high quality compared to other competing protein meal sources [3].

Despite the high quality of soybean protein, raw soybeans and soybean meal cannot be directly included into animal feed mixtures due to anti-nutritional compounds that interfere with digestion, cause pancreatic inflammation, and effectively reduce animal weight gain. Soybean seeds contain

two major classes of proteinaceous anti-nutritional factors [4], which were named after the scientists that discovered them: 1) Kunitz-trypsin inhibitors [5] (KTI) inactivate the digestive enzyme trypsin; and 2) Bowman-Birk inhibitors [6] (BBI), which inactivate both trypsin and chymotrypsin animal digestive enzymes [7]. In soybean, three KTI isoforms (KTI1, KTI2 and KTI3) have been reported [8–10]. There are at least 11 different *BBI* encoding genes present in the soybean genome, however expression analysis of RNAseq data reveals only three of them are expressed at appreciable levels in developing seeds (Figure S1).

Experiments have conclusively shown that ingestion of high levels of active protease inhibitors are responsible for reduced animal weight gain, and can even result in hypoglycemia, pancreatic hypertrophy, or liver damage [11]. To avoid these issues and permit inclusion of soybean meal at high levels in animal feed, KTI and BBI anti-nutritional proteins in typical soybean seeds must be inactivated by heat-treatment, which has been shown to reduce, but not eliminate ANF activity [7]. Heat inactivation allows soybean meal to be included at moderately high levels in chicken and swine feed mixtures. While this step meets the needs of the industry, such treatment is energy intensive, adds expense, and requires a balance between sufficiently reducing ANF activity and negatively impacting the quality of seed proteins [3,11].

One success story in genetically reducing soybean's trypsin inhibitor activity was the identification and analysis of a soybean accession (PI 157740) which features marked reduced (~40%) in trypsin inhibitor activity. This genetic source has been examined through both genetic analysis and animal feeding trials. Raw extruded protein meal lacking KTI3 protein was found to result in superior animal weight gain compared with raw soybeans [12], but not a degree that could eliminate the need for heat inactivation. In our previous research, a soybean line (PI 68679) with reductions in trypsin inhibition due to a mutation affecting the KTI1 gene was identified [13]. Using hybridization, the *KTI-1* mutation (PI 68679) was combined with the previously identified *KTI-3* mutation (PI 542044), to yield soybean lines that lack both KTI1 and KTI3 protein. These soybean lines have lower trypsin inhibitor activity than the currently available *KTI-3* mutant [13]. Unfortunately, all lines which have reduced KTI levels unexpectedly had unacceptable levels of trypsin inhibitor activity, due to overaccumulation of BBI proteins in seeds of all mutant lines (*KTI1*, *KTI3*, and *KTI1/KTI3*).

To develop soybean lines which require extremely low or no heat-inactivation processing, we hypothesize that both *KTI* and *BBI* gene expression in seeds must be reduced or ideally eliminated. Naturally occurring *BBI* mutants have been identified in perennial soybean collections [14]. However, to our knowledge no naturally occurring *G. max* mutant or germplasm entry has been reported with substantive reduced BBI accumulation or activity. However, one report indicated that transgenic expression of a modified protein can lower BBI activity and resulted in viable plants which were able to produce seed [15].

CRISPR/Cas9 gene editing [16,17], is now a routine practice for inducing targeted loss of function mutations in soybean [17–19]. This method could be leveraged to induce mutations of soybean seed *BBI* genes and decrease or eliminate their accumulation in seeds. In this study, we report on the creation of an allelic series of loss-of-function mutations affecting the major *BBI* seed expressed genes in soybean, and mutation of two genes was shown to lead to substantial reductions in both trypsin and chymotrypsin inhibitor activities due to loss of function mutations affecting *BBI* genes.

2. Results

2.1. Generation of *BBI* Mutants by CRISPR/Cas9 Gene Editing

At least 11 *BBI* encoding genes exist in the soybean genome, however transcriptome analysis implicates only one major and two minor seed-specific *BBI* genes in developing soybean seeds (Supplementary Figure S1) [20]. Because *BBI* genes share extensive sequence homology we designed target sites that enabled simultaneous knockout of all three seed-expressed genes, and two CRISPR/Cas9 expression cassettes were designed, which differed in use of either single or double guideRNA (Supplementary Figure S2). Utilizing these two CRISPR/Cas9 expression cassettes, we generated several independent transgenic events by *Agrobacterium*-mediated soybean transformation (Figure 1A). Six transgenic events were grown in the greenhouse and leaf tissue from these plants

were utilized for the isolation of genomic DNA. Sanger sequencing of PCR products generated by utilizing primers that are specific for the most abundantly expressed *BBi* gene (Glyma.16g208900) identified several mutations in this gene. Figure 1A shows a schematic summary of mutations identified in these six independent events leading to deletions, insertions, and frameshift mutations on amino acid sequence. (Figure 1B, Figure S3). Similarly, we also sequenced PCR products generated by utilizing primers that are specific for the second most abundantly expressed *BBi* gene (Glyma.14G117700). This analysis revealed several mutations that introduced deletions, insertions, and frameshift mutations on amino acid sequence (Figure 2A and 2B; Figure S3)




Figure 1. CRISPR-Cas9 mutations identified in Glyma.16g208900, the most highly expressed BBi isoform. A) Summary of mutations identified in six independent events. B) Effect of CRISPR–CAS9 induced deletions, insertions, and frameshift mutations on amino acid sequence.




Figure 2. CRISPR-Cas9 mutations identified in Glyma.14G117700, the second most highly expressed BBi isoform. A) Summary of mutations identified in six independent events. B) Effect of CRISPR–CAS9 induced deletions, insertions, and frameshift mutations on amino acid sequence.

2.2. SDS-PAGE Analysis of Seed Proteins of CRISPR/Cas9 Transgenic Plants

First, we wanted to compare the total seed protein profile between the wildtype and CRISPR/Cas9 transgenic plants. For this purpose, we first isolated total seed proteins from the wildtype and the six independent transgenic events and resolved them by SDS-PAGE (Figure 3A). A visual examination of the total seed protein profile between wild type and transgenic events did not reveal any drastic alterations between them (Figure 3A). All soybean seeds examined accumulated the two most abundant seed storage proteins (7S β -conglycinin and 11S glycinin). However, the accumulation of the β -subunit of β -conglycinin, a 52 kDa glycoprotein, in all transgenic events were noticeably lower when compared to the wild type (Figure 3A). To substantiate these changes, we performed immunoblot analysis using antibodies raised against the 7S β -conglycinin and 11S glycinin antibodies. The 7S β -conglycinin antibody reacted against all three subunits. The accumulation pattern of the α' and α subunits of β -conglycinin was similar between the wild type and the six CRISPR/Cas9 transgenic events (Figure 3B). In contrast, the accumulation of the β -subunit of β -conglycinin was much lower in all transgenic events when compared to the wild type, and for one transgenic event (Figure 3B, lane 7) only an extremely faint band was detected. We also performed immunoblot analysis to quantify any changes in the 11S glycinin accumulation between wild type and CRISPR/Cas9 transgenic events. The glycinin antibody recognized three glycinin species (A3; A1a, A1b, A2, and A4; A5). In contrast to the β -subunit of β -conglycinin, the accumulation of 11S glycinin showed little variation in their abundance between the wild type and the CRISPR/Cas9 transgenic events (Figure 3C).




Figure 3. Proteomic analysis of total seed proteins from CRISPR-Cas9 BBi mutant and control lines.

(A) Sodium dodecyl sulfate polyacrylamide gel electrophoresis of total seed proteins visualized with Coomassie Blue; (B) Immunoblot using β -conglycinin specific antibodies; (C) Immunoblot using glycinin specific antibodies. Samples in order are 'Maverick' (Mav., non-transgenic control); 2= Event 2; 3=Event 3; 4=Event 4; 5=Event 5; 7=Event 7; 8=Event 8.

2.3. Immunoblot Analysis Reveal the Absence of BBi in the Seeds of Transgenic BBi Mutants

To examine the accumulation of BBi in seeds of the CRISPR/Cas9 transgenic events we performed western blot analysis utilizing BBi-specific peptide-antibodies [21]. A protein fraction enriched for protease inhibitors obtained by calcium fractionation method [22] was utilized for the immunoblot analysis and transferred to a nitrocellulose membrane. The nitrocellulose membrane when challenged with soybean BBi-specific antibodies failed to detect the accumulation of this protein in all transgenic events. In contrast, the BBi antibody reacted against a 11-12 kDa protein in the wild-type plant (Figure 4B). We also examined the accumulation of KTi, which also contributes significantly to the overall protease inhibitor of soybean seed, by immunoblot analysis, via antibodies raised against soybean KTi [9]. Immunoblot analysis clearly demonstrated normal accumulation of KTi proteins in all the plants examined in our study (Figure 4C).




Figure 4. Proteomic analysis of soluble seed proteins from CRISPR-Cas9 BBi mutant and control lines. (A) Sodium dodecyl sulfate polyacrylamide gel electrophoresis of calcium fractionated soluble seed proteins visualized with Coomassie Blue; (B) Immunoblot using Bowman-Birk inhibitor (BBi) specific antibodies; (C) Immunoblot using KTi specific antibodies (KTi). Samples in order are 'Maverick' (Mav., non-transgenic control); 2= Event 2; 3=Event 3; 4=Event 4; 5=Event 5; 7=Event 7; 8=Event 8.

2.4. Chymotrypsin and Trypsin Inhibitor Activities Are Drastically Reduced in CRISPR/Cas9 gene Edited Transgenic BBi Mutants

Since our immunoblot analysis clearly showed the absence of BBi protein accumulation in the CRISPR/Cas9 gene edited transgenic *BBi* mutants we also wanted to examine protease inhibitor activities in these plants. The chymotrypsin inhibitor activity (CIU) was measured using N-Succinyl-Ala-Ala-Pro-Phe p-nitroanilide as a substrate and is expressed as CIU /mg seed powder. The wild-type soybean seed contained 17.2 CIU while in transgenic seeds the activity ranged from 2.89-15.9 CIU. Out of the six transgenic events examined in this study, four had drastic reductions; we saw a 2.8-6.0-fold reduction in CIU compared to wild type (Table 1). Since soybean BBi proteins can inhibit both chymotrypsin and trypsin, we also measured the trypsin inhibitor activity (TIU) using the same seed extracts (Table 1), with N α -Benzoyl-DL-arginine 4-nitroanilide hydrochloride as a substrate. The trypsin inhibitor activity in the wild-type soybean seeds averaged 57.25 TIU/mg seed powder, whereas we saw reduced activities in mutant lines ranging from a low of 14.46 (Event 8) to a high of 35.7 TIU (Event 5) for different transgenic *BBi* events. This corresponds to a 1.6-4.0-fold reduction in TIU activities. TIU activity in all examined transgenic *BBi* mutants were significantly lower than in wild type seeds. Moreover, in four of the transgenic events (Events 2, 4, 7 and 8; Table 1) we observed drastic reductions in the TIU activities ranging from 25.3-41.9% of the TIU activities found in wild type seeds. Interestingly, the reduction in TIU activity in these *BBi* mutants is more significant than we observed for a naturally occurring soybean frameshift *KTi-3* mutant (Table 1). CIU and TIU activities in seeds of mutant and the control line were fully concordant when activities were expressed as TIU/mg seed powder or as TIU/mg seed protein (Table 1).

Table 1. Analysis of trypsin inhibitor activity and chymotrypsin inhibitor activities in seeds from transgenic CRISPR-Cas9BBi mutagenized and control plants. Tukey HSD indicates the results of a Tukey-Shapiro Honest Square Difference test; presence of the same letter indicates insignificant differences between genotypes.

Genotype	n=	Seed powder		Seed Protein	
		Mean \pm SD CIU/mg	Tukey HSD	Mean \pm SD CIU/mg	Tukey HSD
'Maverick' (WT)	3	17.2 \pm 1.91	A	56.93 \pm 6.32	A
KTi-3	3	18.55 \pm 0.42	A	60.97 \pm 1.39	A
Event 5	3	15.93 \pm 1.66	A	56.45 \pm 5.89	A
Event 3	3	15.35 \pm 1.63	A	51.34 \pm 5.44	A
Event 4	3	6.2 \pm 1.77	B	21.15 \pm 6.03	B
Event 2	3	4.19 \pm 1.03	B	15.28 \pm 3.77	B
Event 7	3	3.9 \pm 1.24	B	16.85 \pm 5.38	B
Event 8	3	2.89 \pm 0.04	B	11.54 \pm 0.17	B

Genotype	n=	Seed powder		Seed Protein	
		Mean \pm SD TIU/mg	Tukey HSD	Mean \pm SD TIU/mg	Tukey HSD
'Maverick' (WT)	3	57.25 \pm 0.35	A	189.48 \pm 1.17	A
KTi-3	3	36.45 \pm 1.63	B	119.8 \pm 5.35	BC
Event 5	3	35.7 \pm 1.27	B	126.55 \pm 4.51	B
Event 3	3	31.96 \pm 1	C	106.88 \pm 3.35	C
Event 4	3	24 \pm 0.59	D	81.87 \pm 2.01	D
Event 2	3	21.38 \pm 0.06	DE	77.97 \pm 0.21	D

Event 7	3	19.75±1.89	E	85.35±8.15	D
Event 8	3	14.46±1.71	F	57.74±6.82	E

2.5. qRT-PCR Analysis Confirms the Absence of Expression of the Abundant BBi Genes in CRISPR/Cas9 Gene Edited Transgenic BBi Mutants

We also examined the expression of the two most abundantly expressed BBi genes (Glyma.16G208900, Glyma.14G117700) along with few other BBi genes (Glyma.09G158500, Glyma.09G158600, Glyma.09G158700) in the CRISPR/Cas9 gene edited transgenic BBi mutants. Total RNA was isolated from mid-stage (35 days after flowering) developing T1 soybean seeds and used for qRT-PCR analysis. The gene specific primers used in our study are shown in Table 3. In the wild-type soybean seeds, we could detect the expression of most of these BBi genes. At this stage of seed development, we observed high level expression of Glyma.16G208900, Glyma.14G117700 and Glyma.09G158700 (Figure 5). In contrast, the expression of these abundantly expressed BBi genes was essentially not detected in two of the transgenic events (event # 2 and 4), while several fold lower expression was detected in one of the mutants (event # 8).




Figure 5. RT-PCR analysis of T1 seeds with CRISPR-Cas9 BBi mutant and control lines.

3. Discussion

In this study, we have generated several stable transgenic soybean lines that exhibit the lowest levels of trypsin and chymotrypsin inhibitor activities that have been reported in the literature. Several of our CRISPR-Cas9 BBi lines have much lower trypsin inhibitor activity (62% to 75% lower) when compared to control plants. The relative abundance of KTi proteins in soybean seeds is higher than BBi [23]. It has been reported that BBi proteins are more heat stable than KTi protein since the BBi contains multiple cysteine residues leading to conformational change of BBi during heating [24]. Since our CRISPR-Cas9 BBi lines contain undetectable levels of BBi proteins, as evidenced by immunoblot analysis, one would expect these newly developed BBi mutant lines may require only minimal heat treatment.

Until now, no soybean null mutants with reduced or ablated BBi proteins have been identified in *G. max* germplasm. However, BBi mutants have been identified in several perennial *Glycine spp.* [14]. Unfortunately, substantial hybridization barriers exist between perennial *Glycine spp.* and *G. max* which has hampered the development of *G. max* BBi mutants by traditional breeding. Attempts have been made to lower the protease inhibitor activity in soybean seeds through overexpression of a mutated BBi gene [25]. The transgenic seeds generated in that study revealed a 20-50% reduction in

protease inhibitory activity. In contrast, we have developed several CRISPR-Cas9 mutated *B*B*i* lines that showed remarkable reduction (ranging from 63-84%) in chymotrypsin inhibitor activity when compared to the wild-type seeds. Even though our CRISPR-Cas9 constructs were targeted to knockout seed specific *B*B*i* genes, its pronounced effect on lowering the trypsin inhibitor is striking. This drastic reduction in trypsin inhibitor activity is not mediated by a reduction in the accumulation of KTi proteins in the CRISPR-Cas9 *B*B*i* lines. Our immunoblot analysis (Figure 4) clearly demonstrates that KTi proteins accumulate to the similar levels in *B*B*i* mutant lines as in wild type seeds. Thus, despite the presence of functional versions of all three KTi seed expressed genes, we saw a significant reduction in the trypsin inhibitor activity. This reduction is solely because of *B*B*i* mutations, and unexpectedly revealed that *B*B*i* contributes significantly to overall trypsin inhibitor activity in soybean seeds.

Our immunoblot analysis (Figure 4) and qRT-PCR analysis (Figure 5) confirmed that the gRNA we employed in our study to knockout the seed specific *B*B*i* genes was very effective. In addition to eliminating the expression of the two most abundant *B*B*i* genes (Glyma.16G208900 and Glyma.14G117700) other *B*B*i* genes (Glyma.09G158500, Glyma.09G158600, Glyma.09G158700) were also eliminated in some the CRISPR-Cas9 *B*B*i* induced mutant lines. This was an expected result as *B*B*i* genes share extensive sequence homology and constructs were designed to target multiple *B*B*i* genes. The full number of mutated genes is unknown and will likely require creation and analysis of segregating populations to discern the number of mutated genes and their relative contributions to trypsin and chymotrypsin inhibition activities. One intriguing prospect would be the combination of null mutations for both seed expressed KTi genes and our newly identified null mutations affecting all seed expressed *B*B*i* genes.

Our study also reveals proteome alteration due to knock out of *B*B*i* genes. One-dimensional gel analysis of total seed proteins from *B*B*i* mutant lines reveals lower accumulation of the β -subunit of β -conglycinin. The accumulation of this protein is influenced by nitrogen and sulfur status of the plant [26,27]. The accumulation of β subunit of β -conglycinin, a protein that is deficient in sulfur containing amino acids, is elevated when the plants are grown under limited sulfur supply [28–30]. In the current study, there is a drastic reduction in the accumulation of B*B*i in all CRISPR-Cas9 mutated *B*B*i* lines. B*B*i contains 14 cysteine residues and acts as a main sink for sulfur storage in the seed [31,32]. In the absence of B*B*i accumulation, cysteine that would be normally incorporated in B*B*i will be freely available. We speculate that the reduction in the accumulation of the β -subunit of β -conglycinin could be related to an increase in the availability of cysteine in the seeds, which would be incorporated into other, normally lower abundance seed expressed proteins. Our suggestion for this possibility is strengthened by previous reports that has shown an increased availability of sulfur amino acids in transgenic soybean plants reduces the accumulation of β subunit of β -conglycinin [33]. However, this hypothesis remains to be rigorously tested through greenhouse and/or field studies. Although we clearly demonstrate the creation of an allelic series of *B*B*i* null mutations through CRISPR Cas9 mutagenesis, we acknowledge we do not currently know what impact, if any, such mutations (or a combination of mutations) will have on field agronomic performance or seed yield. Similar work with *K*T*i-3* mutant lines did not identify any deleterious impacts on whole plant agronomic traits or seed yield [34].

Our results open the possibility that breeding lines could be developed that require substantially less heat-treatment, which would save both money and energy which is used to heat treat soybeans. One study performed with *K*T*i-3* and lectin free seed demonstrated that partial loss of trypsin/chymotrypsin inhibitors resulted in superior animal weight gain performance as compared to conventional soybean meal, however both were inferior to heat processed soybean meal [11]. Given our finding that our *B*B*i* mutants had reduced trypsin inhibitor activity equivalent to the *K*T*i-3* mutation (Table 1), the *B*B*i* allelic series we have created may open the door to even greater reductions on trypsin and chymotrypsin inhibitor content in animal feed mixtures and for human food products, such as tofu and soymilk, while also reducing the energy required for heat processing.

4. Materials and Methods

4.1. Chemicals

Most chemicals and reagents used in this study were of analytical grade. Chemicals for electrophoresis including acrylamide, bis acrylamide, SDS, TEMED, ammonium persulfate were purchased from GE healthcare (Piscataway, NJ). Tris-HCL, trypsin, chymotrypsin, β -mercaptoethanol, N α -Benzoyl-DL-arginine p-nitroanilide hydrochloride, and N-Succinyl-Ala-Ala-Pro-Phe p-nitroanilide were purchased from Sigma-Aldrich (St. Louis, MO).

4.2. Analysis of Expression of BBi Genes Using Public Data Repositories

Soybase.org is a repository for data from multiple experimental datasets for soybean. We identified protein coding genes in the 'Williams 82' genome soybean with the annotation PFAM PF00228 (Bowman-Birk serine protease inhibitor family) as targets. Previously collected data from the 'RNA-Seq Atlas' [20] was used to prioritize target BBi genes for CRISPR-Cas9 construct, gene sequencing and RT-PCR analysis. In developing seeds, two genes, Glyma.14g117700 and Glyma.16g208900 represent between 17.0-31.3% and 68.5-82.8% of the total detectable BBi gene expression from seeds samples between 21 and 42 days after flowering.

4.3. Generation of CRISPR/CAS9 induced BBi Knockout Soybean Mutants

For CRISPR/Cas9-mediated mutagenesis of the BBi genes in soybean, two single gRNAs were designed to target the two major expressed BBi genes and three other BBi genes. gRNA1: (5'-GAACAACATGGTGGTGCTAAAGG-3') was a common target in five BBi genes (Glyma.09G158500; Glyma.14G117700; Glyma.09G158600; Glyma.09G158700; Glyma.16G202800) and gRNA2 (5'-CACATGCAGAGATCACAGCATGG-3', reverse complimentary of the sense strand sequence 5'-CCATGCTGTGATCTCTGCATGTG-3') was a common target in only 3 BBi genes (Glyma.09G158600; Glyma.09G158700; Glyma.16G202800). Two plasmid vectors were constructed: the first plasmid contained only gRNA1 and the second plasmid contained both gRNA1 and gRNA2. For cloning of the BBi gRNAs in the T-DNA vector, a two-step Golden Gate Assembly was performed firstly in the module vector pMOD_B2103 and then subsequently in the destination T-DNA vector pTRANS230d as described previously [18,35]. Recombinant pMOD_B2103_BB_i gRNAs plasmids harboring BBi gRNAs were verified by Sanger sequencing of the gRNA cassette with primers flanking the gRNAs: the forward primer TC320 (5'-CTAGAAGTAGTCAAGGCGGC-3') and reverse primer TC089R (5'-GGAACCCTAATTCCTTATCTGG-3'), respectively. The second Golden Gate Assembly reaction was composed of the recombinant pMOD_B2103_BB_i gRNAs, pMOD_A0521, pMOD_C2906, and a destination T-DNA vector pTRANS230d. Final CRISPR/Cas9_BB_i gRNA constructs in the destination T-DNA vector were designated as pTRANS230/BB_i_1gRNA and pTRANS230/BB_i_2gRNA, respectively. The CRISPR constructs were mobilized into a disabled *Agrobacterium tumefaciens* strain EHA105 by heat shock procedure.

4.4. Soybean Transformation

The soybean genotype 'Maverick' [36] was transformed using *Agrobacterium*-mediated transformation of half-seed explants. The transformation protocol was modified from Zeng et al., [37]. The process began by germinating sterilized seeds for 16 hours in Petri dishes on germination media. Then, half seed explants were made from the seeds and treated with an *Agrobacterium* culture [38]. After 5 days of co-cultivation, the explants were transferred to shoot induction medium. Subsequently, the explants underwent two rounds of subcultures before being moved into shoot elongation medium. Elongated shoots were then cut from the explants and used for rooting. After 2 weeks, the rooted plants were transferred into soil. To identify putative transgenic soybean plants, leaf paginating assay was performed with 100 mg L⁻¹ glufosinate-ammonium solution on leaves at three stages. Herbicide resistant plants were cultivated in greenhouses under controlled conditions, with a 16-h day at 28°C and 8-h night at 24°C until T1 seeds were harvested.

4.5. Protein Extraction and 1-D Electrophoresis

Total soybean seed protein extraction and subsequent separation of proteins by 1-D SDS-PAGE were carried out as previously described [39]. Briefly, dry soybean seeds were ground to a fine powder with a mortar and pestle and extracted with 1 mL of sodium dodecyl sulfate (SDS)-sample buffer (60 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol and 5% 2-mercaptoethanol). Tubes were vortexed (Vortex Genie, Scientific Industries, San Diego, CA) for 10 min at room temperature, and then heated to 100° C for 5 min, followed by centrifugation (Eppendorf, Beckman Coulter, Inc., Indianapolis, IN) at 15800xg for 5 min. The supernatant is the total seed protein fraction. This fraction was resolved by 13.5% SDS-PAGE gels using a Hoefer SE 260 minigel apparatus (Amersham Biosciences, Piscataway, NJ). Electrophoretic separation of proteins was achieved with 20 mA per gel (constant current). SDS-PAGE separated proteins were visualized using Coomassie Blue R-250 solution (0.3 % Coomassie Brilliant Blue R-250, 45 % methanol, and 10 % glacial acetic acid).

4.6. Real-Time qRT-PCR Analysis of *BBI* Genes in Developing Soybean Seeds

Total RNA from 35-day after anthesis soybean seeds were extracted using TRIzol reagent (Invitrogen) according to the manufacturer's instructions. Real-time qRT-PCR was conducted using a LightCycler 480 II instrument (Roche, Indianapolis, IN). Each qRT-PCR reactions were conducted with two replications in 20 µL reaction using the QuantiNova SYBR Green RT-PCR Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. The cycling condition of absolute quantification analysis were 10 min at 50 °C then 2 min 95°C for cDNA synthesis followed by 40 cycles of 20 sec at 95 °C and 40 sec at 60 °C, with the data acquired at 60 °C in green channel. The reference gene (F-box protein, Glyma.12G051100) was amplified by RT-PCR from Maverick soybean seed total RNA with gene specific primer set (Supplementary Table S1). The amplified reference gene RT-PCR fragment was purified from agarose gel and standard was calculated based on purified fragment concentration and molecular weight. The reference gene was subjected to a 10-fold serial dilution (1.00E+1 to 1.00E-10) with DEPC. For absolute quantification analysis, five standard dilutions from 1.00E-5 to 1.00E-9 for standard curve were used for quantifying both target and reference gene expression in cDNA samples.

4.7. Screening of Putative Transgenic Plants and Identification of CRISPR-Cas9 induced Mutations

Twenty seeds from 10 individual T0 *BBI* events were planted in the Sears Plant greenhouse in 2-gallon pots containing Pro-Mix® HP Mycorrhizae™ (Premier Tech, Quakertown, PA). The plants were fertilized with Osmocote Plus (Scotts) according to the manufacturer's recommendations. Greenhouse settings were 16-h daylength and 30° C/18° C day/night temperatures. For every plant which germinated, fresh tissue was collected, and genomic DNA isolated from ~20-30 mg lyophilized leaf tissue using a Maxwell Promega robot with the Maxwell RSC PureFood GMO and Authentication Kit (Promega).

PCR primer pairs were designed to bear at least two gene-specific SNP differences as compared to homeologous genes. Primers used for PCR amplification and sequencing are detailed in supplementary Table S1. We blasted all primers against the unmasked *Glycine max* 'Williams 82' genomic sequence (E-value cut off = 10.0) to ensure primer specificity (www.soybase.org). PCR amplification was performed using Titanium taq according to manufacturer's recommendations (Takara) in a PTC-200 thermocycler (MJ Research/Bio-Rad), using these conditions: 95°C for an initial 5 minute denaturation, followed by 40 cycles of 95°C for 30 seconds, followed by 60°C for 30 seconds, and an extension step at 72°C for 1 minute/kilobase of target sequence. PCR products were separated on a 1% agarose gel and compared to DNA ladder size standards to ensure appropriate size. PCR products were purified through use of a QIAquick PCR purification kit (Qiagen). After PCR purification, products were Sanger sequenced at the DNA Core Facility of the University of Missouri-Columbia.

PCR product sequencing traces were imported into Geneious Primer software, version 11.0.11+9 (Invitrogen), trimmed, aligned to gene sequences from the G275_W82.a2.v1 assembly (<http://www.phytozome.net/soybean>) using the Geneious Primer software, version 11.0.11+9 (Invitrogen). Disagreements were manually assessed in comparison to the G275_W82.a2.v1 assembly

[40]. Putative indels and polymorphisms were verified by a minimum of two independent PCR reactions.

4.8. Immunoblot Analysis

Total seed proteins isolated from wild-type soybean cultivar 'Maverick' and T1:2 seeds of BBi mutated CRISPR/CAS9 lines were resolved on a 13.5% SDS-PAGE gels. Resolved proteins were transferred to nitrocellulose membranes and incubated with TBS (10 mM Tris-HCl, pH 7.5, 500 mM NaCl) containing 5% non-fat dry milk for 1 h at room temperature. Following this step, the nitrocellulose membranes were washed 3X with TBST (15 min each) and incubated over-night with 1:10,000 diluted antibodies raised either against purified soybean Kunitz trypsin or Bowman-Birk protease inhibitors. 7S β -conglycinin and 11S glycinin specific antibodies were used at 1:50,000 dilution. Specifically bound primary antibodies were detected using anti-rabbit IgG-horseradish peroxidase conjugates and a SuperSignal West Pico kit (Pierce, Rockford, IL, U.S.A.) as previously described [21].

4.9. Trypsin and Chymotrypsin Inhibitor Assay

Trypsin inhibitor activity in different BBi mutated CRISPR/CAS9 lines was measured following the procedure described by [41]. Briefly, to 20 mg of dry soybean seed powder was placed in a 2 mL Eppendorf centrifuge. To this 1 mL of 10 mM NaOH solution was added and vigorously agitated on a vortex mixer for 10 min at room temperature. The resulting slurry was clarified by centrifugation at 16,000xg for 10 min. Clear supernatant was utilized for measuring KTi activity. Trypsin inhibitor activity was expressed as trypsin units inhibited (TUI) per mg of sample, and results were expressed as the mean \pm SD from three biological replicates.

Chymotrypsin inhibitor activity in soybean seed extracts were measured N-Succinyl-Ala-Ala-Pro-Phe p-nitroanilide (AAPF) as the substrate. Soybean seed extract, which resulted in 40 to 60% chymotrypsin inhibition range, was added to a 1.5 mL Eppendorf tube containing 900 μ L of assay buffer (100 mM-Tris-HCL, pH 8.0), 8 μ L of α -chymotrypsin (Sigma-Aldrich Company) dissolved in 1 mM HCl solution (0.1 mg/mL) and seed extract (8 to 10 μ L) and incubated at 37 C for 5 min. Following this step, 80 μ L of AAPF (1 mg/mL) was added and incubated for additional 10 min at 37 C. The assay was stopped by adding 500 μ L of 30% acetic acid. Absorbance at 410 nm was measured, and chymotrypsin inhibitor units were calculated as the amount of inhibitor that reduced the absorbance at 410 nm by 0.1 optical density. The results are expressed as the mean \pm SD from three biological replicates. Trypsin and Chymotrypsin inhibitor assay data was visualized and compared using JMP statistical software 16.0 (SAS Institute Inc., Cary, NC, USA). One-way ANOVA was performed and significant differences ($\alpha=0.05$) between means were determined using the Tukey-Kramer HSD test.

Supplementary Materials: The following supporting information can be downloaded at: www.mdpi.com/xxx/s1, Figure S1. Gene expression of *BBi* genes in soybean. Figure S2. CRISPR-Cas9 cassettes targeting *BBi* genes. Figure S3. Sequencing traces for Glyma.16g208900 and Glyma.14G117700. **Table S1.** Primers used for sequencing analysis and RT-PCR.

Author Contributions: HBK, WSK and JDG conceptualized the original project. Experimental work was performed by WSK, SK, JDG, JL, MRJ and HBK. RMS and JL contributed new reagents and CRISPR/Cas9 constructs. The original manuscript was written by HBK and JDG. All authors were involved in reviewing and editing the manuscript. The project administration was led by HBK. All authors have read and agreed to the published version of the manuscript.

Funding: Funding for this research was provided by United Soybean Board (2020-152-0114 and 2335-203-0101) and Agricultural Research Service, USDA (8042-21220-234-00D).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data supporting the findings of this study are provided in the manuscript and supplementary information.

Acknowledgments: The authors would like to thank Nathan Oehrle for his excellent technical assistance. Mention of a trademark, vendor, or proprietary product does not constitute a guarantee or warranty of the product by the USDA and does not imply its approval to the exclusion of other products or vendors that may also be suitable. The US Department of Agriculture, Agricultural Research Service, Midwest Area, is an equal opportunity, affirmative action employer and all agency services are available without discrimination.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Bullock, D.G. Crop rotation. *Critical Reviews in Plant Sciences* **1992**, *11*, 309-326.
- Mohammad Sohidel, I.; Imam, M.; Md. Rafiqul, I.; Md. Kamrul, H.; Hafeez, A.S.M.G.; Md. Moaz, H.; Hirofumi, S.; Akihiro, U.; Liyun, L.; Misbah, N., et al. Soybean and Sustainable Agriculture for Food Security. In *Soybean*, Takuji, O., Yoshihiko, T., Norikuni, O., Takashi, S., Sayuri, T., Eds. IntechOpen: Rijeka, 2022; 10.5772/intechopen.104129p. Ch. 1.
- Ruiz, N.; Parsons, C.; Stein, H.; Coon, C.; Eys, J.; Miles, R. A review: 100 years of soybean meal. *ADM: Chicago, IL, USA* **2020**.
- Vorster, J.; van der Westhuizen, W.; du Plessis, G.; Marais, D.; Sparvoli, F.; Cominelli, E.; Camilli, E.; Ferrari, M.; Le Donne, C.; Marconi, S., et al. In order to lower the antinutritional activity of serine protease inhibitors, we need to understand their role in seed development. *Front Plant Sci* **2023**, *14*, 1252223.
- Kunitz, M. CRYSTALLINE SOYBEAN TRYPSIN INHIBITOR : II. GENERAL PROPERTIES. *Journal of General Physiology* **1947**, *30*, 291-310, doi:10.1085/jgp.30.4.291.
- Bowman, D.E. Fractions derived from soy beans and navy beans which retard tryptic digestion of casein. *Proceedings of the Society for Experimental Biology and Medicine* **1944**, *57*, 139-140.
- DiPietro, C.M.; Liener, I.E. Heat inactivation of the Kunitz and Bowman-Birk soybean protease inhibitors. *Journal of Agricultural and Food Chemistry* **1989**, *37*, 39-44.
- Jofuku, K.D.; Schipper, R.D.; Goldberg, R.B. A frameshift mutation prevents Kunitz trypsin inhibitor mRNA accumulation in soybean embryos. *The Plant Cell* **1989**, *1*, 427-435.
- Krishnan, H.B. Characterization of a soybean [*Glycine max* (L.) Merr.] mutant with reduced levels of Kunitz trypsin inhibitor. *Plant Science* **2001**, *160*, 979-986.
- Orf, J.; Hymowitz, T. Inheritance of the Absence of the Kunitz Trypsin Inhibitor in Seed Protein of Soybeans 1. *Crop Science* **1979**, *19*, 107-109.
- Palacios, M.; Easter, R.; Soltwedel, K.; Parsons, C.; Douglas, M.; Hymowitz, T.; Pettigrew, J. Effect of soybean variety and processing on growth performance of young chicks and pigs. *Journal of animal science* **2004**, *82*, 1108-1114.
- Bernard, R.L.; Hymowitz, T.; Cremeens, C.R. Registration of 'Kunitz' Soybean. *Crop Science* **1991**, *31*, 232-233.
- Gillman, J.D.; Kim, W.-S.; Krishnan, H.B. Identification of a new soybean kunitz trypsin inhibitor mutation and its effect on bowman- birk protease inhibitor content in soybean seed. *Journal of agricultural and food chemistry* **2015**, *63*, 1352-1359.
- Kollipara, K.P.; Singh, R.J.; Hymowitz, T. Inheritance of Protease Inhibitors in *Glycine tomentella* Hayata (2n = 38), a Perennial Relative of Soybean. *Journal of Heredity* **1996**, *87*, 461-463, doi:10.1093/oxfordjournals.jhered.a023038.
- Livingstone, D.; Beilinson, V.; Kalyaeva, M.; Schmidt, M.A.; Herman, E.M.; Nielsen, N.C. Reduction of protease inhibitor activity by expression of a mutant Bowman-Birk gene in soybean seed. *Plant Mol Biol* **2007**, *64*, 397-408, doi:10.1007/s11103-007-9163-x.
- Cong, L.; Ran, F.A.; Cox, D.; Lin, S.; Barretto, R.; Habib, N.; Hsu, P.D.; Wu, X.; Jiang, W.; Marraffini, L.A. Multiplex genome engineering using CRISPR/Cas systems. *Science* **2013**, *339*, 819-823.
- Mali, P.; Yang, L.; Esvelt, K.M.; Aach, J.; Guell, M.; DiCarlo, J.E.; Norville, J.E.; Church, G.M. RNA-guided human genome engineering via Cas9. *Science* **2013**, *339*, 823-826.
- Liu, J.; Gunapati, S.; Mihelich, N.T.; Stec, A.O.; Michno, J.M.; Stupar, R.M. Genome Editing in Soybean with CRISPR/Cas9. *Methods Mol Biol* **2019**, *1917*, 217-234, doi:10.1007/978-1-4939-8991-1_16.
- Sun, X.; Hu, Z.; Chen, R.; Jiang, Q.; Song, G.; Zhang, H.; Xi, Y. Targeted mutagenesis in soybean using the CRISPR-Cas9 system. *Scientific reports* **2015**, *5*, 1-10.
- Severin, A.J.; Woody, J.L.; Bolon, Y.-T.; Joseph, B.; Diers, B.W.; Farmer, A.D.; Muehlbauer, G.J.; Nelson, R.T.; Grant, D.; Specht, J.E., et al. RNA-Seq Atlas of *Glycine max*: A guide to the soybean transcriptome. *BMC Plant Biology* **2010**, *10*, 160.

21. Krishnan, H.B.; Kim, S.; Pereira, A.E.; Jurkevich, A.; Hibbard, B.E. Adenantha pavonina, a potential plant-based protein resource: Seed protein composition and immunohistochemical localization of trypsin inhibitors. *Food Chem X* **2022**, *13*, 100253.
22. Krishnan, H.B.; Oehrle, N.W.; Natarajan, S.S. A rapid and simple procedure for the depletion of abundant storage proteins from legume seeds to advance proteome analysis: a case study using Glycine max. *Proteomics* **2009**, *9*, 3174-3188.
23. Dipietro, C.; Liener, I. Soybean protease inhibitors in foods. *Journal of Food Science* **1989**, *54*, 606-609.
24. He, H.; Li, X.; Kong, X.; Hua, Y.; Chen, Y. Heat-induced inactivation mechanism of soybean Bowman-Birk inhibitors. *Food chemistry* **2017**, *232*, 712-720.
25. Livingstone, D.; Beilinson, V.; Kalyaeva, M.; Schmidt, M.A.; Herman, E.M.; Nielsen, N.C. Reduction of protease inhibitor activity by expression of a mutant Bowman-Birk gene in soybean seed. *Plant Molecular Biology* **2007**, *64*, 397-408.
26. Sexton, P.; Naeve, S.; Paek, N.; Shibles, R. Sulfur availability, cotyledon nitrogen: sulfur ratio, and relative abundance of seed storage proteins of soybean. *Crop science* **1998**, *38*, 983-986.
27. Paek, N.C.; Imsande, J.; Shoemaker, R.C.; Shibles, R. Nutritional control of soybean seed storage protein. *Crop Science* **1997**, *37*, 498-503.
28. Holowach, L.P.; Thompson, J.F.; Madison, J.T. Storage protein composition of soybean cotyledons grown in vitro in media of various sulfate concentrations in the presence and absence of exogenous L-methionine. *Plant Physiology* **1984**, *74*, 584-589.
29. Gayler, K.R.; Sykes, G.E. Effects of nutritional stress on the storage proteins of soybeans. *Plant Physiology* **1985**, *78*, 582-585.
30. Fujiwara, T.; Hirai, M.Y.; Chino, M.; Komeda, Y.; Naito, S. Effects of sulfur nutrition on expression of the soybean seed storage protein genes in transgenic petunia. *Plant Physiology* **1992**, *99*, 263-268.
31. Clarke, E.; Wiseman, J. Developments in plant breeding for improved nutritional quality of soya beans I. Protein and amino acid content. *The Journal of Agricultural Science* **2000**, *134*, 111-124.
32. BIRK, Y. The Bowman-Birk inhibitor. Trypsin-and chymotrypsin-inhibitor from soybeans. *International journal of peptide and protein research* **1985**, *25*, 113-131.
33. Kim, W.-S.; Sun-Hyung, J.; Oehrle, N.W.; Jez, J.M.; Krishnan, H.B. Overexpression of ATP sulfurylase improves the sulfur amino acid content, enhances the accumulation of Bowman-Birk protease inhibitor and suppresses the accumulation of the β -subunit of β -conglycinin in soybean seeds. *Scientific reports* **2020**, *10*, 1-13.
34. Orf, J.H. Modifying soybean composition by plant breeding. In Proceedings of Soybean Utilization Alternatives Conference, University of Minnesota.
35. Čermák, T.; Curtin, S.J.; Gil-Humanes, J.; Čegan, R.; Kono, T.J.Y.; Konečná, E.; Belanto, J.J.; Starker, C.G.; Mathre, J.W.; Greenstein, R.L., et al. A Multipurpose Toolkit to Enable Advanced Genome Engineering in Plants. *Plant Cell* **2017**, *29*, 1196-1217.
36. Sleper, D.A.; Nickell, C.D.; Noel, G.R.; Cary, T.R.; Thomas, D.J.; Clark, K.M.; Rao Arelli, A.P. Registration of 'Maverick' Soybean. *Crop Science* **1998**, *38*, 549-550.
37. Zeng, P.; Vadnais, D.; Zhang, Z.; Polacco, J. Refined glufosinate selection in Agrobacterium-mediated transformation of soybean [Glycine max (L.) Merrill]. *Plant cell reports* **2004**, *22*, 478-482.
38. Paz, M.M.; Martinez, J.C.; Kalvig, A.B.; Fonger, T.M.; Wang, K. Improved cotyledonary node method using an alternative explant derived from mature seed for efficient Agrobacterium-mediated soybean transformation. *Plant cell reports* **2006**, *25*, 206-213.
39. Song, B.; Oehrle, N.W.; Liu, S.; Krishnan, H.B. Development and Characterization of a Soybean Experimental Line Lacking the α' Subunit of β -Conglycinin and G1, G2, and G4 Glycinin. *J Agric Food Chem* **2018**, *66*, 432-439, doi:10.1021/acs.jafc.7b05011.
40. Schmutz, J.; Cannon, S.B.; Schlueter, J.; Ma, J.; Mitros, T.; Nelson, W. Genome sequence of the palaeopolyploid soybean. *Nature* **2010**, *463*, 178-183.
41. Kim, S.; Krishnan, H.B. Chapter Seven - A fast and cost-effective procedure for reliable measurement of trypsin inhibitor activity in soy and soy products. In *Methods in Enzymology*, Jez, J., Ed. Academic Press: 2023; Vol. 680, pp. 195-213.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.