

Article

Not peer-reviewed version

CO₂-Breathing Prior to Simulated Diving Increases Decompression Sickness Risk in a Mouse Model: The Microbiota Trail Is Not Forgotten

Lucille Daubresse , Aurélie Portas , [Alexandrine Bertaud](#) , Marion Marlinge , Sandrine Gaillard , Jean-Jacques Risso , Céline Ramdani , Jean-Claude Rostain , Nabil Adjirou , Anne-Virginie Desruelle , Jean-Eric Blatteau , [Régis Guieu](#) * , [Nicolas Vallée](#) *

Posted Date: 21 June 2024

doi: 10.20944/preprints202406.1553.v1

Keywords: Bubble formation; Dysbiosis; Auto-immune; Auto-inflammation; Akkermansia; Rhizobium; Gut-Brain axis; Hypercapnia



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Article

CO₂-Breathing Prior to Simulated Diving Increases Decompression Sickness Risk in a Mouse Model: The Microbiota Trail Is Not Forgotten

Running Title: Impact of CO₂ on Experimental DCS

Lucille Daubresse ¹, Aurélie Portas ², Alexandrine Bertaud ³, Marion Marlinge ³, Sandrine Gaillard ², Jean-Jacques Risso ⁴, Céline Ramdani ⁴, Jean-Claude Rostain ³, Nabil Adjirou ³, Anne-Virginie Desruelle ⁴, Jean-Eric Blatteau ^{1,†}, Régis Guieu ^{3,†} and Nicolas Vallee ^{4,*,†}

¹ Hôpital d'instruction des Armées, Service de Médecine hyperbare, 83000 Toulon, France

² Université de Toulon, La Garde, France

³ Center for CardioVascular and Nutrition Research (C2VN), Aix-Marseille Univ, INSERM, INRAE, Faculté de Médecine, 27 Boulevard Jean-Moulin, 13005 Marseille, France

⁴ Military Institute of Biomedical Research (IRBA), Subaquatic Operational Research Team (ERRSO), 83000 Toulon, France

* Correspondence: author: nicolas.vallee5@hotmail.fr

† The last 3 authors contributed equally to this work and share last authorship.

Abstract: Decompression sickness (DCS) with neurological disorders is the leading cause of diving accidents. Exposure to high levels of CO₂ during diving is a safety concern for occupational , groups at risk of DCS. However, the effects of prior exposure to CO₂ have never been evaluated. The purpose of this study was to evaluate the effect of CO₂ breathing prior to a provocative dive on the occurrence of DCS in mice. 50 mice were exposed to a maximum CO₂ concentration of 70 hPa for one hour at atmospheric pressure. Another 50 mice breathing air under similar conditions served as controls. In the AIR group (controls), 22 out of 50 mice showed post-dive symptoms compared to 44 out of 50 in the CO₂ group. We found that CO₂ breathing is associated with a decrease in body temperature in mice and that CO₂ exposure dramatically increases the incidence of DCS. More unexpectedly, it appears that a lower temperature of the animals even before exposure to the accident-prone protocol leads to an unfavorable prognosis. This study also suggests that the composition of the microbiota may influence thermogenesis and thus accidentology. Depending on prior exposure, some bacteria identified in this work could be perceived as protective or pathogenic.

Keywords: bubble formation; dysbiosis; auto-immune; auto-inflammation; Akkermansia; Rhizobium; gut-brain axis; hypercapnia

1. Introduction

Exposure to high levels of CO₂ is a growing safety concern for professional divers at risk of decompression sickness [1–3]. In fact, pre-dive exposure to CO₂ is suspected in a recent series of unusual decompression sickness (DCS) cases with neurological symptoms [4]. Although there is a tendency for hypercapnia during diving [2] and an increase in osteo-articular accidents among hyperbaric workers breathing a CO₂ enriched gas mixture (1.8 to 2.3%) [5]; There is a paucity of information on the effects of CO₂ inhalation prior to hyperbaric exposure and its incidence in DCS.

At atmospheric pressure, the increase in blood CO₂ levels, endogenous or exogenous, defines hypercapnia, which in humans corresponds to an arterial CO₂ partial pressure (PaCO₂) greater than 45 mmHg (60 hPa). This causes a very small decrease in plasma pH, which limits the association of oxygen with hemoglobin at the pulmonary level and promotes its dissociation at the systemic level.

In response, the body increases its respiratory rate and essentially mobilizes its bicarbonate buffering systems according to Le Chatelier's principle [6,7] ($\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3 \rightleftharpoons \text{HCO}_3^- + \text{H}^+$) to eliminate excess CO_2 and keep the blood pH stable.

According to the classical scheme, DCS results from circulating bubbles in the bloodstream [8]. The tissue gas load in the body that occurs during compressed air ventilation during diving is proportional to the duration and depth of the dive. Bubble formation occurs during desaturation, *i.e.* ascent to the surface to sea level, when gas elimination by perfusion and ventilation is too inefficient. In the bloodstream, the presence of bubbles causes cell destruction by increasing shear forces [9] but also platelet activation, possibly due to direct bubble/platelet contact [10–12] ultimately causing inflammation, vasoconstriction, circulatory stasis, and interstitial edema, leading to tissue death [13]. Disorders of the spinal cord and brain are the most common and serious symptoms. They can lead to paralysis, with a risk of long-term neurological sequelae of about 20 to 30 percent [14]. Decompression sickness with neurological disorders is the leading cause of diving accidents and is responsible for increased morbidity.

Circulating bubbles are essentially composed of inspired gases, such as nitrogen [15] but also metabolized gases. A porcine model of a desaturation accident specifies that the composition of the bubbles includes 5 to 9% CO_2 , 1 to 4% O_2 , while the remainder of the bubble is inert gas.[15]. A Japanese study conducted on rabbits highlights that the CO_2 level in the bubble seems to decrease first and then gradually increase, while the O_2 level evolves in the opposite way [16]. The concentration of CO_2 in the bubble and its evolution with time would play an important role in the growth and formation of bubbles during decompression [17,18]. We are not currently aware of any studies that have examined the effect of adding CO_2 on the incidence of DCS. The suggestion [19] that CO_2 retention increases the likelihood of DCS is based on the assumption that tissue gas uptake is enhanced by elevated pCO_2 (CO_2 partial pressure) during deep work and is proportionally slower at normal pCO_2 during decompression. In fact, such CO_2 toxicity is often associated with oxygen toxicity of the nervous system [20–22] and possibly blood pH [18]. Because CO_2 is a highly diffusible gas, we hypothesize that its increase in blood concentration could significantly contribute to increased bubble growth and therefore DCS.

Inflammation, or even the autoimmune reaction [23,24], is often transcribed in articles relating to DCS [25–28]. Differences have thus been observed in animals all from the same strain [26] or in specially selected strains [23]. This naturally led the teams to focus on the predispositions that the host microbiota can induce [23,28–31]. The microbiota is indeed known for its interactions with the host, for example by modulating immune responses [23,25,32–35] or/and thermoregulation [36–40]. Microbiota is sensitive to temperature fluctuation [37] and can confer host plasticity in thermoregulation [33,36]. As example, disruption of the equilibrium of the gut microbiota by cold and high-fat diet can compromise the integrity of the intestinal barrier that induces increased permeability, facilitating the translocation of harmful substances and bacteria into the bloodstream and ultimately triggering systemic inflammation [41].

On the other hand, increases in CO_2 are also known to modulate the microbiota [42] as well as the inflammatory response [42,43] and thermoregulation [17,42,44]. Thermogenesis is the cornerstone of metabolism [45]. Thermoregulation is closely related to the metabolic function and health of the host [32]. Hypercapnia elicits hypothermia in a great number of vertebrates, including mammals [46]. This mechanism would include peripheral and central vasomotion through NO activation, and also hyperventilation which would lead to an increase in pulmonary perspiration among other things, and therefore to cooling of the body, especially during exercise [17,47,48].

This work is the first to clearly establish the influence of exposure to a high concentration of CO_2 on the risk of DCS. This study was performed in a mouse model with clinical and biological evaluation, including thermal measurements and characterization of the native fecal microbiota. The hyperbaric protocol was preceded by exposure of the mice to a maximum CO_2 concentration of 70 hPa, *i.e.* 7% at atmospheric pressure, for one hour. This CO_2 value corresponds to the maximum concentration that does not cause clinical or social effects in the animals [49–51].

2. Materials and Methods

The experimental design can be followed in Figure 1. On the day of the experiment, the temperature of each mouse was measured. Mice were then immediately identified, weighed, and their feces collected for microbiota analysis. Next, the mice were exposed to either CO2 or air. After one hour of exposure, the cages were transferred to the hyperbaric chamber. After the accident-prone protocol, a clinical examination supplemented by temperature measurement was performed. Blood punctures were then performed under anesthesia for biochemical analysis. Details follow.




Figure 1. Flow Chart of Experimental Design - Measurement of Temperature, Identification, Weighing, and Feces Collection for Microbiota Analysis in Mice.

2.1. Ethic Statements

All procedures involving animals comply with European (Directive 2010/63/EU) and French (Decree 2013/118) legislation. The protocol was approved by the Animal Ethics Committee (CE14, University of Aix-Marseille) on May 6, 2022 under number APAFIS#34720-2022011812031043. The population consisted of 100 6-week-old male C57Bl/6n mice (Charles River strain, The Jackson Laboratory, Italy). Animals were housed 5 per cage (as delivered) in an accredited animal facility with controlled temperature (22 ± 1°C) and day/night cycle (12 hours of light per day, 7:00 am - 7:00 pm), with ad libitum access to water and food (A03, UAR).

2.2. Animal Batches

The animals remained grouped in batches of 5, as they were when delivered from the breeder, to maintain social grouping. Each batch of 5 mice was assigned to one of two groups. The first group consisted of 50 mice exposed to a CO₂ enriched atmosphere (CO₂), while the control group (AIR) consisted of 50 mice breathing air in a similar condition. After one hour of exposure, the animal cages were transferred to the hyperbaric chamber.

2.3. Evaluation of Superficial Temperature by Infrared Imaging

The surface temperatures of the mice were measured twice by infrared thermography (camera: model Flir E52, Sweden), at the very beginning of the experiment and before anesthesia. Infrared videos were taken in a room at 23°C while the mice were moving quietly in their cages. More specifically, the animals were filmed in their original cage before being weighed and tagged, then transferred to a temporary cage, and then all returned to the original cage. The same strategy was used before anesthesia. The images were then analyzed to evaluate the averaged temperature using the processing software of the camera manufacturer (Flir ResearchIR), according to our previous study [26].

2.4. CO₂ Exposition

A constant flow of CO₂, controlled by an Arduino system designed by our engineer, perfused the polycarbonate cage to maintain a 7% CO₂ atmosphere for 1 hour. The same cage perfused with air served as a control. The animals were then immediately transferred to the hyperbaric chamber.

2.5. Hyperbaric Exposure

Batches of 10 mice (5 per cage and 5 per group) from the AIR and CO₂ pools were subjected to the hyperbaric protocol in an 80-liter tank equipped with two observation ports. Mice were allowed to move freely within their cages.

The air compression protocol involved two ramps of pressure increase, first at 0.1 atm/min to a pressure of 2 atm, followed by 1 atm/min to 10 atm. 10 atm (equivalent to a depth of 90 msw) is the pressure at which animals were maintained for 45 minutes prior to decompression. The decompression rate was 1 atm/s to 1 atm.

In the hyperbaric chamber, CO₂ and water vapor produced by mice were trapped with soda lime (<300 ppm) and seccagel (relative humidity: 40-60%). The gases were mixed by an electric fan. The day-night cycle was maintained throughout.

2.6. Physical Examination

After exiting the hyperbaric chamber, a 30-minute period was devoted to the identification of clinical signs of a desaturation accident, with the addition of grip tests at 15 and 30 minutes [26,28,52,53]. Manifestations of DCS include itching, respiratory dysfunction (gasping), neurological symptoms (tremors, convulsions, motor and/or sensory deficits including claudication and paralysis), marked asthenia (decrease of gripping duration, prostrate mouse without exploratory posture), and in the most severe form, it may result in death of the animal [26,28,52,53].

2.7. Anaesthesia and Biopsy

After physical examination, animals were first anesthetized by administration of gaseous isoflurane (4.0% in oxygen flow at 2.0 l/min) in an individual cage for 2 min and then through a face mask (anesthesia box: Harvard Apparatus, Les Ulis, France). Intraperitoneal injection of ketamine (100 mg.kg⁻¹) and xylazine (10 mg.kg⁻¹) completed the anesthesia. The degree of anesthesia was determined by testing the absence of withdrawal reflexes in response to pinching of the distal hind limbs.

Immediately after anesthesia, blood samples were collected by intracardiac puncture for assays of glial-derived peptide S100 (PS100), copeptin, neuron-specific enolase (NSE), bicarbonate, and interleukin-6 (IL-6). Blood was collected in a sterile 1 ml syringe containing lithium heparin, immediately placed on ice, and then processed according to the manufacturer's instructions. Finally, animals were euthanized with a lethal dose of anesthetic.

2.8. Blood Biochemical Analysis

Plasma was diluted volume-to-volume to obtain sufficient sample volume for all biochemical analyses. All analyses were performed in duplicate. The coefficient of variation tolerated by the analyzers must be less than 10% for the measurement to be considered valid.

Glial-derived peptide S100 (PS100) analysis was performed by electrochemiluminescence on the Cobas e411 analyzer (Roche®, Meylan, France; detection limit 5 pg/mL). 20 µL of plasma was incubated with 2 different antibodies specifically directed against PS100: a monoclonal and biotinylated antibody and a second antibody that binds to ruthenium to form a sandwich complex. Magnetic microparticles coated with streptavidin were added to bind the immune complex through biotin and to keep it in the measuring cell despite a magnetic field washing step. A potential difference was then applied to induce luminescence due to excitation of ruthenium (quantified by a photomultiplier), this signal being proportional to the concentration of PS100 present in the sample.

Copeptin (50 µL per assay/detection limit 0.88 pmol/L) and NSE (70 µL per assay/detection limit 4 ng/mL) were analyzed on a Kryptor Gold instrument (Thermoscientific®) using TRACE® (Time-Resolved Amplified Cryptate Emission) technology. Briefly, the analyzer measures the fluorescence at 620 and 665 nm emitted by an immune complex-containing donor/acceptor system (and europium cryptate and cyanine, respectively) after an excitation step at 337 nm.

Bicarbonate (HCO_3^-) (5.7 µL per assay / detection limit 10 µmol/L) and IL-6 (50 µL per assay / detection limit 2.7 pg/mL) were determined on Atellica Solution (Siemens®), on the CH930 module for the former (colorimetric enzymatic method based on 2 chemical reactions catalyzed by phosphoenolpyruvate carboxylase PEPC and malate dehydrogenase MDH, (the reduction in absorbance is proportional to the concentration of the analyte) and the IM module (chemiluminescence using a mouse monoclonal antibody labeled with an acridinium ester molecule and monoclonal antibody coated magnetic microparticles, both directed against IL-6) for the latter.

2.9. Stools Bacterial Content: 16s rDNA Extraction and Sequencing

While the mouse clung to the grid during marking, feces falling onto a new surgical field were collected with disinfected tweezers, placed in a 1.5 ml DNA-free Eppendorf tube, and stored at -80°C until genetic analysis. DNA for metagenomic identification of microbiota was extracted from stool using the Maxwell RSC Fecal Microbiome DNA Kit (Maxwell RSC, Promega, France) according to the supplier's recommendations to obtain a 100 µl eluate. The concentration of the extract was adjusted to 20 ng/µl in ultrapure water prior to amplification on the Azenta Genewiz platform (France). The 16S-EZ workflow involves proprietary multiplexed PCR amplification of the V3 and V4 hypervariable regions of the 16S rDNA gene. A limited cycle 2nd round PCR adds sample-specific barcodes to each sample to allow multiplexing of multiple samples on the same sequencing run. Final libraries are pooled and undergo final quality control prior to sequencing. Sequencing is performed using Illumina chemistry in the 2x250bp paired-end configuration to ensure coverage of all V3 and V4 sequences. Subsequent sequencing data undergo quality control prior to 16S-EZ bioinformatics analysis.

2.10. Bioinformatic Processing

Initial quality control of reads in 'fastq' files was performed using FastQC [54]. Forward and reverse primers were trimmed from the paired-end reads. Sequences were quality filtered, denoised, and merged into amplicon sequence variants (ASVs) using the DADA2 [55] within R [56] (v. 4.1.0). Briefly, sequences were filtered based on visual assessment of error rates according to the quality

score ≥ 25 and after chimeras were detected and removed using the 'removeBimeraDenovo' script in DADA2. Each representative 16s rDNA ASV sequence was associated with Blast+ (v. 2.10) using FROGS v.4.0.1 (Escudié et al., 2018) and the naive Bayesian classifier method [57] against SILVA 138 database [58]. Based on the initial assignment, we organized our taxonomic profile to retain the taxonomy of ASVs with greater than 95% identity and 75% sequence coverage. Below these thresholds, if an ASV had a bootstrap confidence score $<65\%$ at any level, it was binned as unknown. More specifically, if the bootstrap confidence score associated with the ASV was less than 65% at any level of classification (e.g. genus, family, etc.), then that ASV was labeled as belonging to an unknown species per example. After processing, ASV with a frequency lower than 0.005% were removed [59]. The raw sequence data generated in this study have been deposited in the NCBI Sequence Read Archive under the accession number BioProject PRJNA1070862.

2.11. Statistical Analyses

Prior to analysis, normality was tested and data normalization (n) was applied when necessary.

The association between qualitative (gas exposure, clinical status) and quantitative variables (weight, temperature, bacterial species abundances...) was studied using Partial Least Square Regression - Discriminant Analysis (PLS-DA). It is based on the Partial Least Squares method and allows to handle multicollinear data, missing values and data sets with few observations and many variables. PLS-DA was set with press minimum as the stop criterion, and the jackknife resampling technique was selected for cross-validation. Variable importance for projection and discriminative effects of gas or clinical status were evaluated.

Kruskal-Wallis tests supplemented with Dunn's post-hoc (with Bonferroni correction) were performed for qualitative or non-parametric data.

The software used was XLStat Biomed from Addinsoft. The maximum alpha level accepted is 5%.

3. Results

3.1. Clinical Status of the Animals Following the Dive

Initial results confirm that the protocol was accident-prone as expected.

Clinical observation after the simulated dive was the main criterion for defining DCS. During the thirty-minute observation period following exit from the hyperbaric chamber, 66% (out of 100 mice) of the total animals showed symptoms of decompression sickness, including convulsions and paralysis in the most severe cases. Two mice died, one per group. There was a significant difference (KW; $p < 0.001$) between the group of mice that breathed air before the simulated dive and the group that breathed 21% O₂ and 7% CO₂ (Figure 2). In the AIR group (control), 22 out of 50 mice showed post-dive symptoms compared to 44 out of 50 in the CO₂ group.




Figure 2. Decompression sickness occurrences 30 minutes after pathogenic decompression as a function of previous gas exposure (AIR or CO₂). **A, Histogram:** Proportion of mice with DCS

symptoms. The DCS status was attributed when the mice presented neurological signs in the form of paresis or paralysis of at least one limb, convulsions and/or reduced performance in the motor test for forelimbs, and also respiratory troubles, even death. **B, Radar Chart:** percentage of mice displaying a type of symptom in a population, considering that a mouse can present the whole panel of symptoms. * denotes $p < 0.050$, ** denotes $p < 0.010$, *** denotes $p < 0.001$.

Knowing that weight can influence accidents, we tested it: no significant difference between the groups was found (KW; $p=0.239$).

3.2. General Description of the Data

A PLS-DA was conducted to model the interaction between the 42 variables studied (see Figure 4 for the complete list), with clinical assessment as an additional qualitative variable. Previous exposure to CO₂ or its control in air was integrated as the only explanatory qualitative variable. The model showed a good fit, with R²X approaching 1 as the number of components increased. R²Y, while slightly slower, also demonstrated a good fit. However, the model had limited predictive ability, indicated by a low Q² value. However, the Receiver Operator Characteristic (ROC) curve analysis shows an area under the curve (AUC) of 0.879 with specificity and sensitivity values of 79% and 80%, respectively, resulting in an accuracy of 80%.

Projection (Figure 3 left) onto components 1 and 2 explains 35.9% of the model, with component 1 alone expected to explain 31%. The confidence ellipses (Fisher test, 95% CI) surrounding the observations of symptomatic and asymptomatic animals on the PLS-DA indicate a slight discrimination of these groups on the T1 axis. The confidence ellipses surrounding the animals according to the inhaled gas have a profile very similar to that of the clinic.




Figure 3. Partial Least Square – Discriminant Analysis (PLS-DA) plot conducted on clinic (Symptomatic or Asymptomatic, blue and red gradient line) as dependant variable) and as explanatory variable gas exposure before the hyperbaric protocol (CO₂ green or Air, brown), micro-organisms genus detected in stools (red), mice superficial temperature before and after the protocol (black), plasma analysis performed at the end of the protocol (orange). (Right) Correlation map based from observations (green dots) according to their qualitative or quantitative explanatory characteristics. (Left) Confusion matrix summarize the observations according to the clinic status of mice (Symptomatic in blue or Asymptomatic in red), according to the gas breathed before the dive (Air in brown or CO₂ in green), with their confidence ellipse (Fisher, Confidence Interval 95%).

The correlations (Figure 3 right) between the components and the variables are generally good, with a considerable number of variables projected onto the disk between 0.8 and 1, including, for example, the nature of the gas, which is essentially projected onto axis 1, while a number of bacteria are projected onto the t2 axis. In general, the correlations are weak in the center. For example, we can see the weight of the animals close to the center. In this particular case, we had previously selected animals of low weight to overcome this well-known problem in diving, where high weight increases accidents.

Looking specifically at the correlations and the projection of the clinical axis (Sympto/Asympto, green line), we note with interest a positive correlation between an asymptomatic state and prior exposure to air (as a control), as well as a higher temperature of the mice upon exiting the hyperbaric chamber. This supports the diagnosis. To a lesser extent, and because it is clearly on the Sympto/Asympto axis, the PS100 marker could be considered. It may also appear that a higher temperature of the mice before the start of the experiment is positively correlated with being asymptomatic. The opposite would be true in the case of prior exposure to CO₂, which would be clinically deleterious and would be associated with an increase in bicarbonate and NSE.

The study of the classification of the variables of importance for the projection (VIP) of the model axes confirms the relationship between the previous exposure to the gas (VIP1=3.354) and the clinic, accompanied by the temperature on leaving the hyperbaric chamber (VIP1=2.757), also the temperature before the start of the experiment (VIP1=1.554). PS100 (VIP1=1.260), NSE (VIP1=0.853) and bicarbonates (VIP1=0.744) are also important data. These values decrease slightly on projection axes 2 and 3. This is followed by *Muribaculaceae* (*unidentified genus*) (VIP1=1.090 ; VIP2=1.017), *Lachnospiraceae_GCA-9000066575* (VIP1=0.892 ; VIP2=0.881), and *Christensenellaceae_R-7 group* (VIP1=0.846 ; VIP2=0.941), in decreasing order of importance on projection axis 1. Globally, microbes are more abundant on axes 2 and 3. However, the importance of these bacterial genera, at t1 and t2, underlies the link between the initial presence of these bacteria and the clinic, taking into account the value of the contribution of axis 2 (4.9%) compared to axis 1 (31%) for the construction of the model.

The discriminant analysis based on the values of the variables of importance in the projection according to the discrimination (VIDs) specifies the interest of the parameters according to the clinical state (Figure 4). Thus, the Asympto state is more clearly associated with a positive correlation with previous exposure to air (rather than CO₂), high surface temperature, PS100 and IL-6 values, and also with the presence of *Oscillospiraceae_intestinimoneas*, *Nostocaceae_Tolypothrix*, *Akkermansiaceae_Akkermansia* et cetera. Negative correlations compared to the Asympto status can also be seen in the figure, which ultimately bring us closer to the Sympto status. On the other hand, the symptomatic status is positively correlated with previous ventilation of CO₂, increased bicarbonate and NSE levels (which would support the clinic), but also with the initial presence of unidentified genera of *Muribaculaceae*, *Roseburia* and *Lachnospiraceae* in the stool. The ratio of aerobic/anaerobic strains seems less interesting for model projection, which does not mean that it is not clinically significant. Post-hoc tests then specify the significance of each criterion.




Figure 4. Classification of the Variables of Importance in projection according to the Discrimination (VIDs) which specify the interest of the parameters according to the clinical state. According to PLS-DA, clinical trend arrows appear in gradient color. The more positive the values (top), the closer they are to the asymptomatic clinical status. The more negative the values (bottom), the more the mice expressed DCS symptoms. VIDs $\pm$ Standard Deviation.

3.3. Two-Way Analysis

There are a number of variables that catch our attention when reading the PLS-DA.

3.3.1. Body Temperature

Infrared thermograms performed after the hyperbaric protocol (KW; $p < 0.001$) show differences between the clinical groups. In this case, symptomatic animals previously exposed to CO₂ had a lower mean surface temperature ($\sim 30.9^{\circ}\text{C}$ vs. $\sim 32.0^{\circ}\text{C}$) than mice in other groups (Figure 5).




Figure 5. Surface temperature measured by infrared (from thermograms) at the very first manipulation, or after the hyperbaric protocol. The clinical group is represented by colors, and the

gas group is represented by shape. Please refer to Figure 1 for the tracking timeline. Kruskal-Wallis (KW) analysis with post-hoc dunn test (Bonferroni correction: 0,0083).

Even before the start of the protocol, the temperature of the animals (Figure 5) would tend to be determinative to the extent that the symptomatic animals had different temperatures than the uninjured animals (KW; $p = 0.046$). It should be noted that the initial temperature of symptomatic AIR mice tended to be lower ($\sim 0.4^{\circ}\text{C}$) than that of asymptomatic AIR mice (post hoc Dunn test; $p=0.0087$). This point is consistent with PLS-DA, i.e. with high VIP and VIDs scores.

3.3.2. Plasma Analysis

Plasma variables such as IL-6, NSE, Copeptin, Bicarbonate, PS100 measured after the accident-prone protocol do not show significant differences between the groups (Figure 6). There is a trend for NSE, especially between the Asympto_CO2 mice with lower levels than the Asympto_AIR mice.




Figure 6. Plasmatic analyses carried out at the very end of the protocol, i.e. after the provocative hyperbaric protocol. The clinical group is represented by colors, and the gas group is represented by shape. Please refer to Figure 1 for the tracking timeline. Normalized data. Kruskal-Wallis (KW) analysis with post-hoc dunn test (Bonferroni correction: 0,0083).

3.4. Initial Fecal Content Analysis

16s rDNA analysis was performed on mouse feces collected at the very beginning of the protocol. Approximately 97% (min 4%_max 99%) of ASVs were not classified/identified, so these results must be interpreted with caution (see Limitations section). Finally, 520 ASVs from the stools of 88 mice allowed the identification of about thirty genera, distributed in about 13 families and 6 classes, with the presence of unknowns (Figure 7). Only about fifteen species appeared to be correctly identified, prompting analyses at the genus level. The distribution of relative abundances (Figure 7, right side) according to Bray-Curtis dissimilarity agglomerative hierarchical clustering (ACH; Figure 7, left side) on the genera allows to separate groups without directly corresponding to our clinical groups. The same is true for richness indices based on ASVs (Figure 7, middle). A plot of diversity according to the Shannon index is available in the supplementary data (Supplementary Data, Figure 1). These results should be viewed in the context of the contribution of bacteria to the PLS-DA model.




Figure 7. The dendrogram clustering illustrates the Bray-Curtis dissimilarity of the taxonomic profile, which was performed using the ASV abundance table. The scale bar of the dendrogram represents the degree of dissimilarity (%) between the prokaryotic communities. The number of ASVs was calculated from the 16S rDNA ASV table for each sample. The clinical group is represented by colors, and the gas group is represented by shape. The bar chart illustrates the cluster from the Bray-Curtis dendrogram. Each bar represents the relative abundance of the family in the sample under study.

Cluster I of the ACH with a high proportion of symptomatic animals (8/10) presents a relatively high richness (Shannon diversity index close to 3.5), but with a lot of Clostridia. This cluster also has a very low proportion of aerobic genera (min 0% _max 2%), while the average for all animals is 35±35%. On the other hand, cluster II, consisting of 8 symptomatic mice (/11), has a very high aerobic presence (min 97%_max 100%). This cluster, whose Shannon index is close to 3.0, shows a particular dominance of 2 families, *Sphingomonadaceae* (*Alphaproteobacteria*) and *Nostocaceae* (*Cyanobacteria*). Cluster IV, with 5 symptomatic mice (/10), has the lowest microbial diversity (Shannon close to 1) and a median aerobic rate of 16% (min 0%_max 66%). Cluster VIII, with 8 symptomatic animals (/10), has a dominance of *Rhizobiaceae* (*Alphaproteobacteria*) and *Cyanobacteria Nostocaceae* (Shannon close to 2.1) and a median aerobic rate of 84% (min 16%_max 98%).

Overall, Shannon, Coverage or leveness Pielou, and dominance dmn (the McNaughton's dominance index) indices (Supplementary Data, Figure 2) organized by clinical and gas groups remain non-significant (KW; $p>0.05$), although at the post-hoc level trends (Dunn; $p < 0.05$) appear between air-exposed mice (symptomatic versus asymptomatic). Asymptomatic animals appear to have lower Shannon diversity indices.

A more specific analysis of genera reveals differences between groups, in relative or raw abundances (Figure 8 & Supplementary data Table S1).

Significant differences, both in relative and raw abundances, are highlighted in the genera *Blautia* and *GCA-900066575* of the family *Lachnospiraceae*, with differences at the post-hoc level. There are much higher values of *Lachnospiraceae Blautia* in the Sympto_Air compared to the other groups, especially compared to the Sympto_CO2 where *Blautia* is almost absent.

There are significantly higher values of Lachnospiraceae GCA-900066575 in Sympto_Air compared to Asympto_Air. A similar trend is found at the post-hoc level for *Candidatus Saccharimonas*, with higher values in Sympto_Air compared to Asympto_Air.

From a less strict statistical point of view, we see trends (KW with $p < 0.010$) in Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium, whose values are higher in Asympto_CO2 compared to Asympto_AIR and Sympto_AIR. The genus Nostoc PCC-73102 has slightly higher values in Asympto_Air compared to Sympto_Air (without relative abundance playing a role). Roseburia has higher abundance values in Sympto_AIR compared to Asympto_AIR and Sympto_CO2. The genus Anaerostipes has higher relative (and raw) abundance values in Asympto_CO2 compared to Asympto_Air.




Figure 8. Heat tree constructed using 16S rDNA data at the genus level. Each connected node represents a different taxonomic rank. The nodes are colored based on the Kruskal-Wallis differences between the 4 conditions (Asympto_AIR, Asympto_CO2, Sympto_AIR and Sympto_CO2).

4. Discussion

This study is the first to demonstrate that prior exposure to a CO₂-enriched atmosphere increases the risk of DCS in mice. Although thermograms taken after the hyperbaric protocol seem to support the diagnosis, more generally the temperature of the animals is associated with the clinic without us knowing if it is the consequence or even the cause. According to the PLS-DA, the variation of bicarbonate levels in the blood confirms the low adaptive CO₂ exfiltration after the provocative protocol, especially after a previous CO₂ exposure. This would require the use of a buffering system.[60].

The trend of differences in NSE levels between CO₂- and AIR-exposed mice may suggest the involvement of different pathogenic pathways, where neurons seem to be less affected in Asympto_CO2 mice.

A specific bacterial community, based on diversity, does not appear to contribute collectively to preventing or increasing DCS risk. However, some bacteria, for example from the *Lachnospiraceae*

family, appear to play a role depending on whether or not there is prior exposure to CO₂. Although secondary, this suggests that more than one pathway may alter the outcome.

4.1. Effects of CO₂ and Diving on Circulating Markers of Neuronal Stress

Neuron-specific enolase (NSE) and the glial-derived peptide S100 calcium-binding protein B (S100B) are commonly described to assess the presence and severity of neurological injury. S100B is known to rise in the peripheral circulation in association with both oxidative [61] and heat stress [62], and a relationship is described with core body temperature (T_c) response [63]. NSE is a glycolytic enzyme found in brain neurons and peripheral nervous tissue that is elevated during cerebral inflammation or ischemia. These proteins were measured in neurological diving accident victims (symptomatic) and compared with asymptomatic divers (same diving profile). In fact, the level of NSE is higher in the symptomatic group than in the control group [64] and S100B level has no change [64]. In rat studies, there appears to be no change in plasma S100B concentrations after a low bubble grade dive, and it is increased after a high bubble grade dive [65,66]. Interestingly, S100B protein levels increase with apnea-induced hypoxic stress [67].

In our study, S100B remains non-contributory. S100B does not appear to be sensitive to cold (see below). Differences in NSE levels between CO₂- and AIR-exposed mice may indicate that neurons in asymptomatic mice appear to be less affected, but we cannot explain why this difference is not found between the symptomatic and asymptomatic AIR groups. Caution should therefore be exercised in interpreting these data. However, one avenue would be worth exploring given the high rates of NSE expressed in patients with chronic hypercapnia and CO₂ retention [68].

4.2. Effects of CO₂ and Diving on Plasma Markers, Thermoregulation and Vasomotion

A high level of bicarbonates over time, as supported by the PLS-DA, could affect cerebral vasomotor function and blood flow. Hypothetically, this vasodilation could make the brain of mice previously exposed to CO₂ more susceptible to bubbles (see moderation), both by increasing gas saturation and by making it more accessible to bubbles of systemic origin.

Temperature

A1>In the present study, the Thermograms performed after the hyperbaric protocol seem to confirm the diagnosis, as expected according to previous studies.[26]. Symptomatic mice have a lower surface temperature (~1.1 °C). To explain this low temperature in symptomatic mice, it would be simplistic to exclude a pro-inflammatory state (usually associated with temperature elevation) as a hypothesis, especially in CO₂ exposed mice. This should probably affect vasomotion, the quality of tissue perfusion, i.e. gas balance and bubble growth, according to the most common theory on DCS.[8].

More unexpectedly, it should be noted that the initial temperature (before any exposure) of symptomatic AIR mice tends to be lower (~0.4°C) than that of asymptomatic AIR mice. This suggests that initial temperature is critical for prognosis, and that a lower baseline temperature currently predicts a state of susceptibility to DCS. Considering that hypercapnia induces hypothermia, but sufficient heat production is a prerequisite for inflammation, a special study should be devoted to. Again, we might wonder about the links between NO/vasodilation/thermogenesis [69] and now CO₂ in the DCS.

4.3. Impact of the Microbiota in the Stress Response

The gut microbiota has been of interest to hyperbaric teams for several years. It has been documented that the core of the intestinal contents is hypoxic, but there is a gradient of oxygen concentration near the intestinal epithelium [70] and that oxygen or hypoxia can reshape its community [71,72]. For example, an increase in the frequency of the Clostridium group is shown after a safety dive in men [31].

In this study, the collection of feces at the beginning of the protocol and the analysis of 16sDNA were conducted to investigate whether the microbial profile could play a critical role in the response to the accident-prone protocol. While no specific bacterial spectrum was identified, it was found that certain bacteria, when examined individually, appeared to have an impact on clinical outcomes depending on whether or not there was prior exposure to CO₂. This suggests that while CO₂ may play an important role in symptoms, the gut microbiota may also have an indirect influence, although this relationship is less pronounced according to the PLS-DA analysis.

The use of richness indices, such as the Shannon index, did not allow for the definition of a specific risk fingerprint. This may be attributed to the use of an inbred animal strain, C57Bl6/J, which exerts a selective pressure on the gut microbiota, resulting in low diversity as reported by the Shannon index. This limitation or advantage of the study is noteworthy because reducing diversity enhances the contrast, making it easier to identify the microorganisms involved in pathology. This finding supports the notion of inter- and intra-individual variability that has been described in both animal and human diving studies [23,29,30]. However, it should be noted that some bacterial genera demonstrated intergroup differences in abundance, as indirectly suggested in previous studies [73–75]. For example, the low representation of *Lachnospiraceae_GCA-900066575* in this study appears to have a beneficial effect after a risky dive, as the Asympto_AIR animals showed significantly less abundance of this bacterium compared to their symptomatic counterparts (i.e., those not exposed to CO₂).

Certain bacteria require further caution or additional targeted studies. For instance, *Candidatus_Saccharimonas* is significantly more abundant in Sympto_Air animals compared to Asympto_Air animals. However, its role as a bacterium promoting decompression sickness (DCS) may be debated, since its contribution to the model was very weak. Similarly, *Nostoc_PCC-73102* demonstrated a slightly greater contribution to the model but only displayed a trend towards being pathogenic. The higher abundance of *Lachnospiraceae_Roseburia* in Sympto_AIR mice suggests a potential promotion of DCS, while its absence may be beneficial in the event of a provocative dive. Paradoxically, this same low abundance may favor DCS in the presence of prior CO₂ inhalation. Similar trends were observed for *Lachnospiraceae_Blautia*, with higher levels of significance in their abundance ($p < 0.002$), but low contribution values to the model. Furthermore, a higher initial proportion of *Lachnospiraceae_Anaerostipes* tends to be beneficial in the case of prior exposure to CO₂, despite its relatively low contribution to the model. These results contribute to the ongoing debate regarding the role of *Lachnospiraceae* in physiology, as they are known to be major producers of short-chain fatty acids [76]. Further investigations are needed to clarify the specific actions of the *Lachnospiraceae* family observed in this study. We carried out tests at the level of the *Lachnospiraceae* family, which did not reveal any significant differences. This implies a more specific action that remains to be clarified. In fact, the effect of short chain fatty acids (SCFA) by fermentation has already been mentioned in DCS, as a protector but also as a promoter of accidents [29,74,77–79].

Recent work suggests that the microbiota plays an increasingly important role in thermoregulation through its metabolic derivatives [38] or through its serotonergic effects via modulation of the gut-brain axis [32,33]. For instance, the inhibition of serotonin reuptake with fluoxetine has been shown to reduce DCS [27,28] and it would be interesting to explore this question further through the influence of the microbiota. The influence of temperature in the DCS is discussed earlier in this document, and we could now add a possible role for bacteria. It should be remembered that the fecal samples were taken at the very beginning of the protocol. Recent studies have shown that cold exposure can lead to dramatic changes in the composition and function of the gut microbiota, with acute responses, and *Akkermansia muciniphila* may be the most appropriate strain [39,40]. Looking at the VIDs ranking, the amount of *Akkermansia* (*Akkermansia muciniphila* is the only one identified in this genus) would be positively correlated with lower accident rates or, on the contrary, negatively correlated with prior exposure to CO₂.

Among the noteworthy findings, a higher initial proportion of *Rhizobiaceae_Allorhizobium-Neorhizobium-Parhizobium-Rhizobium* (*Rhizobium* sp. is the only one identified in this genus) would tend to be beneficial in cases of prior CO₂ exposure, despite its relatively low contribution to the

model. *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium* is related to nitrogen fixation by plants. Therefore, it could come from the diet of mice. These bacteria have been proposed to be involved in the development of multiple sclerosis, where their consumption from root bulbs containing Rhizobiaceae acts as exogenous antigens that trigger an immune response [34]. These bacteria are also known to neutralize the immune response [35], and their role in DCS warrants further exploration based on the findings of this study and others [23,25].

4.4. Context

This study was particularly motivated by an unusual accident where high levels of CO₂ were detected. When fresh air is around 0.04% CO₂, wearing face masks for more than 5 min carries a possible chronic exposure to carbon dioxide of 1.41% to 3.2% of the inhaled air [80–82]. In this context, relatively speaking, this work could open the question of wearing a face mask. In fact, in a situation such as the COVID epidemic, where the rules regarding the wearing of masks can be particularly strict, special attention should be paid to professional divers.

4.5. Moderation, Study Limits

The mouse is probably not the best model for DCS, and as with any model, its limitations must be considered when extrapolating to humans. This applies to the hyperbaric chamber diving protocol and also to the amount of CO₂, which may seem extreme. However, this allows reliable clinical signs to be produced in animals.

In this study, it was unfortunate that before-and-after comparisons could not be made for blood analysis and fecal bacterial identification. Exposure to CO₂ significantly reduced the number of asymptomatic animals that could be sampled. Accidentology reduced blood perfusion (access to punction) and also fecal excretion.

In fact, we have favored the symptomatology to the detriment of the technique of ultrasound bubble detection in this animal. Clinical evaluation remains essential, as it is well known that the presence of bubbles does not necessarily correlate with the reporting of DCS symptoms. We assume that we did not quantify circulating bubbles in this study, whereas their diameter is influenced by gases dissolved in the tissue, such as CO₂ [83]. Because we cannot exclude the involvement of CO₂ in the growth of the bubbles, especially in symptomatic mice previously exposed to CO₂, this point deserves clarification.

We mention the influence of the genetic background of the mice on the selection of the microbiota. To the extent that we have used a single animal strain, it would be interesting to combine several strains to verify their impact on microbial communities and accidentology, and then try to identify a profile, as could be done in rats [23,73,84].

Regarding the microbiota, the low identification rate could be a limitation of this study. Approximately 97% (min 4%_max 99%) of the ASVs were not classified/identified, which requires a cautious interpretation of these results. The remaining 97% could represent, for example, microbes, but also their degradation products or even food waste.

Certain bacterial strains have emerged from this work. This study is not specifically dedicated to their identification or their mode of action, so caution should be exercised in their interpretation. This work could be directed by the use of antibiotics or, on the contrary, by carrying out microbiota transplants, for example.

5. Conclusions

This work is the first to clearly demonstrate the influence of exposure to a high concentration of CO₂ on DCS risk. Exposure of mice to a maximum CO₂ concentration of 70 hPa for one hour prior to the hyperbaric protocol dramatically increased the incidence of DCS. This was confirmed by temperature measurements at the end of the procedure. More unexpectedly, it seems that a lower temperature of the animals even before exposure to the accident-prone protocol leads to an unfavorable prognosis. This study also suggests that the microbiota may partially influence

thermogenesis and thus accidentology. For example, the low representation of *Lachnospiraceae_GCA-9000066575* could be perceived as protective in the AIR condition. However, depending on prior exposure, certain high abundances of bacteria identified in this work, such as *Lachnospiraceae_Blautia*, could be perceived as favoring DCS or, on the contrary, as protective if the risky dive follows exposure to CO₂. Studies should further explore this issue in order to improve prevention in diving.

5.1. Grants and Funding

The last 3 authors are equally placed. The work should be attributed to Aix-Marseille University and to the “Institut de Recherches Biomédicales des Armées” laboratories. It is supported by Grant No. SAN-1-0516 from the Direction Générale de l’Armement of the French army, Paris, France.

5.2. Caption for the Figures

Figure 1: Flow Chart of Experimental Design - Measurement of Temperature, Identification, Weighing, and Feces Collection for Microbiota Analysis in Mice

Figure 2: Decompression sickness occurrences 30 minutes after pathogenic decompression as a function of previous gas exposure (Air or CO₂). **A, Histogram:** Proportion of mice with DCS symptoms. The DCS status was attributed when the mice presented neurological signs in the form of paresis or paralysis of at least one limb, convulsions and/or reduced performance in the motor test for forelimbs, and also respiratory troubles, even death. **B, Radar Chart:** percentage of mice displaying a type of symptom in a population, considering that a mouse can present the whole panel of symptoms. * denotes $p < 0.050$, ** denotes $p < 0.010$, *** denotes $p < 0.001$.

Figure 3: Partial Least Square – Discriminant Analysis (PLS-DA) plot conducted on clinic (Symptomatic or Asymptomatic, green line) as dependant variable) and as explanatory variable gas exposure before the hyperbaric protocol (CO₂ or Air, blue), micro-organisms genus detected in stools (red), mice superficial temperature before and after the protocol (black), plasma analysis performed at the end of the protocol (orange). (Left) Confusion matrix summarize the observations according to the clinic status of mice (Symptomatic in red or Asymptomatic in green), according to the gas breathed before the dive (Air in blue or CO₂ in orange), with their confidence ellipse (Fisher, Confidence Interval 95%). (Right) Correlation map based from observations (green dots) according to their qualitative or quantitative explanatory characteristics.

Figure 4: Classification of the Variables of Importance in projection according to the Discrimination (VIDs) which specify the interest of the parameters according to the clinical state. The more positive the values (top), the closer they are to the asymptomatic clinical status. The more negative the values (bottom), the more the mice expressed DCS symptoms. VIDs $\pm$ Standard Deviation.

Figure 5: Surface temperature measured by infrared, at the very first manipulation, or after the hyperbaric protocol. Normalized data. Kruskal-Wallis (KW) analysis with post-hoc dunn test (Bonferroni correction: 0,0083).

Figure 6: Plasmatic analyses carried out at the very end of the protocol, i.e. after the provocative hyperbaric protocol. Normalized data. Kruskal-Wallis (KW) analysis with post-hoc dunn test (Bonferroni correction: 0,0083).

Figure 7: The dendrogram clustering illustrates the Bray-Curtis dissimilarity of the taxonomic profile, which was performed using the ASV abundance table. The scale bar of the dendrogram represents the degree of dissimilarity (%) between the prokaryotic communities. The number of ASVs was calculated from the 16S rDNA ASV table for each sample. The clinical group is represented by colors, and the gas group is represented by shape. The bar chart illustrates the cluster from the Bray-Curtis dendrogram. Each bar represents the relative abundance of the family in the sample under study.

Figure 8: Heat tree constructed using 16S rDNA data at the genus level. Each connected node represents a different taxonomic rank. The nodes are colored based on the Kruskal-Wallis differences between the 4 conditions (Asympto_AIR, Asympto_CO₂, Sympto_AIR and Sympto_CO₂).

Author Contributions: The last 3 authors are equally placed. L. D., A. B., M. M., A.-V. D., A.P., J.-E. B., R.G. and N.V. conception and design of research; L. D., A. B., M. M., S.G. and N.V. performed the experiments; N. A. developed the equipment. L. D., A.-V. D., A.P. and N.V. analysed data; L. D., A.-V. D., A.P. and N.V. interpreted the results of the experiments; A.P., and N.V. prepared the figures; N.V. drafted, edited the manuscript; L. D., A. B., A.-V. D., A.P., J.-J. R., C. R., J.-C. R. J.-E. B., R.G. and N.V. revised the manuscript; R.G. and N.V. approved the final version of the manuscript.

Conflicts of Interest: All other authors declare no competing interests. The last 3 authors are equally placed.

References

1. Andicochea, C.T.; Henriques, M.E.; Fulkerson, J.; Jay, S.; Chen, H.; Deaton, T. Elevated environmental carbon dioxide exposure confounding physiologic events in aviators? *Military medicine* **2019**, *184*, e863-e867.
2. Doolette, D.J.; Mitchell, S.J. Hyperbaric conditions. *Comprehensive Physiology* **2010**, *1*, 163-201.
3. Gennser, M.; Blogg, S.L. Oxygen or carbogen breathing before simulated submarine escape. *Journal of applied physiology* **2008**, *104*, 50-56.
4. Daubresse, L.; Amara, J.J.; Grau, M.M.; De Maistre, S.; Guillaume, C.G.; Lehot, H.H.; Texier, G.G.; Blatteau, J.-E. Case series of unusual neurologic symptoms after scuba diving training sessions, in the context of SARS COV2 preventive measures and restrictions. In Proceedings of the EUBS, 2022.
5. Mano, Y.; D'Arrigo, J. Relationship between CO2 levels and decompression sickness: implications for disease prevention. *Aviation, space, and environmental medicine* **1978**, *49*, 349-355.
6. Le Chatelier, H.L. Sur un énoncé général des lois des équilibres chimiques. *Comptes Rendus Académie des Sciences* **1884**, *99*, 786-789.
7. Caldwell, H.G.; Carr, J.M.; Minhas, J.S.; Swenson, E.R.; Ainslie, P.N. Acid-base balance and cerebrovascular regulation. *The Journal of Physiology* **2021**, *599*, 5337-5359.
8. Brubakk, A.O.; Ross, J.A.; Thom, S.R. Saturation diving; physiology and pathophysiology. *Comprehensive physiology* **2011**, *4*, 1229-1272.
9. Pickles, D.; Ogston, D.; MacDonald, A. Effects of gas bubbling and other forms of convection on platelets in vitro. *Journal of Applied Physiology* **1989**, *67*, 1250-1255.
10. Giry, P.B.; Porlier, G.; Eastman, D.; Radomski, M.W. Dive-induced modifications in platelet kinetics in rats. *Undersea Biomed Res* **1977**, *4*, 147-157.
11. Pontier, J.M.; Jimenez, C.; Blatteau, J.E. Blood platelet count and bubble formation after a dive to 30 msw for 30 min. *Aviation, space, and environmental medicine* **2008**, *79*, 1096-1099.
12. Thorsen, T.; Klausen, H.; Lie, R.T.; Holmsen, H. Bubble-induced aggregation of platelets: effects of gas species, proteins, and decompression. *Undersea Hyperb Med* **1993**, *20*, 101-119.
13. Brubakk, A.O.; Neuman, T.S.; Elliott, D.H. *Bennett and Elliott's Physiology and Medicine of Diving*; Saunders: 2003; pp. 578-650.
14. Gempp, E.; Blatteau, J.E. Risk factors and treatment outcome in scuba divers with spinal cord decompression sickness. *J Crit Care* **2010**, *25*, 236-242, doi:10.1016/j.jcrc.2009.05.011.
15. Lillo, R.; Maccallum, M.; Caldwell, J. Intravascular bubble composition in guinea pigs: a possible explanation for differences in decompression risk among different gases. *Undersea biomedical research* **1992**, *19*, 375-386.
16. Ishiyama, A. Analysis of gas composition of intravascular bubbles produced by decompression. *The Bulletin of Tokyo Medical and Dental University* **1983**, *30*, 25-35.
17. Hayashi, K.; Honda, Y.; Miyakawa, N.; Fujii, N.; Ichinose, M.; Koga, S.; Kondo, N.; Nishiyasu, T. Effect of CO2 on the ventilatory sensitivity to rising body temperature during exercise. *Journal of Applied Physiology* **2011**, *110*, 1334-1341.
18. Hodes, R.; Larrabee, M.G. The relation between alveolar carbon dioxide tension and susceptibility to decompression sickness. *American Journal of Physiology-Legacy Content* **1946**, *147*, 603-615.
19. Camporesi, E.M.; Bosco, G. Ventilation, Gas Exchange and Exercise Under Pressure. In *Bennett and Elliott's Physiology and Medicine of Diving*; Elsevier Science Limited: 2003; pp. 77-114.
20. Lanphier, E.H. Carbon dioxide poisoning. *Case histories of diving and hyperbaric accidents* **1988**, 199-213.
21. Eynan, M.; Arieli, Y.; Taran, B.; Yanir, Y. Symptoms of central nervous system oxygen toxicity during 100% oxygen breathing at normobaric pressure with increasing inspired levels of carbon dioxide: a case report. *Diving and Hyperbaric Medicine* **2020**, *50*, 70.
22. Eynan, M.; Arieli, R.; Adir, Y. Response to CO2 in novice closed-circuit apparatus divers and after 1 year of active oxygen diving at shallow depths. *Journal of applied physiology* **2005**, *98*, 1653-1659.
23. Vallée, N.; Dugrenot, E.; Desruelle, A.-V.; Richard, S.; Coupé, S.; Ramdani, C.; Guieu, R.; Risso, J.-J.; Gaillard, S.; Guerrero, F. Highlighting of the interactions of MYD88 and NFKB1 SNPs in rats resistant to decompression sickness: toward an autoimmune response. *Frontiers in Physiology* **2023**, *14*.

24. Blatteau, J.E.; Gaillard, S.; De Maistre, S.; Richard, S.; Louges, P.; Gempp, E.; Druelles, A.; Lehot, H.; Morin, J.; Castagna, O.; et al. Reduction in the Level of Plasma Mitochondrial DNA in Human Diving, Followed by an Increase in the Event of an Accident. *Front Physiol* **2018**, *9*, 1695, doi:10.3389/fphys.2018.01695.
25. Lambrechts, K.; de Maistre, S.; Abraini, J.H.; Blatteau, J.-E.; Risso, J.-J.; Vallée, N. Tirofiban, a Glycoprotein IIb/IIIa Antagonist, Has a Protective Effect on Decompression Sickness in Rats: Is the Crosstalk Between Platelet and Leukocytes Essential? *Frontiers in Physiology* **2018**, *9*, doi:10.3389/fphys.2018.00906.
26. Desruelle, A.V.; Louge, P.; Richard, S.; Blatteau, J.E.; Gaillard, S.; De Maistre, S.; David, H.; Risso, J.J.; Vallee, N. Demonstration by Infra-Red Imaging of a Temperature Control Defect in a Decompression Sickness Model Testing Minocycline. *Front Physiol* **2019**, *10*, 933, doi:10.3389/fphys.2019.00933.
27. Cosnard, C.; De Maistre, S.; Abraini, J.H.; Chazalviel, L.; Blatteau, J.E.; Risso, J.J.; Vallee, N. Thirty-five Day Fluoxetine Treatment Limits Sensory-Motor Deficit and Biochemical Disorders in a Rat Model of Decompression Sickness. *Front Physiol* **2017**, *8*, 604, doi:10.3389/fphys.2017.00604.
28. Vallee, N.; Lambrechts, K.; De Maistre, S.; Royal, P.; Mazella, J.; Borsotto, M.; Heurteaux, C.; Abraini, J.; Risso, J.J.; Blatteau, J.E. Fluoxetine Protection in Decompression Sickness in Mice is Enhanced by Blocking TREK-1 Potassium Channel with the "spadin" Antidepressant. *Front Physiol* **2016**, *7*, 42, doi:10.3389/fphys.2016.00042.
29. Oya, M.; Tokunaga, T.; Tadano, Y.; Ogawa, H.; Fujii, S.; Murakami, W.; Tamai, K.; Ikomi, F.; Morimoto, Y. The composition of the human fecal microbiota might be significantly associated with fecal SCFA levels under hyperbaric conditions. *Bioscience of Microbiota, Food and Health* **2021**, *40*, 168-175.
30. Hagberg, M.; Ornhagen, H. Incidence and risk factors for symptoms of decompression sickness among male and female dive masters and instructors--a retrospective cohort study. *Undersea Hyperb Med* **2003**, *30*, 93-102.
31. Oya, M.; Tadano, Y.; Takihata, Y.; Murakami, W.; Fujii, S.; Tamai, K.; Morimoto, Y.; Ikomi, F.; Tokunaga, T. Effects of hyperbaric conditions on fecal microbiota. *Bioscience of microbiota, food and health* **2019**, *38*, 35-39.
32. Bo, T.; Tang, L.; Xu, X.; Liu, M.; Wen, J.; Lv, J.; Wang, D. Role of gut microbiota in the postnatal thermoregulation of Brandt's voles. *Cell Reports* **2023**, *42*.
33. Li, S.; Li, X.; Wang, K.; Li, Y.; Nagaoka, K.; Li, C. Gut microbiota intervention attenuates thermogenesis in broilers exposed to high temperature through modulation of the hypothalamic 5-HT pathway. *Journal of Animal Science and Biotechnology* **2023**, *14*, 159.
34. Boucher, J. The cause of multiple sclerosis is autoimmune attack of adenosyltransferase thereby limiting adenosylcobalamin production. *Medical Hypotheses* **2017**, *109*, 29-37.
35. Phan, H.T.; Ho, T.T.; Chu, H.H.; Vu, T.H.; Gresch, U.; Conrad, U. Neutralizing immune responses induced by oligomeric H5N1-hemagglutinins from plants. *Veterinary research* **2017**, *48*, 1-13.
36. Khakisahneh, S.; Zhang, X.-Y.; Nouri, Z.; Wang, D.-H. Gut microbiota and host thermoregulation in response to ambient temperature fluctuations. *Msystems* **2020**, *5*, 10.1128/msystems.00514-00520.
37. Zhang, L.; Yang, Z.; Yang, F.; Wang, G.; Zeng, M.; Zhang, Z.; Yang, M.; Wang, Z.; Li, Z. Gut microbiota of two invasive fishes respond differently to temperature. *Frontiers in Microbiology* **2023**, *14*, 1087777.
38. Tang, Y.; Wang, Y.-D.; Wang, Y.-Y.; Liao, Z.-Z.; Xiao, X.-H. Skeletal muscles and gut microbiota-derived metabolites: novel modulators of adipocyte thermogenesis. *Frontiers in Endocrinology* **2023**, *14*.
39. Chevalier, C.; Stojanović, O.; Colin, D.J.; Suarez-Zamorano, N.; Tarallo, V.; Veyrat-Durebex, C.; Rigo, D.; Fabbiano, S.; Stevanović, A.; Hagemann, S. Gut microbiota orchestrates energy homeostasis during cold. *Cell* **2015**, *163*, 1360-1374.
40. Worthmann, A.; John, C.; Rühlemann, M.C.; Baguhl, M.; Heinsen, F.-A.; Schaltenberg, N.; Heine, M.; Schlein, C.; Evangelakos, I.; Mineo, C. Cold-induced conversion of cholesterol to bile acids in mice shapes the gut microbiome and promotes adaptive thermogenesis. *Nature medicine* **2017**, *23*, 839-849.
41. Tian, Y.; Fu, M.; Su, J.; Yan, M.; Yu, J.; Wang, C.; Niu, Z.; Du, Y.; Hu, X.; Zheng, J. Gut microbiota dysbiosis and intestinal barrier impairment in diarrhea caused by cold drink and high-fat diet. *Toxicology* **2024**, *502*, 153728.
42. Lv, R.; Liu, X.; Zhang, Y.; Dong, N.; Wang, X.; He, Y.; Yue, H.; Yin, Q. Pathophysiological mechanisms and therapeutic approaches in obstructive sleep apnea syndrome. *Signal transduction and targeted therapy* **2023**, *8*, 218.
43. Campaña-Duel, E.; Ceccato, A.; Morales-Quinteros, L.; Camprubi-Rimblas, M.; Artigas, A. Hypercapnia and its relationship with respiratory infections. *Expert Review of Respiratory Medicine* **2024**.
44. Ruiz, A. Carbon dioxide response curves during hypothermia. *Pflügers Archiv* **1975**, *358*, 125-133.
45. Santiago-Marrero, I.; Liu, F.; Wang, H.; Arzola, E.P.; Xiong, W.-C.; Mei, L. Energy expenditure homeostasis requires ErbB4, an obesity risk gene, in the paraventricular nucleus. *Eneuro* **2023**, *10*.
46. Branco, L.; Wood, S.C. Role of central chemoreceptors in behavioral thermoregulation of the toad, *Bufo marinus*. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **1994**, *266*, R1483-R1487.

47. Tsuji, B.; Hayashi, K.; Kondo, N.; Nishiyasu, T. Characteristics of hyperthermia-induced hyperventilation in humans. *Temperature* **2016**, *3*, 146-160.
48. White, M.D. Components and mechanisms of thermal hyperpnea. *Journal of Applied Physiology* **2006**, *101*, 655-663.
49. Sunanaga, J.; Deng, B.-S.; Zhang, W.; Kanmura, Y.; Kuwaki, T. CO₂ activates orexin-containing neurons in mice. *Respiratory physiology & neurobiology* **2009**, *166*, 184-186.
50. Hacquemand, R.; Buron, G.; Pourié, G.; Karrer, M.; Jacquot, L.; Brand, G. Effects of CO₂ inhalation exposure on mice vomeronasal epithelium. *Cell biology and toxicology* **2010**, *26*, 309-317.
51. Rodrigues, L.T.C.; da Silva, E.N.; José de Anchieta, C.; Gargaglioni, L.H.; Dias, M.B. Glutamate metabotropic receptors in the lateral hypothalamus/perifornical area reduce the CO₂ chemoreflex. *Respiratory physiology & neurobiology* **2019**, *260*, 122-130.
52. Blatteau, J.E.; Barre, S.; Pascual, A.; Castagna, O.; Abraini, J.H.; Risso, J.J.; Vallee, N. Protective effects of fluoxetine on decompression sickness in mice. *PLoS One* **2012**, *7*, e49069, doi:10.1371/journal.pone.0049069.PONE-D-12-20978 [pii].
53. Vallee, N.; Meckler, C.; Risso, J.J.; Blatteau, J.E. Neuroprotective role of the TREK-1 channel in decompression sickness. *J Appl Physiol (1985)* **2012**, *112*, 1191-1196, doi:10.1152/japplphysiol.01100.2011.
54. Andrews, S. FastQC: a quality control tool for high throughput sequence data. Available online. Retrieved May **2010**, *17*, 2018.
55. Callahan, B.J.; McMurdie, P.J.; Rosen, M.J.; Han, A.W.; Johnson, A.J.A.; Holmes, S.P. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods* **2016**, *13*, 581-583.
56. R Core Team, R. R: A language and environment for statistical computing. **2013**.
57. Wang, Q.; Garrity, G.M.; Tiedje, J.M.; Cole, J.R. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and environmental microbiology* **2007**, *73*, 5261-5267.
58. Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F.O. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic acids research* **2012**, *41*, D590-D596.
59. Bokulich, N.A.; Subramanian, S.; Faith, J.J.; Gevers, D.; Gordon, J.I.; Knight, R.; Mills, D.A.; Caporaso, J.G. Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nature methods* **2013**, *10*, 57-59.
60. Pinard, E.; Rigaud, A.; Riche, D.; Naquet, R.; Seylaz, J. Continuous determination of the cerebrovascular changes induced by bicuculline and kainic acid in unanaesthetized spontaneously breathing rats. *Neuroscience* **1987**, *23*, 943-952.
61. Bailey, D.M.; Evans, K.A.; McEneny, J.; Young, I.S.; Hullin, D.A.; James, P.E.; Ogoh, S.; Ainslie, P.N.; Lucchesi, C.; Rockenbauer, A. Exercise-induced oxidative–nitrosative stress is associated with impaired dynamic cerebral autoregulation and blood–brain barrier leakage. *Experimental physiology* **2011**, *96*, 1196-1207.
62. Watson, P.; Shirreffs, S.M.; Maughan, R.J. Blood-brain barrier integrity may be threatened by exercise in a warm environment. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **2005**, *288*, R1689-R1694.
63. Stacey, M.J.; Hill, N.E.; Parsons, I.T.; Wallace, J.; Taylor, N.; Grimaldi, R.; Shah, N.; Marshall, A.; House, C.; O'Hara, J.P. Relative changes in brain and kidney biomarkers with Exertional Heat Illness during a cool weather marathon. *Plos one* **2022**, *17*, e0263873.
64. Gempp, E.; Louge, P.; De Maistre, S.; Emile, L.; Blatteau, J.E. Neuron-specific enolase and S100B protein levels in recreational scuba divers with neurological decompression sickness. *Diving Hyperb Med* **2014**, *44*, 26-29.
65. Havnes, M.B.; Hjelde, A.; Brubakk, A.O.; Møllerlækken, A. S100B and its relation to intravascular bubbles following decompression. *Diving Hyperb Med* **2010**, *40*, 210-212.
66. Havnes, M.B.; Kerlefsen, Y.; Møllerlækken, A. S100B and NSE serum concentrations after simulated diving in rats. *Physiological Reports* **2015**, *3*, e12546.
67. Linér, M.H.; Andersson, J. Hypoxic syncope in a competitive breath-hold diver with elevation of the brain damage marker S100B. *Aviation, space, and environmental medicine* **2009**, *80*, 1066-1068.
68. Li, J.; Kong, X.; Shu, W.; Zhang, W. The association of serum neuron-specific enolase with other disease markers in chronic obstructive pulmonary disease: A case-control study. *Pakistan Journal of Medical Sciences* **2018**, *34*, 1172.
69. Blatteau, J.E.; Brubakk, A.O.; Gempp, E.; Castagna, O.; Risso, J.J.; Vallee, N. Sildenafil pre-treatment promotes decompression sickness in rats. *PLoS One* **2013**, *8*, e60639, doi:10.1371/journal.pone.0060639_PONE-D-12-40063 [pii].
70. Espey, M.G. Role of oxygen gradients in shaping redox relationships between the human intestine and its microbiota. *Free Radical Biology and Medicine* **2013**, *55*, 130-140.

71. Albenberg, L.; Esipova, T.V.; Judge, C.P.; Bittinger, K.; Chen, J.; Laughlin, A.; Grunberg, S.; Baldassano, R.N.; Lewis, J.D.; Li, H. Correlation between intraluminal oxygen gradient and radial partitioning of intestinal microbiota. *Gastroenterology* **2014**, *147*, 1055-1063. e1058.
72. Moreno-Indias, I.; Torres, M.; Montserrat, J.M.; Sanchez-Alcoholado, L.; Cardona, F.; Tinahones, F.J.; Gozal, D.; Poroyko, V.A.; Navajas, D.; Queipo-Ortuño, M.I. Intermittent hypoxia alters gut microbiota diversity in a mouse model of sleep apnoea. *European Respiratory Journal* **2015**, *45*, 1055-1065.
73. Desruelle, A.V.; de Maistre, S.; Gaillard, S.; Richard, S.; Tardivel, C.; Martin, J.C.; Blatteau, J.E.; Boussuges, A.; Rives, S.; Risso, J.J.; et al. Cecal Metabolomic Fingerprint of Unscathed Rats: Does It Reflect the Good Response to a Provocative Decompression? *Front Physiol* **2022**, *13*, 882944, doi:10.3389/fphys.2022.882944.
74. de Maistre, S.; Gaillard, S.; Martin, J.C.; Richard, S.; Boussuges, A.; Rives, S.; Desruelle, A.V.; Blatteau, J.E.; Tardivel, C.; Risso, J.J.; et al. Cecal metabolome fingerprint in a rat model of decompression sickness with neurological disorders. *Sci Rep* **2020**, *10*, 15996, doi:10.1038/s41598-020-73033-z.
75. Vallee, N.; Dugrenot, E.; Desruelle, A.V.; Tardivel, C.; Martin, J.C.; Guernec, A.; Boussuges, A.; Rives, S.; Risso, J.J.; Guerrero, F. Evidence of a hormonal reshuffle in the cecal metabolome fingerprint of a strain of rats resistant to decompression sickness. *Sci Rep* **2021**, *11*, 8317, doi:10.1038/s41598-021-87952-y.
76. Vacca, M.; Celano, G.; Calabrese, F.M.; Portincasa, P.; Gobetti, M.; De Angelis, M. The controversial role of human gut lachnospiraceae. *Microorganisms* **2020**, *8*, 573.
77. de Maistre, S.; Vallee, N.; Gaillard, S.; Duchamp, C.; Blatteau, J.E. Stimulating fermentation by the prolonged acceleration of gut transit protects against decompression sickness. *Sci Rep* **2018**, *8*, 10128, doi:10.1038/s41598-018-28510-x.
78. de Maistre, S.; Vallee, N.; Gempp, E.; Lambrechts, K.; Louge, P.; Duchamp, C.; Blatteau, J.E. Colonic Fermentation Promotes Decompression sickness in Rats. *Sci Rep* **2016**, *6*, 20379, doi:10.1038/srep20379.
79. de Maistre, S.; Vallee, N.; Gempp, E.; Louge, P.; Duchamp, C.; Blatteau, J.E. Gut fermentation seems to promote decompression sickness in humans. *J Appl Physiol (1985)* **2016**, *121*, 973-979, doi:10.1152/jappphysiol.00503.2016.
80. Brooks, J.P.; Layman, J.; Willis, J. Physiologic effects of surgical masking in children versus adults. *PeerJ* **2023**, *11*, e15474.
81. Kisielinski, K.; Hirsch, O.; Wagner, S.; Wojtasik, B.; Funken, S.; Klosterhalfen, B.; Kanti Manna, S.; Prescher, A.; Sukul, P.; Sönnichsen, A. Physio-metabolic and clinical consequences of wearing face masks—Systematic review with meta-analysis and comprehensive evaluation. *Frontiers in Public Health* **2023**, *11*, 749.
82. Kisielinski, K.; Giboni, P.; Prescher, A.; Klosterhalfen, B.; Graessel, D.; Funken, S.; Kempinski, O.; Hirsch, O. Is a mask that covers the mouth and nose free from undesirable side effects in everyday use and free of potential hazards? *International journal of environmental research and public health* **2021**, *18*, 4344.
83. Gireesan, S.; Pandit, A.B. Modeling the effect of carbon-dioxide gas on cavitation. *Ultrasonics sonochemistry* **2017**, *34*, 721-728.
84. Lautridou, J.; Buzzacott, P.; Belhomme, M.; Dugrenot, E.; Lafere, P.; Balestra, C.; Guerrero, F. Evidence of Heritable Determinants of Decompression Sickness in Rats. *Med Sci Sports Exerc* **2017**, *49*, 2433-2438, doi:10.1249/MSS.0000000000001385.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.