

Case Report

Not peer-reviewed version

Advancing Physiatriic Care: A Comprehensive Case Study on Laser Therapy and Hydrotherapy Integration Post-Femoral Head and Neck Resection

Miriam Caramico and [Stefania De Dominicis](#) *

Posted Date: 22 July 2024

doi: 10.20944/preprints202407.1702.v1

Keywords: orthopedic rehabilitation physical therapy; orthopedic surgery; osteoarthritis; musculoskeletal system; photobiomodulation; balance and stability; orthopedic pain



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Case Report

Advancing Physiatric Care: A Comprehensive Case Study on Laser Therapy and Hydrotherapy Integration Post-Femoral Head and Neck Resection

Miriam Caramico and Stefania De Dominicis *

Dept. of Veterinary Medicine, University of Teramo 1; sdedominicis@unite.it

* Correspondence: mcaramico@unite.it Tel + 39 3288992240

Simple Summary: Hip dysplasia is a common hereditary condition in dogs and cats characterized by inadequate hip joint development. This condition can be managed conservatively or surgically. It has been observed that after surgical treatment, physiotherapy is crucial for these patients. Physiotherapy helps in faster recovery and reduces additional complications in gait biomechanics. The rehabilitation plan may vary depending on the case, but the main goals are pain and inflammation control and muscle mass gain. Laser therapy and the use of an aquatic treadmill are effective methods used in achieving these goals.

Abstract: Hip dysplasia is a prevalent hereditary condition affecting dogs and cats, characterized by inadequate development of the hip joint. This condition leads to pain due to capsular distension, microfractures, and joint incongruity in young animals, ultimately progressing to osteoarthritis in adult and elderly animals. The pursuit of pain relief and improved quality of life for individuals with osteoarthritis has been a focal point in several studies. In this context, veterinary physiotherapy has gained prominence in small animal practice, serving as both palliative care and a supportive modality in the clinical or post-surgical management of degenerative joint diseases. Laser therapy and hydrotherapy, recognized for their analgesic effects and capacity to promote muscle mass gain, have garnered attention for their potential efficacy in treating joint diseases. This study aims to evaluate the therapeutic effects of super pulsed laser therapy and controlled low-impact exercises (water treadmill) on a patient recovering from surgery involving bilateral femoral head and forehead resection. We observed positive outcomes, including pain resolution, improved muscle mass, and enhanced range of motion, contributing to the patient's overall recovery.

Keywords: orthopedic rehabilitation physical therapy; orthopedic surgery; osteoarthritis; musculoskeletal system; photobiomodulation; balance and stability; orthopedic pain

1. Introduction

Coxofemoral dysplasia (CFD) is a prevalent orthopedic disease characterized by joint incongruity and subsequent osteoarthritis and pain, influenced by complex interactions of multiple genes and environmental factors impacting disease susceptibility [1,3,8,37]. Although the coxofemoral joint is normal at birth, the imbalance between skeletal and muscular system growth rates during development imposes excessive loads on the femoral head, altering acetabular conformation and leading to irregular joint surface remodeling [6,36,38].

CFD can progress to painful secondary osteoarthritis and affect the dog's behavior, manifesting as chronic lameness, muscle atrophy, and exercise reluctance [2,23,50]. While medical and surgical management can alleviate pain, they do not address underlying skeletal muscle conditions, which necessitates physiotherapy for effective treatment.

Clinical signs of CFD include pain, lameness, stiff gait, and muscle atrophy, with diagnosis relying on comprehensive medical history, clinical evaluation, and imaging [6,10,23]. Manifestations

may remain latent for extended periods, becoming apparent with the onset of degenerative joint disease.

The pathophysiology involves muscle contractures, joint inflammation, and increased intra-articular pressure, contributing to pain and joint dysfunction [3,6,26,28]. Contractures of the adductor muscles, notably the pectineus, can cause microfractures during growth remodeling, exacerbating the condition.

In patients with osteoarthritis, increased intra-articular pressure and subchondral bone ischemia are prominent contributors to pain [3,6,26]. Understanding the multifaceted etiology and clinical manifestations of CFD is essential for targeted therapeutic interventions (Figure 1).

The selection of an appropriate treatment strategy for coxofemoral dysplasia (CFD) hinges upon various factors, including the patient's age, the severity of dysplasia, and the presence of any concurrent pathologies. The primary objectives are to alleviate pain, enhance limb function, and optimize the patient's overall quality of life.

As illustrated in Table 1 below, conventional surgical interventions are inadequate in resolving the complex anatomical alterations associated with CFD. Instead, comprehensive symptom management is achievable through physiotherapy (rehabilitation) complemented by nutraceutical integration. Unlike surgical approaches, physiotherapy addresses the underlying muscular, tendinous, and ligamentous changes associated with CFD [19,29,32,41] (Table 1).

The physiotherapy regimen employed encompassed a multifaceted approach incorporating several specialized techniques tailored to the needs of the patient. These techniques included:

Manual Treatment: Application of massage and stretching protocols to promote joint mobility and tissue flexibility.

Balance and Proprioceptive Exercises: Activities designed to enhance proprioception and stability, crucial for improving limb function and gait.

Therapeutic Exercises: Prescribed exercises targeting specific muscle groups to promote strength and endurance.

Aquatic Treadmill Therapy: Utilization of an aquatic treadmill to facilitate low-impact exercise, fostering muscle conditioning and joint mobilization.

Super-Pulsed Class I Laser Therapy: Utilization of the ACTIVet PRO® device from Multi Radiance Medical (Solon, OH, USA) for laser therapy sessions. This therapy is known for its analgesic and anti-inflammatory effects, aiding in pain management and tissue healing. (table2)

The integration of these modalities within the physiotherapy protocol contributed synergistically to the patient's rehabilitation and recovery process (Table 2).

2. Materials and Methods

2.1. Patient Details

A 7-year-old male Canine Setter was the subject of this study (Figures 2–5).

2.2. History

The canine was discovered abandoned near the municipal kennel and subsequently rescued. Following medical examination and X-rays, the dog was diagnosed with severe coxofemoral dysplasia, necessitating femoral head ostectomy due to joint dislocation. The initial surgery took place in March 2022, followed by a subsequent procedure in August 2022.

2.3. Assessment

In October 2022, the patient presented with pelvic extension pain, gluteal muscle atrophy, quadriceps and femoral biceps contractures, iliopsoas muscle contracture, lumbar thoracic spine pain, and joint capsule retraction. The dog exhibited difficulty in sitting and standing. Goniometry measurements indicated left coxofemoral joint flexion of 48° and extension of 158°, and right coxofemoral joint flexion of 47° and extension of 157° (normal flexion 50°, extension 162°). Thigh perimetry measurements were 28 cm (right) and 29 cm (left).

2.4. Methods

The treatment plan comprised 20 physiotherapy sessions utilizing the ACTIVet PRO® laser (Multi Radiance Medical, Solon, OH, USA) for pain and osteoarthritis management, aquatic treadmill for muscular reinforcement, balance and proprioception exercises, therapeutic exercises, and core strengthening. Additionally, Alevica® was administered for pain control, and Chondrostress® was prescribed to enhance cartilage hydration and synovial fluid viscosity in degenerative diseases such as coxofemoral dysplasia.

Rehabilitation sessions included tecar therapy, super-pulsed laser, and aquatic treadmill exercises administered twice weekly for two months, followed by once-weekly sessions for the final two weeks. Laser therapy aimed to expedite healing, control inflammation and pain, and alleviate muscle contractures. Specific laser treatments targeted thoracolumbar region pain, chronic pain at the surgery site, and pectineus muscle contracture.

Hydrotherapy sessions progressed through different water levels: hip joint level for the first two weeks, mid-femur level for subsequent sessions, and knee-deep water for the remaining sessions. Significant improvements were observed in muscle contracture and joint pressure after the fourth session.

Physiotherapy continued for two months to restore muscle mass, improve joint amplitude, and achieve total pain control. Condrostress® supplement C was continued indefinitely to manage degenerative joint disease.

3. Results

In the last evaluation, the patient exhibited no signs of pain and demonstrated ease in sitting and standing. The post-surgery treatment effectively restored the patient’s function and quality of life. Radiographic assessment revealed no developmental signs of osteoarthritis and an increase in muscle mass. Thigh perimetry measurements showed improvement, with the right thigh measuring 30 cm and the left thigh measuring 31 cm.

3.1 Figures and Tables




Figure 1. Hyp dysplasia.



Figure 2. Pre surgery x ray.




Figure 3. Pre surgery x ray.



Figure 4. X-ray after the first surgery.



Figure 5. X-ray after the second surgery.



Figure 6. Post-physiotherapy X-ray.



Figure 7. Use of images authorized by the guardian.

Table 1. comparative treatment.

Treatment x symptoms	Resection of the head and femoral strain	Denervation	Prosthes is	Rehabilitation
Pain in extension, adduction and flexion of the joint	-	+	+	+
Intermittent claudication paresis of hind limbs	-/+	+	+	+
Intra-articular pressure	+	-	+	+
Pectineus and iliopsoas contracture	-	-	-	+
Hypotrophy of the gluteus, quadriceps and biceps femoris	-	-/+	-	+
Pain in the sacral loin and lumbar thoracic spine.	-	-	-	+
Muscle and joint capsule pain.	-	+/-	-/+	+
Inflammation and retraction of the joint capsule.	+	-	+	+

Table 2. This table shows the characteristics of the Laser Active Pro.

Feature	Specification
Laser radiation	905 nm
Broadband infrared radiation	860 nm
Visible red light radiation	660 nm
Laser radiation peak pulse power	50W
Laser pulse duration	110±20 ns
Broadband infrared radiation average power	250
Red light average power	100 mW
Radiation aperture	4±0.4 cm²
Magnetic induction	35±10 mT

Table 3. This table shows the perimetry pre and post treatment.





Table 4. This table shows the goniometry pre and post treatment.



4. Discussion

The evolution of rehabilitation has brought forth substantial scientific evidence supporting the effectiveness of non-surgical interventions for treating hip dysplasia, particularly in cases where there is no joint luxation. This case report underscores the necessity of rehabilitative measures even following corrective surgery, given that surgical techniques such as osteotomy, denervation, or prosthesis placement do not address soft tissue alterations (e.g., muscles, joint capsule). Consequently, it is essential to integrate pre- and post-surgery physiotherapy into all hip dysplasia treatment protocols to facilitate enhanced and accelerated patient recovery.

5. Conclusions

Recent studies demonstrate the efficacy of various physical therapeutic modalities in the conservative or post-surgical treatment of canine coxofemoral dysplasia (FCD) [28,32,48]. Physiatry exerts biological effects through physical mechanisms, eliciting biochemical responses in the body. Laser therapy is recommended for its analgesic effects [12,39,40], and low-impact exercises on a hydro treadmill [17,18,20,47] are beneficial for individuals with osteoarthritis.

Physical exercises on an aquatic treadmill contribute to weight control, a crucial factor in therapeutic management to avoid joint overload [21,24]. The physical properties of water decrease the relative weight of the animal due to buoyancy, and the increased density challenges muscles more than air [20,42]. These properties make hydro treadmill and swimming ideal exercises for animals with arthropathy, providing greater energy expenditure and less joint impact [11], and they are particularly important in the multimodal treatment of canine obesity in dogs with osteoarthritis [11,21,42,56].

Regular treadmill exercise enhances the release of beta-endorphins, improves self-esteem, and aids in pain relief in humans [11]. Further studies are necessary to elucidate the effects of aquatic physical exercises on dogs in relation to the release of endogenous substances and well-being. Nonetheless, the thermal effect of heated water is known to promote relaxation and improve joint mobility in dogs [57].

Laser therapy is effective in controlling joint pain and chronic disease in both humans [12,15,16] and dogs [38,39]. Its analgesic effects occur through photo biomodulation of the inflammatory response, increasing cellular ATP synthesis and transforming prostaglandins and prostacyclin, thereby exerting anti-edematous and anti-inflammatory effects [16]. Laser therapy also increases the release of endorphins, providing relief from non-inflammatory pain [45].

To promote pain relief, improve joint movement amplitude, and increase muscle mass through physiatry modalities, it is essential to employ appropriate assessment methods for each parameter. In this case report, subjective assessments by the owner, goniometry, and perimetry of the coxofemoral joint and associated muscles were utilized [22,44–46].

In this specific case, the owner reported substantial improvement, indicating that the patient is no longer hesitant to walk, sit, or stand

And in the measurements [13]

1. Goniometry of the left coxofemoral joint pre-treatment: flexion 48° extension 158° (normal flexion 50° extension 162°) Post treatment flexion 50° extension 160°
2. Goniometry of the right coxofemoral joint pre-treatment: flexion 47° extension 157° (normal flexion 50° extension 162°) Post treatment flexion 50° extension 159°
3. Pre-treatment perimeter left thigh 28 cm left thigh 29 cm
4. Post treatment Right thigh perimetry 30 cm; left thigh 31 cm (Tables 3 and 4)

Based on our findings, we conclude that hydrotherapy and laser therapy offer effective means to address muscle contractures, joint extension, and support strength, ultimately enhancing the quality of life for our patients. Importantly, these therapies pose minimal risk or contraindication, as the use of super pulsed laser does not result in burns or abrasions on the patient's skin and does not require shaving. To further elucidate these benefits, continued studies and comparative analyses with cases of hip dysplasia, with or without surgical intervention, are warranted [27–29].

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Acknowledgments: Thank you for the support from the University of Teramo and for the support and assistance from Multi Radiance. I thank the reviewers from Multi Radiance for their valuable assistance in reviewing this manuscript

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Bettini CM, et al. Incidence of coxofemoral dysplasia in Border Collie dogs. *Unipar Veterinary and Zoological Sciences Archive*. 2007;10(1):21-25.
2. McLaughlin Junior R, Tomlinson J. Radiographic diagnosis of canine hip dysplasia. *Veterinary medicine*. 1996;91(2):36-47.
3. Minto BW, et al. Clinical evaluation of acetabular denervation in dogs with coxofemoral dysplasia treated at the FMVZ Veterinary Hospital - Botucatu - SP. *Veterinary and Animal Sciences*. 2012;19(1):91-98.
4. Mueller M, et al. Effects of radial shock wave therapy on limb function of dogs with osteoarthritis of the hip. *Veterinary record*. 2007;160(1):762–765.
5. Morgan JP, Wind A, Davidson AP. Elbow dysplasia. In: *Hereditary diseases of bones and joints in dogs: osteochondrosis, hip dysplasia, elbow dysplasia*. Hanover: Schlütersche; 2000. p. 41-94.
6. Mostafa AA, Berry CR, Nahla MA. Quantitative evaluation of hip morphology to improve the identification of coxofemoral dysplasia in German Shepherd dogs. *Am J Vet Res*. 2023 Feb;7:1-10. PMID: 36729899.
7. Piermattei D. Arthroplasty for excision of the coxofemoral joint in dogs and cats. *Vet Comp Orthop Traumatol*. 2011;24(1):89. PMID: 21274500.
8. Rocha FPC, et al. Coxofemoral dysplasia in dogs. *Revista Cientí Eletrônica de Medicina Veterinária*. 2008;4(11):1-7.
9. Off W, Matis U. Arthroplasty for excision of the coxofemoral joint in dogs and cats. Results of clinical, radiographic and gait analysis of the Department of Surgery, Faculty of Veterinary Medicine, Ludwig-Maximilians-University of Munich, Germany. *Vet Comp Orthop Traumatol*. 2010;23(5):297-305. PMID: 20945541.
10. Virag Y, Gumpenberger M, Tichy A, Lutonsky C, Peham C, Bockstahler B. Center of pressure and reaction forces of the soil in Labradors and Golden Retrievers with and without coxofemoral dysplasia at 4, 8 and 12 months of age. *Front Vet Sci*. 2022 Dec 22;9:1087693. PMID: 36619950; PMCID: PMC9814507.
11. Reusing MSO, Villanova Júnior JA, Weber SH. Efeitos terapêuticos do exercício em esteira aquática e da laserterapia de baixa intensidade em cães com displasia coxofemoral = Therapeutic effects of underwater treadmill exercise and low-level laser therapy in dogs with hip dysplasia. 2019. Available at: link. Accessed July 16, 2023.

12. Ali Q, et al. Low Energy Laser Effects in Relieving Pain in Patients with Knee Osteoarthritis. *International Journal of Medicine and Public Health*. 2018;2(1):6-10.
13. Roush, J.K.; Unis, M.D.; Wodiske, T.; Baker, S.G. Comparison of four commercial devices to measure limb circumference in dogs. *Veter- Comp. Orthop. Traumatol*. **2010**, *23*, 406–410, <https://doi.org/10.3415/vcot-10-03-0032>.
14. Barr, A.R.S.; Denny, H.R.; Gibbs, C. Clinical hip dysplasia in growing dogs: the long-term results of conservative management. *J. Small Anim. Pr*. **1987**, *28*, 243–252, <https://doi.org/10.1111/j.1748-5827.1987.tb03879.x>.
15. Bender, T.; Nagy, G.; Barna, I.; Tefner, I.; Kádas, .; Géher, P. The effect of physical therapy on beta-endorphin levels. *Eur. J. Appl. Physiol*. **2007**, *100*, 371–382, <https://doi.org/10.1007/s00421-007-0469-9>.
16. Bjordal, J.M.; Couppé, C.; Chow, R.T.; Tunér, J.; Ljunggren, E.A. A systematic review of low level laser therapy with location-specific doses for pain from chronic joint disorders. *Aust. J. Physiother*. **2003**, *49*, 107–116, [https://doi.org/10.1016/s0004-9514\(14\)60127-6](https://doi.org/10.1016/s0004-9514(14)60127-6).
17. Bockstahler, B.; Kräutler, C.; Holler, P.; Kotschwar, A.; Vobornik, A.; Peham, C. Pelvic Limb Kinematics and Surface Electromyography of the Vastus Lateralis, Biceps Femoris, and Gluteus Medius Muscle in Dogs with Hip Osteoarthritis. *Veter- Surg*. **2011**, *41*, 54–62, <https://doi.org/10.1111/j.1532-950x.2011.00932.x>.
18. Bockstahler, B.A.; Prickler, B.; Lewy, E.; Holler, P.J.; Vobornik, A.; Peham, C. Hind limb kinematics during therapeutic exercises in dogs with osteoarthritis of the hip joints. *Am. J. Veter- Res*. **2012**, *73*, 1371–1376, <https://doi.org/10.2460/ajvr.73.9.1371>.
19. Brown, D.C.; Boston, R.C.; Coyne, J.C.; Farrar, J.T. Ability of the Canine Brief Pain Inventory to detect response to treatment in dogs with osteoarthritis. *J. Am. Veter- Med Assoc*. **2008**, *233*, 1278–1283, <https://doi.org/10.2460/javma.233.8.1278>.
20. Cartlidge H. Hydrotherapy for the osteoarthritic dog: why might it help and is there any evidence? *The Veterinary Nurse*. 2015;6(10):600-606.
21. Chauvet, A.; Laclair, J.; A Elliott, D.; German, A.J. Incorporation of exercise, using an underwater treadmill, and active client education into a weight management program for obese dogs. *Can. Vet. J*. **2011**, *52*, 491–6.
22. Collard F, et al. Canine hip denervation: Comparison between clinical outcome and gait analysis. *Revue de Medecine Veterinaire*. 2010;161(6):277.
23. Crestani MV, Telöken MA, Gusmão PDF. Impacto femoroacetabular: uma das condições precursoras da osteoartrose do quadril. *Rev Bras Ortop*. 2006;41(8):285-293.
24. Denning, W.E.; Bressel, E.; Dolny, D.G. Underwater Treadmill Exercise as a Potential Treatment for Adults With Osteoarthritis. *Int. J. Aquat. Res. Educ*. **2010**, *4*, 9, <https://doi.org/10.25035/ijare.04.01.09>.
25. Diniz R. Hidroterapia. In: Lopes RS, Diniz R, editors. *Fisiatria em Pequenos Animais*. São Paulo: Editora Inteligente; 2018. p. 156-162.
26. Doust S, et al. Preliminary study of the hip dysplasia incidence based on clinical and radiographical examination in large breed dogs referred to veterinary teaching hospital of Ferdowsi University of Mashhad. *Journal of Veterinary Research*. 2018;73(1).
27. Duff, R.; Campbell, J. Long term results of excision arthroplasty of the canine hip. *Veter- Rec*. **1977**, *101*, 181–184, <https://doi.org/10.1136/vr.101.10.181>.
28. Edge-Hughes, L. Hip and Sacroiliac Disease: Selected Disorders and Their Management with Physical Therapy. *Clin. Tech. Small Anim. Pr*. **2007**, *22*, 183–194, <https://doi.org/10.1053/j.ctsap.2007.09.007>.
29. Ginja, M.; Silvestre, A.; Gonzalo-Orden, J.; Ferreira, A. Diagnosis, genetic control and preventive management of canine hip dysplasia: A review. *Veter- J*. **2009**, *184*, 269–276, <https://doi.org/10.1016/j.tvjl.2009.04.009>.
30. Greshake, R.J.; Ackerman, N. ULTRASOUND EVALUATION OF THE COXOFEMORAL JOINTS OF THE CANINE NEONATE. *Veter- Radiol. Ultrasound* **1993**, *34*, 99–104, <https://doi.org/10.1111/j.1740-8261.1993.tb01516.x>.
31. Gusi, N.; Tomas-Carus, P.; Häkkinen, A.; Häkkinen, K.; Ortega-Alonso, A. Exercise in waist-high warm water decreases pain and improves health-related quality of life and strength in the lower extremities in women with fibromyalgia. *Arthritis Care Res*. **2006**, *55*, 66–73, <https://doi.org/10.1002/art.21718>.
32. Harper TA. Conservative Management of Hip Dysplasia. *Veterinary Clinics: Small Animal Practice*. 2017;47(4):807-821.
33. Henrigson, B.; Norberg, I.; Olssons, S. On the Etiology and Pathogenesis of Hip Dysplasia: a Comparative Review. *J. Small Anim. Pr*. **1966**, *7*, 673–688, <https://doi.org/10.1111/j.1748-5827.1966.tb04393.x>.
34. Honmura, A.; Yanase, M.; Obata, J.; Haruki, E. Therapeutic effect of Ga-Al-As diode laser irradiation on experimentally induced inflammation in rats. *Lasers Surg. Med*. **1992**, *12*, 441–449, <https://doi.org/10.1002/lsm.1900120414>.
35. Jaegger, G.; Marcellin-Little, D.J.; Levine, D. Reliability of goniometry in Labrador Retrievers. *Am. J. Veter- Res*. **2002**, *63*, 979–986, <https://doi.org/10.2460/ajvr.2002.63.979>.
36. Cardinet, G.H.; Kass, P.H.; Wallace, L.J.; Guffy, M.M. Association between pelvic muscle mass and canine hip dysplasia. *J. Am. Veter- Med Assoc*. **1997**, *210*, 1466–1473, <https://doi.org/10.2460/javma.1997.210.10.1466>.

37. Kyriazis A. Canine hip dysplasia. Part I: Aetiopathogenesis & diagnostic approach. *Hellenic Journal of Companion Animal Medicine*. 2016;5(1):37.
38. König HE, Liebich H-G. *Anatomia dos Animais Domésticos: Texto e Atlas Colorido*. Artmed Editora; 2016.
39. Lewis G, Holt S. The Effects of Low-level Laser Therapy on the Gait of the Osteoarthritic Canine Hindlimb. *BVNA Congress 2018*. 2018.
40. Matera JM, Tatarunas AC, Oliveira SM. Uso do laser arseneto de gálio (904nm) após excisão artroplástica da cabeça do fêmur em cães. *Acta Cirúrgica Brasileira*. 2003;18(2):102-106.
41. McCarthy, G.; O'donovan, J.; Jones, B.; McAllister, H.; Seed, M.; Mooney, C. Randomised double-blind, positive-controlled trial to assess the efficacy of glucosamine/chondroitin sulfate for the treatment of dogs with osteoarthritis. *Veter- J*. **2007**, *174*, 54–61, <https://doi.org/10.1016/j.tvjl.2006.02.015>.
42. Mendes S, Coutinho I, Rebelo P. Hidroterapia canina. *Revista Portuguesa de Ciências Veterinárias*. 2015;110(595-596):160-164.
43. Millis, D.L.; Levine, D. The Role of Exercise and Physical Modalities in The Treatment of Osteoarthritis. *Veter- Clin. North Am. Small Anim. Pr*. **1997**, *27*, 913–930, [https://doi.org/10.1016/s0195-5616\(97\)50086-6](https://doi.org/10.1016/s0195-5616(97)50086-6).
44. A Morales, J.; Ruiz-Gómez, M.J.; Gil-Carmona, L.; Souvirón, A.; Martínez-Morillo, M. He-Ne laser has no effect on cell cycle phases of human colon adenocarcinoma cells. *Rev. Esp. Fisiol*. **1995**, *51*, 43–7.
45. Pluhar, G.E.; Nielsen, C. Diagnosis and treatment of hind limb muscle strain injuries in 22 dogs. *Veter-Comp. Orthop. Traumatol*. **2005**, *18*, 247–253, <https://doi.org/10.1055/s-0038-1632962>.
46. Pereira, F.C.; Parisi, J.R.; Maglioni, C.B.; Machado, G.B.; Barragán-Iglesias, P.; Silva, J.R.T.; Silva, M.L. Antinociceptive effects of low-level laser therapy at 3 and 8 j/cm² in a rat model of postoperative pain: possible role of endogenous Opioids. *Lasers Surg. Med*. **2017**, *49*, 844–851, <https://doi.org/10.1002/lsm.22696>.
47. Prankel, S. Hydrotherapy in practice. *Practice* **2008**, *30*, 272–277, <https://doi.org/10.1136/inpract.30.5.272>.
48. Raghuvir H, et al. Treatment of canine hip dysplasia: A review. *J Anim Sci Adv*. 2013;3(12):589-597.
49. Remedios, A.M.; Fries, C.L. Treatment of canine hip dysplasia: a review. *Can. Vet. J*. **1995**, *36*, 503–9.
50. Syrcle J. Hip Dysplasia: Clinical Signs and Physical Examination Findings. *Veterinary Clinics: Small Animal Practice*. 2017;47(4):769-775.
51. Tai, G.; Tai, M.; Zhao, M. Electrically stimulated cell migration and its contribution to wound healing. *Burn. Trauma* **2018**, *6*, 20, <https://doi.org/10.1186/s41038-018-0123-2>.
52. Upariputti, R.; Vijarnsorn, M.; Niyom, S.; Boonyong, S. Effect of interferential current therapy on ground reaction force in dogs with hip osteoarthritis: A randomized placebo controlled cross-over clinical trial. *Thai J. Veter- Med*. **2018**, *48*, 111–116, <https://doi.org/10.56808/2985-1130.2879>.
53. Vince, K.J. Canine hip dysplasia: surgical treatment for the military working dog. *Army Med. Dep. J*. **2007**, 44–50.
54. Wallace, L.J. Pectineus Tendon Surgery for the Management of Canine Hip Dysplasia. *Veter- Clin. North Am. Small Anim. Pr*. **1992**, *22*, 607–621, [https://doi.org/10.1016/s0195-5616\(92\)50059-6](https://doi.org/10.1016/s0195-5616(92)50059-6).
55. Weigel, J.P.; Cartee, R.E.; Marich, K.W. Preliminary study on the use of ultrasonic transmission imaging to evaluate the hip joint in the immature dog. *Ultrasound Med. Biol*. **1983**, *9*, 371–378, [https://doi.org/10.1016/0301-5629\(83\)90090-x](https://doi.org/10.1016/0301-5629(83)90090-x).
56. Wong E. *Swim to Recovery: Canine Hydrotherapy Healing*. Veloce Publishing Ltd; 2011.
57. Zink C, Carr BJ. What is a canine athlete? *Canine sports medicine and rehabilitation*. 2018;1-22.
58. Zink C, Van Dyke JB. *Canine sports medicine and rehabilitation*. John Wiley & Sons; 2018.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.