

Brief Report

Not peer-reviewed version

Novel *De Novo* *RAI1*:c.2736delC Variant Identified in A Child with Features of Smith-Magenis: Insight from Whole Genomes Joint Analysis

[Mario Cuk](#) , [Busra Unal](#) , [Nives Jandric](#) , Connor P. Hayes , McKenzie Walker , [Kristina Crkvenac Gornik](#) , [Feruza Abraamyan](#) , [Arezou A. Ghazani](#) *

Posted Date: 26 July 2024

doi: 10.20944/preprints202407.2052.v1

Keywords: Smith-Magenis syndrome; retinoic acid-induced 1 protein; neurodevelopmental disorders; joint whole genome sequencing analysis



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Brief Report

Novel *De Novo* *RAI1*:c.2736delC Variant Identified in A Child with Features of Smith-Magenis: Insight from Whole Genomes Joint Analysis

Mario Cuk ^{1,†}, Busra Unal ^{2,†}, Nives Jandric ¹, Connor P. Hayes ², McKenzie Walker ²,
Feruza Abraamyan ², Kristina Crkvenac Gornik ³ and Arezou A. Ghazani ^{2,4,5,*}

¹ Department of Pediatrics, University Hospital Centre Zagreb and University of Zagreb, School of Medicine, Zagreb, Croatia

² Division of Genetics, Brigham and Women's Hospital, Boston, MA, USA

³ Department of Laboratory Diagnostics, Division of Cytogenetics, University Hospital Centre Zagreb, Zagreb, Croatia

⁴ Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA

⁵ Harvard Medical School, Boston, MA, USA

* Correspondence: aghazani@bwh.harvard.edu

† These authors contributed equally.

Abstract: Smith-Magenis syndrome is a complex neurobehavioral genetic disorder. The spectrum of SMS phenotype is broad but commonly includes craniofacial, neurobehavioral, and otolaryngologic features. While the etiology of SMS is commonly attributed to one copy interstitial deletion in the 17p11.2 region (90-95% of cases), variants identified by sequence analysis in *RAI1* have also been reported in 5-10% of cases. The phenotypic spectrum of patients with *RAI1* sequence alterations is not clear. In this study, we report a 9-year-old male with global cognitive and psychomotor developmental delay, musculoskeletal and cardiovascular abnormalities, and dysmorphic craniofacial features. Whole genome quad analysis of the proband, unaffected parent, and unaffected brother identified a novel *de novo* *RAI1*:c.2736delC variant. This is the first report of this variant in the literature. This report also highlights the details of the genome analysis and this patient's phenotypic spectrum.

Keywords: Smith-Magenis syndrome; retinoic acid-induced 1 protein; neurodevelopmental disorders; joint whole genome sequencing analysis

1. Introduction

Smith-Magenis syndrome (SMS; OMIM #182290) is a complex neurobehavioral disorder with an estimated prevalence of 1:15,000. This rare syndrome is characterized by neurodevelopmental delay, coarse facial features evolving with age, and maladaptive behaviors that include prolonged temper tantrums and self-injurious behaviors, including hand biting, trichotillomania, and polyembolokoilomania [1]. During infancy, hypotonia, hyporeflexia, and feeding difficulties are commonly seen neurological findings [2–5]. The maladaptive behavior phenotype is typically recognized after 18 months of age. The classic neurological features include intellectual disability, delayed speech, sensory integration problems, decreased pain sensitivity, and abnormalities in sleep patterns [1]. Additional system involvements, such as musculoskeletal and cardiovascular, are also reported as part of SMS [2–6].

The underlying genomic etiology of SMS is interstitial deletion at 17p11.2 in 90-95% of patients. This genomic location encompasses the retinoic acid-induced 1 (*RAI1*) gene. *RAI1* is the key gene in SMS and plays a regulatory role in healthy skeletal and nervous system development, behavioral maturation, and circadian rhythm. Pathogenic *de novo* single nucleotide variants (SNVs) in *RAI1* are

reported in 5-10% of patients with SMS [2,5–7]. In this category, missense, nonsense, and splice site variants in *RAI1* have been reported [4]. The diagnosis of the SMS syndrome is based on clinical molecular diagnostics tests. The first-tier molecular test is chromosomal microarray analysis (CMA), and if negative, a sequencing-based test is typically performed. Genome studies have reported SMS-like features, including ID, sleep disturbances, and self-injuries behaviors in patients without an SMS molecular diagnosis who harbored deleterious variants in *KMTD2*, *MECP2*, *KDM5C*, *IQSEC2*, and *DEAF1* genes [8,9].

Here, we report a 9-year-old male with global cognitive and psychomotor developmental delay and facial, musculoskeletal, and cardiovascular abnormalities. The patient's initial chromosomal microarray and karyotype tests were normal. The quad joint WGS analysis of the proband, parents, and brother detected a pathogenic novel *de novo* variant, which has not been reported previously. This report highlights the importance of sequencing analysis in patients clinically suspected of SMS and with negative conventional diagnostic tests. Additionally, sharing the genotype and detailed phenotype of these relatively rare SMS cases helps understand the genetic and phenotypic spectrum of the disease.

2. Material and Methods

2.1. Participants

The index patient, a 9-year-old male, his parents, and his younger brother were enrolled in the CROseq genome program. The CROseq Genome Program is a collaborative research program between Brigham and Women's Hospital (BWH) (Boston, USA), and the Department of Pediatrics, University Hospital Centre Zagreb (Zagreb, Croatia), supported by the Mila Za Sve Foundation (Rijeka, Croatia). This program aims to investigate the genomic etiology of disease in patients with complex phenotypes and negative conventional genetic tests. Patient consent, enrollment, and clinical assessment were performed at the University Hospital Center Zagreb.

2.2. Clinical Assessment of Brain Structure and Activity

The brain abnormalities were assessed by MRI scanning (MAGNETOM Trio, A Tim System 3T eco, Zagreb, Croatia). The patient was under general anesthesia during the scan. Age-adjusted repetition time and time-to-echo were applied to create multiplanar T1- and T2-weighted images. The electroencephalogram (Nihon Kohden, Neurofax EEG-1200K; Zagreb, Croatia) examination was performed to evaluate the brain activity. During the record, the sampling rate was 200-10000 Hz. The patient's forehead is cleaned with alcohol to lower the impedance less than 5 Ω for each electrode. The electrodes were set up in FPz-F9 and FPz-AF7 of the international 10-20 system, which focuses on the brain in the same hemisphere.

2.3. Traditional Genetic Testing

Karyotyping was performed to analyze chromosomal abnormalities according to standard procedures at University Hospital Centre Zagreb. Briefly, the lymphocytes were purified from the peripheral blood sample. Cells were arrested at metaphase and fixed before being harvested for slide preparation. The number and morphology of chromosomes were assessed in at least 20 metaphases. The karyotype result was reported per The International System for Human Cytogenomic Nomenclature (ISCN) 2020.

Chromosome microarray (CMA) was performed according to standard procedures at University Hospital Centre Zagreb. Briefly, genomic DNA was isolated with the FlexiGene DNA Kit (Qiagen) and CMA was performed using Agilent 4x180k aCGH+SNP array (Agilent Technologies, Santa Clara, CA, USA). The genomic DNA of the test and control samples were labeled with fluorescent dyes and hybridized to the array. The array was scanned with a DNA Microarray Scanner (Agilent Technologies, Santa Clara, CA, USA). Data were obtained using the "CytoGenomics 5.1.1.15" software (Agilent Technologies, Santa Clara, CA, USA) to generate the final plots.

2.4. Sample Preparation and Whole Genome Sequencing (WGS)

DNA extraction and WGS were performed at the Medical College of Wisconsin (Milwaukee, USA). Genomic DNA was isolated from peripheral blood samples (2ml) with a purity ratio of 1.75 – 2.0. After robotic DNA library construction, sequencing was conducted on the NovaSeq 6000 platform, with an average depth of 40X.

2.5. Quad Joint Whole Genome Analysis

Genome analyses were performed at the BWH, USA. The quad whole genome analysis was performed on proband, unaffected parents, and the unaffected proband's brother. Following the quad analysis, a phenotype-based analysis was performed using human phenotype ontology (HPO) terms of the patient's indications. Variants were prioritized by the gene's association with the phenotype and variant type. The HPO terms were: Neurodevelopmental delay (HP:0012758), moderate intellectual disability (HP:0002342), delayed speech and language development (HP:0000750), attention deficit hyperactivity disorder (HP:0007018), bicuspid aortic valve (HP:0001647) and stenosis (HP:0001650), failure to thrive (HP:0001508), abnormal facial shape (HP:0001999), unilateral ptosis (HP:0007687), frontal bossing (HP:0002007), relative macrocephaly (HP:0004482), abnormality of mitochondrial metabolism (HP:0003287).

All variants across the genome were included in the investigation. Following the technical assessment, medium, high, and very high-quality variants were further evaluated, and low-quality variants were excluded. The variant frequencies were annotated by gnomAD exome and genome population allele frequencies. The in silico prediction algorithms for missense variants included CADD, REVEL, Polyphen, SIFT, MutationTaster, Mutation Assessor, FATHMM, FITCONS, GENOCANYON, dbSNV ADA, and dbSNV RF. The spliceAI prediction score was used in the evaluation of splice variants.

Zygosity analysis was performed by identifying and evaluating compound heterozygous and homozygous variants in the proband. The recessive variants in the proband inherited from unaffected parents and related to the patient's HPOs were assessed. Separately, *de novo* analysis was performed to assess variants present in the proband and related to the patient's HPOs that are absent in the unaffected parents and the brother.

The variant classification was performed according to ACMG-AMP guidelines (25741868). In the attribution of PM2 and BS1 criteria, the gene-specific threshold was applied based on gnomAD (v2.1.1) aggregated allele frequency. The aggregated in silico prediction score was indicative of a deleterious effect if it was higher than 0.7; scores less than 0.15 were assigned for a benign effect. The genes' tolerance to the loss of function variants was evaluated by using gene constraint metrics of pLI=1 and/or o/e<0.35.

3. Results

3.1. Clinical Description

The proband is a 9-year-old male presented with indications of global cognitive and psychomotor developmental delay, musculoskeletal and cardiovascular abnormalities, and dysmorphic craniofacial features. During his infancy, the patient had generalized hypotonia and feeding difficulties. His brain ultrasound, which was performed at four months of age, was normal. Following intensive physical therapy, he started walking at two years old. The psychomotor assessment revealed that the patient has severe psychomotor retardation. The brain MRI performed at 3.5 years of age showed perinatal hypoxic-ischemic lesions. The electroencephalogram was unremarkable (data not shown).

The detailed clinical evaluation at 9 years of age indicated brachycephaly, broad face, hypertelorism, wide nasal bridge, short and full-tipped nose, midface retrusion, deeply set ears, tented and down-turned upper lip, everted upper lip vermillion, micrognathia, and low hairline (Figure 1). He exhibited global cognitive and psychomotor developmental delay, severe ID, and developmental language disorder. The pervasive developmental disorder was evident with the lack

of speech development and poor contact. Behavioral phenotypes included aggression toward others self-injurious episodes, and polyembolokoilamania. He did not acquire sphincter control and exhibited incontinence. He had sleep disturbances, including night awakening, with difficulty falling back asleep. There were skeletal abnormalities including mild scoliosis, thoracic kyphosis, short palm and foot, brachydactyly, and partial syndactyly of both hands' second and third fingers (Figure 1). He reportedly had a bicuspid aortic valve with mild stenosis and congenital left eyelid ptosis.



Figure 1. Dysmorphic features of the proband at the age of 9 years. A) Brachycephaly, broad face, hypertelorism, wide nasal bridge, midface retrusion, deeply set ears, tented and down-turned upper lip, everted upper lip vermillion, light-colored hair; B) Brachycephaly, prominent forehead, short and full-tipped nose, deeply set ears, midface retrusion, micrognathia, tented and down-turned upper lip, everted upper lip vermillion, light-colored hair; C) Short foot, brachydactyly; D) Short palm, brachydactyly.

3.2. Genomic Findings

CMA and Karyotype Analysis

Karyotype testing was 46, XY. CMA analysis of the proband detected a maternally inherited 2.1 Mb duplication in 22q11.21 (Supplementary Figure S1 and S2). This region contains OMIM genes *SLC25A1*, *CDC45*, *GP1BB*, *TBX1*, *TXNRD2*, *COMT*, *TANGO2*, *RTN4R*, *SCARF2*, *PI4KA*, *SERPIND1*, *SNAP29*, and *LZTR1*. The mother is unaffected, therefore, this 22q11.21 gain was considered unrelated to the patient’s clinical findings of SMS. The patient was enrolled in the CROseq genome program, and a quad WGS joint analysis was performed.

Quad WGS Joint Analysis

After WGS, a quad-WGS analysis of the proband, unaffected brother, and parents was performed. HPO-based analysis revealed 10,290 variants in 2,567 genes. Compound heterozygous analysis interrogated possible causative variants inherited from the unaffected parents. Quad analysis did not identify deleterious homozygous or compound heterozygous candidate variants in the proband.

De novo analysis interrogated variants related to the HPO that are present in the proband, but absent in the parents and the unaffected sibling. This analysis identified the variant NM_030665.4 (*RAI1*):c.2736delC (p.Gly913Alafs*37), located in exon 3 of *RAI1* (Figure 2). This variant is a 1bp deletion that results in a frameshift and a premature stop codon. Due to its location, this variant likely results in a premature translational stop signal and nonsense-mediated mRNA decay. The gene constraint metrics for *RAI1* are pLI = 1, o/e = 0.04. The variant has not been reported in gnomAD, ClinVar, or any published literature. The phasing analysis by quad joint WGS showed that the parents and younger brother did not harbor this variant. No other deleterious variants in genes related to the patient’s HPO list were identified.




Figure 2. Lollipop plot of the pathogenic and likely pathogenic *RAI1* frameshift variants. The plot shows the p.G913fs variant identified in the proband in this study and the distribution of frameshift *RAI1* variants reported in the ClinVar database to date (n=30). The Y axis shows the variant number, and the X axis illustrates the position of amino acid residues along the *RAI1* protein. It includes exon 3 and exon 4, the green box denotes the PHD-like zinc-binding domain.

Discussion

Smith-Magenis Syndrome, also known as 17p11.2 microdeletion syndrome, was first discovered in 1986 [10]. Since then, the genomic and clinical spectrum of more than 500 SMS patients has been reported [11]. Most cases of SMS are caused by microdeletion in 17p11.2, while 5-10% of cases are due to pathogenic *RAI1* sequence variants [2]. When the phenotypic findings suggest SMS, the first molecular testing approach is generally CMA, followed by a single gene or multipanel sequence analysis that includes *RAI1*.

The *RAI1* has a regulatory role in embryogenesis through the transcriptional regulation of many genes [12–14]. These include the skeletal development genes *PSTPIP2* and *ANGH* [15]; lipid metabolism genes *LIPE*, *HMGCS1*, and *INSIG1* [15]; neurological development genes *ZIC1*, *PSEN2*, *RXR*, *CLN8*, *SMA4*, *NF1*, and *KMT2A*; behavioral function gene *SCN12A* 19236431 [15]; circadian activity genes *NR1D2*, *PER2*, *PER3*, *CRY1*, and *ARNTL* 22578325; cellular growth and cell cycle

regulation genes *SPTBN1*, *POLDIP3*, *PPP1R14D*, *GLI3*, *KMT2A*, and *ADD3* [15]; and insulin regulation genes *INSIG1*, *PIK3R1*, *ZNF236*, and *LIPE* [15]. As a result, patients with SMS commonly have moderate to severe neurodevelopmental and behavioral abnormalities, metabolic problems, sleep problems, and skeletal abnormalities [3].

Here, we report a patient with a spectrum of Smith-Magenis syndrome and the pathogenic *de novo* novel NM_030665.4 (*RAI1*):c.2736delC variant. This variant is located in exon 3 (of 6 exons). Most reported pathogenic *RAI1* variants are located in exon 3 [4] which encodes nearly 98% of the protein. There are reports of mutational hot spots in this exon [16]. The *RAI1*-encoded nuclear protein has a zinc finger homology structural domain, which is indicative of the transcription regulatory activity of *RAI1* [17]. The truncating variants lead to aberrant cytoplasmic subcellular localization and, therefore, the inability of transcription activation [18]. To date, there are 78 frameshift, 29 nonsense, and 4 missense variants reported as pathogenic or likely pathogenic in ClinVar. Most of these variants, including the (*RAI1*):c.2736delC variant in our case, are located in exon 3 (Figure 2).

The broad phenotypical variation of SMS stems from the genetic pathophysiology of the genomic region. Patients with the 17p11.2 microdeletion are more likely to have severe cognitive impairment, hearing loss, cardiac abnormalities, and hypotonia. These features are associated with the deletion of genes residing within the region [19]. Patients with normal copy17p number who carry a deleterious sequence variant in *RAI1* reportedly exhibit polyembolokoilamania (inserting foreign objects into orifices), skin picking, and self-hugging behavior [19]. Consistent with these reports, our patient exhibited these behaviors. Our patient also exhibited severe intellectual disability and cardiac abnormalities, as well as hypotonia during infancy. The joint analysis did not reveal any additional alterations in the genome consistent with these phenotypes.

In conclusion, we report a detailed clinical description of a patient with SMS and the novel pathogenic *RAI1* sequence variant. Studies reporting clinical findings of SMS and pathogenic *RAI1* variants help expand the understanding of the genetic pathophysiology of this complex syndrome.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org. Figure S1: Chromosomal microarray of the proband. Interstitial duplication of the long arm of chromosome 22 in the region 22q11.21 (18996947_21086225), 2.1 Mb in size (arr[GRCh38] 22q11.21(18996947-21086225)x3 mat). Figure S2: Chromosomal microarray of the mother. Interstitial duplication of the long arm of chromosome 22 in the q11.21 region (18996947-21454704), 2.46 Mb in size (arr[GRCh38]22q11.21 (18996947_21454704)x3).

Funding: This investigation was done in collaboration with Brigham and Women's Hospital, Inc/University Hospital Centre Zagreb, with funding support provided by Mila za Sve Foundation.

Institutional Review Board Statement: Institutional Review Board at BWH and University Hospital Centre Zagreb approved this study (Class: 8.1-21/6-2; Reg. No.: 02/21 AG). Written informed consent was obtained from all the participants at the University Hospital Centre Zagreb, Croatia. This study is compliant with the General Data Protection Regulation (GDPR), approved by BWH, Mila Za Sve Foundation, and the University Hospital Centre Zagreb, Croatia.

Data Availability Statement: Data for this manuscript are subject to BWH institutional and GDPR privacy policies and restricted from inclusion in repositories. They may be available upon request.

Acknowledgments: We thank our patient and their family for their participation in the CROseq Genome Program and for permitting us to anonymously share the clinical details.

References

1. Elsea SH, Girirajan S. Smith-Magenis syndrome. *Eur J Hum Genet.* Apr 2008;16(4):412-21. doi:10.1038/sj.ejhg.5202009
2. Gropman AL, Duncan WC, Smith AC. Neurologic and developmental features of the Smith-Magenis syndrome (del 17p11.2). *Pediatr Neurol.* May 2006;34(5):337-50. doi:10.1016/j.pediatrneurol.2005.08.018
3. Adam MP, Feldman J, Mirzaa GM, et al. GeneReviews. 1993.
4. Rinaldi B, Villa R, Sironi A, Garavelli L, Finelli P, Bedeschi MF. Smith-Magenis Syndrome-Clinical Review, Biological Background and Related Disorders. *Genes (Basel).* Feb 11 2022;13(2)doi:10.3390/genes13020335

5. Yeetong P, Vilboux T, Ciccone C, et al. Delayed diagnosis in a house of correction: Smith-Magenis syndrome due to a de novo nonsense RAI1 variant. *Am J Med Genet A*. Sep 2016;170(9):2383-8. doi:10.1002/ajmg.a.37602
6. Carmona-Mora P, Encina CA, Canales CP, et al. Functional and cellular characterization of human Retinoic Acid Induced 1 (RAI1) mutations associated with Smith-Magenis Syndrome. *BMC Mol Biol*. Aug 25 2010;11:63. doi:10.1186/1471-2199-11-63
7. Slager RE, Newton TL, Vlangos CN, Finucane B, Elsea SH. Mutations in RAI1 associated with Smith-Magenis syndrome. *Nat Genet*. Apr 2003;33(4):466-8. doi:10.1038/ng1126
8. Loviglio MN, Beck CR, White JJ, et al. Identification of a RAI1-associated disease network through integration of exome sequencing, transcriptomics, and 3D genomics. *Genome Med*. Nov 01 2016;8(1):105. doi:10.1186/s13073-016-0359-z
9. Berger SI, Ciccone C, Simon KL, et al. Exome analysis of Smith-Magenis-like syndrome cohort identifies de novo likely pathogenic variants. *Hum Genet*. Apr 2017;136(4):409-420. doi:10.1007/s00439-017-1767-x
10. Smith AC, McGavran L, Robinson J, et al. Interstitial deletion of (17)(p11.2p11.2) in nine patients. *Am J Med Genet*. Jul 1986;24(3):393-414. doi:10.1002/ajmg.1320240303
11. Wolters PL, Gropman AL, Martin SC, et al. Neurodevelopment of children under 3 years of age with Smith-Magenis syndrome. *Pediatr Neurol*. Oct 2009;41(4):250-8. doi:10.1016/j.pediatrneurol.2009.04.015
12. Tahir R, Kennedy A, Elsea SH, Dickinson AJ. Retinoic acid induced-1 (Rai1) regulates craniofacial and brain development in *Xenopus*. *Mech Dev*. Aug 2014;133:91-104. doi:10.1016/j.mod.2014.05.004
13. Bi W, Yan J, Shi X, et al. Rai1 deficiency in mice causes learning impairment and motor dysfunction, whereas Rai1 heterozygous mice display minimal behavioral phenotypes. *Hum Mol Genet*. Aug 01 2007;16(15):1802-13. doi:10.1093/hmg/ddm128
14. Falco M, Amabile S, Acquaviva F. gene mutations: mechanisms of Smith-Magenis syndrome. *Appl Clin Genet*. 2017;10:85-94. doi:10.2147/TACG.S128455
15. Girirajan S, Truong HT, Blanchard CL, Elsea SH. A functional network module for Smith-Magenis syndrome. *Clin Genet*. Apr 2009;75(4):364-74. doi:10.1111/j.1399-0004.2008.01135.x
16. Dubourg C, Bonnet-Brilhault F, Toutain A, et al. Identification of Nine New RAI1-Truncating Mutations in Smith-Magenis Syndrome Patients without 17p11.2 Deletions. *Mol Syndromol*. Feb 2014;5(2):57-64. doi:10.1159/000357359
17. Bi W, Saifi GM, Shaw CJ, et al. Mutations of RAI1, a PHD-containing protein, in nondeletion patients with Smith-Magenis syndrome. *Hum Genet*. Nov 2004;115(6):515-24. doi:10.1007/s00439-004-1187-6
18. Carmona-Mora P, Canales CP, Cao L, et al. RAI1 transcription factor activity is impaired in mutants associated with Smith-Magenis Syndrome. *PLoS One*. 2012;7(9):e45155. doi:10.1371/journal.pone.0045155
19. Girirajan S, Vlangos CN, Szomju BB, et al. Genotype-phenotype correlation in Smith-Magenis syndrome: evidence that multiple genes in 17p11.2 contribute to the clinical spectrum. *Genet Med*. Jul 2006;8(7):417-27. doi:10.1097/01.gim.0000228215.32110.89

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.